



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>







Y2A9B1J 34A1

Entered according to Act of Congress, in the year 1885, by
LEA BROTHERS & CO.,
in the Office of the Librarian of Congress at Washington. All rights reserved.

WESTCOTT & THOMSON,
Stereotypers and Electrotypers, Philada.

WILLIAM J. DORNAN,
Printer, Philada.

W. E. Sander -
Oct. 1847.



AMERICAN PUBLISHER'S NOTICE.

THE Author has carefully revised this edition of his Treatise on Obstetrics, and presented the subject in its latest aspect from a British standpoint. It is well known, however, that American opinions and practice differ somewhat from those of Great Britain, and require that certain teachings should be remodelled in accordance with our usages and experience. We differ in the form of decubitus for the application of the forceps; in the models of instruments in use; in the measure of fear of the Cæsarean operation when based upon our more favorable results; on the question of the use of stimulants for wet-nurses and convalescent parturient women, etc.

During the several months that have elapsed since the publication of the English edition large additions have been made to our obstetrical statistics, and some changes effected in the management of cases requiring surgical assistance. The American Editor feels that he has been called upon to bring up his work to the latest period, and has therefore added the new forms of Cæsarean operation devised in Germany, the experiences in Berlin with Hicks' method of treating placenta prævia, the latest American statistics in the Cæsarean section and laparo-elytomy, the latest Porro statistics of the world, etc. He has also added articles on spondylolisthesis (a pregnant case of which is now in this city), the rational treatment of rupture of the uterus, etc. In all cases the additions have been distinguished by enclosure in brackets [—].

PHILADELPHIA, *March, 1885.*

AUTHOR'S PREFACE

TO THE

FOURTH AMERICAN EDITION.

IN preparing a new edition of his work on Midwifery for his American readers, the Author has gratefully to acknowledge the kind reception it continues to receive from the profession. A comparatively short time having elapsed since the last edition was published, there are naturally not many changes to make. The whole work, however, has been carefully revised, and the chapter on "Conception and Generation" has been in great part rewritten, so as to incorporate the most recent advances in Embryology. The Author has to acknowledge the kind assistance he has received in this subject from his late colleague, Dr. W. TYRRELL BROOKES, formerly of the Physiological Laboratory in King's College, now of Oxford. Several new illustrations have been added, and it is hoped that the work may thus prove more worthy of being used as a guide in the anxieties and emergencies of obstetric practice.

31 GEORGE STREET, HANOVER SQUARE, W., LONDON,
August, 1884.



PREFACE TO THE FIRST EDITION.

THOSE who have studied the progress of Midwifery know that there is no department of medicine in which more has been done of late years, and none in which modern views of practice differ more widely from those prevalent only a short time ago. The Author's object has been to place in the hands of his readers an epitome of the science and practice of midwifery which embodies all recent advances. He is aware that on certain important points he has recommended practice which not long ago would have been considered heterodox in the extreme, and which even now will not meet with general approval. He has, however, the satisfaction of knowing that he has only done so after very deliberate reflection, and with the profound conviction that such changes are right and that they will stand the test of experience. He has endeavored to dwell especially on the practical part of the subject, so as to make the work a useful guide in this most anxious and responsible branch of the profession. It is admitted by all that emergencies and difficulties arise more often in this than in any other branch of practice; and there is no part of the practitioner's work which requires more thorough knowledge or greater experience. It is, moreover, a lamentable fact that students generally leave their schools more ignorant of obstetrics than of any other subject. So long as the absurd regulations exist which oblige the lecturer on midwifery to attempt the impossible task of teaching obstetrics in a short three months' course—an absurdity which has over and over again been pointed out—such must of necessity be the case. This must be the Author's excuse for dwelling on many topics at greater length than some will doubtless think their importance merits, since he desires to place in the hands of his students a work which may in some measure supply the inevitable defects of his lectures.

Many of the illustrations are copied from previous authors, while some are original. The following quotation from the preface to Tyler Smith's *Manual of Obstetrics* will explain why the source of the copied wood-cuts has not been in each instance acknowledged: "When I began to publish, I determined to give the authority for every wood-cut copied from other works. I soon found, however, that obstetric authors of all countries, from the time of Mauriceau downward, had copied each other so freely without acknowledgment as to render it difficult or impossible to trace the originals."

The Author has to express his acknowledgments to many friends for their kind assistance by the loan of illustrations and otherwise, and more especially to his colleague, DR. HAYES, for his valuable aid in passing the work through the press.

31 GEORGE STREET, HANOVER SQUARE,
March, 1876.

CONTENTS.

PART I.

ANATOMY AND PHYSIOLOGY OF THE ORGANS CONCERNED IN PARTURITION.

CHAPTER I.

ANATOMY OF THE PELVIS.

	PAGE
Its importance—Formation of the Pelvis—The os innominatum: its three divisions—Separation between the True and False Pelvis—The Sacrum and Coccyx—Mechanical relations of the Sacrum—Pelvic articulations and ligaments—Movements of the Pelvic Joints—The Pelvis as a whole—Differences in the Two Sexes—Measurements of the Pelvis—Its diameters, planes, and axes—Development of the Pelvis—Soft parts in connection with the Pelvis . . .	33

CHAPTER II.

THE FEMALE GENERATIVE ORGANS.

Division according to Function: 1. External or Copulative; 2. Internal or Formative Organs—Mons Veneris—Labia majora and minora—The Clitoris—The Vestibule and Orifice of Urethra—Passing of the female catheter—Orifice of Vagina—The Hymen—The glands of the Vulva—The Perineum—The Vagina—The Uterus: its position and anatomy—The Ligaments of the Uterus—The Parovarium—The Fallopian Tubes—The Ovaries—The Graafian Follicles and the Ova—The Mammary Glands	49
--	----

CHAPTER III.

OVULATION AND MENSTRUATION.

Functions of the Ovary—Changes in the Graafian Follicle: 1. Maturation; 2. Escape of the Ovum—Formation of the Corpus Luteum—Quality and source of the Menstrual blood—Theory of Menstruation—Purpose of the Menstrual loss—Vicarious Menstruation—Cessation of Menstruation	81
--	----

PART II.

PREGNANCY.

CHAPTER I.

CONCEPTION AND GENERATION.

	PAGE
The Semen—Site and mode of Impregnation—Changes in the Ovum—Cleavage of the Yolk—The Decidua and its formation—Formation of the Amnion—The Umbilical Vesicle and Allantois—The Liquor Amnii and its uses—The Chorion—The Placenta: its formation, anatomy, and functions	95

CHAPTER II.

THE ANATOMY AND PHYSIOLOGY OF THE FŒTUS.

Appearance of the Fœtus at various stages of development—Anatomy of the Fœtal Head—The Sutures and Fontanelles—Influence of Sex and Race on the Fœtal Head—Position of the Fœtus in utero—Functions of the Fœtus—The Fœtal Circulation	118
--	-----

CHAPTER III.

PREGNANCY.

Changes in the form and dimensions of the Uterus—Changes in the Cervix—Changes in the texture of the Uterine Tissues, the Peritoneal, Muscular, and Mucous Coats—General modifications in the Body produced by Pregnancy .	132
--	-----

CHAPTER IV.

SIGNS AND SYMPTOMS OF PREGNANCY.

Signs of a fruitful Conception—Cessation of Menstruation—Sympathetic Disturbances: Morning Sickness, etc.—Mammary Changes—Enlargement of the Abdomen—Quickening—Intermittent Uterine Contractions—Vaginal Signs of Pregnancy—Ballotement, etc.—Auscultatory Signs of Pregnancy—Fœtal Pulsations—Uterine Souffle, etc.	143
---	-----

CHAPTER V.

THE DIFFERENTIAL DIAGNOSIS OF PREGNANCY—SPURIOUS PREGNANCY—THE DURATION OF PREGNANCY—SIGNS OF RECENT PREGNANCY.

Adipose enlargement of the Abdomen—Distension of the Uterus by retained Menses, etc.—Congestive enlargement of the Uterus—Ascites—Uterine and Ovarian Tumors—Spurious Pregnancy: its Causes, Symptoms, and Diagnosis—The Duration of Pregnancy—Sources of Fallacy—Methods of Predicting Date of Delivery—Protraction of Pregnancy—Signs of recent Delivery . .	157
--	-----

CHAPTER VI.

ABNORMAL PREGNANCY, INCLUDING MULTIPLE PREGNANCY, SUPER-FŒTATION, EXTRA-UTERINE FŒTATION, AND MISSED LABOR.

	PAGE
Plural Births, their frequency: Relative frequency in different Countries; Causes, etc.—Super-fœtation and Super-fecundation—Nature—Explanation—Objections to admission of such cases—Their possibility admitted—Extra-uterine Pregnancy—Classification—Causes—Tubal Pregnancies—Changes in the Fallopian Tubes—Condition of Uterus—Progress and Termination—Diagnosis—Treatment—Abdominal Pregnancy: Description; Diagnosis; Treatment—Missed Labor: its Symptoms, Causes, and Treatment	166

CHAPTER VII.

DISEASES OF PREGNANCY.

Some only Sympathetic, others Mechanical or Complex in their Origin—Derangements of the Digestive Organs: Excessive Nausea and Vomiting; Diarrhœa; Constipation; Hemorrhoids; Ptyalism; Dyspnœa, etc.—Palpitation—Syncope—Anæmia and Chlorosis—Albuminuria	198
--	-----

CHAPTER VIII.

DISEASES OF PREGNANCY (*continued*).

Disorders of the Nervous System: Insomnia; Headaches and Neuralgia; Paralysis; Chorea; Disorders of the Urinary Organs; Retention of Urine; Irritability of the Bladder; Incontinence of Urine; Phosphatic Deposits; Leucorrhœa; Effects of Pressure; Laceration of Veins; Displacements of the Gravid Uterus; Prolapse, Anteversion, Retroversion—Diseases co-existing with Pregnancy: Eruptive Fevers—Small-pox, Measles, Scarlet Fever, Continued Fever; Phthisis; Cardiac Disease; Syphilis; Icterus; Carcinoma; Pregnancy complicated with Ovarian and Fibroid Tumors	211
--	-----

CHAPTER IX.

PATHOLOGY OF THE DECIDUA AND OVUM.

Pathology of the Decidua—Hydorrhœa Gravidarum—Pathology of the Chorion; Vesicular Degeneration, Myxoma Fibrosum—Pathology of the Placenta; Blood Extravasations, Fatty Degeneration, etc.—Pathology of the Umbilical Cord—Pathology of the Amnion, Hydramnios; Deficiency of Liquor Amnii, etc.—Pathology of the Fœtus: Blood Diseases transmitted through the Mother, Small-pox, Measles, and Scarlet Fever, Intermittent Fevers, Lead-poisoning, Syphilis—Inflammatory Diseases—Dropsies—Tumors—Wounds and Injuries of the Fœtus—Intra-uterine Amputations—Death of the Fœtus	226
---	-----

CHAPTER X.

ABORTION AND PREMATURE LABOR.

PAGE

Importance and Frequency—Definition and Classification—Frequency—Recurrence—Causes—Causes referable to Fœtus—Changes in a Dead Ovum retained in Utero—Extravasations of Blood—Moles, etc.—Causes depending on Maternal State—Syphilis: Causes acting through Nervous System, Physical Causes, etc.—Causes depending on Morbid States of Uterus—Symptoms—Preventive Treatment—Prophylactic Treatment—Treatment when Abortion is inevitable—After-treatment	242
---	-----

PART III.

LABOR.

CHAPTER I.

THE PHENOMENA OF LABOR.

Causes of Labor—Mode in which the Expulsion of the Child is effected—The Uterine Contraction—Mode in which the Dilatation of the Cervix is effected—Rupture of the Membranes—Character and Source of Pains during Labor—Effect of Pains on Mother and Fœtus—Division of Labor into Stages—Preparatory Stage—False Pains—First Stage—Second Stage—Third Stage—Mode in which the Placenta is expelled—Duration of Labor	255
---	-----

CHAPTER II.

MECHANISM OF DELIVERY IN HEAD PRESENTATIONS.

Importance of Subject—Frequency of Head Presentations—The Different Positions of the Head—First Position—Division of Mechanical Movements into Stages—Flexion—Descent and Levelling Movement—Rotation—Extension—External Rotation—Second Position—Third Position—Fourth Position—Caput Succedaneum—Alteration in Shape of Head from Moulding	268
--	-----

CHAPTER III.

MANAGEMENT OF NATURAL LABOR.

Preparatory Treatment—Dress of Patient during Pregnancy—The Obstetric Bag—Duties on first visiting Patient—False Pains—Their Character and Treatment—Vaginal Examination—The Position of Patient—Artificial Rupture of Membranes—Treatment of Propulsive Stage—Relaxation of the Perineum—Treatment of Lacerations—Expulsion of Child—Promotion of Uterine Contraction—Ligature of the Cord—Management of the Third Stage of Labor—Application of the Binder—After-treatment	280
--	-----

CONTENTS.

xiii

CHAPTER IV.

ANÆSTHESIA IN LABOR.

PAGE

Agents employed—Chloral: its Object and Mode of Administration—Ether— Chloroform: its Use, Objections to, and Mode of Administration	295
---	-----

CHAPTER V.

PELVIC PRESENTATIONS.

Frequency—Causes—Prognosis to Mother and Child—Diagnosis by Abdominal Palpation and by Vaginal Examination—Differential Diagnosis of Breech, Knee, and Feet—Mechanism—Treatment—Management of Impacted Breech Presentations	299
--	-----

CHAPTER VI.

PRESENTATIONS OF THE FACE.

Erroneous Views formerly held on the Subject—Frequency—Mode of Production —Diagnosis—Mechanism—Four Positions of the Face—Description of Deliv- ery in First Face Position—Mento-posterior Position in which Rotation does not take place—Prognosis—Treatment—Brow Presentations	310
---	-----

CHAPTER VII.

DIFFICULT OCCIPITO-POSTERIOR POSITIONS.

Causes of Face to Pubes Delivery—Mode of Treatment—Upward Pressure on Forehead—Downward Traction on Occiput—Use of Forceps—Peculiarities of Forceps Delivery	319
--	-----

CHAPTER VIII.

PRESENTATIONS OF SHOULDER, ARM, OR TRUNK—COMPLEX PRESENTATIONS—PRO- LAPSE OF THE FUNIS.

Position of the Fœtus—Division into Dorso-anterior and Dorso-posterior Positions —Causes—Prognosis and Frequency—Diagnosis—Mode of distinguishing Po- sition of Child—Differential Diagnosis of Shoulder, Elbow, and Hand—Mech- anism—The Two Possible Modes of Delivery by the Natural Powers—Spon- taneous Version—Spontaneous Evolution—Treatment—Complex Presentation: Foot or Hand, with Head, Hand, and Feet together—Dorsal Displacement of the Arm—Prolapse of the Umbilical Cord—Frequency—Prognosis—Causes— Diagnosis—Postural Treatment—Artificial Reposition—Treatment when Re- position fails	322
--	-----

CHAPTER IX.

PROLONGED AND PRECIPITATE LABORS.

PAGE

Evil Effects of Prolonged Labor—Influence of the Stage of Labor in Protraction— Delay in the First Stage rarely serious—Temporary Cessation of Pains—Symptoms of Protraction in the Second Stage—State of the Uterus in Protracted Labor—Cases of Protraction due to morbid condition of the expulsive powers— Causes of Protraction—Treatment—Oxytocic Remedies—Ergot of Rye, etc.— Manual Pressure—Instrumental Delivery (case of Princess Charlotte of Wales) —Precipitate Labor—Its Causes and Treatment	337
--	-----

CHAPTER X.

LABOR OBSTRUCTED BY FAULTY CONDITION OF THE SOFT PARTS.

Rigidity of the Cervix: its Causes, Effects, and Treatment—Ante-partum Hour-glass Contraction—Bands and Cicatrices in the Vagina—Extreme Rigidity of the Perineum—Labor complicated with Tumor—Vaginal Cystocele—Calculus—Hernial Protrusions—Œdema of Vulva—Hæmatic Effusions, etc.	351
--	-----

CHAPTER XI.

DIFFICULT LABOR DEPENDING ON SOME UNUSUAL CONDITION OF THE FŒTUS.

Plural Births, Treatment of—Locked Twins—Conjoined Twins—Intra-uterine Hydrocephalus: its Dangers, Diagnosis, and Treatment—Other Dropsical Effusions—Fœtal Tumors—Excessive Development of Fœtus	363
---	-----

CHAPTER XII.

DEFORMITIES OF THE PELVIS.

Classification—Causes of Pelvic Deformity—Rickets and Osteo-malacia—The Equally-enlarged Pelvis—The Equally-contracted Pelvis—The Undeveloped Pelvis—Masculine or Funnel-shaped Pelvis—Contraction of Conjugate Diameter of the Brim—Figure-of-eight Deformity—Spondylolisthesis—Spondylolizema—Narrowing of the Oblique Diameters—Obliquely-contracted Pelvis—Kyphotic Pelvis—Robert's Pelvis—Deformity from Old-standing Hip-joint Disease—Deformity from Tumors, Fractures, etc.—Effects of Contracted Pelvis on Labor—Risks to the Mother and Child—Mechanism of Delivery in Head Presentation: <i>a</i> , in Contracted Brim; <i>b</i> , in Generally-contracted Pelvis—Diagnosis—External Measurements—Internal Measurements—Mode of Estimating the Conjugate Diameter of the Brim—Mode of Diagnosing the Oblique Pelvis—Treatment—The Forceps—Turning—Craniotomy—The Induction of Premature Labor—Induction of Abortion	375
--	-----

CHAPTER XIII.

HEMORRHAGE BEFORE DELIVERY: PLACENTA PRÆVIA.

Definition—Causes—Symptoms—Sources and Causes of Hemorrhage—Prognosis— Treatment	400
---	-----

CONTENTS.

xv

CHAPTER XIV.

HEMORRHAGE FROM SEPARATION OF A NORMALLY-SITUATED PLACENTA.

	PAGE
Causes and Pathology—Symptoms and Diagnosis—Prognosis—Treatment	411

CHAPTER XV.

HEMORRHAGE AFTER DELIVERY.

Its Frequency—Generally a Preventible Accident—Causes—Nature's Method of controlling Hemorrhages—Uterine Contraction—Thrombosis—Secondary Causes of Hemorrhage—Irregular Uterine Contraction—Placental Adhesions—Constitutional Predisposition to Flooding—Symptoms—Preventive Treatment—Curative Treatment—Secondary Treatment—Secondary Post-partum Hemorrhage—Its Causes and Treatment	415
---	-----

CHAPTER XVI.

RUPTURE OF THE UTERUS, ETC.

Its Fatality—Seat of Rupture—Causes, Predisposing and Exciting—Symptoms—Prognosis—Treatment: when the Fœtus remains in Utero; when the Fœtus has escaped from the Uterus—Lacerations of the Cervix—Recapitulation—Lacerations of the Vagina—Vesico- and Recto-vaginal Fistulæ—Their Mode of Formation—Treatment	431
---	-----

CHAPTER XVII.

INVERSION OF THE UTERUS.

Division into Acute and Chronic Forms—Description—Symptoms—Diagnosis—Mode of Production—Treatment	442
---	-----

PART IV.

OBSTETRIC OPERATIONS.

CHAPTER I.

INDUCTION OF PREMATURE LABOR.

History—Objects—May be performed either on account of the Mother or Child—Modes of inducing Labor—Puncture of Membranes—Administration of Oxytocics—Means acting Indirectly on the Uterus—Dilatation of Cervix—Separation of Membranes—Vaginal and Uterine Douches—Introduction of Flexible Catheter	449
--	-----

CHAPTER II.

TURNING.

PAGE

History—Turning by External Manipulation—Object and Nature of the Operation—Cases Suitable for the Operation—Statistics and Dangers—Method of Performance—Cephalic Version—Method of Performance—Podalic Version—Position of Patient—Administration of Anæsthetics—Period when the Operation should be Undertaken—Choice of Hand to be used—Turning by Bi-polar Method—Turning when the Hand is introduced into the Uterus—Turning in Abdomino-anterior Positions—Difficult Cases of Arm Presentation . . . 457

CHAPTER III.

THE FORCEPS.

Frequent Use of the Forceps in Modern Practice—Description of the Instrument—The Short Forceps—Its Varieties—The Long Forceps—Suitable to all Cases alike—Action of the Instrument—Its Power as a Tractor, Lever, and Compressor—Preliminary Considerations before Operation—Use of Anæsthetics—Description of the Operation—Low Forceps Operation—High Forceps Operation—Possible Dangers of Forceps Delivery—Possible Risks to the Child . . . 472

CHAPTER IV.

THE VECTIS—THE FILLET.

Nature of the Vectis—Its Use as a Lever or Tractor—Cases in which it is Applicable—Its Use as a Rectifier of Malpositions—The Fillet—Nature of the Instrument—Objection to its Use . . . 495

CHAPTER V.

OPERATIONS INVOLVING DESTRUCTION OF THE FŒTUS.

Their Antiquity and History—Division of Subject—Nature of Instruments employed—Perforator—Crotchet—Craniotomy Forceps—Cephalotribe—Forceps-saw—Écraseur—Basilyst—Cases requiring Craniotomy—Method of Perforation—Extraction of the Head—Comparative merits of Cephalotripsy and Craniotomy—Extraction by the Craniotomy Forceps—Extraction of the Body—Embryotomy—Decapitation and Evisceration. . . 497

CHAPTER VI.

THE CÆSAREAN SECTION—PORRO'S OPERATION—SYMPHYSEOTOMY.

History of the Operation—Statistics—Results to Mother and Child—Causes requiring the Operation—Post-mortem Cæsarean Section—Causes of Death after the Cæsarean Section—Preliminary Preparation—Description of the Operation—New Forms of Operating—Subsequent Management—Porro's Operation—Substitutes for the Cæsarean Section—Symphyseotomy . . . 511

CHAPTER VII.

LAPARO-ELYTROTOMY.

PAGE

History—Nature of the Operation—Advantages over the Cæsarean Section—Cases suitable for the Operation—Anatomy of the Parts concerned in the Operation— Method of Performance—Subsequent Treatment	529
---	-----

CHAPTER VIII.

THE TRANSFUSION OF BLOOD.

History—Nature and Object of the Operation—Use of Blood taken from the Lower Animals—Difficulties from Coagulation of Fibrin—Modes of Obviating them —Immediate Transfusion—Addition of Chemical Agents to prevent Coagula- tion—Defibrination of the Blood—Statistical Results—Possible Dangers of the Operation—Cases suitable for Transfusion—Description of the Operation— Schäfer's Directions for Immediate Transfusion—Effects of Successful Trans- fusion—Secondary Effects of Transfusion	534
--	-----

PART V.

THE PUERPERAL STATE.

CHAPTER I.

THE PUERPERAL STATE AND ITS MANAGEMENT.

Importance of Studying the Puerperal State—The Mortality of Childbirth— Alterations in the Blood after Delivery—Condition after Delivery—Nervous Shock—Fall of the Pulse—The Secretions and Excretions—Secretion of Milk —Changes in the Uterus after Delivery—The Lochia—The After-pains—Man- agement of Women after Delivery—Treatment of Severe After-pains—Diet and Regimen	546
--	-----

CHAPTER II.

MANAGEMENT OF THE INFANT, LACTATION, ETC.

Commencement of Respiration after the Birth of the Child—Apparent Death of the New-born Child—Its Treatment—Washing and Dressing the Child—Ap- plication of the Child to the Breast—The Colostrum and its Properties—Secre- tion of Milk—Importance of Nursing—Selection of a Wet-nurse—Manage- ment of Lactation—Diet and Regimen of Nursing Women—Period of Weaning —Disorders of Lactation—Means of Arresting the Secretion of Milk—Defective Secretion of Milk—Depressed Nipples—Fissures and Excoriations of the Nip- ples—Excessive Flow of Milk—Mammary Abscess—Hand-feeding—Causes of Mortality in Hand-feeding—Various Kinds of Milk—Method of Hand-feeding	557
--	-----

CHAPTER III.

PUERPERAL ECLAMPSIA.

PAGE

Its Doubtful Etiology—Premonitory Symptoms—Symptoms of the Attack—Condition between the Attacks—Relation of the Attacks to Labor—Results to Mother and Child—Pathology—Treatment—Obstetric Management	573
---	-----

CHAPTER IV.

PUERPERAL INSANITY.

Classification—Proportion of Various Forms—Insanity of Pregnancy—Predisposing Causes—Period of Pregnancy at which it Occurs—Type of Insanity—Prognosis—Transient Mania during Delivery—Puerperal Insanity (Proper)—Type of Insanity—Causes—Theory of its Dependence on a Morbid State of the Blood—Objections to the Theory—Prognosis—Post-mortem Signs—Duration—Insanity of Lactation—Type—Symptoms—Of Mania—Of Melancholia—Treatment—Question of Removal to Asylum—Treatment during Convalescence	582
---	-----

CHAPTER V.

PUERPERAL SEPTICÆMIA.

Differences of Opinion—Confusion from this Cause—Modern View of this Disease—History—Its Mortality in Lying-in Hospitals—Numerous Theories as to its Nature—Theory of Local Origin—Theory of an Essential Zymotic Fever—Theory of its Identity with Surgical Septicæmia—Nature of this View—Channels through which Septic Matter may be Absorbed—Character and Origin of Septic Matter often Obscure—Division into Auto-genetic and Hetero-genetic Cases—Sources of Self-infection—Sources of Hetero-genetic Infection—Influence of Cadaveric Poison—Infection from Erysipelas—Infection from other Zymotic Diseases—Infection from Sewer Gas—Contagion from other Puerperal Patients—Mode in which the Poison may be Conveyed to the Patient—Conduct of the Practitioner in relation to the Disease—Nature of the Septic Poison—Local Changes resulting from the Absorption of Septic Material—Channels through which Systemic Infection is Produced—Pathological Phenomena observed after General Blood-Infection—Four Principal Types of Pathological Change—Intense Cases without Marked Post-mortem Signs—Cases characterized by Inflammation of the Serous Membranes—Cases characterized by the Impaction of Infected Emboli and Secondary Inflammation and Abscess—Description of the Disease—Duration—Varieties of Symptoms in Different Cases—Symptoms of Local Complications—Treatment	593
--	-----

CHAPTER VI.

PUERPERAL VENOUS THROMBOSIS AND EMBOLISM.

Puerperal Thrombosis and its Results—Conditions which favor Thrombosis—Conditions which favor Coagulation in the Puerperal State—Distinction between Thrombosis and Embolism—Is Primary Thrombosis of the Pulmonary Arteries possible?—History—Symptoms of Pulmonary Obstruction—Is Recovery possible—Causes of Death—Post-mortem Appearances—Treatment—Puerperal Pleuro-pneumonia: its Causes and Treatment	621
--	-----

CONTENTS.

xix

CHAPTER VII.

PUERPERAL ARTERIAL THROMBOSIS AND EMBOLISM.

Causes—Symptoms—Treatment	PAGE 633
--	---------------------

CHAPTER VIII.

OTHER CAUSES OF SUDDEN DEATH DURING LABOR AND THE PUERPERAL STATE.

Organic and Functional Causes—Idiopathic Asphyxia—Pulmonary Apoplexy— Cerebral Apoplexy—Syncope—Shock and Exhaustion—Entrance of Air into the Veins	635
--	------------

CHAPTER IX.

**PERIPHERAL VENOUS THROMBOSIS (SYNS.: CRURAL PHLEBITIS; PHLEGMASIA DO-
LENS; ANASARCA SEROSA; OEDEMA LACTEUM; MILK LEG, ETC.).**

Nature—Symptoms—History and Pathology—Anatomical Form of the Thrombi in the Veins—Detachment of Emboli—Treatment	637
---	------------

CHAPTER X.

PELVIC CELLULITIS AND PELVIC PERITONITIS.

Two Forms of Disease—Variety of Nomenclature—Importance of Differential Diagnosis—Etiology—Connection with Septicæmia—Seat of Inflammation— Relative Frequency of the Two Forms of Disease—Symptomatology—Results of Physical Examination—Terminations—Prognosis—Treatment	644
---	------------

INDEX	653
------------------------	------------



ILLUSTRATIONS.

- PLATE I.—Section of a Frozen Body in the last months of Pregnancy (after Braune). Illustrating the Relations of the Uterus to the surrounding Parts, and the attitude of the Fœtus, which is lying in the second Cranial Position *Frontispiece*
- PLATE II.—Section of a Frozen Body at the termination of the First Stage of Labor (after Braune). Membranes unbroken; Cervix fully dilated; and the Head (in the Second Position) in the Pelvic Cavity *Frontispiece*
- PLATE III.—Illustrations of the Corpora Lutea of Menstruation and Pregnancy (after Dalton) *Facing page 81*

FIG.	PAGE
1. Os Innominatum	34
2. Sacrum and Coccyx	35
3. Section of Pelvis and Heads of Thigh-bones, showing the Suspensory Action of the Sacro-iliac Ligaments. (After Wood.)	37
4. Outlet of Pelvis	39
5. The Female Pelvis	40
6. The Male Pelvis	40
7. Brim of Pelvis, showing Antero-posterior, Oblique, and Conjugate Diameters	41
8. Transverse Section of Pelvis, showing the Diameters	42
9. Planes of the Pelvis, with Horizon	43
10. Axes of the Pelvis	44
11. Representing General Axis of the Parturient Canal, including the Uterine Cavity and Soft Parts	45
12. Side View of Pelvis	46
13. Pelvis of a Child	47
14. External Genitals of Virgin with Diaphragmatic Hymen. (After Sappey.)	49
15. Vascular Supply of Vulva. (After Kobelt.)	53
16. Right Half of Virgin Vagina with Walls held apart, showing the abundant transverse Rugæ, the greater depth of the Vagina above than below, and the Hymeneal Segment. (After Hart.)	54
17. Longitudinal Section of Body, showing Relations of Generative Organs	55
18. Transverse Section of Body, showing Relations of the Fundus Uteri	56
19. Transverse Section of Uterus	57
20. Uterus and Appendages in an Infant. (After Farre.)	58
21. Portion of Interior of Cervix. (Enlarged nine diameters.)	59
22. Muscular Fibres of unimpregnated Uterus. (After Farre.)	60
23. Developed Muscular Fibres from the Gravid Uterus. (After Wagner.)	60
24. Lining Membrane of Uterus, showing network of Capillaries and Orifices of Uterine Glands. (After Farre.)	62
25. The Course of the Glands in the fully-developed Mucous Membrane of the Uterus. (After Williams.)	62

FIG.	PAGE
26. Vertical Section through the Mucous Membrane of the Human Uterus. (After Turner.)	63
27. Villi of Os Uteri stripped of Epithelium. (After Tyler Smith and Hassall.)	64
28. Villi of Uterus, covered with Pavement Epithelium, and containing Looped Vessels. (After Tyler Smith and Hassall.)	64
29. Bifid Uterus. (After Farre.)	66
30. Partitioned Uterus. (Kussmaul.)	68
31. Adult Parovarium, Ovary, and Fallopian Tube. (After Kobelt.)	69
32. Posterior View of Muscular and Vascular Arrangements. (After Rouget.)	70
33. Fallopian Tube laid open. (After Richard.)	72
34. Ovary enlarged under Menstrual Nisus	73
35. Longitudinal Section of Adult Ovary. (After Farre.)	74
36. Section through the Cortical part of the Ovary. (After Turner.)	75
37. Vertical section through the Ovary of the Human Fœtus. (After Foulis.)	76
38. Diagrammatic Section of Graafian Follicle	77
39. Bulb of Ovary	78
40. Mammary Gland	79
41. Section of Ovary, showing Corpus Luteum three weeks after Menstruation. (After Dalton.)	83
42. Corpus Luteum at the fourth month of Pregnancy. (After Dalton.)	84
43. Corpus Luteum of Pregnancy at Term. (After Dalton.)	84
44. Section of Parts of three Seminiferous Tubules of a Rat. (From a preparation by Mr. A. Frazer.)	96
45. Ovum of Rabbit containing Spermatozoa	97
46. Formation of the "Polar Globule".	99
47. Segmentation of the Yolk	99
48. Formation of the Blastodermic Membrane. (After Joulin.)	100
49. Aborted Ovum (of about forty days), showing the Triangular Shape of the Decidua (which is laid open), and the Aperture of the Fallopian Tube. (After Coste.)	102
50. } Formation of the Decidua. (After Dalton.)	102
51. }	
52. }	
53. An Ovum removed from the Uterus, and part of the Decidua Vera cut away. (After Coste.)	103
54. Diagram of Area Germinativa, showing the Primitive Trace and Area Pellicula	105
55. Development of the Amnion	106
56. Development of the Umbilical Vesicle and Amnion	107
57. An Embryo of about twenty-five days laid open. (After Coste.)	107
58. Development of the Chorion	108
59. Placental Villus, greatly magnified. (After Joulin.)	113
60. Terminal Villus of Fœtal Tuft, minutely injected. (After Farre.)	113
61. Diagram representing a Vertical Section of the Placenta. (After Dalton.)	114
62. Diagram illustrating the Mode in which a Placental Villus derives a Covering from the Vascular System of the Mother. (After Priestley.)	115
63. The Extremity of a Placental Villus. (After Goodsir.)	115
64. Anterior and Posterior Fontanelles	121
65. Bi-parietal Diameter, Sagittal and Lambdoidal Sutures, with Posterior Fontanelle	121
66. Diameters of the Fœtal Skull	122


FIG.	PAGE
67. Mode of Ascertaining the Position of the Fœtus by Palpation	124
68. Diagram illustrating the Effect of Gravity on the Fœtus. (After Duncan.)	126
69. Illustrating the greater Mobility of the Fœtus and the larger relative amount of Liquor Amnii in Early Pregnancy. (After Duncan.)	126
70. Diagram of Fœtal Heart. (After Dalton.)	129
71. Diagram of Heart of Infant. (After Dalton.)	131
72. Relations of Pregnant Uterus at six months. (After Martin.)	133
73. Size of Uterus at various Periods of Pregnancy	135
74. } Supposed Shortening of the Cervix at the third, sixth, eighth, and ninth	
75. } months of Pregnancy, as figured in Obstetric Works	136
76. }	
77. }	
78. Cervix of a Woman Dying in the Eighth Month of Pregnancy. (After Duncan.)	137
79. Appearance of the Areola in Pregnancy	147
80. Illustrating the Cavity between the Decidua Vera and the Decidua Reflexa during the early Months of Pregnancy. (After Coste.)	171
81. Tubal Pregnancy, with the Corpus Luteum in the Ovary of the opposite side.	174
82. Tubal Pregnancy. (From a specimen in the Museum of King's College.) .	175
83. Extra-uterine Pregnancy at term of the Tubo-ovarian Variety. (After a case of Dr. A. Sibley Campbell's.)	177
84. Uterus and Fœtus in a case of Abdominal Pregnancy	184
85. Lithopædion. (From a preparation in the Museum of the Royal College of Surgeons.)	185
86. Contents of the Cyst in Dr. Oldham's case of Missed Labor	194
87. Hypertrophied Decidua laid open, with the Ovum attached to its Fundal Portion. (After Duncan.)	227
88. Imperfectly developed Decidua Vera, with the Ovum. (After Duncan.) . .	228
89. Hydatidiform Degeneration of the Chorion	230
90. Double Placenta, with Single Cord	233
91. Fatty Degeneration of the Placenta	234
92. Knots in the Umbilical Cord	235
93. Intra-uterine Amputation of both Arms and Legs	240
94. An Apoplectic Ovum, with Blood effused in masses under the Fœtal Surface of the Membranes	245
95. Blighted Ovum, with Fleishy Degeneration of the Membranes	246
96. Mode in which the Placenta is Naturally Expelled. (After Duncan.) . .	266
97. Attitude of Child in First Position. (After Hodge.)	270
98. First Position: Movement of Flexion	271
99. First Position: Occiput in Cavity of Pelvis. (After Hodge.)	273
100. First Position: Occiput at Outlet of Pelvis. (After Hodge.)	274
101. First Position: Head Delivered. (After Hodge.)	275
102. External Rotation of Head in First Position. (After Hodge.)	275
103. Third Position of Occiput at Brim of Pelvis.	276
104. Fourth Position of Occiput at Pelvic Brim	278
105. Examination during the First Stage of Labor	282
106. Mode of effecting Relaxation of the Perineum	288
107. Usual Method of Removing the Placenta by Traction on the Cord	291
108. Illustrating Expression of the Placenta	293
109. First, or left Sacro-anterior position of the Breech	303
110. Passage of the Shoulders and partial Rotation of the Thorax	304
111. Descent of the Head	304

FIG.	PAGE
112. Third Position in Face Presentation	313
113. Rotation forward of Chin	314
114. Passage of the Head through the External Parts in Face Presentation . . .	315
115. Illustrating the Position of the Head when Forward Rotation of the Chin does not take place	315
116. Dorso-anterior Presentation of the Arm	323
117. Dorso-posterior Presentation of the Arm	324
118. Spontaneous Evolution. (After Chiara.)	329
119. Dorsal Displacement of the Arm	331
120. Dorsal Displacement of the Arm in Footling Presentation. (After Barnes.)	331
121. Prolapse of the Umbilical Cord	332
122. Postural Treatment of Prolapse of the Cord	334
123. Braun's Apparatus for Replacing the Cord	336
124. Labor complicated by Ovarian Tumor	358
125. Twin Pregnancy, Breech and Head presenting	363
126. Head Locking, both Children presenting Head first. (After Barnes.) . .	365
127. Head Locking, first Child coming Feet first: Impaction of Heads from wedging in Brim. (After Barnes.)	366
128. Labor impeded by Hydrocephalus	371
129. Adult Pelvis retaining its Infantile Type	378
130. Scolio-rachitic Pelvis	379
131. Rickety Pelvis, with backward depression of Symphysis Pubis	380
132. Flatness of Sacrum, with narrowing of the Pelvic Cavity	381
133. Pelvis deformed by Spondylolisthesis. (After Kilian.)	381
134. [" " " (After Neugebauer.)]	382
135. Osteo-malacic Pelvis	385
136. Extreme degree of Osteo-malacic Deformity	385
137. Obliquely-contracted Pelvis. (After Duncan.)	386
138. Kyphotic Pelvis	387
139. Robert's, or Double Obliquely-contracted Pelvis. (After Duncan.) . . .	387
140. Bony Growth from Sacrum obstructing the Pelvic Cavity	388
141. Greenhalgh's Pelvimeter	393
142. Section of Fœtal Cranium, showing its Conical Form	396
143. Showing the greater Breadth of the Bi-parietal Diameter of the Fœtal Cranium. (After Simpson.)	396
144. Showing the greater Space for the Bi-parietal Diameter at the side of the Pelvis in certain Cases of Deformity. (After Simpson.)	397
145. Irregular Contraction of the Uterus, with Encystment of the Placenta . .	418
146. Illustrating the Dangerous Thinning of the Lower Segment of Uterus, owing to non-descent of Head in a case of Intra-uterine Hydrocephalus. (After Bandl.)	434
147. Partial Inversion of the Fundus	443
148. Illustrating the Commencement of Inversion at the Cervix. (After Duncan.)	446
149. Barnes's Bag for Dilating the Cervix	453
150. First Stage of Bi-polar Version. (After Barnes.)	463
151. Second Stage of Bi-polar Version. (After Barnes.)	464
152. Third Stage of Bi-polar Version. (After Barnes.)	464
153. Fourth Stage of Bi-polar Version. (After Barnes.)	465
154. Seizure of the Feet when the Hand is introduced into the Uterus	466
155. Drawing down of the Feet and Completion of Version	467
156. Showing the Completion of Version. (After Barnes.)	469
157. Showing the use of the Right Hand in Abdomino-anterior positions . . .	470

FIG.	PAGE
158. Denman's Short Forceps	473
159. Ziegler's Forceps	473
160. Simpson's Forceps	474
161. Tarnier's Forceps	476
162. Simpson's Axis-Traction Forceps	476
163. Position of Patient for Forceps Delivery, and Mode of Introducing the Lower Blade	480
164. Introduction of the Upper Blade	481
165. Forceps in position; Traction in the Axis of the Brim, downward and backward	482
166. Last Stage of Extraction; the Handles of the Forceps turned upward toward the Mother's Abdomen	483
167. Hodge Forceps	488
168. Wallace Forceps	488
169. Davis Forceps	488
170. Elliott Forceps	490
171. Sawyer Forceps	490
172. Application of the Forceps at the Inferior Strait	491
173. Application of the Forceps with the Head at the Superior Strait, the Left Blade held in Place by an Assistant	493
174. Direction of the Forceps as the Head is being Delivered	494
175. Vectis with Hinged Handle	496
176. Wilmot's Fillet	497
177. }	
178. } Various Forms of Perforators	499
179. }	
180. }	
181. } Crotchets	499
182. Craniotomy Forceps	499
183. Simpson's Cranioclast	500
184. Hicks's Cephalotribe	501
185. Perforation of the Skull	505
186. Fœtal Head crushed by Cephalotribe	507
187. Professor Simpson's Basilyst	508
188. Straight Craniotomy Forceps	509
189. Curved Craniotomy Forceps	510
190. Method of Transfusion by Aveling's Apparatus	541
191. Schäfer's Canula for Immediate Transfusion	541
192. Section of a Uterine Sinus from the Placental Site nine weeks after delivery. (After Williams.)	551
193. }	
194. }	
195. }	
196. }	
197. }	
198. }	
199. Hayes's Tube for Intra-uterine Injections	615
200. }	
201. }	




PLATE I.



SECTION OF A FROZEN BODY IN THE LAST MONTH OF PREGNANCY (AFTER BRAUNE), ILLUSTRATING THE RELATIONS OF THE UTERUS TO THE SURROUNDING PARTS, AND THE ATTITUDE OF THE FETUS, WHICH IS LYING IN THE SECOND CRANIAL POSITION.

PLATE II.



SECTION OF A FROZEN BODY AT THE TERMINATION OF THE FIRST STAGE OF LABOR (AFTER DRAUNE).
 THE BAG OF MEMBRANES IS STILL UNBROKEN, THE CERVIX IS FULLY DILATED, AND
 THE HEAD (IN THE SECOND POSITION) IS IN THE PELVIC CAVITY.



THE
SCIENCE AND PRACTICE
OF
MIDWIFERY.

PART I.

ANATOMY AND PHYSIOLOGY OF THE ORGANS
CONCERNED IN PARTURITION.

CHAPTER I.

ANATOMY OF THE PELVIS.

THE *pelvis* is the bony basin situated between the trunk and the lower extremities. To the obstetrician its study is of paramount importance, for it not only contains, in the unimpregnated state, all the organs connected with the function of reproduction, but through its cavity the foetus has to pass in the process of parturition. An accurate knowledge, therefore, of its anatomical formation may be said to be the very alphabet of obstetrics, without which no one can practise midwifery, either with satisfaction to himself or safety to his patient.

In a treatise on obstetrics, however, any detailed account of the purely descriptive anatomy of the pelvis would be out of place. A knowledge of that must be taken for granted, and it is only necessary to refer to those points which have a more or less direct bearing on the study of its obstetrical relations.

The *pelvis* is formed of four bones. On either side are the *ossa innominata*, joined together by the *sacrum*; to the inferior extremity of the sacrum is attached the *coccyx*, which is, in fact, its continuation.

The *os innominatum* (Fig. 1) is an irregularly shaped bone originally formed of three distinct portions, the *ilium*, the *ischium*, and the *pubes*, which remain separated from each other up to and beyond the period of puberty. They are united at the acetabulum by a Y-shaped cartilaginous junction, which does not, as a rule, become ossified until about the twentieth year. The consequence is that the pelvis, during the period of growth, is subject to the action of various mechanical influences to a

far greater extent than in adult life; and these, as we shall presently see, have an important effect in determining the form of the bones. The external surface and borders of the os innominatum are chiefly of obstetric interest from giving attachment to muscles, many of which have an important accessory influence on parturition, such as the muscles forming the abdominal wall, which are attached to its crest, and those closing its outlet and forming the perineum, which are attached to the tuberosity of the ischium. On the anterior and posterior extremities of the crest of the ilium are two prominences (the anterior and posterior spinous processes) which are points from which certain measurements are sometimes taken. The internal surface of the upper fan-shaped portion of

FIG. 1.



Os Innominatum.

the os innominatum gives attachment to the iliacus muscle, and contributes to the support of the abdominal contents; along with its fellow of the opposite side it forms the *false* pelvis. The false is separated from the *true* pelvis by the ilio-pectineal line, which, with the upper margin of the sacrum, forms the brim of the pelvis. This is of especial obstetric importance, as it is the first part of the pelvic cavity through which the child passes, and that in which osseous deformities are most often met with. At one portion of the ilio-pectineal line, corresponding with the junction of the ilium and pubes, is situated a prominence which is known as the ilio-pectineal eminence.

Internal Surface.—The internal smooth surface of the innominate bone below the linea ilio-pectinea forms the greater portion of the pelvis proper. In front, with the corresponding portions of the opposite bone, it forms the arch of the pubes, under which the head of the child passes in labor.

Behind this we observe the oval obturator foramen, and below that the tuberosity and spine of the ischium, the latter separating the great and lesser sciatic notches, and giving attachment to ligaments of importance. The rough articulating surface posteriorly, by which the junction with the sacrum is effected, may be noted, and above this the prominence

to which the powerful ligaments joining the sacrum and os innominatum are attached.

The *sacrum* (Fig. 2) is a triangular and somewhat spongy bone forming the continuation of the spinal column and binding together the *ossa innominata*. It is originally composed of five separate portions, analogous to the *vertebræ*, which ossify and unite about the period of puberty, leaving on its internal surface four prominent ridges at the points of junction. The upper of these is sometimes so well marked as to be mistaken, on vaginal examination, for the promontory of the sacrum itself.

The base of the sacrum is about $4\frac{1}{2}$ inches in width, and its sides rapidly approximate until they nearly meet at its apex, giving the whole bone a triangular or wedge shape. The anterior and posterior surfaces also approximate in the same way, so that the bone is much thicker at the base than at the apex. The sacrum, in the erect position of the body, is directed from above downward and from before backward. At its upper edge it is joined, the lumbo-sacral cartilage intervening, with the fifth lumbar vertebra. The point of junction, called the promontory of the sacrum, is of great importance, as on its undue projection many deformities of the brim of the pelvis depend. The anterior surface of the bone is concave, and forms the curve of the sacrum—more marked in some cases than in others. There is also more or less concavity from side to side. On it we observe four apertures on each side, the intervertebral foramina, giving exit to nerves. The posterior surface is convex, rough and irregular for the attachment of ligaments and muscles, and showing a ridge of vertical prominences corresponding to the spinous processes of the *vertebræ*.

Mechanical Relations of the Sacrum.—The sacrum is generally described as forming a keystone to the arch constituted by the pelvic bones, and transmitting the weight of the body, in consequence of its wedge-like shape, in a direction which tends to thrust it downward and backward, as if separating the *ossa innominata*. Dr. Duncan,¹ however, has shown, from a careful consideration of its mechanical relations, that it should rather be regarded as a strong transverse beam, curved on its anterior surface, the extremities of which are in contact with the corresponding articular surfaces of the *ossa innominata*. The weight of the body is thus transmitted to the innominate bones, and through them to the acetabula and the femora (Fig. 3). There counter-pressure is applied, and the result is, as we shall subsequently see, an important modifying influence on the development and shape of the pelvis.

The *coccyx* (Fig. 2) is composed of four small separate bones, which eventually unite into one, but not until late in life. The uppermost of

FIG. 2.



Sacrum and Coccyx.

¹ *Researches in Obstetrics*, p. 67.

these articulates with the apex of the sacrum. On its posterior surface are two small cornua, which unite with corresponding points at the tip of the sacrum. The bones of the coccyx taper to a point. To it are attached various muscles which have the effect of imparting considerable mobility. During labor, also, it yields to the mechanical pressure of the presenting part, so as to increase the antero-posterior diameter of the pelvic outlet to the extent of an inch or more.

Ossification of Coccyx.—If, through disease or accident, as sometimes happens, the articular cartilages of the coccyx become prematurely ossified, the enlargement of the pelvic outlet during labor may be prevented, and considerable difficulty may thus arise. This is most apt to happen in aged primiparæ or in women who have followed sedentary occupations; and not infrequently, under such circumstances, the bone fractures under the pressure to which it is subjected by the presenting part.

Pelvic Articulations.—The pelvic bones are firmly joined together by various articulations and ligaments. The latter are arranged so as to complete the canal through which the fœtus has to pass, and which is in great part formed by the bones. On its internal surface, where the absence of obstruction is of importance, they are everywhere smooth; while externally, where strength is the desideratum, they are arranged in larger masses, so as to unite the bones firmly together. The pelvic articulations have been generally described as symphyses or amphiarthrodia—a term which is properly applied to two articulating surfaces united by fibrous tissue in such a way as to prevent any sliding motion. It is certain, however, that this is not the case with the joints of the female pelvis during pregnancy and parturition. Lenoir found that in 22 females between the ages of 18 and 35 there was a distinct sliding motion. Therefore, the pelvic articulations are, strictly speaking, to be considered examples of the class of joints termed arthrodia.

Lumbo-sacral Joint.—The last lumbar vertebra is united to the sacrum by ligamentous union similar to that which joins the vertebræ to each other. The intervening fibro-cartilage forms a disk which is thicker in front than behind, and this, in connection with a similar peculiarity of the fifth lumbar vertebra, tends to increase the sloped position of the sacrum and the angle which it forms with the vertebral column. It constitutes the most prominent portion of the promontory of the sacrum, and is the part on which the finger generally impinges in vaginal examinations. The anterior common vertebral ligament passes over the surface of the joints, and we also find the ligamenta subflava and the interspinous ligaments, as in the other vertebræ. The articular processes are joined together by a fibrous capsule, and there is also a peculiar ligament, the lumbo-sacral, extending from the transverse process of the vertebra on each side, and attaching itself to the sides of the sacrum and the sacro-iliac synchondrosis.

Ligaments of Coccyx.—The sacrum is joined to the coccyx, and, in some cases at least, the separate bones of the coccyx to each other, by small cartilaginous disks like that connecting the sacrum with the last lumbar vertebra. They are further united by anterior and posterior common ligaments, the latter being much the thicker and more marked. In the adult female a synovial membrane is found between the sacrum

and coccyx, and it is supposed that this is formed under the influence of the movements of the bones on each other.

Sacro-iliac Synchondrosis.—The opposing articular surfaces of the sacrum and ilium are each covered by cartilages, that of the sacrum being the thicker. These are firmly united, but, in the female, according to Mr. Wood,¹ they are always more or less separated by an intervening synovial membrane. Posterior to these cartilaginous convex surfaces there are strong interosseous ligaments passing directly from bone to bone, filling up the interspace between them and uniting them firmly. There are also accessory ligaments, such as the superior and anterior sacro-iliac, which are of secondary consequence. The posterior sacro-iliac ligaments, however, are of great obstetric importance. They are the very strong attachments which unite the rough surfaces on the posterior iliac tuberosities to the posterior and lateral surfaces of the sacrum. They pass obliquely downward from the former points, and suspend, as it were, the sacrum from them. According to Duncan, the sacrum has nothing to prevent its being depressed by the weight of the body but these ligaments, and it is mainly through them that the weight

FIG. 3.



Section of Pelvis and Heads of Thigh-bones, showing the Suspensory Action of the Sacro-iliac Ligaments. (After Wood.)

of the body is transmitted to the sacro-cotyloid beams and the heads of the femora.

Sacro-sciatic Ligaments.—The sacro-sciatic ligaments are instrumental in completing the canal of the pelvis. The greater sacro-sciatic ligament is attached by a broad base to the posterior-inferior spine of the ilium and to the posterior surfaces of the sacrum and coccyx. Its fibres unite into a thick cord, cross each other in an X-like manner, and again

¹ Todd's *Cyclopædia of Anatomy and Physiology*, article "Pelvis," p. 123.

expand at their insertion into the tuberosity of the ischium. The lesser sacro-sciatic ligament is also attached with the former to the back parts of the sacrum and coccyx, its fibres passing to their much narrower insertion at the spine of the ischium, and converting the sacro-sciatic notch into a complete foramen.

Obturator Membrane.—The obturator membrane is the fibrous aponeurosis that closes the large obturator foramen. Joulin¹ supposes that, along with the sacro-sciatic ligaments, it may, by yielding somewhat to the pressure of the foetal head, tend to prevent the contusion to which the soft parts would be subjected if they were compressed between two entirely osseous surfaces.

Symphysis Pubis.—The junction of the pubic bones in front is effected by means of two oval plates of fibro-cartilage, attached to each articular surface by nipple-shaped projections, which fit into corresponding depressions in the bones. There is a greater separation between the bones in front than behind, where the numerous fibres of the cartilaginous plates intersect and unite the bones firmly together. At the upper and back part of the articulation there is an interspace between the cartilages which is lined by a delicate membrane. In pregnancy this space often increases in size, so as to extend even to the front of the joint. The juncture is further strengthened by four ligaments—the anterior, the posterior, the superior, and the subpubic. Of these the last is the largest, connecting together the pubic bones and forming the upper boundary of the pubic arch.

Movements of Pelvic Joints.—The close apposition of the bones of the pelvis might not unreasonably lead to the supposition that no movement took place between its component parts; and this is the opinion which is even yet held by many anatomists. It is tolerably certain, however, that even in the unimpregnated condition there is a certain amount of mobility. Thus, Zaglas has pointed out² that in man there is a movement in an antero-posterior direction of the sacro-iliac joints which has the effect, in certain positions of the body, of causing the sacrum to project downward to the extent of about a line, thus narrowing the pelvic brim, tilting up the point of the bone, and thereby enlarging the outlet of the pelvis. This movement seems habitually brought into play in the act of straining during defecation.

Observations in the Lower Animals.—During pregnancy in some of the lower animals there is a very marked movement of the pelvic articulations, which materially facilitates the process of parturition. This, in the case of the guinea-pig and cow, has been especially pointed out by Dr. Matthews Duncan.³ In the former, during labor the pelvic bones separate from each other to the extent of an inch or more. In the latter, the movements are different, for the symphysis pubis is fixed by bony ankylosis, and is immovable; but the sacro-iliac joints become swollen during pregnancy, and extensive movements in an antero-posterior direction take place in them which materially enlarge the pelvic canal during labor.

¹ *Traité d'Accouchements*, p. 11.

² *Monthly Journal of Med. Science*, Sept., 1851.

³ *Researches in Obstetrics*, p. 19.

Mode in which the Movements are Effected.—It is extremely probable that similar movements take place in women, both in the symphysis pubis and in the sacro-iliac joints, although to a less marked extent. These are particularly well described by Dr. Duncan. They seem to consist chiefly in an elevation and depression of the symphysis pubis, either by the ilia moving on the sacrum, or by the sacrum itself undergoing a forward movement on an imaginary transverse axis passing through it, thus lessening the pelvic brim to the extent of one or even two lines, and increasing, at the same time, the diameter of the outlet by tilting up the apex of the sacrum. These movements are only an exaggeration of those which Zaglas describes as occurring normally during defecation. The instinctive positions which the parturient woman assumes find an explanation in these observations. During the first stage of labor, when the head is passing through the brim, she sits or stands or walks about, and in these erect positions the symphysis pubis is depressed and the brim of the pelvis enlarged to its utmost. As the

FIG. 4.



Outlet of Pelvis.

head advances through the cavity of the pelvis, she can no longer maintain her erect position, and she lies down and bends her body forward, which has the effect of causing a nutatory motion of the sacrum, with corresponding tilting up of its apex and an enlargement of the outlet.

Alterations in the Pelvic Joints during Pregnancy.—These movements during parturition are facilitated by the changes which are known to take place in the pelvic articulations during pregnancy. The ligaments and cartilages become swollen and softened, and the synovial membranes existing between the articulating surfaces become greatly augmented in size and distended with fluid. These changes act by forcing the bones apart, as the swelling of a sponge placed between them might do after it had imbibed moisture. The reality of these alterations receives a clinical illustration from those cases, which are far from uncommon, in which these changes are carried to so extreme an extent that the power of progression is materially interfered with for a considerable time after delivery.

Pelvis as a Whole.—On looking at the pelvis as a whole, we are at once struck with its division into the true and false pelvis. The latter portion (all that is above the brim of the pelvis) is of comparatively little obstetric importance, except in giving attachments to the accessory muscles of parturition, and need not be further considered. The brim of the pelvis is a heart-shaped opening, bounded by the sacrum behind,


FIG. 5.



The Female Pelvis.

the linea ilio-pectinea on either side, and the symphysis of the pubes in front. All below it forms the cavity, which is bounded by the hollow of the sacrum behind, by the inner surfaces of the innominate bones at the sides and in front, and by the posterior surface of the symphysis pubis. It is in this part of the pelvis that the changes in direction

FIG. 6.




The Male Pelvis.

which the foetal head undergoes in labor are imparted to it. The lower border of this canal, or pelvic outlet (Fig. 4), is lozenge-shaped, is bounded by the ischiatic tuberosities on either side, the tip of the coccyx behind, and the under surface of the pubic symphysis in front. Posteriorly to the tuberosities of the ischia the boundaries of the outlet are completed by the sacro-sciatic ligaments.

Difference in the Two Sexes.—There is a very marked difference between the pelvis in the male and the female, and the peculiarities of the latter all tend to facilitate the process of parturition. In the female pelvis (Fig. 5) all the bones are lighter in structure, and have the points for muscular attachments much less developed. The iliac bones are more spread out, hence the greater breadth which is observed in the female figure, and the peculiar side-to-side movement which all females have in walking. The tuberosities of the ischia are lighter in structure and farther apart, and the rami of the pubes also converge at a much less acute angle. This greater breadth of the pubic arch gives one of the most easily appreciable points of contrast between the male and the female pelvis; the pubic arch in the female forms an angle of from 90° to 100° , while in the male (Fig. 6) it averages from 70° to 75° . The obturator foramina are more triangular in shape.

The whole cavity of the female pelvis is wider and less funnel-shaped than in the male, the symphysis pubis is not so deep, and, as the promontory of the sacrum does not project so much, the shape of the pelvic brim is more oval than heart-shaped. These differences between the male and female pelvis are probably due to the presence of the female

FIG. 7.




Brim of Pelvis, showing Antero-posterior, Oblique, and Conjugate Diameters.

genital organs in the true pelvis, the growth of which increases its development in width. In proof of this, Schroeder states that in women with congenitally defective internal organs, and in women who have had both ovaries removed early in life, the pelvis has always more or less of the masculine type.

Measurements of the Pelvis.—The measurements of the pelvis that are of most importance from an obstetric point of view are taken between various points directly opposite to each other, and are known as the *diameters* of the pelvis. Those of the true pelvis are the diameters which it is especially important to fix in our memories, and it is customary to describe three in works on obstetrics—the antero-posterior or conjugate, the oblique, and the transverse—although of course the measurements may be taken at any opposing points in the circumference of the bones. The *antero-posterior* (sacro-pubic), at the brim (Fig. 7), is taken

from the upper part of the posterior surface of the symphysis pubis to the centre of the promontory of the sacrum ; in the cavity, from the centre

FIG. 8.



Transverse Section of Pelvis, showing the Diameters.

of the symphysis pubis to a corresponding point in the body of the third piece of the sacrum ; and at the outlet (coccy-pubic), from the lower border of the symphysis pubis to the tip of the coccyx. The *oblique*, at the brim, is taken from the sacro-iliac joint on either side to a point of the brim corresponding with the ilio-pectineal eminence (that starting from the right sacro-iliac joint being called the right oblique, that from the left the left oblique); in the cavity a similar measurement is made at the same level as the conjugate, while at the outlet an oblique diameter is not usually measured. The *transverse* is taken at the brim, from a point midway between the sacro-iliac joint and the ilio-pectineal eminence to a corresponding point at the opposite side of the brim ; in the cavity from points in the same plane as the conjugate and oblique diameters ; and at the outlet from the centre of the inner border of one ischial tuberosity to that of the other. The measurements given by various writers differ considerably and vary somewhat in different pelves. Taking the average of a large number, the following may be given as the standard measurements of the female pelvis :


	Antero-posterior (in.).	Oblique (in.).	Transverse (in.).
Brim	4.25	4.8	5.2
Cavity	4.7	5.2	4.75
Outlet	5.0	4.2

Differences in Various Parts of Pelvis.—It will be observed that the lengths of the corresponding diameters at different places vary greatly ; thus, while the transverse is longest at the brim, the oblique is longest in the cavity and the antero-posterior at the outlet. It will be subsequently seen that this fact is of great practical importance in studying the mechanism of delivery, for the head in its descent through the pelvis alters its position in such a way as to adapt itself to the longest diameter of the pelvis ; thus, as it passes through the cavity it lies in the oblique diameter, and then rotates so as to be expelled in the antero-posterior diameter of the outlet.

Diameters as Altered by Soft Parts.—In thinking of these measurements of the pelvis it must not be forgotten that they are taken in the dried bones, and that they are considerably modified during life by the soft parts. This is especially the case at the brim, where the projection of the psoas and iliacus muscles lessens the transverse diameter about half an inch, while the antero-posterior diameter of the brim and all the diameters of the cavity are lessened by a quarter of an inch. The right oblique diameter of the brim is, even in the dried pelvis, found to be on an average slightly longer than the left, probably on account of the increased development of the right side of the pelvis from the greater use made of the right leg; but in addition to this the left oblique diameter is somewhat lessened during life by the presence of the rectum on the left side. The advantage gained by the comparatively frequent passage of the head through the pelvis in the right oblique diameter is thus explained.

Other Measurements.—There are one or two other measurements of the true pelvis which are sometimes given, but which are of secondary importance. One of these, the sacro-cotyloid diameter, is that between the promontory of the sacrum and a point immediately above the coty-

FIG. 9.



Planes of the Pelvis with Horizon.

- A B. Horizon. C D. Vertical line.
A B I. Angle of inclination of pelvis to horizon, equal to 60°.
B I C. Angle of inclination of pelvis to spinal column, equal to 150°.
C I J. Angle of inclination of sacrum to spinal column, equal to 130°.
E F. Axis of pelvic inlet. L M. Mid plane in the middle line.
N. Lowest point of mid plane of ischium.


loid cavity, and averages from 3.4 to 3.5 inches. Another, called by Wood the lower or inclined conjugate diameter, is that between the centre of the lower margin of the symphysis pubis and the promontory

of the sacrum, and averages half an inch more than the antero-posterior diameter of the brim. These measurements are chiefly of importance in relation to certain pelvic deformities.

External Measurements.—The external measurements of the pelvis are of no real consequence in normal parturition, but they may help us, in certain cases, to estimate the existence and amount of deformities. Those which are generally given are: Between the anterior-superior iliac spines, 10 inches; between the central points of the crests of the ilia, $10\frac{1}{2}$ inches; between the spinous process of the last lumbar vertebra and the upper part of the symphysis pubis (external conjugate), 7 inches.

Planes of the Pelvis.—By the planes of the pelvis are meant imaginary levels at any portion of its circumference. If we were to cut out a piece of cardboard so as to fit the pelvic cavity, and place it either at the brim

FIG. 10.




or elsewhere, it would represent the pelvic plane at that particular part, and it is obvious that we may conceive as many planes as we desire. Observation of the angle which the pelvic planes form with the horizon shows the great obliquity at which the pelvis is placed in regard to the spinal column. Thus the angle A B I (Fig. 9) represents the inclination to the horizon of the plane of the pelvic brim, I B, and is estimated to be about 60° , while the angle which the same plane forms with the vertebral column is about 150° . The plane of the outlet forms, with the coccyx in its usual position, an angle with the horizon of about 11° , but which varies greatly with the movements of the tip of coccyx and the degree to which it is pushed back during parturition. These figures must only be taken as giving an approximate idea of the inclination of the pelvis to the spinal column, and it must be remembered that the degree of inclination varies considerably in the same female at different

times, in accordance with the position of the body. During pregnancy especially, the obliquity of the brim is lessened by the patient throwing herself backward in order to support more easily the weight of the gravid uterus. The height of the promontory of the sacrum above the upper margin of the symphysis pubis is on an average about $3\frac{3}{4}$ inches, and a line passing horizontally backward from the latter point would impinge on the junction of the second and third coccygeal bones.

Axes of the Parturient Canal.—By the axis of the pelvis is meant an imaginary line which indicates the direction which the foetus takes during its expulsion. The axis of the brim (Fig. 10) is a line drawn perpendicular to its plane, which would extend from the umbilicus to about the apex of the coccyx; the axis of the outlet of the bony pelvis intersects this, and extends from the centre of the promontory of the sacrum to midway between the tuberosities of the ischia. The axis of the entire pelvic canal is represented by the sum of the axes of an indefinite number of planes at different levels of the pelvic cavity, which forms an irregular parabolic line, as represented in the diagram (Fig. 10, A D).

FIG. 11.



Representing General Axis of Parturient Canal, including the Uterine Cavity and Soft Parts.

It must be borne in mind, however, that it is not the axis of the bony pelvis alone that is of importance in obstetrics. We must always, in considering this subject, remember that the general axis of the parturient canal (Fig. 11) also includes that of the uterine cavity above and of the soft parts below. These are variable in direction according to circumstances; and it is only the axis of that portion of the parturient canal extending between the plane of the pelvic brim and a plane between the

lower edge of the pubic symphysis and the base of the coccyx that is fixed. The axis of the lower part of the canal will vary according to the amount of distension of the perineum during labor; but when this is stretched to its utmost, just before the expulsion of the head, the axis of the plane between the edge of the distended perineum and the lower border of the symphysis looks nearly directly forward. The axis of the uterine cavity generally corresponds with that of the pelvic brim, but it may be much altered by abnormal positions of the uterus, such as anteversion from laxity of the abdominal walls. The foetus, under such circumstances, will not enter the brim in its proper axis, and difficulties in the labor arise. A knowledge of the general direction of the parturient canal is of great importance in practical midwifery in guiding us to the introduction of the hand or instruments in obstetric operations, and in showing us how to obviate difficulties arising from such accidental deviations of the uterus as have been just alluded to.

Cavity of the Pelvis.—The arrangements of the bones in the interior of the pelvic canal (Fig. 12) are important in relation to the mechanism of delivery. A line passing between

FIG. 12.



Side View of Pelvis.

the spine of the ischium and the iliopectineal eminence divides the inner surface of the ischial bone into two smooth plane surfaces, which have received the name of the planes of the ischium. Two other planes are formed by the inner surfaces of the pubic bones in front and by the upper portion of the sacrum behind, both having a direction downward and backward. In studying the mechanism of delivery, it will be seen that many obstetricians attribute to these planes, in conjunction with the spines of the ischia, a very important influence in effecting rotation of the foetal head from the oblique to the antero-posterior diameter of the pelvis.

Development of the Pelvis.—The peculiarities of the pelvis during infancy and childhood are of interest as leading to a knowledge of the manner in which the form observed during adult life is impressed upon it. The sacrum in the pelvis of the child (Fig. 13) is less developed transversely, and is much less deeply curved, than in the adult. The pubes are also much shorter from side to side, and the pubic arch is an acute angle. The result of this narrowness of both the pubes and sacrum is that the transverse diameter of the pelvic brim is shorter instead of longer than the antero-posterior. The sides of the pelvis have a tendency to parallelism, as well as the antero-posterior walls; and this is stated by Wood to be a peculiar characteristic of the infantile pelvis. The iliac bones are not spread out as in adult life, so that the centres of the crests of the ilia are not more distant from each other than the anterior-superior spines. The cavity of the true pelvis is small, and the tuberosities of the ischia are proportionately nearer to

each other than they afterward become; the pelvic viscera are consequently crowded up into the abdominal cavity, which is, for this reason, much more prominent in children than in adults. The bones are soft and semi-cartilaginous until after the period of puberty, and yield readily to the mechanical influences to which they are subjected; and the three divisions of the innominate bone remain separate until about the twentieth year.

As the child grows older the transverse development of the sacrum increases, and the pelvis begins to assume more and more of the adult shape. The mere growth of the bones, however, is not sufficient to account for the change in the shape of the pelvis, and it has been well

FIG. 13.



Pelvis of a Child.

shown by Duncan that this is chiefly produced by the pressure to which the bones are subjected during early life. The iliac bones are acted upon by two principal and opposing forces. One is the weight of the body above, which acts vertically upon the sacral extremity of the iliac beam through the strong posterior sacro-iliac ligaments, and tends to throw the lower or acetabular ends of the sacro-cotyloid beams outward. This outward displacement, however, is resisted, partly by the junction between the two acetabular ends at the front of the pelvis, but chiefly by the opposing force, which is the upward pressure of the lower extremities through the femurs. The result of these counteracting forces is that the still soft bones bend near their junction with the sacrum, and thus the greater transverse development of the pelvic brim characteristic of adult life is established. In treating of pelvic deformities it will be seen that the same forces applied to diseased and softened bones explain the peculiarities of form that they assume.

Pelvis in Different Races.—The researches that have been made on the differences of the pelvis in different races prove that these are not so great as might have been expected. Joulin pointed out that in all human pelvises the transverse diameter was larger than the antero-posterior, while the reverse was the case in all the lower animals, even in the highest simiæ. This observation has been more recently confirmed

by Von Franque,¹ who has made careful measurements of the pelvis in various races. In the pelvis of the gorilla, the oval form of the brim, resulting from the increased length of the conjugate diameter, is very marked. In certain races there is so far a tendency to animality of type that the difference between the transverse and conjugate diameters is much less than in European women, but is not sufficiently marked to enable us to refer any given pelvis to a particular race. Von Franque makes the general observation that the size of the pelvis increases from south to north, but that the conjugate diameter increases in proportion to the transverse in southern races.

Soft Parts in connection with Pelvis.—In closing the description of the pelvis the attention of the student must be directed to the muscular and other structures which cover it. It has already been pointed out that the measurements of the pelvic diameters are considerably lessened by the soft parts, which also influence parturition in other ways. Thus, attached to the crests of the ilia are strong muscles which not only support the enlarged uterus during pregnancy, but are powerful accessory muscles in labor; in the pelvic cavity are the obturator and pyriformis muscles lining it on either side; the pelvic cellular tissue and fasciæ; the rectum and bladder; the vessels and nerves, pressure on which often gives rise to cramps and pains during pregnancy and labor; while below the outlet of the pelvis is closed and its axis directed forward by the numerous muscles forming the floor of the pelvis and perineum. The structures closing the pelvis have been accurately described by Dr. Berry Hart,² who points out that they form a complete diaphragm stretching from the pelvis to the sacrum, in which are three "faults" or "slits" formed by the orifices of the urethra, vagina, and rectum. The first of these is a mere capillary slit, the last is closed by a strong muscular sphincter, while the vagina, in a healthy condition, is also a mere slit, with its walls in accurate apposition. Hence it follows that none of these apertures impairs the structural efficiency of the pelvic floor or the support it gives to the structures above it.

CHAPTER II.

THE FEMALE GENERATIVE ORGANS.

Division according to Function.—The reproductive organs in the female are conveniently divided, according to their function, into—1. The external or copulative organs, which are chiefly concerned in the act of insemination, and are only of secondary importance in parturition; they include all the organs situated externally which form the vulva; and the vagina, which is placed internally and forms the canal

¹ Scanzoni's *Beiträge*, 1867.

² *The Structural Anatomy of the Female Pelvic Floor*.

of communication between the uterus and the vulva. 2. The internal or formative organs: they include the ovaries, which are the most important of all, as being those in which the ovule is formed; the Fallopian tubes, through which the ovule is carried to the uterus; and the uterus, in which the impregnated ovule is lodged and developed.

1. The external organs consist of—

Mons Veneris.—The *mons veneris* (Fig. 14, *f*) is a cushion of adipose and fibrous tissue which forms a rounded projection at the upper part of the vulva. It is in relation above with the lower part of the hypogastric region, from which it is often separated by a furrow, and below it is continuous with the labia majora on either side. It lies over the symphysis and horizontal rami of the pubes. After puberty it is covered

FIG. 14.



External Genitals of Virgin with Diaphragmatic Hymen. (After Sappey.)

a. Labium majus. b. Labium minus. c. Præputium clitoridis. d. Glans clitoridis.
e. Vestibule just above urethral orifice. f. Mons veneris.

with hair. On its integument are found the openings of numerous sweat and sebaceous glands.

Labia Majora.—The *labia majora* (Fig. 14, *a*) form two symmetrical sides to the longitudinal aperture of the vulva. They have two surfaces—one external, of ordinary integument, covered with hair, and another internal, of smooth mucous membrane, in apposition with the correspond-

ing portion of the opposite labium, and separated from the external surface by a free convex border. They are thicker in front, where they run into the mons veneris, and thinner behind, where they are united, in front of the perineum, by a thin fold of integument called the fourchette, which is almost invariably ruptured in the first labor. In the virgin the labia are closely in apposition and conceal the rest of the generative organs. After childbearing they become more or less separated from each other, and in the aged they waste and the internal nymphæ protrude through them. Both their cutaneous and mucous surfaces contain a large number of sebaceous glands, opening either directly on the surface or into the hair-follicles. In structure the labia are composed of connective tissue, containing a varying amount of fat, and parallel with their external surface are placed tolerably close plexuses of elastic tissue, interspersed with regularly arranged smooth muscular fibres. These fibres are described by Broca as forming a membranous sac resembling the dartos of the scrotum, to which the labia majora are analogous. Toward its upper and narrower end this sac is continuous with the external inguinal ring, and in it terminate some of the fibres of the round ligament. The analogy with the scrotum is further borne out by the occasional hernial protrusion of the ovary into the labium, corresponding to the normal descent of the testis in the male.

Labia Minora.—The *labia minora*, or *nymphæ* (Fig. 14, *b*), are two folds of mucous membrane, commencing below, on either side, about the centre of the internal surface of the labium externum; they converge as they proceed upward, bifurcating as they approach each other. The lower branch of this bifurcation is attached to the clitoris (Fig. 14, *c*), while the upper and larger unites with its fellow of the opposite side, and forms a fold round the clitoris known as its prepuce. The nymphæ are usually entirely concealed by the labia majora, but after childbearing and in old age they project somewhat beyond them; then they lose their delicate pink color and soft texture, and become brown, dry, and like skin in appearance. This is especially the case in some of the negro races, in whom they form long projecting folds called the apron.

The surfaces of the nymphæ are covered with tessellated epithelium, and over them are distributed a large number of vascular papillæ, somewhat enlarged at their extremities, and sebaceous glands, which are more numerous on their internal surfaces. The latter secrete an odorous, cheesy matter, which lubricates the surface of the vulva and prevents its folds adhering to each other. The nymphæ are composed of trabeculæ of connective tissue containing muscular fibres.

The Clitoris.—The *clitoris* (Fig. 14, *d*) is a small erectile tubercle situated about half an inch below the anterior commissure of the labia majora. It is the analogue of the penis in the male, and is similar to it in structure, consisting of two corpora cavernosa, separated from each other by a fibrous septum. The crura are covered by the ischio-cavernous muscles, which serve the same purpose as in the male. It has also a suspensory ligament. The corpora cavernosa are composed of a vascular plexus with numerous traversing muscular fibres. The arteries are derived from the internal pudic artery, which gives a branch, the cavernous, to each half of the organ; there is also a dorsal artery dis-

tributed to the prepuce. According to Gussenbauer, these cavernous arteries pour their blood directly into large veins, and a finer venous plexus near the surface receives arterial blood from small arterial branches. By these arrangements the erection of the organ which takes place during sexual excitement is favored. The nervous supply of the clitoris is large, being derived from the internal pudic nerve, which supplies branches to the corpora cavernosa, and terminates in the glands and prepuce, where Paccinian corpuscles and terminal bulbs are to be found. On this account the clitoris has been supposed by some to be the chief seat of voluptuous sensation in the female.

The Vestibule.—The *vestibule* (Fig. 14, e) is a triangular space, bounded at its apex by the clitoris and on either side by the folds of the nymphæ. It is smooth, and, unlike the rest of the vulva, is destitute of sebaceous glands, although there are several groups of muciparous glands opening on its surface. At the centre of the base of the triangle, which is formed by the upper edge of the opening of the vagina, is a prominence, distant about an inch from the clitoris, on which is the orifice of the urethra. This prominence can be readily made out by the finger, and the depression upon it—leading to the urethra—is of importance as our guide in passing the female catheter. This little operation ought to be performed without exposing the patient, and it is done in several ways. The easiest is to place the tip of the index finger of the left hand (the patient lying on her back) on the apex of the vestibule, and slip it gently down until we feel the bulb of the urethra and the dimple of its orifice, which is generally readily found. If there is any difficulty in finding the orifice, it is well to remember that it is placed immediately below the sharp edge of the lower border of the symphysis pubis, which will guide us to it. The catheter (and a male elastic catheter is always the best, especially during labor, when the urethra is apt to be stretched) is then passed under the thigh of the patient and directed to the orifice of the urethra by the finger of the left hand, which is placed upon it. We must be careful that the instrument is really passed into the urethra, and not into the vagina. It is advisable to have a few feet of elastic tubing attached to the end of the catheter, so that the urine can be passed into a vessel under the bed without uncovering the patient. If the patient be on her side, in the usual obstetric position, the operation can be more readily performed by placing the tip of the finger in the vagina and feeling its upper edge. The orifice of the urethra lies immediately above this, and if the catheter be slipped along the palmar surface of the finger, it can generally be inserted without much trouble. If, however, as is often the case during labor, the parts are much swollen, it may be difficult to find the aperture, and it is then always better to look for the opening than to hurt the patient by long-continued efforts to feel it.

The Urethra.—The *urethra* is a canal $1\frac{1}{2}$ inches in length, and it is intimately connected with the anterior wall of the vagina, through which it may be felt. It is composed of muscular and erectile tissue, and is remarkable for its extreme dilatability—a property which is turned to practical account in some of the operations for stone in the female bladder.

Orifice of the Vagina.—The orifice of the vagina is situated immediately below the bulb of the urethra. In virgins it is a circular opening, but in women who have borne children or practised sexual intercourse it is, in the undistended state, a fissure, running transversely and at right angles to that between the labia.¹ In virgins it is generally more or less blocked up by a fold of mucous membrane, containing some cellular tissue and muscular fibres, with vessels and nerves, which is known as the *hymen*. This is most often crescentic in shape, with the concavity of the crescent looking upward; sometimes, however, it is circular with a central opening, or cribriform; or it may even be entirely imperforate, and this gives rise to the retention of the menstrual secretion. These varieties of form depend on the peculiar mode of development of the fold of vaginal mucous membrane which blocks up the orifice of the vagina in the foetus, and from which the hymen is formed. The density of the membrane also varies in different individuals. Most usually it is very slight, so as to be ruptured in the first sexual approaches, or even by some accidental circumstance, such as stretching the limbs, so that its absence cannot be taken as evidence of want of chastity. A knowledge of this fact is of considerable importance from a medico-legal point of view. Sometimes it is so tough as to prevent intercourse altogether, and may require division by the knife or scissors before this can be effected; and at others it rather unfolds than ruptures, so that it may exist even after impregnation has been effected, and it has been met with intact in women who have habitually led unchaste lives. In a few rare cases it has even formed an obstacle to delivery, and has required incision during labor.

Carunculæ Myrtiformes.—The *carunculæ myrtiformes* are small fleshy tubercles, varying from two to five in number, situated around the orifice of the vagina, and which are generally supposed to be the remains of the ruptured hymen. Schroeder, however, maintains that they are only formed after childbearing in consequence of parts of the hymen having been destroyed by the injuries received during the passage of the child.

Vulvo-vaginal Glands.—Near the posterior part of the vaginal orifice, and below the superficial perineal fascia, are situated two conglomerate glands which are the analogues of Cowper's glands in the male. Each of these is about the size and shape of an almond, and is contained in a cellular fibrous envelope. Internally they are of a yellowish-white color, and are composed of a number of lobules separated from each other by prolongations of the external envelope. These give origin to separate ducts which unite into a common canal, about half an inch in length, which opens in front of the attached edge of the hymen in virgins, and in married women at the base of one of the *carunculæ myrtiformes*. According to Huguier, the size of the glands varies much in different women, and they appear to have some connection with the ovary, as he has always found the largest gland to be on the same side as the largest ovary. They secrete a glairy, tenacious fluid, which is ejected in jets during the sexual orgasm, probably through the spasmodic action of the perineal muscles. At other times their secretion serves the purpose of

¹ Hart, *op. cit.*


lubricating the vulva, and thus preserves the sensibility of its mucous membrane.

Fossa Navicularis.—Immediately behind the hymen in the unmarried, and between it and the perineum, is a small depression called the *fossa navicularis*, which disappears after childbearing.

Perineum.—The *perineum* separates the orifice of the vagina from that of the rectum. It is about $1\frac{1}{2}$ inches in breadth, and is of great obstetric interest, not only as supporting the internal organs from below, but because of its action in labor. It is largely stretched and distended by the presenting part of the child, and if unusually tough and unyielding may retard delivery, or it may be torn to a greater or less extent, thus giving rise to various subsequent troubles.

Vascular Supply of the Vulva.—The structures described above together form the *vulva*, and they are remarkable for their abundant vas-

FIG. 15.




Vascular Supply of Vulva. (After Kobelt.)

a. Bulb of vestibule. b. Muscular tissue of vagina. c, d, e, f. The clitoris and muscles. g, h, i, k, l, m, n. Veins of the nymphæ and clitoris communicating with the epigastric and obturator veins.

cular and nervous supply. The former constitutes an erectile tissue, similar to that which has already been described in the clitoris, and which is especially marked about the bulb of the vestibule (Fig. 15). From this point and extending on either side of the vagina there is a well-marked plexus of convoluted veins, which, in their distended state, are likened by Dr. Arthur Farre to a filled leech. The erection of the erectile tissue, as well as that of the clitoris, is brought about under excitement, as in the male, by the compression of the efferent veins by the contraction of the ischio-cavernous muscles, and by that of a thin layer of muscular tissues surrounding the orifice of the vagina and described as the constrictor vaginæ.

The Vagina.—The *vagina* is the canal which forms the communication between the external and internal generative organs, through which the semen passes to reach the uterus, the menses flow, and the fœtus is expelled. Roughly speaking, it lies in the axis of the pelvis, but its opening is placed anterior to the axis of the pelvic outlet, so that its lower portion is curved forward, so as to lie parallel to the pelvic brim. It is narrow below, but dilated above, where the cervix uteri is inserted into it, so that it is more or less conoidal in shape. Under ordinary circumstances, especially in the virgin, the anterior and posterior walls lie in close contact with each other (see Plate I.), and there is, strictly speaking, no vaginal canal, although they are capable of wide distension, as in copulation and during the passage of the fœtus. The anterior wall of the vagina is shorter than the posterior, the former measuring on an average $2\frac{1}{2}$ inches, the latter 3 inches; but the length of the canal varies greatly in different subjects and under certain circumstances. In front the vagina is closely connected with the base of the bladder, so that when the vagina is prolapsed, as often occurs, it drags the bladder with it (Fig. 17); behind, it is in relation with the rectum, but less intimately; laterally with the broad ligaments and pelvic fascia; and superiorly with the lower portion of the uterus and folds of peritoneum both before and behind. The vagina is composed of mucous, muscular, and cellular coats. The mucous lining is thrown into numerous folds. These start from longitudinal ridges which exist on both the anterior and posterior walls, but most distinctly on the anterior. They are very numerous in the young and unmarried, and greatly increase the sensitive surface of the vagina (Fig. 16). After childbearing, and in the aged, they become

FIG. 16.



Right Half of Virgin Vagina, with Walls held apart, showing the abundant *Transverse Rugæ*, the greater depth of the Vagina above than below, and the Hymenal Segment. (After Hart.)

atrophied, but they never completely disappear, and toward the orifice of the vagina, where they exist in greatest abundance, they are always to be met with. The whole of the mucous membrane is lined with tessellated epithelium, and it is covered with a large number of papillæ,

either conical or divided, which are highly vascular and project into the epithelial layer. Unlike the vulvar mucous membrane, that of the vagina seems to be destitute of glands. Beneath the epithelial layer is a submucous tissue containing a large number of elastic and some muscular fibres, derived from the muscular walls of the vagina. These are strong and well developed, especially toward the ostium vaginae. They consist of two layers—an internal longitudinal and an external circular—with oblique decussating fibres connecting the two. Below they are attached to the ischio-pubic rami, and above they are continuous with the muscular coat of the uterus. The muscular tissue of the vagina

FIG. 17.



Longitudinal Section of Body, showing Relations of Generative Organs.


increases in thickness during pregnancy, but to a much less degree than that of the uterus. Its vascular arrangements, like those of the vulva, are such as to constitute an erectile tissue. The arteries form an intricate network around the tube, and eventually end in a submucous capillary plexus from which twigs pass to supply the papillæ; these again give origin to venous radicles which unite into meshes freely interlacing with each other and forming a well-marked venous plexus.

2. *The Internal Organs of Generation.*—The internal organs of generation consist of the uterus, the Fallopian tubes, and the ovaries; and in connection with them we have to study the various ligaments and folds of peritoneum which serve to maintain the organs in position, along with certain accessory structures. Physiologically, the most important of all the generative organs are the ovaries, in which the ovules

are formed, and which dominate the entire reproductive life of the female. The Fallopian tubes, which convey the ovule to the uterus, and the uterus itself—whose main function is to receive, nourish, and eventually expel the impregnated product of the ovary—may be said to be, in fact, accessory to these viscera. Practically, however, as obstetricians, we are chiefly concerned with the uterus, and may conveniently commence with its description.

The Uterus.—The *uterus* is correctly described as a pyriform organ, flattened from before backward, consisting of the body, with its rounded fundus, and the cervix, which projects into the upper part of the vaginal canal. In the adult female it is deeply situated in the pelvis, being placed between the bladder in front and the rectum behind, its fundus being below the plane of the pelvic brim (Fig. 18). It only assumes this position, however, toward the period of puberty; and in the *fœtus* it is placed much higher, and lies, indeed, entirely within the cavity of the abdomen. It is maintained in this position partly by being slung

FIG. 18.




Transverse Section of the Body, showing Relations of the Fundus Uteri.

m. Pubes. *a a* (in front). Remainder of hypogastric arteries. *a a* (behind). Spermatic vessels and nerves. *B.* Bladder. *L L.* Round ligaments. *U.* Fundus uteri. *t, t.* Fallopian tubes. *o, o.* Ovaries. *r.* Rectum. *g.* Right ureter, resting on the psoas muscle. *e.* Utero-sacral ligaments. *e.* Last lumbar vertebra.

by its ligaments, which we shall subsequently study, and partly by being supported from below by the pelvic cellular tissue and the fleshy column of the vagina. The result is that the uterus, in the healthy female, is a perfectly movable body, altering its position to suit the condition of the surrounding viscera, especially the bladder and rectum, which are subjected to variations of size according to their fulness or emptiness. When from any cause—as, for example, some peri-uterine inflammation producing adhesions to the surrounding textures—the mobility of the organ is interfered with, much distress ensues, and if pregnancy supervenes more or less serious consequences may result. Generally speaking,

the uterus may be said to lie in a line roughly corresponding with the axis of the pelvic brim, its fundus being pointed forward, and its cervix lying in such a direction that a line drawn from it would impinge on the junction between the sacrum and coccyx. According to some authorities, the uterus in early life is more curved in the anterior direction, and is, in fact, normally in a state of ante flexion. Sappey holds that this is not necessarily the case, but that the amount of anterior curvature depends on the emptiness or fulness of the bladder, on which the uterus, as it were, moulds itself in the unimpregnated state. It is believed also that the body of the uterus is very generally twisted somewhat obliquely, so that its interior surface looks a little toward the right side, this probably depending on the presence and frequent distension of the rectum in the left side of the pelvis. The anterior surface of the uterus is convex, and is covered in three-fourths of its extent by the peritoneum, which is intimately adherent to it. Below the reflexion of the membrane it is loosely connected by cellular tissue to the bladder, so that any downward displacement of the uterus drags the bladder along with it. The posterior surface is also convex, but more distinctly so than the anterior, as may be observed in looking at a transverse section of the organ (Fig. 19). It is also covered by peritoneum, the

FIG. 19.




Transverse Section of Uterus.

reflexion of which on the rectum forms the cavity known as Douglas's pouch. The fundus is the upper extremity of the uterus, lying above the points of entry by the Fallopian tubes. It is only slightly rounded in the virgin, but becomes more decidedly and permanently rounded in the woman who has borne children.

Its Surfaces.—Until the period of puberty the uterus remains small and undeveloped (Fig. 20); after that time it reaches the adult size, at which it remains until menstruation ceases, when it again atrophies. If the woman has borne children it always remains larger than in the nullipara. In the virgin adult the uterus measures $2\frac{1}{2}$ inches from the orifice to the fundus, rather more than half being taken up by the cervix. Its greatest breadth is opposite the insertion of the Fallopian tubes; its greatest thickness, about 11 or 12 lines, opposite the centre of its body. Its average weight is about 9 or 10 drachms. Independently of pregnancy, the uterus is subject to great alterations of size toward the menstrual period, when, on account of the congestion then present, it enlarges, sometimes, it is said, considerably. This fact should be borne in mind, as this periodical swelling might be taken for an early pregnancy.

Regional Divisions.—For the purpose of description the uterus is conveniently divided into the *fundus*, with its rounded upper extremity, situated between the insertions of the Fallopian tubes; the *body*, which is bounded above by the insertions of the Fallopian tubes and below by the upper extremity of the cervix, and which is the part chiefly con-

FIG. 20.



Uterus and Appendages in an Infant. (After Farre.)

cerned in the reception and growth of the ovum; and the *cervix*, which projects into the vagina, and dilates during labor to give passage to the child. The cervix is conical in shape, measuring 11 to 12 lines transversely at the base, and 6 or 7 in the antero-posterior direction; while at the apex it measures 7 to 8 transversely, and 5 antero-posteriorly. It projects about 4 lines into the canal of the vagina, the remainder of the cervix being placed above the reflexion of the vaginal mucous membrane. It varies much in form in the virgin and nulliparous married woman and in the woman who has borne children; and the differences are of importance in the diagnosis of pregnancy and uterine disease. In the virgin it is regularly pyramidal in shape. At its lower extremity is the opening of the external os uteri, forming a small transverse fissure, sometimes difficult to feel, and generally described as giving a sensation to the examining finger like the extremity of the cartilage at the tip of the nose. It is bounded by two lips, the anterior of which is apparently larger on account of the position of the uterus. The surface of the cervix and the borders of the os are very smooth and regular.

Changes after Childbirth.—In women who have borne children these parts become considerably altered. The cervix is no longer conical, but is irregular in form and shortened. The lips of the os uteri become fissured and lobulated, on account of partial lacerations which have occurred during labor. The os is larger and more irregular in outline, and is sometimes sufficiently patulous to admit the tip of the finger. In old age the cervix atrophies, and after the change of life it not uncom-

monly entirely disappears, so that the orifice of the os uteri is on a level with the roof of the vagina.

Internal Surface of the Uterus.—The internal surface of the uterus comprises the cavities of the body and cervix—the former being rather less than the latter in length in virgins, but about equal in women who have borne children—separated from each other by a constriction forming the upper boundary of the cervical canal. The cavity of the body is triangular in shape, the base of the triangle being formed by a line joining the openings of the Fallopian tubes, its apex by the upper orifice of the cervix, or internal os, as it is sometimes called. In the virgin its boundaries are somewhat convex, projecting inward. After child-bearing they become straight or slightly concave. The opposing surfaces of the cavity are always in contact in the healthy state, or are only separated from each other by a small quantity of mucus.

Cavity of the Cervix.—The cavity of the cervix is spindle-shaped or fusiform, narrower above and below at the internal and external os uteri, and somewhat dilated between these two points. It is flattened

FIG. 21.



Portion of Interior of Cervix. (Enlarged nine diameters.) (After Tyler Smith and Hassall.)

from before backward, and its opposing surfaces also lie in contact, but not so closely as those of the body. On the mucous lining of the anterior and posterior surfaces is a prominent perpendicular ridge, with a lesser one at each side, from which transverse ridges proceed at more or less acute angles. They have received the name of the *arbor vitæ*. According to Guyon, the perpendicular ridges are not exactly opposite, so that they fit into each other, and serve more completely to fill up the cavity of the cervix, especially toward the internal os (Fig. 21). The

arbor vitæ is most distinct in the virgin, and atrophies considerably after childbearing.

The superior extremity of the cervical canal forms a narrow isthmus separating it from the cavity of the body, and measuring about $\frac{3}{8}$ ths of an inch in diameter. Like the external os, it contracts after the cessation of menstruation, and in old age sometimes becomes entirely obliterated.

Structure of the Uterus.—The uterus is composed of three principal structures—the peritoneal, muscular, and mucous coats. The peritoneum forms an investment to the greater part of the organ, extending downward in front to the level of the os internum, and behind to the

top of the vagina, from which points it is reflected upward on the bladder and rectum respectively. At the sides the peritoneal investment is not so extensive, for a little below the level of the Fallopian tubes the peritoneal folds separate from each other, forming the broad ligaments (to be afterward described); here it is that the vessels and nerves supplying the uterus gain access to it. At the upper part of the organ the peritoneum is so closely adherent to the muscular tissue that it cannot be separated from it; below, the connection is more loose. The mass of the uterine tissue, both in the body and cervix, consists of unstripped muscular fibres, firmly united together by nucleated connective tissue and elastic fibres. The muscular fibre-cells are large and fusiform, with very attenuated extremities, generally containing in their centre a distinct nucleus. These cells, as well as their nuclei, become greatly enlarged during pregnancy (Fig. 23); according to Stricker, this is only

FIG. 22.




Muscular Fibres of Unimpregnated Uterus.
(After Farre.)

- a. Fibres united by connective tissue.
b. Separate fibres and elementary corpuscles.

toneum is so closely adherent to the muscular tissue that it cannot be separated from it; below, the connection is more loose. The mass of the uterine tissue, both in the body and cervix, consists of unstripped muscular fibres, firmly united together by nucleated connective tissue and elastic fibres. The muscular fibre-cells are large and fusiform, with very attenuated extremities, generally containing in their centre a distinct nucleus. These cells, as well as their nuclei, become greatly enlarged during pregnancy (Fig. 23); according to Stricker, this is only

FIG. 23.



Developed Muscular Fibres from the Gravid Uterus. (After Wagner.)

the case with the muscular fibres which play an important part in the expulsion of the fœtus, those of the outermost and innermost layers not sharing in the increase of size.¹ In addition to these developed fibres there are, especially near the mucous coat, a number of round elementary corpuscles, which are believed by Dr. Farre² to be the elementary form of the muscular fibres, and which he has traced in various intermediate

¹ *Comparative Histology*, vol. iii. Syd. Soc. Trans., p. 477.

² *The Uterus and its Appendages*, p. 632.

states of development. Dr. John Williams¹ believes that a great part of the muscular tissue of the uterus, rather more indeed than three-fourths of its thickness, is an integral part of the mucous membrane, analogous to the muscularis mucosæ of the mucous membrane of the alimentary canal. This he describes as being separated from the rest of the muscular tissue by a layer of rather loose connective tissue, containing numerous vessels. In early fetal life, and in the uteri of some of the lower animals, this appearance is very distinct; in the adult female uterus, however, it cannot be readily made out.

Arrangement of the Muscular Fibres.—On examining the uterine tissue in an unimpregnated condition no definite arrangement of its muscular fibres can be made out, and the whole seem blended in inextricable confusion. By observation of their relations when hypertrophied during pregnancy, Helié² has shown that they may, speaking roughly, be divided into three layers—an external; a middle, chiefly longitudinal; and an internal, chiefly circular. Into the details of their distribution, as described by him, it is needless to enter at length. Briefly, however, he describes the external layer as arising posteriorly at the junction of the body and cervix, and spreading upward and over the fundus. From this are derived the muscular fibres found in the broad and round ligaments, and more particularly described by Rouget. The middle layer is made up of strong fasciculi, which run upward, but decussate and unite with each other in a remarkable manner, so that those which are at first superficial become most deeply seated, and vice versa. The muscular fasciculi which form this coat curve in a circular manner round the large veins, so as to form a species of muscular canal through which they run. This arrangement is of peculiar importance, as it affords a satisfactory explanation of the mechanism by which hemorrhage after delivery is prevented. The internal layer is mainly composed of circular rings of muscular fibres, beginning round the openings of the Fallopian tubes, and forming wider and wider circles which eventually touch and interlace with each other. They surround the internal os, to which they form a kind of sphincter. In addition to these circular fibres on the internal uterine surface, both anteriorly and posteriorly, there is a well-marked triangular layer of longitudinal fibres, the base being above and the apex below, which sends muscular fasciculi into the mucous membrane.

Its Mucous Membrane.—The anatomy of the lining membrane of the uterus has been the subject of considerable discussion. Its existence has been denied by many authorities, most recently by Snow Beck,³ who maintains that it is in no sense a mucous membrane, but only a softened portion of true uterine tissue. It is, however, pretty generally admitted by the best authorities that it is essentially a mucous membrane, differing from others only in being more closely adherent to the subjacent structures in consequence of not possessing any definite connective-tissue framework.

It is a pale pink membrane of considerable thickness, most marked at

¹ "On the Structure of the Mucous Membrane of the Uterus," *Obstet. Journ.*, 1875.

² *Recherches sur la Disposition des Fibres musculaires de l'Utérus*, Paris, 1869.

³ *Obst. Trans.*, vol. xiii. p. 294.

the centre of the body, where it forms from one-eighth to one-fourth of the thickness of the whole uterine walls. At the internal os uteri it

FIG. 24.



Lining Membrane of Uterus, showing Network of Capillaries and Orifices of Uterine Glands. (After Farre.)


From the body.

From orifice of Fallopian tube.

terminates by a distinct border, which separates it from the mucous membrane lining the cervical cavity.

The Utricular Glands.—On the surface of the mucous membrane may be observed a multitude of little openings, about $\frac{1}{30}$ th of a line in width (Fig. 24). These are the orifices of the utricular glands, which are found in immense numbers all over the cavity of the uterus, and very closely agglomerated together. They are little

FIG. 25.



The Course of the Glands in the fully-developed Mucous Membrane of the Uterus—viz. just before the onset of a Menstrual Period. (After Williams.)

cul-de-sacs, narrower at their mouths than in their length, the blind extremities of which are found in the subjacent tissues (Fig. 26). Williams describes them as running obliquely toward the surface at the lower third of the cavity, perpendicularly at its middle, while toward the fundus they are at first perpendicular, and then oblique in their course (Fig. 25). By others they are described as being often twisted and corkscrew-like. One or more may unite to form a common orifice, several of which may open together in little pits or depressions on the surface of the mucous membrane. These glands are composed of structureless membrane lined with epithelium, the precise character of which is doubtful. By some it is described as columnar, by others tessellated, and by some again as ciliated. The most generally received opinion is that it is columnar, but not ciliated; therein differing from the epithelium covering the surface of the membrane, which is undoubtedly ciliated, the movements of the cilia being from within outward. Williams, however, has observed cilia in active movement on the columnar epithelium lining the glands, and also states that at the deep-seated extremities of the glands, which penetrate between the muscular fibres for some distance, the columnar epi-

thelium is replaced by rounded cells. The capillaries of the mucous membrane run down between the tubes, forming a lacework on their surfaces and round their orifices. No true papillæ exist in the membrane lining the uterine cavity. The mucous membrane of the uterus is peculiar in being always in a state of change and alteration, being thrown off at each menstrual period in the form of débris in consequence of fatty degeneration of its structures, and re-formed afresh by proliferation of the cells of the muscular and connective tissues, probably from below upward, the new membrane commencing at the internal os. Hence its appearance and structure vary considerably

FIG. 28.



Vertical Section through the Mucous Membrane of the Human Uterus. (After Turner.)


e. Columnar epithelium; the cilia are not represented. *g.g.* Utricular glands. *ct. ct.* Interglandular connective tissue. *v.v.* Blood-vessels. *m.m.* Muscularis mucosæ (43°).

according to the time at which it is examined. The subject, however, will be more particularly studied in connection with menstruation.

Mucous Membrane of the Cervix.—The mucous membrane of the cervix is much thicker and more transparent than that of the body of the uterus, from which it also differs in certain structural peculiarities. The general arrangements of its folds and surface have already been described. The lower half of the membrane lining the cavity of the cervix, and the whole of that covering its external or vaginal portion, are closely set with a large number of minute filiform or clavate papillæ (Fig. 27). Their structure is similar to that of the mucous membrane itself, of which they seem to be merely elevations. They each contain a vascular loop (Fig. 28), and they are believed by Kilian and Farre to be mainly concerned in giving sensibility to this part of the generative

tract. All over the interior of the cervix, both on the ridges of the mucous membrane and between their folds, are a very large number of

FIG. 27.



Villi of Os Uteri stripped of Epithelium. (After Tyler Smith and Hassall.)

mucous follicles, consisting of a structureless membrane lined with cylindrical epithelium and intimately united with connective tissue.

FIG. 28.



Villi of Uterus covered with Pavement Epithelium, and containing Looped Vessels. (After Tyler Smith and Hassall.)

They cease at the external orifice of the cervix, and they secrete the thick, tenacious, and alkaline mucus which is generally found filling the cervical cavity. The transparent follicles, known as the "*ovula Nabothii*," which are sometimes found in considerable numbers in the cavity of the cervix, consist of mucous follicles, the mouths of which have become obstructed and their canals distended by mucous secretion. The lower third of the cervical canal, as well as the exterior of the cervix, is covered with pavement epithelium; while on its upper portion is found a columnar and ciliated epithelium similar to that lining the uterine cavity.

Bandl¹ describes the cervical mucous membrane as extending much higher in the virgin than in women who have borne children, being traceable in the former nearly to the middle of the body of the uterus. During the first pregnancy he believes that the upper portion of the cervix is taken up into the body of the uterus, its mucous membrane never regaining the arrangement peculiar to that of the cervical canal.

Vessels of the Uterus.—The arteries of the uterus are derived from the internal iliac and from the ovarian. They enter the uterus between the folds of the broad ligaments, and, penetrating its muscular coat, anastomose freely with each other and with the corresponding vessels of the opposite side. Their walls are thick and well developed, and they are remarkable for their very tortuous course, forming spiral curves, especially in the upper part of the uterus. They end in minute capillaries which form the fine meshes surrounding the glands, and in the cervix give off the loops entering the papillæ. Beneath the uterine mucous membrane these capillaries form a plexus, terminating in veins without valves, which unite with each other to form the large veins traversing the substance of the uterus, known during pregnancy as the uterine sinuses, the walls of which are closely adherent to the uterine tissues. These veins, freely anastomosing with each other, pass outward to the folds of the broad ligaments, where they unite to form, with the ovarian and vaginal veins, a large and well-developed venous network, known as the *pampiniform plexus*.

The Lymphatics of the Uterus.—The lymphatics of the uterus are large and well developed, and they have recently, and with much probability, been supposed to play an important part in the production of certain puerperal diseases. A more minute knowledge than we at present possess of their course and distribution will probably throw much light on their influence in this respect. According to the researches of Leopold,² who has studied their minute anatomy carefully, they originate in lymph-spaces between the fine bundles of connective tissue forming the basis of the mucous lining of the uterus. Here they are in intimate contact with the utricular glands and the ultimate ramifications of the uterine blood-vessels. As they pass into the muscular tissue they become gradually narrowed into lymph-vessels and spaces, which have a very complicated arrangement, and which eventually unite together in the external muscular layer, especially on the sides of the uterus, to form large canals which probably have valves. Immediately under this peritoneal covering these lymph-vessels form a large and characteristic

¹ *Arch. f. Gyn.*, B. xiv. S. 237.

² *Ibid.*, Bd. vi. Heft i.

network covering the anterior and posterior surfaces of the uterus, and present, in various parts of their course, large ampullæ. They then spread over the Fallopian tubes. The lymphatics of the body of the uterus unite with the lumbar glands, those of the cervix with the pelvic glands.

The Nerves of the Uterus.—The distribution and arrangement of the nerves of the uterus have been the subject of much controversy. They are derived mainly from the ovarian and hypogastric plexuses, inosculating freely with each other between the folds of the broad ligament, from which they enter the muscular tissue of the uterus, generally, but not invariably, following the course of the arteries. They are chiefly derived from the sympathetic, but, as the hypogastric plexus is connected with the sacral nerves, it is probable that some fibres from the cerebro-spinal system are distributed to the cervix. It is now generally admitted that nervous filaments are distributed to the cervix, even as far as the external os, although their existence in this situation has been denied by Jobert and other writers. The ultimate distribution of the nerves is not yet made out. Polle describes a nerve-filament as entering the papillæ of the cervical mucous membrane along with the capillary loop, and Frankenhauser says the nerve-fibres surround the muscles of the uterus in the form of plexuses and terminate in the nuclei of the muscle-cells.

Anomalies of the Uterus.—Various abnormal conditions of the uterus and vagina are occasionally met with, which it is necessary to mention, as they may have an important practical bearing on parturition. The most frequent of these is the existence of a double, or partially double,

FIG. 29.



Bifid Uterus. (After Farre.)

uterus (Fig. 29), similar to that found normally in many of the lower animals. This abnormality is explained by the development of the organ during fetal life. The uterus is formed out of structures existing only in early fetal life, known as the Wolffian bodies. These consist of a number of tubes, situated on either side of the vertebral column, and opening externally into an excretory duct. Along their external border a hollow canal is formed, termed the canal of Müller, which, like the excretory ducts, proceeds to the common cloaca of the digestive and urinary organs, which then exists. The canal of Müller unites with

its fellow of the opposite side to form the uterus and Fallopian tubes in the female, and subsequently the central partition at their point of junction disappears. If, however, the progress of development be in any way checked, the central partition may remain. Then we have produced either a complete double uterus or the uterus bicornis, which is bifid at its upper extremity only; or a double vagina, each leading to a separate uterus.

Pregnancy in Cases of Bifid Uterus.—If pregnancy occur in any of these anomalous uteri—and many such cases are recorded—serious troubles may follow. It may happen that one horn of the double uterus is not sufficiently large to admit of pregnancy going on to term, and rupture may occur. It is supposed that some cases, presumed to be tubal gestation, were really thus explicable. Impregnation may also occur in the two cornua at different times, leading to superfœtation. It is, however, quite possible that impregnation may occur in one horn of a bifid uterus, and labor be completed without anything unusual being observed. A remarkable case of this sort has been recorded by Dr. Ross of Brighton,¹ in which a patient miscarried of twins on July 16, 1870, and on October 31, fifteen weeks later, was delivered of a healthy child. Careful examination showed the existence of a complete double uterus, each side of which had been impregnated. Curiously enough, this patient had formerly given birth to six living children at term, nothing remarkable having been observed in her labors. It can only rarely happen that, under such circumstances, so favorable a result will follow, and more or less difficulty and danger may generally be expected. Occasionally the vagina only is double, the uterus being single. Dr. Matthews Duncan has recorded some cases of this kind² in which the vaginal septum formed an obstacle to the birth of the child, and required division.

Ligaments of the Uterus.—The various folds of peritoneum which invest the uterus serve to maintain it in position, and they are described as its ligaments. They are the broad, the vesico-uterine, and sacro-uterine ligaments; the round ligaments are not peritoneal folds like the others.


[Within a few years we have had no less than five reports of cases of labor in New York and Philadelphia in which it was claimed that tubal fetal cysts had discharged their contents through the uterus and vagina. This is certainly not in accordance with what is usually taught and believed in regard to the termination of Fallopian or interstitial pregnancies. The illustration here introduced is in evidence that a uterus may be duplex and still bear the outward form of a single organ. Cases of this type have been distinctly recognized during life by Drs. Baer and Drysdale of Philadelphia; and the patient of the former bore twins, whose placentæ were discharged separately and at an interval of a quarter of an hour. Two of the reports already referred to are illustrated by drawings representing a normal uterus and a dilated Fallopian tube. How such a cyst is to discharge its contents other than by the usual way of bursting into the pelvic cavity I cannot comprehend, particularly when we consider that in what is called *interstitial* pregnancy the thickest

¹ *Lancet*, August, 1871.

² *Researches in Obstetrics*, p. 443.

part of the cyst-wall is next to the uterine cavity, and that rupture takes place on the remote side. Where the uterus has but one cornu, or where one cornu is in a rudimentary state, and pregnancy occurs, we may fall into the error of supposing that it is of the Fallopian tube. Such a cornu may be emptied *per vias naturales*, but the usual terminus is by a rupture of the sac. Dr. Sanger of Leipzig calls this form "*gynatretic*" pregnancy, and has collected 21 cases which ended fatally in the first six months by rupture, and 3 in which a *lithopædion* formed, one of which was successfully operated upon by Koeberle.^[1] The impregnated cornu has been removed, successfully in two cases under Salin of Stockholm and Sanger, respectively, and unsuccessfully under Litzmann of Kiel. Sanger believes it possible to distinguish during life a Fallopian pregnancy from one of a rudimentary cornu, but this is not the

FIG. 30.



Partitioned Uterus. (Kussmaul.)

general opinion, as even after death certain anatomical points must be relied upon. We must note where the round ligament is given off, it being between the cyst and the uterus in a Fallopian pregnancy, and at the distal side of an impregnated cornu. It is to be hoped that the five cases of New York and Philadelphia under dispute may some day be examined by autopsy, and the true character of the pregnancies determined. If a Fallopian foetal cyst can discharge itself into the uterus, as claimed, it is strange that an impregnated rudimentary cornu almost universally fails to do so, but ends fatally by rupture.—E.D.]

The Broad Ligaments.—The *broad ligaments* extend from either side of the uterus, where their laminae are separated from each other, transversely across to the pelvic wall, and thus divide the cavity of the pelvis

[¹ *Trans. International Med. Congress of Copenhagen*, Aug. 16, 1884.]

into two parts; the anterior containing the bladder, the posterior the rectum. Their upper borders are divided into three subsidiary folds, the anterior of which contains the round ligament, the middle the Fallopian tube, and the posterior the ovary. The arrangement has received the name of the *ala vespertilionis*, from its fancied resemblance to a bat's wing. Between the folds of the broad ligaments are found the uterine vessels and nerves, and a certain amount of loose cellular tissue con-

FIG. 31.



Adult Parovarium, Ovary, and Fallopian Tube. (After Kobelt.)

tinuous with the pelvic fasciæ. Here is situated that peculiar structure called the organ of Rosenmüller, or the *parovarium* (Fig. 31), which is the remains of the Wolffian body, and corresponds to the epididymus in the male. This may best be seen in young subjects by holding up the broad ligaments and looking through them by transmitted light; but it exists at all ages. It consists of several tubes (eight or ten according to Farre, eighteen or twenty according to Bankes¹), which are tortuous in their course. They are arranged in a pyramidal form, the base of the pyramid being toward the Fallopian tube, its apex being lost on the surface of the ovary. They are formed of fibrous tissue, and lined with pavement epithelium. They have no excretory duct or communication with either the uterus or ovary, and their function, if they have any, is unknown.

Muscular Fibres between its Folds.—A number of muscular fibres are also found in this situation, lying between the meshes of the connective tissue. They have been particularly studied by Rouget, who describes them as interlacing with each other, and forming an open network continuous with the muscular tissues of the uterus (Fig. 32). They are divisible into two layers, the anterior of which is continuous with the muscular fibres of the anterior surface of the uterus, and goes to form part of the round ligament; the posterior arises from the posterior wall of the uterus, and proceeds transversely outward to become attached to the sacro-iliac synchondrosis. A continuous muscular envelope is thus formed, which surrounds the whole of the uterus, Fallopian tubes, and

¹ Bankes, *On the Wolffian Bodies.*

ovaries. Its function is not yet thoroughly established. It is supposed to have the effect of retracting the stretched folds of peritoneum after delivery, and more especially of bringing the entire generative organs into harmonious action during menstruation and the sexual orgasm; in this way explaining, as we shall subsequently see, the mechanism by which the fimbriated extremity of the Fallopian tube grasps the ovary prior to the rupture of a Graafian follicle.

The Round Ligaments.—The round ligaments are essentially muscular in structure. They extend from the upper border of the uterus, with the fibres of which their muscular fibres are continuous, transversely and then obliquely downward, until they reach the inguinal rings, where they blend with the cellular tissue. In the first part of their course the

FIG. 32.



Posterior View of Muscular and Vascular Arrangements. (After Rouget.)

1, 2, 3. Vaginal, cervical, and uterine plexuses. 4. Arteries of body of uterus. 5. Arteries supplying ovary. 6. Muscular broad ligament. 7. Fibres attached to vagina, symphysis pubis, and sacro-iliac joint. 8. Muscular broad ligament from uterus and broad ligaments. 9, 10, 11, 12. Fasciculi attached to ovary and Fallopian tube.

muscular fibres are solely of the unstriped variety, but soon they receive striped fibres from the transversalis muscles and the columns of the inguinal ring, which surround and cover the unstriped muscular tissue. In addition to these structures they contain elastic and connective tissue and arterial, venous, and nervous branches; the former from the iliac or crurae arteries, the latter from the genito-crural nerve. According to Mr. Ruiney, the principal function of these ligaments is to draw the

uterus toward the symphysis pubis during sexual intercourse, and thus to favor the ascent of the semen.

The Vesico-uterine Ligaments.—The *vesico-uterine ligaments* are two folds of peritoneum passing in front from the lower part of the body of the uterus to the fundus of the bladder.


The Utero-sacral Ligaments.—The *utero-sacral ligaments* consist of folds of peritoneum of a crescentic form, with their concavities looking inward: they start from the lower part of the posterior surface of the uterus, and curve backward to be attached to the third and fourth sacral vertebræ. Within their folds exist bundles of muscular fibres, continuous with those of the uterus, as well as connective tissue, vessels, and nerves. The experiments of Savage, as well as of other anatomists, show that these ligaments have an important influence in preventing downward displacement of the womb.

Alterations during Pregnancy.—During pregnancy all these ligaments become greatly stretched and unfolded, rising out of the pelvic cavity and accommodating themselves to the increased size of the gravid uterus; and they again contract to their natural size, possibly through the agency of the muscular fibres contained within them, after delivery has taken place.

The Fallopian Tubes.—The *Fallopian tubes*, the homologues of the vasa deferentia in the male, are structures of great physiological interest. They serve the double purpose of conveying the semen to the ovary and of carrying the ovule to the uterus. From the latter function they may be looked on as the excretory ducts of the ovaries; but, unlike other excretory ducts, they are movable, so that they may apply themselves to the part of the ovaries from which the ovule is to come; and so great is their mobility that there is reason to believe that a Fallopian tube may even grasp the ovary of the opposite side. Each tube proceeds from the upper angle of the uterus at first transversely outward, and then downward, backward, and inward, so as to reach the neighborhood of the ovary. In the first part of its course it is straight; afterward it becomes flexuous and twisted on itself. It is contained in the upper part of the broad ligament, where it may be felt as a hard cord. It commences at the uterus by a narrow opening, admitting only the passage of a bristle, known as the *ostium uterinum*. As it passes through the muscular walls of the uterus the tube takes a somewhat curved course, and opens into the uterine cavity by a dilated aperture. From its uterine attachment the tube expands gradually until it terminates in its trumpet-shaped extremity; just before its distal end, however, it again contracts slightly. The ovarian end of the tube is surrounded by a number of remarkable fringe-like processes. These consist of longitudinal membranous fimbriae, surrounding the aperture of the tube, like the tentacles of a polyp, varying considerably in number and size, and having their edges cut and subdivided. On their inner surface are found both transverse and longitudinal folds of mucous membrane, continuous with those lining the tube itself (Fig. 33). One of these fimbriae is always larger and more developed than the rest, and is indirectly united to the surface of the ovary by a fold of peritoneum proceeding from its external surface. Its under surface is grooved so as to form a channel,

open below. The function of this fringe-like structure is to grasp the ovary during the menstrual nixus; and the fimbria which is attached to the ovary would seem to guide the tentacles to the ovary which they are intended to seize. One or more supplementary series of fimbriae sometimes exist, which have an aperture of communication with the canal of the Fallopian tube beyond its ovarian extremity. His has recently shown that the fimbriated extremity of the tube, after running over the

FIG. 33.



Fallopian Tube laid open. (After Richard.)

a, b. Uterine portion of tube. c, d. Plicae of mucous membrane. e. Tubo-ovarian ligaments and fringes. f. Ovary. g. Round ligaments.

upper part of the ovary, turns down along its free border; so that its aperture lies below it, ready to receive the ovule when expelled from the Graafian follicle.¹

Their Structure.—The tubes themselves consist of peritoneal, muscular, and mucous coats. The peritoneum surrounds the tube for three-fourths of its calibre, and comes into contact with the mucous lining at its fimbriated extremity, the only instance in the body where such a junction occurs. The muscular coat is principally composed of circular fibres, with a few longitudinal fibres interspersed. Its muscular character has been doubted by Robin and Richard, but Farre had no difficulty in demonstrating the existence of muscular fibres, both in the human female and many of the lower animals. According to Robin, the muscular tissue of the Fallopian tubes is entirely distinct from that of the uterus, from which he describes it as being separated by a distinct cellular septum. The mucous lining is thrown into a number of remarkable longitudinal folds, each of which contains a dense and vascular fibrous septum, with small muscular fibres, and is covered with columnar and ciliated epithelium. The apposition of these produces a series of minute capillary tubes, along which the ovules are propelled, the action of the cilia, which is toward the uterus, apparently favoring their progress.

The Ovaries.—The ovaries are the bodies in which the ovules are

¹ His, *Archiv für Anat. und Phys.*, 1881.

formed, and from which they are expelled, and the changes going on in them in connection with the process of ovulation, during the whole period between the establishment of puberty and the cessation of menstruation, have an enormous influence in the female economy. Normally, the ovaries are two in number; in some exceptional cases a supplementary ovary has been discovered; or they may be entirely absent. They are placed in the posterior folds of the broad ligament, usually below the brim of the pelvis, behind the Fallopian tubes, the left in front of the rectum, the right in front of some coils of the small intestine. Their situation varies, however, very much under different circumstances, so that they can scarcely be said to have a fixed and normal position; most probably, however, as has been recently shown by His,¹ they are normally placed close below the brim of the pelvis, with their long diameters almost vertical, and immediately above the aperture of the distal extremity of the Fallopian tubes. In pregnancy they rise into the abdominal cavity with the enlarging uterus; and in certain conditions they are dislocated downward into Douglas's space, where they may be felt through the vagina as rounded and very tender bodies.

Their Connections.—The folds of the broad ligament, between which the ovaries are placed, form for them a kind of loose mesentery. Each of them is united to the upper angle of the uterus by a special ligament called the utero-ovarian. This is a rounded band of organic muscular fibres, about an inch in length, continuous with the superficial muscular fibres of the posterior wall of the uterus, and attached to the inner

FIG. 34.



A A. Ovary enlarged under menstrual nixus. B. Ripe follicle projecting on its surface. a, a, a. Traces of previously-ruptured follicles.

extremity of the ovary. It is surrounded by peritoneum, and through it the muscular fibres, which form an important integral part in the structure of the ovaries, are conveyed to them. The ovary is also attached to the fimbriated extremity of the Fallopian tube in the manner already described.

The ovary is of an irregular oval shape (Fig. 34), the upper border being convex, the lower—through which the vessels and nerves enter—

¹ *Op. cit.*

being straight. The anterior surface, like that of the uterus, is less convex than the posterior. The outer extremity is more rounded and bulbous than the inner, which is somewhat pointed and eventually lost in its proper ligament. By these peculiarities it is possible to distinguish the left from the right ovary after they have been removed from the body. The ovary varies much in size under different circumstances. On an average, in adult life, it measures from one to two inches in length, three-quarters of an inch in width, and about half an inch in thickness. It increases greatly in size during each menstrual period—a fact which has been demonstrated in certain cases of ovarian hernia, in which the protruded ovary has been seen to swell as menstruation commenced; also during pregnancy, when it is said to be double its usual size. After the change of life it atrophies, and becomes rough and wrinkled on its surface. Before puberty, the surface of the ovary is smooth and polished and of a whitish color. After menstruation commences, its surface becomes scarred by the rupture of the Graafian follicles (Fig. 34, *a a*), each of which leaves a little linear or striated cicatrix of a brownish color; and the older the patient the greater is the number of these cicatrices.

Their Structure—The structure of the ovary has been made the subject of many important observations. It has an external covering of epithelium, originally continuous with the peritoneum, called by some the germ-epithelium, in consequence of the ovules being formed from it in early foetal life. In the adult it is separated from the peritoneum at the base of the organ by a circular white line, and it consists of columnar epithelium, differing only from the epithelium lining the Fallopian tubes, with which it is sometimes continuous through the attached fimbria uniting the tube and the ovary, in being destitute of cilia. Immediately beneath this covering is the dense coat known as the *tunica albuginea*, on account of its whitish color. It consists of short connective-tissue fibres, arranged in laminae, among which are interspersed fusiform muscular fibres. At the point where the vessels and nerves enter the ovary this membrane is raised into a ridge, which is continuous with the utero-ovarian ligament, and is called the *hilum*.

The *tunica albuginea* is so intimately blended with the stroma of the ovary as to be inseparable on dissection; it does not, however, exist as a distinct lamina, but is merely the external part of the proper structure of the ovary, in which more dense connective tissue is developed than elsewhere.

The Stroma.—On making a longitudinal section of the ovary (Fig. 35), it will be seen to be composed of two parts, the more internal of which is of a reddish color from the number of vessels that ramify in it, and is called the *medullary* or vascular zone; while the external, of a whitish tint, receives the name of




Longitudinal Section of Adult Ovary. (After Parre.)

the *cortical* or parenchymatous substance. The former consists of loose connective tissue interspersed with elastic, and a considerable number of

muscular fibres. According to Rouget¹ and His,² the muscular structure forms the greater part of the ovarian stroma. The latter describes it as consisting essentially of interwoven muscular fibres, which he terms the "fusiform tissue," and which he believes to be continuous with the muscular layers of the ovarian vessels. The former believes that the muscular fasciculi accompany the vessels in the form of sheaths, as in erectile tissues. Both attribute to the muscular tissues an important influence in the expulsion of the ovules and in the rupture of the Graafian fol-

FIG. 36.



Section through the Cortical Part of the Ovary.

e. Surface epithelium. *s s.* Ovarian stroma. *1 1.* Large-sized Graafian follicles. *2 2.* Middle-sized; and *3 3.* Small-sized Graafian follicles. *o.* Ovule within Graafian follicle. *v c.* Blood-vessels in the stroma. *g.* Cells of the membrana granulosa. (After Turner.)

licles. Waldeyer and other writers, however, do not consider it to be so extensively developed as Rouget and His believe. The cortical substance is the more important, as that in which the Graafian follicles and ovules are formed. It consists of interlaced fibres of connective tissue, containing a large number of nuclei. The muscular fibres of the medullary substance do not seem to penetrate into it in the human female. In it are found the Graafian follicles, which exist in enormous numbers from the earliest periods of life, and in all stages of development (Fig. 36.)


The Graafian Follicles.—According to the researches of Pflüger, Waldeyer, and other German writers, the Graafian follicles are formed in early fetal life by cylindrical inflexions of the epithelial covering of the ovary, which dip into the substance of the gland. These tubular filaments anastomose with each other, and in them are formed the ovules, which are originally the epithelial cells lining the tubes. Portions become shut off from the rest of the filaments and form the Graafian follicles. The ovules, on this view, are highly-developed epithelial cells, originally derived from the surface of the ovary, and

¹ *Journal de Physiol.*, i. p. 737.

² Schultze's *Arch. f. Mikroskop. Anat.*, 1865.

not developed in its stroma. These tubular filaments disappear shortly after birth, but they have recently been detected by Slavyansky¹ in the ovaries of a woman thirty years of age. These observations have been modified by Dr. Foulis.² He recognizes the origin of the ovules from the germ-epithelium covering the surface of the ovary, which is itself derived from the Wolffian body. He believes all the ovules to be formed from the germ-epithelium corpuscles, which become imbedded in the stroma of the ovary, by the outgrowth of processes of vascular connective tissue, fresh germ-epithelial corpuscles being constantly produced on the surface of the organ up to the age of two and a half years, to take the place of those already imbedded in its stroma. He believes the Graafian follicles to be formed by the growth of delicate processes of connective tissue between and around the ovules, but not from tubular inflections of the epithelium covering the gland, as described by Waldeyer (Fig. 37). This view is supported by the researches of Balfour,³ who arrives at the conclusion that the whole egg-containing

FIG. 37.



Vertical Section through the Ovary of the Human Fœtus.

g g. Germ-epithelium, with *o o.* Developing ovules in it. *ss.* Ovarian stroma, containing *c c c.* Fusiform connective-tissue corpuscles. *v v.* Capillary blood-vessels. In the centre of the figure an involution of the germ-epithelium is shown; and at the left lower side a primordial ovule, with the connective-tissue corpuscles ranging themselves round it. (After Foulis.)

part of the ovary is really the thickened germinal epithelium, broken up into a kind of meshwork by growths of vascular stroma. According to this theory, Pflüger's tubular filaments are merely trabeculae of germinal epithelium, modified cells of which become developed into ovules.

The greater portion of the Graafian follicles are only visible with the high powers of the microscope, but those which are approaching maturity are distinctly to be seen by the naked eye. The quantity of these follicles is immense. Foulis estimates that at birth each human ovary contains not less than 30,000. No fresh follicles appear to be formed after birth, and as development goes on some only grow, and, by pres-

¹ *Annales de Gynéc.*, Feb., 1871.


² *Proceedings of the Royal Soc. of Edinb.*, April, 1875, and *Journ. of Anat. and Phys.*, vol. xiii., 1879.

³ F. M. Balfour, "Structure and Development of Vertebrate Ovary," *Journal of Microscopical Science*, vol. xviii., 1878.

sure on the others, destroy them. Of those that grow, of course only a few ever reach maturity; they are scattered through the substance of the ovary, some developing in the stroma, others on the surface of the organ, where they eventually burst, and are discharged into the Fallopian tube.

Structure of the Graafian Follicle.—A ripe Graafian follicle has an external investing membrane (Fig. 38), which is generally described as

FIG. 38.



Diagrammatic Section of Graafian Follicle.

1. Ovum. 2. *Membrana granulosa*. 3. External membrane of Graafian follicle. 4. Its vessels. 5. Ovarian stroma. 6. Cavity of Graafian follicle. 7. External covering of ovary.

consisting of two distinct layers—the external, or *tunica fibrosa*, highly vascular, and formed of connective tissue; the internal, or *tunica propria*, composed of young connective tissue, containing a large number of fusiform or stellate cells and numerous oil-globules. These layers, however, appear to be essentially formed of condensed ovarian stroma. Within this capsule is the epithelial lining called the *membrana granulosa*, consisting of stratified columnar epithelial cells, which, according to Foulis, are originally formed from the nuclei of the fibro-nuclear tissue of the stroma of the ovary. At one part of the circumference of the ovisac is situated the ovule, around which the epithelial cells are congregated in greater quantity, constituting the projection known as the *discus proligerus*. The remainder of the cavity of the follicle is filled with a small quantity of transparent fluid, the *liquor folliculi*, traversed by three or four minute bands, the *retinacula of Barry*, which are attached to the opposite walls of the follicular cavity, and apparently serve the purpose of suspending the ovule and maintaining it in a proper position. In many young follicles this cavity does not at first exist, the follicle being entirely filled by the ovule. According to Waldeyer, the liquor folliculi is formed by the disintegration of the epithelial cells, the fluid thus produced collecting, and distending the interior of the follicle.

The Ovule.—The *ovule* is attached to some part of the internal surface of the Graafian follicle. It is a rounded vesicle about $\frac{1}{120}$ th of an inch in diameter, and is surrounded by a layer of columnar cells, distinct from those of the *discus proligerus*, in which it lies. It is invested by a transparent elastic membrane, the *zona pellucida*, or vitelline mem-

brane. In most of the lower animals the zona pellucida is perforated by numerous very minute pores, only visible under the highest powers of the microscope; in others there is a distinct aperture of a larger size, the micropyle, allowing the passage of the spermatozoa into the interior of the ovule. It is possible that similar apertures may exist in the human ovule, but they have not been demonstrated. Within the zona pellucida some embryologists describe a second fine membrane, the existence of which has been denied by Bischoff. The cavity of the ovule is filled with a viscid yellow fluid, the *yolk*, containing numerous granules. It entirely fills the cavity, to the walls of which it is non-adherent. In the centre of the yolk in young, and at some portion of its periphery in mature ovules, is situated the *germinal vesicle*, which is a clear circular vesicle, refracting light strongly, and about $\frac{1}{80}$ th of a line in diameter. It contains a few granules, and a nucleolus, or *germinal spot*, which is sometimes double.

From within outward, therefore, we find—

1. The *germinal spot*; round this
2. The *germinal vesicle*, contained in
3. The *yolk*, which is surrounded by the
4. *Zona pellucida*, with its layers of columnar epithelial cells.

These constitute the ovule.


The ovule is contained in—

The *Graafian follicle*, and lies in that part of its epithelial lining called the—

Discus proligerus, the rest of the follicle being occupied by the *liquor folliculi*. Round these we have the epithelial lining or *membrana granulosa*, and the external coat, consisting of the *tunica propria* and the *tunica fibrosa*.

The Vessels and Nerves of the Ovary.—The vascular supply of the ovary is complex. The arteries enter at the hilum, penetrating the stroma in a spiral curve, and are ultimately distributed in a rich capil-

FIG. 39.



Bulb of Ovary.


v. Uterus. o. Ovary and utero-ovarian ligament. r. Fallopian tube. 1. Utero-ovarian vein. 2. Pampiniform ovarian plexus. 3. Commencement of spermatic vein.

lary plexus to the follicles. The large veins unite freely with each other, and form a vascular and erectile plexus, continuous with that surrounding the uterus, called the bulb of the ovary (Fig. 39). Lymphatics and nerves exist, but their mode of termination is unknown.

The Mammary Glands.—To complete the consideration of the generative organs of the female we must study the *mammary glands*, which secrete the fluid destined to nourish the child. In the human subject they are two in number, and instead of being placed upon the abdomen, as in most animals, they are situated on either side of the sternum, over the *pectorales majora* muscles, and extend from the third to the sixth rib. This position of the glands is obviously intended to suit the erect position of the female in suckling. They are convex anteriorly, and flattened posteriorly where they rest on the muscles. They vary greatly in size in different subjects, chiefly in proportion to the amount of adipose tissue they contain. In man, and in girls previous to puberty, they are rudimentary in structure, while in pregnant women they increase greatly in size, the true glandular structures becoming much hypertrophied. Anomalies in shape and position are sometimes observed. Supplementary *mammæ*, one or more in number, situated on the upper portion of the *mammæ*, are sometimes met with, identical in structure with the normally situated glands; or, more commonly, an extra nipple is observed by the side of the normal one. In some races, especially the African, the *mammæ* are so enormously developed that the mother is able to suckle her child over her shoulder.

Their Structure.—The skin covering the gland is soft and supple, and during pregnancy often becomes covered with fine white lines, while

FIG. 40.



1. Galactophorous ducts. 2. Lobuli of the mammary gland.

large blue veins may be observed coursing over. Underneath it is a quantity of connective tissue, containing a considerable amount of fat, which extends beneath the true glandular structure. This is composed of from fifteen to twenty lobes, each of which is formed of a number of lobules. The lobules are produced by the aggregation of the terminal acini in which the milk is formed. The acini are minute culs-de-sac opening into little ducts, which unite with each other until they form a large duct for each lobule; the ducts of each lobule unite with each other until they end in a still larger duct common to each of the fifteen or twenty lobes into which the gland is divided, and eventually open on the surface of the nipple. These terminal canals are known as the *galactophorous ducts* (Fig. 40). They become widely dilated as they approach the nipple, so as to form reservoirs in which milk is stored until it is required, but when they actually enter the nipple they again

contract. Sometimes they give off lateral branches, but, according to Sappey, they do not anastomose with each other, as some anatomists have described. These excretory ducts are composed of connective tissue, with numerous elastic fibres, on their external surface. Sappey and Robin describe a layer of muscular fibres, chiefly developed near their terminal extremities. They are lined with columnar epithelium, continuous with that in the acini; and it is by the distension of its cells with fatty matter, and their subsequent bursting, that the milk is formed.

The Nipple.—The *nipple* is the conical projection at the summit of the mamma, and it varies in size in different women. Not very unfrequently, from the continuous pressure to which it has been subjected by the dress, it is so depressed below the surface of the skin as to prevent lactation. It is generally larger in married than in single women, and increases in size during pregnancy. Its surface is covered with numerous papillæ, giving it a rugous aspect, and at their bases the orifices of the lactiferous ducts open. Here are also the openings of numerous sebaceous follicles, which secrete an unctuous material supposed to protect and soften the integument during lactation. Beneath the skin are muscular fibres, mixed with connective and elastic tissues, vessels, nerves, and lymphatics. When the nipple is irritated it contracts and hardens, and by some this is attributed to its erectile properties. The vascularity, however, is not great, and it contains no true erectile tissue; the hardening is, therefore, due to muscular contraction. Surrounding the nipple is the *areola*, of a pink color in virgins, becoming dark from the development of pigment-cells during pregnancy, and always remaining somewhat dark after childbearing. On its surface are a number of prominent tubercles, sixteen to twenty in number, which also become largely developed during gestation. They are supposed by some to secrete milk, and to open into the lactiferous tubes; most probably they are composed of sebaceous glands only. Beneath the areola is a circular band of muscular fibres, the object of which is to compress the lactiferous tubes which run through it, and thus to favor the expulsion of their contents. The mammae receive their blood from the internal mammary and intercostal arteries, and they are richly supplied with lymphatic vessels, which open into the axillary glands. The nerves are derived from the intercostal and thoracic branches of the brachial plexus.

The secretion of milk in women who are nursing is accompanied by a peculiar sensation, as if milk were rushing into the breast, called the "draught," which is excited by the efforts of the child to suck and by various other causes. The sympathetic relations between the mammae and the uterus are very well marked, as is shown in the unimpregnated state by the fact of the frequent occurrence of sympathetic pains in the breast in connection with various uterine diseases, and after delivery by the well-known fact that suction produces reflex contraction of the uterus, and even severe after-pains.

Plate III.



Fig. 1.
A recently ruptured and bloody Graafian
follicle, just developing into a Corpus luteum



Fig. 2.
Corpus luteum ten days after menstruation.



Fig. 3.
Degenerated Graafian follicles
which has never ruptured.



Fig. 4.
Corpus luteum of pregnancy

ILLUSTRATIONS OF THE CORPUS LUTEUM. (AFTER DALTON)

2. Maria 1843 B&L

CHAPTER III.

OVULATION AND MENSTRUATION.

Functions of the Ovary.—The main function of the ovary is to supply the female generative element, and to expel it, when ready for impregnation, into the Fallopian tube, along which it passes into the uterus. This process takes place spontaneously in all viviparous animals, and without the assistance of the male. In the lower animals this periodical discharge receives the name of the œstrum or rut, at which time only the female is capable of impregnation and admits the approach of the male. In the human female the periodical discharge of the ovule, in all probability, takes place in connection with menstruation, which may therefore be considered to be the analogue of the rut in animals. Between each menstrual period Graafian follicles undergo changes which prepare them for rupture and the discharge of their contained ovules. After rupture certain changes occur which have for their object the healing of the rent in the ovarian tissue through which the ovule has escaped, and the filling up of the cavity in which it was contained. This results in the formation of a peculiar body in the substance of the ovary, called the *corpus luteum*, which is essentially modified should pregnancy occur, and is of great interest and importance. During the whole of the child-bearing epoch the periodical maturation and rupture of the Graafian follicles are going on. If impregnation does not take place, the ovules are discharged and lost; if it does, ovulation is stopped, as a general rule, during gestation and lactation.

Theory of Menstruation.—This, broadly speaking, is an outline of the modern theory of menstruation, which was first broached in the year 1821 by Dr. Power, and subsequently elaborated by Negrier, Bischoff, Raciborski, and many other writers. Although the sequence of events here indicated may be taken to be the rule, it must be remembered that it is one subject to many exceptions, for undoubtedly ovulation may occur without its outward manifestation, menstruation, as in cases in which impregnation takes place during lactation or before menstruation has been established, of which many examples are recorded. These exceptions have led some modern writers to deny the ovular theory of menstruation, and their views will require subsequent consideration.

In order to understand the subject properly it will be necessary to study the sequence of events in detail.

Changes in the Graafian Follicle.—The changes in the Graafian follicle which are associated with the discharge of the ovules comprise—1. *Maturation.* As the period of puberty approaches, a certain number of the Graafian follicles, fifteen to twenty in number, increase in size, and come near the surface of the ovary. Amongst these one becomes especially developed, preparatory to rupture, and upon it, for the time being,

all the vital energy of the ovary seems to be concentrated. A similar change in one, sometimes in more than one, follicle takes place periodically during the whole of the childbearing epoch, in connection with each menstrual period, and an examination of the ovary will show several follicles in different stages of development. The maturing follicle becomes gradually larger, until it forms a projection on the surface of the ovary, from five to seven lines in breadth, but sometimes even as large as a nut (Fig. 34). This growth is due to the distension of the follicle by the increase of its contained fluid, which causes it so to press upon the ovarian structures covering it that they become thinned, separated from each other, and partially absorbed, until they eventually readily lacerate. The follicle also becomes greatly congested, the capillaries coursing over it become increased in size and loaded with blood, and, being seen through the attenuated ovarian tissue, give it, when mature, a bright-red color. At this time some of these distended capillaries in its inner coat lacerate, and a certain quantity of blood escapes into its cavity. This escape of blood takes place before rupture, and seems to have for its principal object the increase of the tension of the follicle, of which it has been termed the menstruation. Pouchet was of opinion that the blood collects behind the ovule, and carries it up to the surface of the follicle. 2. *Escape of the Ovule.* By these means the follicle is more and more distended, until at last it ruptures (Plate III. Fig. 1), either spontaneously or, it may be, under the stimulus of sexual excitement. Whether the laceration takes place during, before, or after the menstrual discharge is not yet positively known; from the results of post-mortem examination in a number of women who died shortly before or after the period, Williams believes that the ovules are expelled before the monthly flow commences.¹ In order that the ovule may escape, the laceration must, of course, involve not only the coats of the Graafian follicles, but also the superincumbent structures.

Laceration seems to be aided by the growth of the internal layer of the follicle, which increases in thickness before rupture, and assumes a characteristic yellow color from the number of oil-globules it then contains. It is also greatly facilitated, if it be not actually produced, by the turgescence of the ovary at each menstrual period, and by the contraction of the muscular fibres in the ovarian stroma. As soon as the rent in the follicular walls is produced, the ovule is discharged, surrounded by some of the cells of the membrana granulosa, and is received into the fimbriated extremity of the Fallopian tube, which grasps the ovary over the site of the rupture. By the vibratile cilia of its epithelial lining it is then conducted into the canal of the tube, along which it is propelled, partly by ciliary action and partly by muscular contraction in the walls of the tube.


Obliteration of the Graafian Follicle.—After the ovule has escaped, certain characteristic changes occur in the empty Graafian follicle, which have for their object its cicatrization and obliteration. There are great differences in the changes which occur when impregnation has followed the escape of the ovule, and they are then so remarkable that they have been considered certain signs of pregnancy. They are, however, differ-

¹ *Proceedings of the Royal Society, 1875.*

ences of degree rather than of kind. It will be well, however, to discuss them separately.

Changes undergone by the Follicle when Impregnation does not Occur.—As soon as the ovule is discharged, the edges of the rent through which it has escaped become agglutinated by exudation, and the follicle shrinks, as is generally believed, by the inherent elasticity of its internal coat, but according to Robin, who denies the existence of this coat, from compression by the muscular fibres of the ovarian stroma. In proportion to the contraction that takes place, the inner layer of the follicle, the cells of which have become greatly hypertrophied and loaded with fat-granules previous to rupture, is thrown into numerous folds (Plate III. Fig. 2). The greater the amount of contraction the deeper these folds become, giving to a section of the follicle an appearance similar to that of the convolutions of the brain (Fig. 41). These folds in the human subject are generally of a bright-yellow color, but in some of the mammalia they are of a deep red. The tint was formerly ascribed by Raciborski to absorption of the coloring matter of the blood-clot contained in the follicular cavity—a theory he has more recently abandoned in favor of the view maintained by Coste that it is due to the inherent color of the cells of the lining membrane of the follicle, which, though not well marked in a single cell, becomes very apparent *en masse*. The existence of a contained blood-clot is also denied by the latter physiologist, except as an unusual pathological condition; and he describes the cavity as containing a gelatinous and plastic fluid, which becomes absorbed as contraction advances. The more recent researches of Dalton,¹ however, show the existence of a central blood-clot in the cavity of the follicle, and he considers its occasional absence to be connected with disturbance or cessation of the menstrual function. The folds into which the membrane has been thrown continue to increase in size, from the proliferation of their cells, until they unite and become adherent, and eventually fill the follicular cavity. By the time that another Graafian follicle is matured and ready for rupture the diminution has advanced considerably, and the empty ovisac is reduced to a very small size. The cavity is now nearly obliterated, the yellow color of the convolutions is altered into a whitish tint, and on section the corpus luteum has the appearance of a compact white stellate cicatrix, which generally disappears in less than forty days from the period of rupture. The tissue of the ovary at the site of laceration also shrinks, and this, aided by the contraction of the follicle, gives rise to one of those permanent pits or depressions which mark the surface of the adult ovary. Slavyansky² has shown that only a few of the immense number of Graafian follicles

FIG. 41.



Section of Ovary, showing Corpus Luteum three weeks after Menstruation. (After Dalton.)

¹ "Report on the Corpus Luteum," *American Gynec. Trans.*, vol. ii., 1878.


² *Archiv. de Phys.*, March, 1874.

undergo these alterations. The greater proportion of them seem never to discharge their ovules, but, after increasing in size, undergo retrogressive changes exactly similar in their nature, but to a much less extent, to those which result in the formation of a corpus luteum. The sites of these may afterward be seen as minute striæ in the substance of the ovary.

Changes undergone by the Follicle when Impregnation has taken place.—Should pregnancy occur, all the changes above described take place; but, inasmuch as the ovary partakes of the stimulus to which all the generative organs are then subjected, they are much more marked and apparent (Plate III. Fig. 4). Instead of contracting and disappearing in a few weeks, the corpus luteum continues to grow until the third or fourth month of pregnancy; the folds of the inner layer of the ovisac become large and fleshy and permeated by numerous capillaries, and ultimately become so firmly united that the margins of the convolutions thin and disappear, leaving only a firm fleshy yellow mass, averaging from 1 to $1\frac{1}{2}$ inches in thickness, which surrounds a central cavity, often containing a whitish fibrillated structure, believed to be the remains of a central blood-clot. This was erroneously supposed by Montgomery to be the inner layer of the follicle itself, and he conceived the yellow substance to be a new formation between it and the external layer, while Robert Lee thought it was placed external to both the external and internal layers.

Between the third and fourth months of pregnancy, when the corpus luteum has attained its maximum of development (Fig. 42), it forms a firm projection on the surface of the ovary, averaging about 1 inch in

FIG. 42.



Corpus Luteum of the Fourth Month of Pregnancy. (After Dalton.)

FIG. 43.



Corpus Luteum of Pregnancy at Term. (After Dalton.)

length and rather more than $\frac{1}{2}$ an inch in breadth. After this it commences to atrophy (Fig. 43), the fat-cells become absorbed, and the capillaries disappear. Cicatrization is not complete until from one to two months after delivery.

Its Value as a Sign of Pregnancy.—On account of the marked appearance of the corpus luteum it was formerly considered to be an infallible

sign of pregnancy; and it was distinguished from the corpus luteum of the non-pregnant state by being called a "true" as opposed to a "false" corpus luteum. From what has been said it will be obvious that this designation is essentially wrong, as the difference is one of degree only. Dalton¹ applies the term "false corpus luteum" to a degenerated condition sometimes met with in an unruptured Graafian follicle consisting in reabsorption of its contents and thickening of its walls (Plate III. Fig. 3). It differs from the "true" corpus luteum in being deeply seated in the substance of the ovary, in having no central clot, and in being unconnected with a cicatrix on the surface of the ovary. Nor do obstetricians attach by any means the same importance as they did formerly to the presence of the corpus luteum as indicating impregnation; for even when well marked, other and more reliable signs of recent delivery, such as enlargement of the uterus, are sure to be present, especially at the time when the corpus luteum has reached its maximum of development; while after delivery at term it has no longer a sufficiently characteristic appearance to be depended on.

Menstruation.—By the term *menstruation* (catamenia, periods, etc.) is meant the periodical discharge of blood from the uterus which occurs, in the healthy woman, every lunar month, except during pregnancy and lactation, when it is, as a rule, suspended.

Period of Establishment.—The first appearance of menstruation coincides with the establishment of puberty, and the physical changes that accompany it indicate that the female is capable of conception and child-bearing, although exceptional cases are recorded in which pregnancy occurred before menstruation had begun. In temperate climates it generally commences between the 14th and 16th years, the largest number of cases being met with in the 15th year. This rule is subject to many exceptions, it being by no means very rare for menstruation to become established as early as the 10th or 11th year, or to be delayed until the 18th or 20th. Beyond these physiological limits a few cases are from time to time met with in which it has begun in early infancy, or not until a comparatively late period of life.

Influence of Climate, Race, etc.—Various accidental circumstances have much to do with its establishment. As a rule, it occurs somewhat earlier in tropical, and later in very cold than in temperate climates. The influence of climate has been unduly exaggerated. It used to be generally stated that in the Arctic regions women did not menstruate until they were of mature age, and that in the tropics, girls of 10 or 12 years of age did so habitually. The researches of Robertson of Manchester² first showed that the generally received opinions were erroneous; and the collection of a large number of statistics has corroborated his opinion. There can be no doubt, however, that a larger proportion of girls menstruate early in warm climates. Joulin found that in tropical climates, out of 1635 cases, the largest proportion began to menstruate between the 12th and 13th years; so that there is an average difference of more than two years between the period of its establishment in the tropics and in temperate countries. Harris³ states that among the

¹ *Op. cit.*, p. 64.

² *Edin. Med. and Surg. Journ.*, 1832.

³ *Amer. Journ. of Obstet.*, 1871, R. P. Harris on early puberty.

Hindoos 1 to 2 per cent. menstruate as early as nine years of age; 3 to 4 per cent. at ten; 8 per cent. at eleven; and 25 per cent. at twelve; while in London or Paris probably not more than one girl in 1000 or 1200 does so at nine years. The converse holds true with regard to cold climates, although we are not in possession of a sufficient number of accurate statistics to draw very reliable conclusions on this point; but out of 4715 cases, including returns from Denmark, Norway and Sweden, Russia, and Labrador, it was found that menstruation was established on an average a year later than in more temperate countries. It is probable that the mere influence of temperature has much to do in producing these differences, but there are other factors, the action of which must not be overlooked. Raciborski attributes considerable importance to the effect of race; and he has quoted Dr. Webb of Calcutta to the effect that English girls in India, although subjected to the same climatic influence as the Indian races, do not, as a rule, menstruate earlier than in England; while in Austria girls of the Magyar race menstruate considerably later than those of German parentage.¹ The surroundings of girls, and their manner of education and living, have probably also a marked influence in promoting or retarding its establishment. Thus, it will commence earlier in the children of the rich, who are likely to have a highly-developed nervous organization, and are habituated to luxurious living and a premature stimulation of the mental faculties by novel-reading, society, and the like; while amongst the hard-worked poor, or in girls brought up in the country, it is more likely to begin later. Premature sexual excitement is said also to favor its early appearance, and the influence of this among the factory-girls of Manchester, who are exposed in the course of their work to the temptations arising from the promiscuous mixing of the sexes, has been pointed out by Dr. Clay.²

[*Precocious Physical Womanhood.*—Within a few years the photographic process has made us familiar with the appearances of several little girls of four, five, and six years of age who were menstruating regularly, had large mammæ, and pubes covered with hair, all natives of our Northern States. In growth and obesity such children are far beyond their years as a general rule, but in mental development and character very child-like. Lately I saw one of six, who, although large, broad, and full-chested, and with arms and thighs like a woman, was playing with a doll and acting like a little girl of her own age; another, of five and a half, was a beautiful miniature of a developing girl of twelve or fifteen; and a third, of four, had breasts as large as an orange. Fortunately, the sexual passion, so general with the precociously-developed male infant, is seldom a marked characteristic in the female, although cases of precocity are much more frequently met with in girls. The nubile period has seldom been tested in these subjects, but occasionally in the lower classes of society, pregnancy has occurred at a very early age. Sue, in his *Essais historiques*,^[3] Paris, 1779, reports a case where the young mother was eight years and ten months old, the fetus being mature, but dead. Menstruation began at two years. Her mammæ

¹ *Op. cit.*, p. 227.
² Vol. ii. p. 344.]

³ *Brit. Record of Obstet. Med.*, vol. i.

and pubes resembled those of a girl of seventeen. The youngest mother in the United States was ten years and thirteen days old, four feet seven inches in height, and 100 pounds in weight. [¹] She commenced to menstruate at one year; her child weighed $7\frac{3}{4}$ pounds. The youngest English mother was born on August 8, 1871; commenced to menstruate when a year old, and bore a child of 7 pounds in weight, after a labor of only 6 hours, in March, 1881. [²] She ceased to menstruate on June 22, 1880, and must have become pregnant when eight years ten and a half months old. She was nine years seven and a half months old when her case was reported, March 24, 1881.

A Southern negress just thirteen years old gave birth to a female child at maturity, who in turn became also pregnant at twelve. This child menstruated at ten years and nine months, and a second child at seven years and nine months. If the first daughter became a mother in due time, then the first-mentioned negress must have become a grandmother before she was twenty-six. [³]

A Spanish girl of Maracaibo is reported by the late Prof. C. D. Meigs to have given birth to a child at twelve, and twins at a second birth before she was fourteen. A quadroon of Nassau, New York, in 1822, performed the Cæsarean operation on herself when in labor with twins at the age of fourteen.—Ed.]

Changes occurring at Puberty.—The first appearance of menstruation is accompanied by certain well-marked changes in the female system, on the occurrence of which we say that the girl has arrived at the period of puberty. The pubes becomes covered with hair, the breasts enlarge, the pelvis assumes its fully-developed form, and the general contour of the body fills out. The mental qualities also alter: the girl becomes more shy and retiring, and her whole bearing indicates the change that has taken place. The menstrual discharge is not established regularly at once. For one or two months there may be only premonitory symptoms—a vague sense of discomfort, pains in the breasts, and a feeling of weight and heat in the back and loins. There then may be a discharge of mucus tinged with blood, or of pure blood, and this may not again show itself for several months. Such irregularities are of little consequence on the first establishment of the function, and need give rise to no apprehension.

Period of Duration and Recurrence.—As a rule, the discharge recurs every twenty-eight days, and with some women with such regularity that they can foretell its appearance almost to the hour. The rule is, however, subject to very great variations. It is by no means uncommon, and strictly within the limits of health, for it to appear every twentieth day, or even with less interval; while in other cases as much as six weeks may habitually intervene between two periods. The period of recurrence may also vary in the same subject. I am acquainted with patients who sometimes only have twenty-eight days, at others as many as forty-eight days, between their periods, without their health in any way suffering. Joulin mentions the case of a lady who only menSTRU-

[¹ *Transylvania Med. Journ.*, vol. vii. p. 447.]

[² *Lancet*, April 9, 1881, p. 601.]

[³ *Am. Journ. Obstetrics*, vol. vi. p. 572, 1873-74.]

ated two or three times in the year, and whose sister had the same peculiarity.

The duration of the period varies in different women, and in the same woman at different times. In this country its average is four or five days, while in France, Dubois and Brierre de Boismont fix eight days as the most usual length. Some women are only unwell for a few hours, while in others the period may last many days beyond the average without being considered abnormal.

Quantity of Blood lost.—The quantity of blood lost varies in different women. Hippocrates puts it at xxviiij , which, however, is much too high an estimate. Arthur Farre thinks that from xiij to xxiiij is the full amount of a healthy period, and that the quantity cannot habitually exceed this without producing serious constitutional effects. Rich diet, luxurious living, and anything that unhealthily stimulates the body and mind will have an injurious effect in increasing the flow, which is, therefore, less in hard-worked country-women than in the better classes and residents in towns.

It is more abundant in warm climates, and our countrywomen in India habitually menstruate over-profusely, becoming less abundantly unwell when they return to England. The same observation has been made with regard to American women residing in the Gulf States, who improve materially by removing to the Lake States. Some women appear to menstruate more in summer than in winter. I am acquainted with a lady who spends the winter in St. Petersburg, where her periods last eight or ten days, and the summer in England, where they never exceed four or five. The difference is probably due to the effect of the overheated rooms in which she lives in Russia.

The daily loss is not the same during the continuance of the period. It generally is at first slight, and gradually increases, so as to be most profuse on the second or third day, and as gradually diminishes. Toward the last days it sometimes disappears for a few hours, and then comes on again, and is apt to recur under any excitement or emotion.

Quality of Menstrual Blood.—As the menstrual fluid escapes from the uterus it consists of pure blood, and if collected through the speculum it coagulates. The ordinary menstrual fluid does not coagulate unless it is excessive in amount. Various explanations of this fact have been given. It was formerly supposed either to contain no fibrin or an unusually small amount. Retzius attributes its non-coagulation to the presence of free lactic and phosphoric acids. The true explanation was first given by Mandl, who proved that even small quantities of pus or mucus in blood were sufficient to keep the fibrin in solution; and mucus is always present to greater or less amount in the secretions of the cervix and vagina, which mix with the menstrual blood in its passage through the genital tract. If the amount of blood be excessive, however, the mucus present is insufficient in quantity to produce this effect, and coagula are then formed.

On microscopic examination the menstrual fluid exhibits blood-corpuscles, mucous corpuscles, and a considerable amount of epithelial scales, the last being the debris of the epithelium lining the uterine

cavity. According to Virchow, the form of the epithelium often proves that it comes from the interior of the utricular glands. The color of the blood is at first dark, and as the period progresses it generally becomes lighter in tint. In women who are in bad health it is often very pale. These differences doubtless depend upon the amount of mucus mingled with it. The menstrual blood has always a characteristic, faint, and heavy odor, which is analogous to that which is so distinct in the lower animals during the rut. Raciborski mentions a lady who was so sensitive to this odor that she could always tell to a certainty when any woman was menstruating. It is attributed either to decomposing mucus mixed with the blood, which, when partially absorbed, may cause the peculiar odor of the breath often perceptible in menstruating women, or to the mixture with the fluid of the sebaceous secretion from the glands of the vulva. It probably gave rise to the old and prevalent prejudices as to the deleterious properties of menstrual blood, which, it is needless to say, are altogether without foundation.

Source of the Blood.—It is now universally admitted that the source of the menstrual blood is the mucous membrane lining the interior of the uterus, for the blood may be seen oozing through the os uteri by means of the speculum, and in cases of prolapsus uteri, while in cases of inverted uterus it may be actually observed escaping from the exposed mucous membrane, and collecting in minute drops upon its surface. During the menstrual nixus the whole mucous lining becomes congested to such an extent that, in examining the bodies of women who have died during menstruation, it is found to be thicker, larger, and thrown into folds, so as to completely fill the uterine cavity. The capillary circulation at this time becomes very marked, and the mucous membrane assumes a deep-red hue, the network of capillaries surrounding the orifices of the utricular glands being especially distinct. These facts have an unquestionable connection with the production of the discharge, but there is much difference of opinion as to the precise mode in which the blood escapes from the vessels. Coste believed that the blood transudes through the coats of the capillaries without any laceration of their structure. Farre inclines to the hypothesis that the uterine capillaries terminate by open mouths, the escape of blood through these, between the menstrual periods, being prevented by muscular contraction of the uterine walls. Pouchet believed that during each menstrual epoch the entire mucous membrane is broken down and cast off in the form of minute shreds, a fresh mucous membrane being developed in the interval between two periods. During this process the capillary network would be laid bare and ruptured, and the escape of blood readily accounted for. Tyler Smith, who adopted this theory, states that he has frequently seen the uterine mucous membrane, in women who have died during menstruation, in a state of dissolution, with the broken loops of the capillaries exposed. The phenomena attending the so-called membranous dysmenorrhœa, in which the mucous membrane is thrown off in shreds or as a cast of the uterine cavity—the nature of which was first pointed out by Simpson and Oldham—have been supposed to corroborate this theory. This view is, in the main, corroborated by the

recent researches of Engelmann,¹ Williams,² and others. Williams describes the mucous lining of the uterus as undergoing a fatty degeneration before each period, which commences near the inner os, and extends over the whole mucous membrane and down to the muscular wall. This seems to bring on a certain amount of muscular contraction, which drives the blood into the capillaries of the mucosa, and these, having become degenerated, readily rupture, and permit the escape of the blood. The mucous membrane now rapidly disintegrates, and is cast off in shreds with the menstrual discharge, in which masses of epithelial cells may always be detected. Engelmann, however, holds that the fatty degeneration is limited to the superficial layers, and that a portion only of the epithelial investment is thrown off. As soon as the period is over the formation of a new mucous membrane is begun, from proliferation of the elements of the muscular coat, and at the end of a week the whole uterine cavity is lined by a thin mucous membrane. This grows until the advent of another period, when the same degenerative changes occur unless impregnation has taken place, in which case it becomes further developed into the decidua.

Theory of Menstruation.—That there is an intimate connection between ovulation and menstruation is admitted by most physiologists, and it is held by many that the determining cause of the discharge is the periodic maturation of the Graafian follicles. There is abundant evidence of this connection, for we know that when, at the change of life, the Graafian follicles cease to develop, menstruation is arrested; and when the ovaries are removed by operation, of which there are now numerous cases on record, or when they are congenitally absent, menstruation does not generally take place. A few cases, however, have been observed in which menstruation continued after double ovariectomy or the removal of the ovaries by Battey's operation; and these have been used as an argument by those physiologists who doubt the ovular theory of menstruation. Slavyanski has particularly insisted on such cases, which, however, are probably susceptible of explanation. It may be that the habit of menstruation may continue for a time even after the removal of the ovaries; and it has not been shown that menstruation has continued permanently after double ovariectomy, although it certainly has occasionally, although quite exceptionally, done so for a time. It is possible, also, that in such cases a small portion of ovarian tissue may have been left unremoved, sufficient to carry on ovulation. Roberts, a traveller quoted by Depaul and Gueniot in their article on "Menstruation" in the *Dictionnaire des Sciences Médicales*, relates that in certain parts of Central Asia it is the custom to remove both ovaries in young girls who act as guards to the harems. These women, known as "hed-jeras," subsequently assume much of the virile type, and never menstruate. The same close connection between ovulation and the rut of animals is observed, and supports the conclusion that the rut and menstruation are analogous. The chief difference between ovulation in man and the lower animals is that in the latter the process is not generally accompanied by a sanguineous flow. To this there are exceptions, for in

¹ *American Journal of Obstetrics*, May, 1875.

² "On the Structure of the Mucous Membrane of the Uterus," *Obst. Journ.*, 1875.

monkeys there is certainly a discharge analogous to menstruation occurring at intervals. Another point of distinction is that in animals connection never takes place except during the rut, and that it is then only that the female is capable of conception; while in the human race conception only occurs in the interval between the periods. This is another argument brought against the ovular theory, because, it is said, if menstruation depends on the rupture of a Graafian follicle and the emission of an ovule, then impregnation should only take place during or immediately after menstruation. Coste explains this by supposing that it is the *maturation* and not the *rupture* of the follicle which determines the occurrence of menstruation, and that the follicle may remain unruptured for a considerable time after it is mature, the escape of the ovule being subsequently determined by some accidental cause, such as sexual excitement. However this may be, there is good reason to believe that the susceptibility to conception is greater during the menstrual epochs. Raciborski believes that in the large proportion of cases impregnation occurs in the first half of the menstrual interval, or in the few days immediately preceding the appearance of the discharge. There are, however, very numerous exceptions, for in Jewesses, who almost invariably live apart from their husbands for eight days after the cessation of menstruation, impregnation must constantly occur at some other period of the interval, and it is certain that they are not less prolific than other people. This rule with them is very strictly adhered to, as will be seen by the accompanying interesting letter from a medical friend who is a well-known member of that community, and which I have permission to publish.¹ This fact is of itself sufficient to disprove the theory

¹ 10 Bernard Street, Russell Square. July 21, 1873.

MY DEAR SIR:

1. To the best of my knowledge and belief, the law which prohibits sexual intercourse among Jews for seven clear days after the cessation of menstruation, is almost universally observed, the exceptions not being sufficient to vitiate statistics. The law has perhaps fewer exceptions on the Continent—especially Russia and Poland, where the Jewish population is very great—than in England. Even here, however, women who observe no other ceremonial law observe this, and cling to it after everything else is thrown overboard. There are doubtless many exceptions, especially among the better classes in England, who keep only three days after the cessation of the menses.

2. The law is—as you state—that should the discharge last only an hour or so, or should there be only one gush or one spot on the linen, the five days during which the period *might* continue are observed; to which must be superadded the seven clear days—twelve days per mensem in which connection is disallowed. Should any discharge be seen in the inter-menstrual period, seven days would have to be kept, but not the five, for such *irregular* discharge.

3. The “bath of purification,” which must contain *at least* eighty gallons, is used on the last night of the seven clear days. It is not used till after a bath for cleansing purposes; and, from the night when such “purifying” bath is used Jewish women are accustomed to calculate the commencement of pregnancy. That you should not have heard it is not strange: its mention would be considered highly indelicate.

4. Jewish women reckon their pregnancy to last nine calendar or ten lunar months—270 to 280 days. There are no special data on which to reckon on an average, nor do I know of any books on the subject, except some Talmudic authorities, which I will look up for you if you desire it. Pray make no apologies for writing to me: any information I possess is at your service.

I am, dear sir, yours very truly,

Dr. Playfair.

A. ASHER.

P. S.—The biblical foundation for the law of seven clear days is Leviticus xv., verse 19 to the end of the chapter—especially verse 28.

advanced by Dr. Avrard,¹ that impregnation is impossible in the latter half of the menstrual interval. This, and the other reasons referred to, undoubtedly throw some doubt on the ovular theory, but they do not seem to be sufficient to justify the conclusion that menstruation is a physiological process altogether independent of the development and maturation of the Graafian follicles. All that they can be fairly held to prove is that the escape of the ovules may occur independently of menstruation, but the weight of evidence remains strongly in favor of the theory which is generally received. It should be stated that Lawson Tait attributes considerable influence in menstruation to the Fallopian tubes themselves; but his views on this point, based on observations made after the removal of the ovaries for certain morbid conditions, cannot yet be taken as proved.

Purpose of the Menstrual Loss.—The cause of the monthly periodicity is quite unknown, and will probably always remain so. Goodman² has suggested what he calls the "cyclical theory of menstruation," which refers the phenomena to a general condition of the vascular system specially localizing itself in the generative organs, and connected with rhythmical changes in their nerve-centres. It does not seem to me, however, that he has satisfactorily proved the recurrence of the conditions which his ingenious theory assumes. The purpose of the loss of so much blood is also somewhat obscure. To a certain extent it must be considered an accident or complication of ovulation, produced by the vascular turgescence. Nor is it essential to fecundation, because women often conceive during lactation, when menstruation is suspended or before the function has become established. It may, however, serve the negative purpose of relieving the congested uterine capillaries which are periodically filled with a supply of blood for the great growth which takes place when conception has occurred. Thus immediately before each period the uterus may be considered to be placed by the afflux of blood in a state of preparation for the function it may be suddenly called upon to perform. That the discharge relieves a state of vascular tension which accompanies ovulation is proved by the singular phenomenon of vicarious menstruation which is occasionally, though rarely, met with. It occurs in cases in which, from some unexplained cause, the discharge does not escape from the uterine mucous membrane. Under such circumstances a more or less regular escape of blood may take place from other sites. The most common situations are the mucous membranes of the stomach, of the nasal cavities, or of the lungs; the skin, not uncommonly that of the mammae, probably on account of their intimate sympathetic relation with the uterine organs; from the surface of an ulcer; or from hemorrhoids. It is a noteworthy fact that in all these cases the discharge occurs in situations where its external escape can readily take place. This strange deviation of the menstrual discharge may be taken as a sign of general ill-health, and it is usually met with in delicate young women of highly mobile nervous constitution. It may, however, begin at puberty, and it has even been observed during the whole sexual life. The recurrence is regular, and always in connection with the men-

¹ *Rev. de. Thérap. Méd.-Chir.*, 1857.

² *American Journal of Obstetrics*, Oct., 1868.

strual nismus, although the amount of blood lost is much less than in ordinary menstruation.

Cessation of Menstruation.—After a certain time changes occur, showing that the woman is no longer fitted for reproduction; menstruation ceases, Graafian follicles are no longer matured, and the ovary becomes shrivelled and wrinkled on its surface. Analogous alterations take place in the uterus and its appendages. The Fallopian tubes atrophy, and are not unfrequently obliterated. The uterus decreases in size. The cervix undergoes a remarkable change, which is readily detected on vaginal examination; the projection of the cervix into the vaginal canal disappears, and the orifice of the os uteri in old women is found to be flush with the roof of the vagina. In a large number of cases there is, after the cessation of menstruation, an occlusion both of the external and internal os; the canal of the cervix between them, however, remains patulous, and is not unfrequently distended with a mucous secretion.

Period of Cessation.—The age at which menstruation ceases varies much in different women. In certain cases it may cease at an unusually early age, as between 30 and 40 years, or it may continue far beyond the average time, even up to 60 years; and exceptional, though perhaps hardly reliable, instances are recorded in which it has continued even to 80 or 90 years. These are, however, strange anomalies, which, like cases of unusually precocious menstruation, cannot be considered as having any bearing on the general rule. Most cases of so-called protracted menstruation will be found to be really morbid losses of blood depending on malignant or other forms of organic disease, the existence of which, under such circumstances, should always be suspected.

In this country menstruation usually ceases between 40 and 50 years of age. Raciborski says that the largest number of cases of cessation are met with in the 46th year. It is generally said that women who commence to menstruate when very young cease to do so at a comparatively early age, so that the average duration of the function is about the same in all women. Cazcaux and Raciborski, whose opinion is strengthened by the observations of Guy in 1500 cases,¹ think, on the contrary, that the earlier menstruation commences the longer it lasts, early menstruation indicating an excess of vital energy which continues during the whole childbearing life. Climate and other accidental causes do not seem to have as much effect on the cessation as on the establishment of the function. It does not appear to cease earlier in warm than in temperate climates. The change of life is generally indicated by irregularities in the recurrence of the discharge. It seldom ceases suddenly, but it may be absent for one or more periods, and then occur irregularly; or it may become profuse or scanty, until eventually it entirely stops. The popular notions as to the extreme danger of the menopause are probably much exaggerated, although it is certain that at that time various nervous phenomena are apt to be developed. So far from having a prejudicial effect on the health, however, it is not an uncommon observation to see an hysterical woman, who has been for years a martyr to uterine and other complaints, apparently take a new lease of life when her uterine functions have ceased to be in active operation;

¹ *Med. Times and Gaz.*, 1845.

and statistical tables abundantly prove that the general mortality of the sex is not greater at this than at any other time.

[The theory that the average duration of menstrual life is thirty years, and that the period of cessation is usually regulated by the age at which menstruation was established, is an error. The age of cessation is much more irregular than that of commencement, and women cease to menstruate at all ages from 28 to 56, and even beyond 60. If careful inquiry is made, especially among women in the higher walks of life, it will be found that those who begin to menstruate at 9, 10, 11, and 12 years of age often continue to do so until long after thirty years have passed, and that it is not uncommon for such women to be among the latest in reaching the menopause. I have known the menopause to be established before 30, and menstruation to be still regular at 62 years of age.—ED.]

PART II.

PREGNANCY.

CHAPTER I.

CONCEPTION AND GENERATION.

GENERATION in the human female, as in all mammals, requires the congress of the two sexes, in order that the semen, the male element of generation, may be brought into contact with the ovule, the female element of generation.

The Semen.—The semen secreted by the testicle of an adult male is a viscid, opalescent fluid, forming an emulsion when mixed with water, and having a peculiar faint odor, which is attributed to the secretions which are mixed with it, such as those from the prostate and Cowper's glands. On analysis it is found to be an albuminous fluid, holding in solution various salts, principally phosphates and chlorides, and an animal substance, spermatin, analogous to fibrin. Examined under a magnifying power of from 400 to 500 diameters, it consists of a transparent and homogeneous fluid, in which are floating a certain number of granules and epithelial cells derived from the secretions mixed with it, and the characteristic sperm-cells and spermatozoa which form its essential constituents. The sperm-cells are those occupying the tubuli semeniferi of the testicle. Several kinds of sperm-cells are described which receive their name from the position they occupy with regard to the lumen of the tubule (Fig. 44). The cells which are next to the wall of the tubule are called the outer or lining cells. They are more or less flattened in form, and are situated on a distinct basement membrane. Internal to this layer is another, consisting of round cells, the nuclei of which are in a state of proliferation; this is the intermediate layer. Between this and the lumen of the tubule are a number of cells, irregular in shape, amongst which are imbedded the heads of the spermatozoa, the tails of which project into the lumen. The spermatozoa are thought to arise from this innermost layer in the following manner: the nuclei of the sperm-cells proliferate, and from their subdivisions arise the heads of the spermatozoa, the bodies of which originate from the protoplasm of the cells. By the decomposition of the substance in which the heads of the spermatozoa are imbedded the contained spermatozoa become liberated, and move about freely in the seminal fluid. As seen under the microscope, the spermatozoa, which exist in healthy semen in enormous numbers, present the appearance of minute particles, not unlike a tadpole in

the tubules measuring about $\frac{1}{100000}$ of an inch in diameter. This tubule is a filamentous expansion of the spermatozoa, and its termination cannot be seen in the microscope. The whole spermatozoa are contained in the sperm. The spermatozoa are motile, sometimes very rapid, sometimes more

FIG. 4.




FIG. 4. Three Seminiferous Tubules of the Rat. *a*, Early development. *b*, More advanced. *c*, Containing fully-developed spermatozoa. *Strands* of interstitial cells and lymph-spaces. (From a preparation by Dr. J. E. Smith.)

are supposed to be the means by which they pass upward to the genital organs. They retain their vitality and power of motion for a considerable time after emission, provided the semen is kept at a temperature similar to that of the body. Under such circumstances have been observed in active motion from forty-eight to seventy-two hours after ejaculation, and they have also been seen alive as long as twenty-four hours after death. In all probability they are active much longer within the generative organs, as Dr. J. E. Smith has observed them in full vitality in bitches and cats several days after copulation. The recent experiments of Dr. J. E. Smith show that they lose their power of motion in the female within twelve hours after coitus, although they doubtless remain in the uterus and Fallopian tubes. Abundant leucocytes and acid vaginal secretions destroy their movements, and cause sterility in the female. On account of their mobility, they were long considered to be independent animalcules—this view, however, has exploded, and has been maintained in modern times by Joulin, and other writers, while Coste, Robin, and others have attributed their motion to that of ciliated epithelium. There is no doubt that the fertilizing power of the semen is due to the spermatozoa, although some of the older physiologists attributed it to the seminal fluid. The former view, however, has been

conclusively proved by the experiments of Prévost and Dumas, who found that on carefully removing the spermatozoa by filtration the semen lost its fecundating properties.

Sites of Impregnation.—There has been great difference of opinion as to the part of the genital tract in which the spermatozoa and the ovule come into contact, and in which impregnation, therefore, occurs. Spermatozoa have been observed in all parts of the female genital organs in animals killed shortly after coitus, especially in the Fallopian tubes, and even on the surface of the ovary. The phenomena of ovarian gestation, and the fact that fecundation has been proved to occur in certain animals within the ovary, tend to support the idea that it may also occur in the human female before the rupture of the Graafian follicle. In order to do so, however, it is necessary for the spermatozoa to penetrate the proper structure of the follicle and the epithelial covering of the ovary, and no one has actually seen them doing so. Most probably the contact of the spermatozoa and the ovule occurs very shortly after the rupture of the follicle, and in the outer part of the Fallopian tubes. Coste maintains that unless the ovule is impregnated it very rapidly degenerates after being expelled from the ovary, partly by inherent changes in the ovule itself, and partly because it then soon becomes invested by an albuminous covering which is impermeable to the spermatozoa. He believes, therefore, that impregnation can only occur either on the surface of the ovary or just within the fimbriated extremity of the tube.

Mode in which the Ascent of the Semen is Effected.—The semen is probably carried upward chiefly by the inherent mobility of the spermatozoa. It is believed by some that this is assisted by other agencies: amongst them are mentioned the peristaltic action of the uterus and Fallopian tubes; a sort of capillary attraction effected when the walls of the uterus are in close contact, similar to the movement of fluid in minute tubes; and also the vibratile action of the cilia of the epithelium of the uterine mucous membrane. The action of the latter is extremely doubtful, for they are also supposed to effect the descent of the ovule, and they can hardly act in two opposite ways. The movement of the cilia being from within outward, it would certainly oppose rather than favor the progress of the spermatozoa. It must, therefore, be admitted that they ascend chiefly through their own powers of motion. They certainly have this power to a remarkable extent, for there are numerous cases on record in which impregnation has occurred without penetration, and even when the hymen was quite entire, and in which the semen has simply been deposited on the exterior of the vulva; in such cases, which are far from uncommon, the spermatozoa must have found their way through the whole length of the vagina. It is probable, however, that under ordinary circumstances the passage of the spermatic fluid into the uterus is facilitated by changes which take place in the cervix during




FIG. 45.
Ovum of Rabbit containing Spermatozoa.
1. Zona pellucida. 2. The germ, consisting of two large cells, several smaller cells, and spermatozoa.

the sexual orgasm, in the course of which the os uteri is said to dilate and close again in a rhythmical manner.¹

Mode of Impregnation.—The precise method in which the spermatozoa effect impregnation was long a matter of doubt. It is now, however, certain that they actually penetrate the ovule and reach its interior. This has been conclusively proved by the observations of Barry, Meissner, and others, who have seen the spermatozoa within the external membrane of the ovule in rabbits (Fig. 45). In some of the invertebrata a canal or opening exists in the zona pellucida, through which the spermatozoa pass. No such aperture has yet been demonstrated in the ovules of mammals, but its existence is far from improbable. According to the observations of Newport, several spermatozoa enter the ovule, and the greater the number that do so the more certain fecundation becomes. After the spermatozoa penetrate the zona pellucida, they disintegrate and mingle with the yolk, having, while doing so, imparted to the ovule a power of vitality and initiated its development into a new being.


Progress of the Impregnated Ovule toward the Uterus.—The length of time which lapses before the fecundated ovule arrives in the cavity of the uterus has not yet been ascertained, and it probably varies under different circumstances. It is known that in the bitch it may remain eight or ten days in the Fallopian tube, in the guinea-pig three or four. In the human female the ovum has never been discovered in the cavity of the uterus before the tenth or twelfth day after impregnation.

Changes which the Ovule undergoes immediately before and after Impregnation.—The changes which occur in the human ovule immediately before and after impregnation, and during its progress through the Fallopian tube, are only known to us by analogy, as, of course, it is impossible to study them by actual observation. We are in possession, however, of accurate information of what has been made out in the lower animals, and it is reasonable to suppose that similar changes occur in man. Immediately after the ovule has passed into the Fallopian tube, it is found to be surrounded by a layer of granular cells, a portion of the lining membrane of the Graafian follicle, which was described as the discus proligerus. As it proceeds along the tube, these surrounding cells disappear—partly, it is supposed, by friction on the walls of the tube, and partly by being absorbed to nourish the ovule. Be this as it may, before long they are no longer observed, and the zona pellucida forms the outermost layer of the ovule. When the ovule has advanced some distance along the tube, it becomes invested with a covering of albuminous material, which is deposited around it in successive layers, the thickness of which varies in different animals. It is very abundant in birds, in whom it forms the familiar white of the egg. In some animals it has not been detected, so that its presence in the human ovule is uncertain. Where it exists it doubtless contributes to the nourishment of the ovule. Coincident with these changes is the disappearance of the germinal vesicle. At the same time the yolk contracts and becomes more solid, retiring from close contact with the zona pellucida, and thus forming a species of cavity between the outer edge of the yolk and

¹ *How do the Spermatozoa enter the Uterus?* by J. Beck, M. D.

the vitelline membrane, which in some animals is filled with a transparent liquid. Coincident with the shrinking of the yolk, a small granular mass of a rounded form is extruded from the yolk into the clear space beneath the zona pellucida. At a later period another similar mass is extruded. These are the *polar globules* (Fig. 46), the origin of which is thought to be in connection with the disappearance of the germinal vesicle and the germinal spot. These changes occur in all ovules, whether they are impregnated or not, but if the ovule is not fecundated, no further alterations occur. Supposing impregnation has taken place, a bright, clear vesicle, called the *vitelline nucleus*, very similar in appearance to a drop of oil, appears in the centre of the yolk. After this occurs the very peculiar phenomenon known as the cleavage of the yolk, which results in the formation of the layer of cells from which the foetus is developed. The segmentation of the yolk (Fig. 47) occupies in mammals the whole of its substance. In birds the cleavage is confined to a small area of the yolk called the *cicatrix* or *blastoderm*. Hence the term *holoblastic* has been applied to the ova of mammals, *mesoblastic* to those of birds. It divides at first into two


FIG. 46.



Formation of the "Polar Globule."

1. Zona pellucida, containing spermatozoa.
2. Yolk. 3 and 4. Germinal vesicle. 5. The polar globule.

FIG. 47.



Segmentation of the Yolk.


- A. Ovum with first embryo cell. B. Division of embryo cell and cleavage of the yolk around it. C, D, E. Further division of the yolk.

halves, and at the same time the vitelline nucleus becomes constricted in its centre, and separates into two portions, one of which forms a centre for each of the halves into which the yolk has divided. Each of these immediately divides into two, as does its contained portion of the vitelline nucleus, and so on in rapid succession until the whole yolk is divided into a number of spheres, each of which consists of a clump of nucleated protoplasm.

By these continuous dichotomous divisions the whole yolk is formed

shape. The head is oval and flattened, measuring about $\frac{1}{10000}$ of an inch in breadth, and attached to it is a delicate filamentous expansion or tail, which tapers to a point so fine that its termination cannot be seen by the highest powers of the microscope. The whole spermatozoon measures from $\frac{1}{400}$ to $\frac{1}{600}$ of an inch in length. The spermatozoa are observed to be in constant motion, sometimes very rapid, sometimes more

FIG. 44.



Section of Parts of three Semeniferous Tubules of the Rat.

a. With the spermatozoa least advanced in development. b. More advanced. c. Containing fully-developed spermatozoa. Between the tubules are seen strands of interstitial cells and lymph-spaces. (From a preparation by Mr. A. Frazer.)

gentle, which is supposed to be the means by which they pass upward through the female genital organs. They retain their vitality and power of movement for a considerable time after emission, provided the semen is kept at a temperature similar to that of the body. Under such circumstances they have been observed in active motion from forty-eight to seventy-two hours after ejaculation, and they have also been seen alive in the testicle as long as twenty-four hours after death. In all probability they continue active much longer within the generative organs, as many physiologists have observed them in full vitality in bitches and rabbits seven or eight days after copulation. The recent experiments of Haussman, however, show that they lose their power of motion in the human vagina within twelve hours after coitus, although they doubtless retain it longer in the uterus and Fallopian tubes. Abundant leucorrheal discharges and acrid vaginal secretions destroy their movements, and may thus cause sterility in the female. On account of their mobility, the spermatozoa were long considered to be independent animalcules—a view which is by no means exploded, and has been maintained in modern times by Pouchet, Joulin, and other writers, while Coste, Robin, Kölliker, etc. liken their motion to that of ciliated epithelium. There can be no doubt that the fertilizing power of the semen is due to the presence of the spermatozoa, although some of the older physiologists assigned it to the spermatic fluid. The former view, however, has been

uterus greatly hypertrophied. The second, the *decidua reflexa*, is closely applied round the ovum, and it is probably formed by the sprouting of the decidua vera around the ovum at the point on which the latter rests, so that it eventually completely surrounds it. As the ovum enlarges, the decidua reflexa is necessarily stretched, until it comes everywhere into contact with the decidua vera, with which it firmly unites. After the third month of pregnancy true union has occurred, and the two layers of decidua are no longer separate. The *decidua serotina*, which is described as a third portion, is merely that part of the decidua vera on which the ovum rests and where the placenta is eventually developed.

Views of William and John Hunter.—It is needless to refer to the various views which have been held by anatomists as to the structure and formation of the decidua. That taught by John Hunter was long believed to be correct, and down to a recent date it received the adherence of most physiologists. He believed the decidua to be an inflammatory exudation, which on account of the stimulus of pregnancy was thrown out all over the cavity of the uterus, and soon formed a distinct lining membrane to it. When the ovum reached the uterine orifice of the Fallopian tube it found its entrance barred by this new membrane, which accordingly it pushed before it. This separated portion formed a covering to the ovum and became the decidua reflexa, while a fresh exudation took place at that portion of the uterine wall which was thus laid bare, and this became the decidua vera. William Hunter had much more correct views of the decidua, the accuracy of which was at the time much contested, but which have recently received full recognition. He describes the decidua in his earlier writings as a hypertrophy of the uterine mucous membrane itself—a view which is now held by all physiologists.

Structure of the Decidua.—When the decidua is first formed, it is a hollow triangular sac lining the uterine cavity (Fig. 49), and having three openings into it—those of the Fallopian tubes at its upper angles, and one, corresponding to the internal os uteri, below. If, as is generally the case, it is thick and pulpy, these openings are closed up and can no longer be detected. In early pregnancy it is well developed, and continues to grow up to the third month of utero-gestation. After that time it commences to atrophy, its adhesion with the uterine walls lessens, it becomes thin and transparent, and is ready for expulsion when delivery is effected. When it is most developed, a careful examination of the decidua enables us to detect in it all the elements of the uterine mucous membrane greatly hypertrophied. Its substance chiefly consists of large round or oval nucleated cells and elongated fibres, mixed with the tubular uterine gland-ducts, which are much elongated, and filled with cylindrical epithelium cells and a small quantity of milky fluid. According to Friedlander, the decidua is divisible into two layers: the inner, being formed by a proliferation of the corpuscles of the sub-epithelial connective tissue of the mucous membrane; the deeper, in contact with the uterine walls, out of flattened or compressed gland-ducts. In an early abortion the extremities of these ducts may be observed by a lens, on the external or uterine surface of the decidua, occupying the summit of minute projections separated from each other by depressions.

the sexual orgasm, in the course of which the os uteri is said to dilate and close again in a rhythmical manner.¹

Mode of Impregnation.—The precise method in which the spermatozoa effect impregnation was long a matter of doubt. It is now, however, certain that they actually penetrate the ovule and reach its interior. This has been conclusively proved by the observations of Barry, Meissner, and others, who have seen the spermatozoa within the external membrane of the ovule in rabbits (Fig. 45). In some of the invertebrata a canal or opening exists in the zona pellucida, through which the spermatozoa pass. No such aperture has yet been demonstrated in the ovules of mammals, but its existence is far from improbable. According to the observations of Newport, several spermatozoa enter the ovule, and the greater the number that do so the more certain fecundation becomes. After the spermatozoa penetrate the zona pellucida, they disintegrate and mingle with the yolk, having, while doing so, imparted to the ovule a power of vitality and initiated its development into a new being.


Progress of the Impregnated Ovule toward the Uterus.—The length of time which lapses before the fecundated ovule arrives in the cavity of the uterus has not yet been ascertained, and it probably varies under different circumstances. It is known that in the bitch it may remain eight or ten days in the Fallopian tube, in the guinea-pig three or four. In the human female the ovum has never been discovered in the cavity of the uterus before the tenth or twelfth day after impregnation.

Changes which the Ovule undergoes immediately before and after Impregnation.—The changes which occur in the human ovule immediately before and after impregnation, and during its progress through the Fallopian tube, are only known to us by analogy, as, of course, it is impossible to study them by actual observation. We are in possession, however, of accurate information of what has been made out in the lower animals, and it is reasonable to suppose that similar changes occur in man. Immediately after the ovule has passed into the Fallopian tube, it is found to be surrounded by a layer of granular cells, a portion of the lining membrane of the Graafian follicle, which was described as the discus proligerus. As it proceeds along the tube, these surrounding cells disappear—partly, it is supposed, by friction on the walls of the tube, and partly by being absorbed to nourish the ovule. Be this as it may, before long they are no longer observed, and the zona pellucida forms the outermost layer of the ovule. When the ovule has advanced some distance along the tube, it becomes invested with a covering of albuminous material, which is deposited around it in successive layers, the thickness of which varies in different animals. It is very abundant in birds, in whom it forms the familiar white of the egg. In some animals it has not been detected, so that its presence in the human ovule is uncertain. Where it exists it doubtless contributes to the nourishment of the ovule. Coincident with these changes is the disappearance of the germinal vesicle. At the same time the yolk contracts and becomes more solid, retiring from close contact with the zona pellucida, and thus forming a species of cavity between the outer edge of the yolk and

¹ *How do the Spermatozoa enter the Uterus?* by J. Beck, M. D.

the vitelline membrane, which in some animals is filled with a transparent liquid. Coincident with the shrinking of the yelk, a small granular mass of a rounded form is extruded from the yelk into the clear space beneath the zona pellucida. At a later period another similar mass is extruded. These are the *polar globules* (Fig. 46), the origin of which is thought to be in connection with the disappearance of the germinal vesicle and the germinal spot. These changes occur in all ovules, whether they are impregnated or not, but if the ovule is not fecundated, no further alterations occur. Supposing impregnation has taken place, a bright, clear vesicle, called the *vitelline nucleus*, very similar in appearance to a drop of oil, appears in the centre of the yelk. After this occurs the very peculiar phenomenon known as the cleavage of the yelk, which results in the formation of the layer of cells from which the fœtus is developed. The segmentation of the yelk (Fig. 47) occupies in mammals the whole of its substance. In birds the cleavage is confined to a small area of the yelk called the cicatrix or blastoderm. Hence the term *holoblastic* has been applied to the ova of mammals, *mesoblastic* to those of birds. It divides at first into two


FIG. 46.



Formation of the "Polar Globule."

1. Zona pellucida, containing spermatozoa. 2. Yelk. 3 and 4. Germinal vesicle. 5. The polar globule.

FIG. 47.



Segmentation of the Yelk.


- A. Ovum with first embryo cell. B. Division of embryo cell and cleavage of the yelk around it. C, D, E. Further division of the yelk.

halves, and at the same time the vitelline nucleus becomes constricted in its centre, and separates into two portions, one of which forms a centre for each of the halves into which the yelk has divided. Each of these immediately divides into two, as does its contained portion of the vitelline nucleus, and so on in rapid succession until the whole yelk is divided into a number of spheres, each of which consists of a clump of nucleated protoplasm.

By these continuous dichotomous divisions the whole yelk is formed

into a granular mass, which from its supposed resemblance to a mulberry has been named the *muriform body*. When the subdivision of the yelk is completed, its separate spheres become converted into cells, consisting of a fine membrane with granular contents. These cells unite by their edges to form a continuous membrane (Fig. 48), which, through the expansion of the muriform body by fluid which forms in its interior, is distended until it forms a lining to the zona pellucida. This is the *blastodermic membrane*, from which the foetus is developed. By this time the ovum has reached the uterus; and before proceeding to consider the further changes which it undergoes, it will be well to study the alteration which the stimulus of impregnation has set on foot in the mucous membrane of the uterus, in order to prepare it for the reception and growth of its contents.

FIG. 48.



Formation of the Blastodermic Membrane from the Cells of the Muriform Body. (After Joulin.)
1. Layer of albuminous material surrounding 2. The zona pellucida.

Changes in the Uterine Mucous Membrane consequent on Pregnancy.—Even before the ovum reaches the uterus the mucous membrane becomes thickened and vascular, so that its opposing surfaces entirely fill the uterine cavity. These changes may be said to be the same in kind—although more marked and extensive in degree—as the alterations which take place in the mucous membrane in connection with each menstrual period. The result is the formation of a distinct membrane, which affords the ovum a safe anchorage and protection until its connections with the uterus are more fully developed. After delivery, this membrane, which is by that time quite altered in appearance, is at least partially thrown off with the ovum; on which account it has received the name of the *decidua* or *caduca*.

Divisions of the Decidua.—The decidua consists of two principal portions, which in early pregnancy are separated from each other by a considerable interspace. One of these, called the *decidua vera*, lines the entire uterine cavity, and is, no doubt, the original mucous lining of the

uterus greatly hypertrophied. The second, the *decidua reflexa*, is closely applied round the ovum, and it is probably formed by the sprouting of the decidua vera around the ovum at the point on which the latter rests, so that it eventually completely surrounds it. As the ovum enlarges, the decidua reflexa is necessarily stretched, until it comes everywhere into contact with the decidua vera, with which it firmly unites. After the third month of pregnancy true union has occurred, and the two layers of decidua are no longer separate. The *decidua serotina*, which is described as a third portion, is merely that part of the decidua vera on which the ovum rests and where the placenta is eventually developed.

Views of William and John Hunter.—It is needless to refer to the various views which have been held by anatomists as to the structure and formation of the decidua. That taught by John Hunter was long believed to be correct, and down to a recent date it received the adherence of most physiologists. He believed the decidua to be an inflammatory exudation, which on account of the stimulus of pregnancy was thrown out all over the cavity of the uterus, and soon formed a distinct lining membrane to it. When the ovum reached the uterine orifice of the Fallopian tube it found its entrance barred by this new membrane, which accordingly it pushed before it. This separated portion formed a covering to the ovum and became the decidua reflexa, while a fresh exudation took place at that portion of the uterine wall which was thus laid bare, and this became the decidua vera. William Hunter had much more correct views of the decidua, the accuracy of which was at the time much contested, but which have recently received full recognition. He describes the decidua in his earlier writings as a hypertrophy of the uterine mucous membrane itself—a view which is now held by all physiologists.

Structure of the Decidua.—When the decidua is first formed, it is a hollow triangular sac lining the uterine cavity (Fig. 49), and having three openings into it—those of the Fallopian tubes at its upper angles, and one, corresponding to the internal os uteri, below. If, as is generally the case, it is thick and pulpy, these openings are closed up and can no longer be detected. In early pregnancy it is well developed, and continues to grow up to the third month of utero-gestation. After that time it commences to atrophy, its adhesion with the uterine walls lessens, it becomes thin and transparent, and is ready for expulsion when delivery is effected. When it is most developed, a careful examination of the decidua enables us to detect in it all the elements of the uterine mucous membrane greatly hypertrophied. Its substance chiefly consists of large round or oval nucleated cells and elongated fibres, mixed with the tubular uterine gland-ducts, which are much elongated, and filled with cylindrical epithelium cells and a small quantity of milky fluid. According to Friedländer, the decidua is divisible into two layers: the inner, being formed by a proliferation of the corpuscles of the sub-epithelial connective tissue of the mucous membrane; the deeper, in contact with the uterine walls, out of flattened or compressed gland-ducts. In an early abortion the extremities of these ducts may be observed by a lens, on the external or uterine surface of the decidua, occupying the summit of minute projections separated from each other by depressions.

If these projections be bisected they will be found to contain little cavities, filled with lactescent fluid, which were first described by Montgomery of Dublin, and are known as *Montgomery's cups*. They are in fact

FIG. 49.



Aborted Ovum of about forty days, showing the Triangular Shape of the Decidua (which is laid open), and the Aperture of the Fallopian Tube. (After Coste.)

the dilated canals of the uterine tubular glands. On the internal surface of such an early decidua a number of shallow depressions may be made out, which are the open mouths of these ducts.

FIG. 50.



Formation of Decidua.
(The decidua is colored black;
the ovum is represented as
engaged between two pro-
jecting folds of membrane.)

FIG. 51.



Projecting Folds of Mem-
brane growing up around
the Ovum.

FIG. 52.




Showing Ovum completely sur-
rounded by the Decidua Re-
flexa.

(After Dalton.)

Formation of the Decidua Reflexa.—When the ovum reaches the uterine cavity, it soon becomes imbedded in the folds of the hypertrophied mucous membrane, which almost entirely fills the uterine cavity. As a

rule, it is attached to some point near the opening of a Fallopian tube, the swollen folds of mucous membrane preventing its descent to the lower part of the uterus; in exceptional circumstances, however—as in women who have borne many children and have a more than usually dilated uterine cavity—it may fix itself at a point much nearer the internal os uteri. According to the now generally accepted opinion of Coste, the mucous membrane at the base of the ovum soon begins to sprout around it, and gradually extends until it eventually covers the ovum (Figs. 50–52) and forms the decidua reflexa. Coste describes, under the name of the *umbilicus*, a small depression at the most prominent part of the ovum, which he considers to be the indication of the point where the closure of the decidua reflexa is effected. There are some objections to this theory, for no one has seen the decidua reflexa incomplete and in the process of formation, and on examining its external surface—that is, the one farthest from the ovum—its microscopical appearance is identi-

FIG. 53.



An Ovum removed from Uterus, and part of the Decidua Vera cut away. (After Coste.)
 a. Decidua vera, showing the follicles opening on its inner surface. b. Inner extremity of Fallopian tube
 c. Flap of decidua reflexa. d. Ovum.

cal with that of the inner surface of the decidua vera. To meet these difficulties, Weber and Goodsir, whose views have been adopted by Priestley, contended that the decidua reflexa is “the primary lamina of the mucous membrane, which, when the ovum enters the uterus, separates in two-thirds of its extent from the layers beneath it, to adhere to the ovum; the remaining third remains attached, and forms a centre of nutrition.” According to this view, the decidua vera would be a subsequent growth over the separated portion, and the decidua serotina the portion of the primary lamina which remained attached. In this way the fact of the opposed surfaces of the decidua vera and reflexa being identical in structure would be accounted for. The difficulty which this theory is thus intended to meet does not seem so great as is supposed; for if, as is likely, it is only the epithelial or internal surface of the mucous membrane which sprouts over the ovum, and not its deeper layers, the facts of the case would be sufficiently met by Coste’s view.

Up to the third month of pregnancy the decidua reflexa and vera are not in close contact, and there may even be a considerable interspace between them, which sometimes contains a small quantity of mucous fluid, called the *hydroperione*. This fact may account for the curious circumstance, of which many instances are on record, that a uterine sound may be passed into a gravid uterus in the early months of pregnancy without necessarily producing abortion, and also for the occasional occurrence of menstruation after conception (Figs. 53 and 80). Eventually, by the growth of the ovum, the decidua reflexa comes closely into contact with the vera, and the two become intimately blended and inseparable.

Decidua at the End of Pregnancy and after Delivery.—As pregnancy advances, the decidua alters in appearance and becomes fibrous and thin. In the later months of utero-gestation fatty degeneration of its structure commences, its vessels and glands are obliterated, and its adhesion to the uterine walls is lessened, so as to prepare it for separation. As we shall subsequently see, this fatty degeneration was assumed by Simpson to be the determining cause of labor at term. After the eighth month, thrombi form in the veins lying underneath the decidua serotina, and at the end of pregnancy they are described by Leopold¹ as having become, to a great extent, obliterated. This, he supposes, may have some effect in inducing the contractions of the uterus in labor.

Views of Robin.—It was long believed that the entire decidua was thrown off after labor, leaving the muscular coat of the uterus bare and denuded, and that a new mucous membrane was formed during convalescence. According to Robin,² whose views are corroborated by Priestley, no such denudation of the muscular tissue of the uterus ever occurs, but a portion of the decidua always remains attached after delivery. After the fourth month of pregnancy they believe that a new mucous membrane is formed under the decidua, which remains in a somewhat imperfect condition till after delivery, when it rapidly develops and assumes the proper functions of the mucous lining of the uterus. Robin also believes that that portion of the decidua which covers the placental site, the so-called decidua serotina, is not thrown off with the membranes, like the decidua vera and reflexa, but remains attached to the uterine walls, a thin layer of it only being expelled with the placenta, on which it may be observed. Duncan³ entirely dissents from these views, and does not admit the formation of a new mucous membrane during the later months of utero-gestation. He believes that the greater part of the decidua is thrown off, but that part remains, and from this the fresh mucous membrane is developed. This view is similar to that of Spiegelberg, who holds that the portion of the decidua that is expelled is the more superficial of the two layers described by Friedlander, composed chiefly of the epithelial elements, while the deeper or glandular layer remains attached to the walls of the uterus. From the epithelium of the glands a new epithelial layer is rapidly developed after delivery. Leopold⁴ has shown that the uterine mucous membrane is completely re-formed within six weeks after delivery, and that its regeneration is sometimes completed as early as the end of the

¹ *Arch. f. Gyn.*, B. xi. H. 3.

² *Mémoires de l'Acad. Imp. de Méd.*, 1861.

³ *Researches in Obstetrics*, p. 186 et seq.

⁴ *Arch. f. Gyn.*, B. xii. H. 2.

third week. This theory bears on the well-known analogy of the uterus after delivery to the stump of an amputated limb—an old simile, principally based on the erroneous theory that the whole muscular tissue of the uterus was laid bare. This, as we have seen, is not the case, but the simile so far holds good in that the mucous lining is deprived of its epithelial covering; and this fact, together with the existence of numerous open veins on the interior of the uterus, readily explains the extreme susceptibility to septic absorption which forms so peculiar a characteristic of the puerperal state.

Changes in the Ovum.—Before we commenced the study of the decidua we had traced the impregnated ovum into the uterine cavity, and described the formation of the blastodermic membrane by the junction of the cells of the muriform body. We must now proceed to consider the further changes which result in the development of the fœtus and of the membranes that surround it. It would be quite out of place in a work of this kind to enter into the subject of embryology at any length, and we must therefore be content with such details as are of importance from a practical point of view.

Division of the Blastodermic Membrane into Layers.—The blastodermic membrane, which forms a complete spherical lining to the ovum between the yolk and the zona pellucida, soon divides into two layers—the most external, called the *epiblast*, and an internal, the *hypoblast*—and between them is subsequently developed a third, known as the *mesoblast*. From these three layers are formed the entire fœtus: the epiblast giving origin to the central nervous system, to the superficial layer of the skin, and aiding in formation of the organs of special sense and of the amnion; the hypoblast forming the epithelial lining membrane of the alimentary and respiratory tracts, and of the tubes and glands in connection with them, and helping in the development of the yolk-sac; the mesoblast giving rise to the skeleton, the muscles, the connective tissues, the vascular system, the genito-urinary organs, and taking part in the formation of all the membranes.

The Area Germinativa.—Almost immediately after the separation of the blastodermic membrane into layers one part of it becomes thickened by the aggregation of cells, and is called the *area germinativa*. This is at first round and then oval in shape, and in its centre the first indication of the embryo may be detected in the form of a narrow straight line, the *primitive trace*. Surrounding it are some cells more translucent than those of the rest of the *area germinativa*, and hence called the *area pellucida* (Fig. 54). In front of the primitive trace two elevated ridges soon arise, the *laminae dorsales*, which include between them a groove—the medullary groove—and gradually unite posteriorly

FIG. 54.




Diagram of Area Germinativa, showing the Primitive Trace and Area Pellucida.

to form a cavity within which the cerebro-spinal axis is subsequently developed. The medullary groove as it grows backward overlaps the primitive trace, which disappears. The embryo is differentiated from the rest of the blastoderm by a fold anteriorly, which is called the cephalic or head fold. Another fold afterward appears posteriorly, which is called the caudal or tail fold. Laterally folds also arise. These folds all tend to grow toward the centre of the under surface of what will be the embryo.

The mesoblastic layer of the blastoderm, except that part which forms the axis of the embryo, splits into an upper layer, the somato-pleure, which is beneath the epiblast, and a lower layer, the splanchno-pleure, which lies upon the hypoblast. The space formed by this cleavage of the mesoblast is called the pleuro-peritoneal cavity. The somato-pleure is engaged in the formation of the body walls of the embryo. The splanchno-pleure forms the walls of the alimentary tract.

Formation of the Amnion.—Processes arise from the somato-pleure anteriorly, posteriorly, and laterally, which gradually arch over the

dorsal surface of the fœtus until they meet each other and form a complete envelope to it. At the ventral surface these processes are separated by the whole length of the embryo, but they here also gradually approach each other, and eventually surround what is subsequently the umbilical cord, and blend with the integument of the fœtus at the point of its insertion. In this way is formed the *amnion* (Fig. 55), consisting of two layers: the internal, derived from the epiblast, is formed of tessellated epithelial cells; the external, arising from the mesoblast, is formed of cells like those of young connective tissue. Before the folds of the amnion unite, the free edge of each is bent outward and



Development of the Amnion.

1. Vitelline membrane. 2. External layer of blastodermic membrane. 3. Internal layers forming the umbilical vesicle. 4. Umbilical vessels. 5. Projections forming amnions. 6. Embryo. 7. Allantois.

spreads around the ovum, immediately within the zona pellucida, forming a lining to it, termed by Turner the *subzonal membrane*, which is connected with the development of the chorion. The amnion is the most internal of the membranes surrounding the fœtus, and will presently be studied more in detail. It soon becomes distended with fluid, the *liquor amnii*, and as this increases in amount it separates the amnion more and more from the fœtus.

Changes in the Hypoblast.—During this time the innermost layer of the blastodermic membrane or hypoblast is also developing two projections at either extremity of the fœtus, and these gradually approach each other anteriorly. As the hypoblast is in contact with the yolk, when these meet they have the effect of dividing the yolk into two portions. One, and the smaller of the two, forms eventually the intestinal canal of the fœtus; the other, and much the larger, contains the greater por-

tion of the yelk, and forms the ephemeral structure known as the *umbilical vesicle*, from which the foetus derives most of its nourishment during the early stage of its existence. Its communication with the abdominal cavity of the foetus is through the constricted portion at the point of

FIG. 56.



1. Exo-chorion. 2. External layer of blastodermic membrane. 3. Umbilical vesicle. 4. Its vessels. 5. Amnion. 6. Embryo. 7. Allantois increasing in size.

division called the *vitelline duct* (Fig. 56). An artery and vein, the *omphalo-mesenteric*, ramify on the vesicle and its duct.

As the amnion increases in size, it pushes back the umbilical vesicle toward the external membrane of the ovum, between which and the amnion it lies (Fig. 57); and when the allantois is developed, it ceases to be of any use, and rapidly shrinks and dwindles away. In most mammals no trace of it can be found after the fourth month of utero-gestation; in some, including the human female, it is said to exist as a minute vesicle at the placental end of the umbilical cord at the full period of pregnancy. The umbilical vesicle is filled with a yellowish fluid, containing many oil- and fat-globules, similar to the yelk of an egg.

The Allantois.—Somewhere about the twentieth day after conception a small vesicle is formed toward the caudal extremity of the foetus, which is called the *allantois*. This membrane in mammals is important, as it forms the greater part of the fetal placenta, a small portion of it remaining inside the body permanently as the bladder. It begins as a diverticulum from the lower part of the intestinal canal. This, at first spherical, rapidly develops and becomes pyriform in shape, while, by a process of constriction similar to that which occurs in the vitellus to form the umbilical vesicle, it becomes divided into two parts, communicating with each other, the smaller of them being eventually developed into the urinary bladder. The larger portion, leaving the abdominal cavity along with the vitelline duct, rapidly grows until it

FIG. 57.



An Embryo of about twenty-five days laid open. (After Coste.)

a. Chorion. b. Amnion. c. Cavity of chorion. d. Umbilical vesicle. e. Pedicle of allantois. f. Embryo.

comes into contact with the most external ovular membrane, the chorion, over the entire inner surface of which it spreads. In this part vessels soon develop—namely, the two umbilical arteries, derived from the abdominal aorta, and two umbilical veins, one of which subsequently disappears; these, along with the vitelline duct and the pedicle of the allantois, form the umbilical cord. The main and very important function of the allantois, therefore, is to carry the fetal vessels up to the inner surface of the subzonal membrane. Besides this purpose, the allantois, at a very early period, may receive the excretions of the fœtus and

FIG. 58.



1. Exo-chorion. 2. External layer of the blastodermic membrane. 3. Allantois. 4. Umbilical vesicle.
5. Amnion. 6. Embryo. 7. Pedicle of allantois.

serve as an excrementitious organ. According to Cazeaux, scarcely a trace of the allantois can be seen a few days after its formation. Its lower part or pedicle, however, long remains distinct, and forms part of the umbilical cord; and traces of it may be found even in adult life in the form of the urachus, which is really the dwindled pedicle and forms one of the ligaments of the bladder.

The Corps Reticulé or Vitriform Body.—Between the chorion and amnion is often found a gelatinous fluid, with minute filamentous processes traversing it, called by Velpeau the *corps reticulé*, which is not met with until the allantois comes into contact with the chorion, and which seems to be formed out of the tissues of that vesicle. It is analogous to the so-called Wharton's jelly found in the umbilical cord. When first formed it is highly vascular, but the vessels entirely disappear after the placenta is formed, and the remainder of the chorionic villi atrophy. Sometimes it exists in considerable quantities, and should the chorion rupture at the end of pregnancy it may escape and give rise to an erroneous impression that the liquor amnii has been discharged.

Recapitulation.—Before proceeding to consider the fœtal envelopes more at length, it may be useful to recapitulate the structures already alluded to as forming the ovum. In this we find—

1. The *embryo* itself.
2. A fluid, the *liquor amnii*, in which it floats.

3. The *amnion*, a purely foetal membrane surrounding the embryo, and containing the liquor amnii.

4. The *umbilical vesicle*, containing the greater portion of the yolk, serving as a source of nutrition to the early embryo through the vitelline duct, and on which ramify the omphalo-mesenteric vessels.

5. The *allantois*, a vesicle proceeding from the caudal extremity of the embryo, spreading itself over the interior of the ovum, and serving as a channel of vascular communication between the chorion and the foetus through the umbilical vessels.

6. An interspace between the outer layer of the ovum and the amnion, in which is contained the *umbilical vesicle* and *allantois* and the *corps réticulé* of Velpeau.

7. The outer layer of the ovum, along with the subzonal membrane, forming the *chorion* and *placenta*.

The Amnion.—The *amnion* is the most internal of the two membranes surrounding the foetus; its origin at an early period of foetal life has already been described. It is a perfectly smooth, transparent, but tough membrane, continuous with the integument of the foetus at the insertion of the umbilical cord, round which it forms a sheath. Soon after it is formed it becomes distended with a fluid, the *liquor amnii*, in which the foetus is suspended and floats. This fluid increases gradually in quantity, distending the amnion as it does so, until this is brought into contact with the inner surface of the chorion, from which it was at first separated by a considerable interspace.

Structure of the Amnion.—The internal surface of the amnion is smooth and glistening, and on microscopic examination it is found to consist of a layer of flattened cells, each containing a large nucleus. These rest on a stratum of fibrous tissue which gives to the membrane its toughness, and by which it is attached to the inner surface of the chorion. It is entirely destitute of vessels, nerves, and lymphatics.

The Liquor Amnii.—The quantity of the liquor amnii varies much at different periods of pregnancy. In the early months it is relatively greater in amount than the foetus, which it outweighs. As pregnancy advances, the weight of the foetus becomes four or five times greater than that of the liquor amnii, although the actual quantity of fluid increases during the whole period of gestation. The amount of fluid varies much in different pregnancies. Sometimes there is comparatively little, while at others the quantity is immense, reaching several pounds in weight, greatly distending the uterus, and thus, it may be, producing difficulty in labor.

Its Quality.—At first the liquid is clear and limpid. As pregnancy advances it becomes more turbid and dense, from the admixture of epithelial débris derived from the cutaneous surface of the foetus. In some cases, without actual disease, it may be dark green in color and thick and tenacious in consistency. It has a peculiar heavy odor, and it consists chemically of water containing albumen, with various salts, principally phosphates and chlorides.

Its Source.—The source of the liquor amnii has been much disputed. Some maintain that it is derived chiefly from the foetus—a view sufficiently disproved by the fact that the liquor amnii continues to increase in amount after the death of the foetus. Burdach believed that it is

secreted by the internal surface of the uterus, and arrives in the cavity of the amnion by transudation through the membrane. Priestley—and this seems the most probable hypothesis—thinks that it is secreted by the epithelial cells lining the membrane, which become distended with fluid, burst, and pour their contents into the amniotic cavity.

Functions and Uses.—The most obvious use of the liquor amnii is to afford a fluid medium in which the fœtus floats, and so is protected from the shocks and jars to which it would otherwise be subjected, and from undue pressure upon the uterine walls. By distending the uterus it saves the uterus from injury, which the movements of the fœtus might otherwise inflict, and the fœtus is thus also enabled to change its position freely. The facility with which version by external manipulation can be effected depends entirely on the mobility of the fœtus in the fluid which surrounds it. Some have also supposed that it prevents the fœtus, in the early months of pregnancy, from forming adhesions to the amnion. In labor it is of great service by lubricating the passages, but chiefly by forming, with the membranes, a fluid wedge which dilates the circle of the os uteri.

The Chorion.—The chorion is the more external of the truly fœtal membranes, although external to it is the decidua, having a strictly maternal origin. It is a perfectly closed sac, its external surface, in contact with the decidua, being rough and shaggy from the development of villi (Fig. 56), its internal smooth and shining. As the ovum passes along the Fallopian tube it receives, as we have seen, an albuminous coating, and this, with the zona pellucida, is developed into a temporary structure, the *primitive chorion*. On its external surface villous prominences soon appear, which have no ascertained structure, and which seem to supply the early ovum with nutriment by endosmotic absorption from the mucous membrane of the uterus. This primitive chorion, however, has not been observed in the human subject, although it may be readily seen in the ova of some of the lower animals, such as the dog and the rabbit. Some twelve days after conception, when the blastodermic membrane is formed, the true chorion appears. This is, in fact, formed by the epiblast layer of the blastodermic membrane, which everywhere lines the zona pellucida or primitive chorion, and, by pressure, causes its absorption and disappearance. On the surface of the true chorion thus formed, which is now the external envelope of the ovum, villi soon appear.

Formation of the Villi.—These villi are hollow projections like the fingers of a glove, which are raised up from the surface of the chorion (the hollows looking into the chorionic cavity), and they cover the whole external surface of the ovum, giving it the peculiar shaggy appearance observed in early abortions. They push themselves into the substance of the decidua, with which they soon become so firmly united that they cannot be separated without laceration. At first they are absolutely non-vascular, but soon the allantois, previously described, reaches the inner surface of the chorion and spreads itself over the whole of it. Each villus now receives a separate artery and vein, the former having a branch to each of the subdivisions into which the villus divides. These vessels are encased in a fine sheath of the allantois,

which enters the villus along with them and forms a lining to it, described by some as the *endo-chorion*, the external epithelial membrane of the villus, derived from the epiblast layer of the blastodermic membrane, being called the *exo-chorion*. The artery and vein lie side by side in the centre of the villus and anastomose at its extremity; each villus thus having a separate circulation.

Growth and Atrophy of the Villi.—As soon as the union of the allantois with the chorion has been effected, the villi grow very rapidly, give off branches, which in their turn give off secondary branches, and so form root-like processes of great complexity. In the early months of gestation they exist equally over the whole surface of the ovum. As pregnancy advances, however, those which are in contact with the decidua reflexa shrivel up, and by the end of the second month disappear, being no longer required for the nutrition of the ovum. The chorion and decidua thus come into close contact, being united together by fibrous shreds, which on microscopic examination are found to consist of atrophied villi. A certain number of villi—viz., those which are in contact with the decidua serotina—instead of dwindling away, increase greatly in size, and eventually develop into the organ by which the fœtus is nourished—the *placenta*.

Form of the Placenta.—This important organ serves the purpose of supplying nutriment to, and aërating the blood of, the fœtus, and on its integrity the existence of the fœtus depends. It is met with in all mammals, but is very different in form and arrangement in different classes. Thus, in the sow, mare, and in the cetacea it is diffused over the whole interior of the uterus; in the ruminants it is divided into a number of separate small masses, scattered here and there over the uterine walls; while in the carnivora and elephant it forms a zone or belt round the uterine cavity. In the human race, as well as in rodentia, insectivora, etc., the placenta is in the form of a circular mass, attached generally to some part of the uterus near the orifices of one Fallopian tube; but it may be situated anywhere in the uterine cavity, even over the internal os uteri. As it is expelled after delivery with the fœtal membranes attached to it, and as the aperture in these corresponds to the os uteri, we can generally determine pretty accurately the situation in which the placenta was placed by examining them after expulsion. The maternal surface of the placenta is somewhat convex, the fœtal concave. Its size varies greatly in different cases, and it is usually largest when the child is big, but not necessarily so. Its average diameter is from 6 to 8 inches, its weight from 18 to 24 oz., but in exceptional cases it has been found to weigh several pounds. Abnormalities of form are not very rare. Thus, the placenta has been found to be divided into distinct parts—a form said by Professor Turner to be normal in certain genera of monkeys; or smaller supplementary placenta (*placentæ succentariæ*) may exist round a central mass. These variations of shape are only of importance in consequence of a risk of part of the detached placenta being left in the uterus after delivery and giving rise to septicæmia or secondary hæmorrhage.

Attachment of the Membranes.—The fœtal membranes cover the whole fœtal surface of the placenta, being reflected from its edges so as to line

the uterine cavity, and being expelled with it after delivery. They also leave it at the insertion of the cord, to which they form a sheath. The cord is generally attached near the centre of the placenta, and from its insertion the umbilical vessels may be seen dividing and radiating over the whole foetal surface.


Its Maternal Surface.—The maternal surface is rough and divided by numerous sulci, which are best seen if the placenta is rendered convex, so as to resemble its condition when attached to the uterus. A careful examination shows that a delicate membrane covers the entire maternal surface, unites the sulci together, and dips down between them. This is, in fact, the cellular layer of the decidua serotina, which is separated and expelled with the placenta, the deeper layer remaining attached to the uterus. Numerous small openings may be seen on the surface, which are the apertures of the veins torn off from the uterus, as also those of some arteries, which, after taking several sharp turns, open suddenly into the substance of the organ.

Minute Structure of the Placenta.—As regards the minute structure of the placenta, it is certain that it consists essentially of two distinct portions—one *foetal*, consisting of the greatly hypertrophied chorion villi, with their contained vessels, which carry the foetal blood so as to bring it into intimate relation with the maternal blood, and thus admit of the necessary changes occurring in it connected with the nutrition of the foetus; and the other *maternal*, formed out of the decidua serotina and the maternal blood-vessels. These two portions are in the human female so intimately blended as to form the single deciduous organ which is thrown off after delivery. These main facts are admitted by all, but considerable differences of opinion still exist among anatomists as to the precise arrangement of these parts. In the following sketch of the subject I shall describe the views most generally entertained, merely briefly indicating the points which are contested by various authorities.

Foetal Portion of the Placenta.—The foetal portion of the placenta consists essentially of the ultimate ramifications of the chorion villi, which may be seen on microscopic examination in the form of club-shaped digitations, which are given off at every possible angle from the stem of a parent trunk, just like the branches of a plant. Within the transparent walls of the villi the capillary tubes of the contained vessels may be seen lying, distended with blood, and presenting an appearance not unlike loops of small intestine. The capillaries are the terminal ramifications of the umbilical arteries and veins, which, after reaching the site of the placenta, divide and subdivide until they at last form an immense number of minute capillary vessels, with their convexities looking toward the maternal portion of the placenta, each terminal loop being contained in one of the digitations of the chorion villi. Each arterial twig is accompanied by a corresponding venous branch, which unites with it to form the terminal arch or loop (Fig. 59). The foetal blood is carried through these arterial twigs to the villi, where it comes into intimate contact with the maternal blood, in consequence of the anatomical arrangements presently to be described; but the two do not directly mix, as the older physiologists believed, for none of the mater-

nal blood escapes when the umbilical cord is cut, nor can the minutest injections through the foetal vessels be made to pass into the maternal

FIG. 59.



Placental Villus, greatly magnified. (After Joulin.)

- 1, 2. Placental vessels, forming terminal loops. 3. Chorion tissue, forming external walls of villus.
4. Tissue surrounding vessels.

vascular system, or vice versa. In addition to the looped terminations of the umbilical vessels, Farre and Schroeder van der Kolk have described

FIG. 60.



- a. Terminal villus of foetal tuft, minutely injected. b. Its nucleated non-vascular sheath. (After Farre.)

another set of capillary vessels in connection with each villus (Fig. 60). This consists of a very fine network covering each villus, and very differ-

ent in appearance from the convoluted vessels lying in its interior, which are the only ones which have been usually described. Dr. Farre believes that these vessels only exist in the early months of pregnancy, and that they disappear as pregnancy advances. Priestley¹ suggests that they may not be vessels at all, but lymphatics, which may possibly absorb nutrient material from the mother's blood and throw it into the foetal vascular system. The existence of lymphatics, or nerves, in the placenta, however, has never been demonstrated, and they are believed not to exist.

Maternal Portion of the Placenta.—As generally described, the maternal portion of the placenta consists of large cavities, or of a single large cavity, which contains the maternal blood, and into which the villi of the chorion penetrate (Fig. 61). Into this maternal part of the viscus the curling arteries of the uterus pour their blood, which is collected from it by the uterine sinuses. The villi of the chorion, therefore, are

FIG. 61.



Diagram representing a Vertical Section of the Placenta. (After Dalton.)
 a, a. Chorion. b, b. Decidua. c, c, c, c. Orifices of uterine sinuses.

suspended in a sac filled with maternal blood, which penetrates freely between them, and with which they are brought into very intimate contact.

Theory of Reid.—Dr. John Reid believed that only the delicate internal lining of the maternal vessels entered the substance of the placenta to form the sac just spoken of. Into this the villi project, pushing before them the membrane forming the limiting wall of the placental sinuses, each of them in this way receiving an investment, just as the fingers of a hand are covered by a glove (Fig. 62).

Theory of Goodsir.—Schroeder van der Kolk and Goodsir (Fig. 63) were of opinion that not only were the maternal blood-vessels continued into the substance of the placenta, but also the processes of the decidua, which accompanied the vessels and were prolonged over each villus, so as to separate it from the limiting membrane of the maternal sinuses.

¹ *The Gravid Uterus*, p. 52.

Each villus would thus be covered by two layers of fine tissue—one from the internal lining membrane of the maternal blood-vessels, the other from the epithelial cells of the decidua.

Theory of Turner.—Turner, whose valuable researches on the comparative anatomy of the placenta have thrown much light on its structure, points out that the placenta of all animals are formed on the same

FIG. 62.





Diagram illustrating the mode in which a Placental Villus derives a Covering from the Vascular System of the Mother. (After Priestley.)

a. Villus having three terminal digitations projecting into b. Cavity of the mother's vessel. c. Dotted lines representing coat of vessel.

FIG. 63.



The Extremity of a Placental Villus.
(After Goodsir.)

a. External membrane of villus (the lining membrane of vascular system of Weber).
b. External cells of villus derived from decidua.
c. c. Nuclei of ditto.
d. The space between the maternal and fetal portions of villus.
e. Its internal membrane.
f. Its internal cells.
g. The loop of umbilical vessels.

fundamental type,¹ in which the *fœtal portion* consists of a smooth, plane-surfaced vascular membrane covered with pavement epithelium, which is brought into contact with the *maternal portion*, consisting of a smooth, plane-surfaced vascular membrane covered with columnar epithelium. The fœtal capillaries are separated from the maternal capillaries only by two opposed layers of epithelium. In various animals the placenta are more or less specialized from the generalized form, in some to a much greater extent than others. In the human placenta the maternal vessels have lost their normal cylindrical form, and are dilated into a system of freely intercommunicating placental sinuses, which are, in fact, maternal capillaries enormously enlarged, with their walls so expanded and thinned out that they cannot be recognized as a distinct layer limiting the sinus. Each fœtal chorion villus projecting into these sinuses is covered with a layer of cells distinct from those of the epithelial layer of the villus, and readily stripped from it. These are maternal in their origin, and are derived from the decidua, which sends prolongations of its tissue into the placenta. These cells, he believes, form a secreting epithelium which separates from the maternal blood a secretion for the nourishment of the fœtus, which is, in its turn, absorbed by the villi of the chorion.

Theory of Ercolani.—A view not very dissimilar to this has been advanced by Professor Ercolani of Bologna, who maintains that the maternal portion of the placenta is a new formation, strictly glandular, and not vascular, in its structure. It is formed, he thinks, by the sub-mucous connective tissue of the decidua serotina, and it dips down into the placenta and forms a sheath to each of the chorion villi, which it

¹ *Introduction to Human Anatomy, Part 2.*

separates from the maternal blood. This new glandular structure he describes as secreting a fluid, termed the "uterine milk," which is absorbed by the villi of the chorion, just as the mother's milk is absorbed by the villi of the intestines, and it is with this fluid alone that the chorion villi are in direct contact. The sheath thus formed to each villus is doubtless analogous to the layer of cells which Goodsir described as encasing each villus, but is attributed to a new structure formed after conception.

Theory of Braxton Hicks.—The existence of the maternal sinus system in the placenta is altogether denied by anatomists of eminence whose views are worthy of careful consideration. Prominent among these is Braxton Hicks,¹ who has written an elaborate paper on the subject. He holds that there is no evidence to prove that the maternal blood is poured out into a cavity in which the chorion villi float, and he believes that the curling arteries, instead of entering the so-called maternal portion of the placenta, terminate in the decidua serotina. The hypertrophied chorion villi at the site of the placenta are firmly attached to the decidual surface, into which their tips are imbedded. The line of junction between the decidual reflexa and serotina forms a circumferential margin to, and limits, the placenta. The arrangement of the foetal portion of the placenta on this view is very similar to that generally described, but the villi are not surrounded by maternal blood at all, and nothing exists between them, unless it be a small quantity of serous fluid. The change in the foetal blood is effected by endosmosis, and Hicks suggests that the follicles of the decidua may secrete a fluid, which is poured into the intervillous spaces for absorption by the villi.

Functions of the Placenta.—It will thus be seen that anatomists of repute are still undecided as to important points in the minute anatomy of the placenta, which further investigation will doubtless clear up. The main functions of the organ are, however, sufficiently clear. During the entire period of its existence it fills the important office of both stomach and lungs to the foetus. Whatever view of the arrangement of the maternal blood-vessels be taken, it is certain that the foetal blood is propelled by the pulsations of the foetal heart into the numberless villi of the chorion, where it is brought into very intimate relation with the mother's blood, gives off its carbonic acid, absorbs oxygen, and passes back to the foetus, through the umbilical vein, in a fit state for circulation. The mode of respiration, therefore, in the foetus is analogous to that in fishes, the chorion villi representing the gills, the maternal blood the water in which they float. Nutrition is also effected in the organ, and, by absorption through the chorion villi, the pabulum for the nourishment of the foetus is taken up. It also probably serves as an emunctory for the products of excretion in the foetus. Picard found that the blood in the placenta contained an appreciably larger quantity of urea than that in other parts of the body, this urea probably being derived from the foetus. Claude Bernard also attributed to it a glyco-genic function,² supposing it to take the place of the foetal liver until that organ was sufficiently developed.

Degenerative Changes Previous to Expulsion.—Finally, we find that

¹ *Obst. Trans.*, vol. xiv.

² *Acad. des Sciences*, April, 1859.

the temporary character of the placenta is indicated by certain degenerative changes which take place in it previous to expulsion. These consist chiefly in the deposit of calcareous patches, on its uterine surface, and in fatty degeneration of the villi and of the decidual layer between the placenta and the uterus. If this degeneration be carried to excess, as is not unfrequently the case, the foetus may perish from want of a sufficient number of healthy villi through which its respiration and nutrition may be effected.

Umbilical Cord.—The *umbilical* cord is the channel of communication between the foetus and placenta, being attached to the former at the umbilicus, to the latter generally near its centre, but sometimes, as in the battledore placenta, at its edge. It varies much in length, measuring on an average from 18 to 24 inches, but in exceptional cases being found as long as 50 or 60, and as short as 5 or 6, inches.

When fully formed it consists of an external membranous layer formed of the amnion, two umbilical arteries, one umbilical vein, and a considerable quantity of a transparent gelatinous substance surrounding the vessels, called Wharton's jelly, which is contained in a fine network of fibres, and is formed out of the tissue of the allantois. At an early period of pregnancy, in addition to these structures, the cord contains the pedicle of the umbilical vesicle, with the omphalo-mesenteric vessels ramifying on it, and two umbilical veins, one of which soon atrophies and disappears. No nerves or lymphatics have been satisfactorily demonstrated in the cord, although such have been described as existing. The vessels of the cord are at first straight in their course, but shortly they become greatly twisted, the arteries being external to the vein, and in nine cases out of ten the twist is from left to right. Various explanations have been given of this peculiarity, none of them entirely satisfactory. Tyler Smith attributed it to the movements of the foetus twisting the cord, its attachment to the placenta being a fixed point; this would not, however, account for the frequency with which the spiral turns occur in one direction. Mr. John Simpson attributed it to the greater pressure of the blood through the right hypogastric artery, on account of that vessel having a more direct relation to the aorta than the left. The umbilical arteries give off no branches, and the vein contains no valves, nor can any vasa vasorum be detected in their coats after they have left the umbilicus. The umbilical arteries increase in size after they leave the cord to divide on the surface of the placenta. This is the only example in the body in which arteries are larger near their terminations than their origin, and the object of this arrangement is probably to effect a retardation of the current of the blood distributed to the placenta. The tortuous course of the vein probably compensates for the absence of valves, and moderates the flow of blood through it. Distinct knots are not unfrequently observed in the cord, but they rarely have the effect of obstructing the circulation through it. They no doubt form when the foetus is very small. They may sometimes also be produced in labor by the child being propelled through a coil of the cord lying circularly round the os uteri. The so-called false knots are merely accidental nodosities due to local enlargements of the vessels.

CHAPTER II.

THE ANATOMY AND PHYSIOLOGY OF THE FÆTUS.

It is obviously impossible to attempt anything like a full account of the development of the various fœtal structures, or of their growth, during intra-uterine life. To do so would lead us far beyond the scope of this work, and would involve a study of complex details only suitable in a treatise on embryology. It is of importance, however, that the practitioner should have it in his power to determine approximately the age of the fœtus in abortions or premature labors, and for this purpose it is necessary to describe briefly the appearance of the fœtus at various stages of its growth.

Appearance of the Fœtus at Various Stages of Development.—1st Month: The fœtus in the first month of gestation is a minute gelatinous and semi-transparent mass, of a grayish color, in which no definite structure can be made out, and in which no head or extremities can be seen. It is rarely to be detected in abortions, being lost in surrounding blood-clots. In the few examples which have been carefully examined it did not measure more than a line in length. It is, however, already surrounded by the amnion, and the pedicle of the umbilical vesicle can be traced into the unclosed abdominal cavity.

2d Month: The embryo becomes more distinctly apparent, and is curved on itself, weighing about 62 grains, and measuring 6 to 8 lines in length. The head and extremities are distinctly visible, the latter in the form of rudimentary projections from the body. The eyes are to be seen as small black spots on the side of the head. The spinal column is divided into separate vertebrae. The independent circulatory system of the fœtus is now beginning to form, the heart consisting of only one ventricle and one auricle, from the former of which both the aorta and pulmonary arteries arise. On either side of the vertebral column, reaching from the heart to the pelvis, are two large glandular structures, the *corpora Wolffiana*, which consist of a series of convoluted tubes opening into an excretory duct running along their external borders, and connected below with the common cloaca of the genito-urinary and digestive tracts. They seem to act as secreting glands, and fulfil the functions of the kidneys before they are formed. Toward the end of the second month they atrophy and disappear, and the only trace of them in the fœtus at term is to be found in the parovarium lying between the folds of the broad ligaments. At this stage of development there are met with in the human embryo, as in that of all mammals, four transverse fissures opening into the pharynx, which are analogous to the permanent branchiae of fishes. Their vascular supply is also similar, as the aorta at this time gives off four branches on each side, each of which forms a branchial arch, and these afterward unite to form the descending aorta. By the end of the sixth week these, as well as the transverse

fissures to which they are distributed, disappear. By the end of the second month the kidneys and supra-renal capsules are forming, and the single ventricle is divided into two by the growth of the inter-ventricular septum. The umbilical cord is quite straight, and is inserted into the lower part of the abdomen. Centres of ossification are showing themselves in the inferior maxillary bones and the clavicle.

3d Month: The embryo weighs from 70 to 300 grains, and measures from $2\frac{1}{2}$ to $3\frac{1}{2}$ inches in length. The forearm is well formed, and the first traces of the fingers can be made out. The head is large in proportion to the rest of the body, and the eyes are prominent. The umbilical vesicle and allantois have disappeared, the greater portion of the chorion villi have atrophied, and the placenta is distinctly formed.

4th Month: The weight is from 4 to 6 oz., and the length about 6 inches. The convolutions of the brain are beginning to develop. The sex of the child can now be ascertained on inspection. The muscles are sufficiently formed to produce distinct movements of the limbs. Ossification is extending, and can be traced in the occipital and frontal bones and in the mastoid processes. The sexual organs are differentiated.

5th Month: Weight, about 10 oz.; length, 9 or 10 inches. Hair is observed covering the head, which forms about one-third of the length of the whole fœtus. The nails are beginning to form, and ossification has commenced in the ischium.

6th Month: Weight, about 1 lb.; length, 11 to $12\frac{1}{2}$ inches. The hair is darker. The eyelids are closed, and the membrana pupillaris exists; eyelashes have now been formed. Some fat is deposited under the skin. The testicles are still in the abdominal cavity. The clitoris is prominent. The pubic bones have begun to ossify.

7th Month: Weight, from 3 to 4 lbs.; length, 13 to 15 inches. The skin is covered with unctuous, sebaceous matter, and there is a more considerable deposit of subcutaneous fat. The eyelids are open. The testicles have descended into the scrotum.

8th Month: Weight, from 4 to 5 lbs.; length, 16 to 18 inches, and the fœtus seems now to grow in thickness rather than in length. The nails are completely developed. The membrana pupillaris has disappeared.

Fœtus at Term.—At the completion of pregnancy the fœtus weighs on an average $6\frac{1}{2}$ lbs. and measures about 20 inches in length. These averages are, however, liable to great variation. Remarkable histories are given by many writers of fœtuses of extraordinary weight, which have been probably greatly exaggerated. Out of 3000 children delivered under the care of Cazeaux at various charities, one only weighed 10 lbs. There are, however, several carefully recorded instances of weight far exceeding this, but they are undoubtedly much more uncommon than is generally supposed. Dr. Ramsbottom mentions a fœtus weighing $16\frac{1}{2}$ lbs.; Cazeaux tells us of one which he delivered by turning which weighed 18 lbs. and measured 2 feet $1\frac{1}{2}$ inches, and the birth of one weighing 21 lbs. has been recently recorded.¹ Such overgrown children are almost invariably stillborn.^[1]

¹ *Brit. Med. Journ.*, Feb. 1, 1879.

[² Probably the largest fœtus on record was that of Mrs. Captain Bates, the Nova

The average size of male children at birth, as in after-life, is somewhat greater than that of female. Thus, Simpson¹ found that out of 100 cases the male children averaged 10 oz. more in weight than the female, and half an inch more in length.

Vernix Caseosa.—A new-born child at term is generally covered to a greater or less extent with a greasy, unctuous material, the *vernix caseosa*, which is formed of epithelial scales and the secretion of the sebaceous glands, and which is said to be of use in labor by lubricating the surface of the child. The head is generally covered with long dark hair, which frequently falls off or changes in color shortly after birth. Dr. Wiltshire² has called attention to an old observation, that the eyes of all new-born children are of a peculiar dark steel-gray color, and that they do not acquire their permanent tint until some time after birth. The umbilical cord is generally inserted below the centre of the body.

Anatomy of the Fœtal Head.—The most important part of the fœtus from an obstetrical point of view is the head, which requires a separate study, as it is the usual presenting part, and the facility of the labor depends on its accurate adaptation to the maternal passages.

The chief anatomical peculiarity of interest in the head of the fœtus at term is that the bones of the skull, especially of its vertex—which, in the vast majority of cases, has to pass first through the pelvis—are not firmly ossified as in adult life, but are joined loosely together by membrane or cartilage. The result of this is, that the skull is capable of being moulded and altered in form to a very considerable extent by the pressure to which it is subjected, and thus its passage through the pelvis is very greatly facilitated. This, however, is chiefly the case with the cranium proper, the bones of the face and of the base of the skull being more firmly united. By this means the delicate structures at the base of the brain are protected from pressure, while the change of form which the skull undergoes during labor implicates a portion of the skull where pressure on the cranial contents is least likely to be injurious.

The divisions between the bones of the cranium are further of obstetric importance in enabling us to detect the precise position of the head during labor, and an accurate knowledge of them is therefore essential to the obstetrician.

The Sutures and Fontanelles.—We talk of them as *sutures* and *fontanelles*, the former being the lines of junction between the separate bones, which overlap each other to a greater or less extent during labor; the latter, membranous interspaces where the sutures join each other.

The principal sutures are: 1st, the *sagittal*, which separates the two parietal bones, and extends longitudinally backward along the vertex

Scotia giantess, a woman of 7 ft. 9 in., whose husband is also of gigantic build, reaching 7 ft. 7 in. in height. This child, born in Ohio, was their second, and was lost in its birth, as no forceps could be procured of sufficient size to grasp the head. The fœtus weighed 23½ lbs. and was 30 in. in length. Their first infant weighed 18 lbs. Dr. George Eddowes of Crewe, England, delivered a woman, on Nov. 12, 1884, of a male child weighing 20 lbs. 2 oz. It matured 23 inches in length and 14½ inches around the chest (*Lancet*, Nov. 22, 1884, p. 941). We have had children born in this city (Philadelphia) at maturity and live that weighed but one pound. The well-remembered "Pincus baby" weighed a pound and an ounce.—Ed.]

¹ *Selected Obst. Works*, p. 327.

² *Lancet*, February 11, 1871.

of the head ; 2d, the *frontal*, which is a continuation of the sagittal, and divides the two halves of the frontal bone, at this time separate from each other ; 3d, the *coronal*, which separates the frontal from the parietal bones, and extends from the squamous portion of the temporal bone across the head to a corresponding point on the opposite side ; and 4th, the *lambdoidal*, which receives its name from its resemblance to the Greek letter Λ , and separates the occipital from the parietal bones on either side. The fontanelles (Fig. 64) are the membranous interspaces where the sutures join—the *anterior* and larger being lozenge-shaped, and formed by the junction of the frontal, sagittal, and two halves of the coronal sutures. It will be well to note that there are, therefore, four lines of sutures running into it, and four angles, of which the anterior, formed by the frontal suture, is most elongated and well marked. The *posterior* fontanelle (Fig. 65) is formed by the junction of the sagittal suture with the two legs of the lambdoidal. It is therefore triangular in shape, with three lines of suture entering it in three angles, and is

FIG. 64.


Anterior and Posterior
Fontanelles.

FIG. 65.


Bi-parietal Diameter. Sagittal and Lambdoidal
Sutures, with Posterior Fontanelle.

much smaller than the anterior fontanelle, forming merely a depression into which the tip of the finger can be placed, while the latter is a hollow as big as a shilling, or even larger. As it is the posterior fontanelle which is generally lowest, and the one most commonly felt during labor, it is important for the student to familiarize himself with it, and he should lose no opportunity of studying the sensations imparted to the finger by the sutures and fontanelles in the head of the child after birth.

The Diameters of the Fœtal Skull.—For the purpose of understanding the mechanism of labor we must study the measurements of the fetal head in relation to the cavity through which it has to pass. They are taken from corresponding points opposite to each other, and are known as the diameters of the skull (Fig. 66). Those of most importance are : 1st. The *occipito-mental*, from the occipital protuberance to the point of the chin, 5.25'' to 5.50''. 2d. The *occipito-frontal*, from the occiput to the centre of the forehead, 4.50'' to 5''. 3d. The *sub-occipito-breg-*

matic, from a point midway between the occipital protuberance and the margin of the foramen magnum to the centre of the anterior fontanelle, 3.25". 4th. The *cervico-bregmatic*, from the anterior margin of the

FIG. 66.



- 1 and 2. Occipito-frontal diameter.
3 and 4. Occipito-mental.
5 and 6. Cervico-bregmatic.
7 and 8. Fronto-mental.

foramen magnum to the centre of the anterior fontanelle, 3.75". 5th. *Transverse*, or *bi-parietal*, between the parietal protuberances, 3.75" to 4". 6th. *Bi-temporal*, between the ears, 3.50". 7th. *Fronto-mental*, from the apex of the forehead to the chin, 3.25".

Alteration of Diameters by Compression and Moulding during Labor.—The length of these respective diameters, as given by different writers, differs considerably, a fact to be explained by the measurements having been taken at different times—by some just after birth, when the head was altered in shape by the

moulding it had undergone; by others when this had either been slight or after the head had recovered its normal shape. The above measurements may be taken as the average of those of the normally shaped head, and it is to be noted that the first two are most apt to be modified during labor. The amount of compression and moulding to which the head may be subjected without proving fatal to the fœtus is not certainly known, but it is doubtless very considerable. Some interesting examples of the extent to which the head may be altered in shape in difficult labors have been given by Barnes,¹ who has shown by tracings of the shape of the head taken immediately after delivery that in protracted labor the occipito-mental and occipito-frontal diameters may be increased more than an inch in length, while lateral compression may diminish the bi-parietal diameter to the same length as the inter-auricular. The fœtal head is movable on the vertical column to the extent of a quarter of a circle; and it seems probable that the laxity of the ligaments admits with impunity a greater circular movement than would be possible in the adult.

Influence of Sex and Race on the Fœtal Head.—On taking the average of a large number of measurements, it is found that the heads of male children are larger and more firmly ossified than those of females, the former averaging about half an inch more in circumference. Sir James Simpson attributed great importance to this fact, and believed that it was sufficient to account for the larger proportion of stillbirths in male than in female children, as well as for the greater difficulty of labor and the increased maternal mortality that are found to attend on male births. His well-known paper on this subject, which has given rise to much controversy, is full of the most elaborate details; and so great did he believe the fœtal influence to be that he calculated that between the years 1834 and 1837 there were lost in Great Britain, as

¹ *Obst. Trans.*, vol. vii.

a consequence of the slightly larger size of the male than of the female head at birth, about 50,000 lives, including those of about 46,000 or 47,000 infants, and of between 3000 and 4000 mothers who died in childbed.¹ It is probable that race and other conditions, such as civilization and intellectual culture, have considerable influence on the size of the foetal skull, but we are not in possession of sufficiently accurate data to justify any very positive opinion on these points.

Position of the Fœtus in Utero.—In the very large majority of cases the foetus lies in utero with head downward, and is so placed as to be adapted in the most convenient way to the cavity in which it is placed. The uterine cavity is most roomy at the fundus, and narrowest at the cervix, and the greatest bulk of the foetus is at the breech, so that the largest part of the child usually lies in the part of the uterus best adapted to contain it. The various parts of the child's body are further so placed, in regard to each other, as to take up the least possible amount of space. (See frontispiece.) The body is bent so that the spine is curved with its convexity outward, this curvature existing from the earliest period of development; the chin is flexed on the sternum; the forearms are flexed on the arms, and lie close together on the front of the chest; the legs are flexed on the thighs, and the thighs drawn up on the abdomen; the feet are drawn up toward the legs; the umbilical cord is generally placed out of reach of injurious pressure, in the space between the arms and the thighs. Variations from this attitude, however, are not uncommon, and are not, as a rule, of much consequence. Although the cranial presentations are much the most common, averaging 96 out of every 100 cases, other presentations are by no means rare, the next most frequent being either that of the breech, in which the long diameter of the child lies in the long diameter of the uterine cavity, or some variety of transverse presentation, in which the long diameter of the foetus lies obliquely across the uterus, and no longer corresponds to its longitudinal axis.

Changes of Fœtal Position during Pregnancy.—It was long believed that the head presentation was only assumed toward the end of pregnancy, when it was supposed to be produced by a sudden movement on the part of the foetus, known as the *culbute*. It is now well known that, in the large majority of cases, the head is lowest during all the latter part of pregnancy, although changes in position are more common than is generally believed to be the case, and presentation of parts other than the head is much more frequent in premature labor than in delivery at term. In evidence of the last statement, Churchill says that in labor at the seventh month the head presents only 83 times out of 100 when the child is living, and that as many as 53 per cent. of the presentations are preternatural when the child is stillborn. The frequency with which the foetus changes its position before delivery has been made the subject of investigation by various German obstetricians, and the fact can be readily ascertained by examination. Valenta² found that out of nearly 1000 cases, carefully and frequently examined by him, in 57.6 per cent. the presentation underwent no change in the latter months of pregnancy, but in the remaining 42.4 per cent. a change


¹ *Selected Obst. Works*, p. 363.

² *Mon. f. Geburt.*, 1866.

could be readily detected. These alterations were found to be most frequent in multiparæ, and the tendency was for abnormal presentations to alter into normal ones. Thus it was common for transverse presentations to alter longitudinally, and but rare for breech presentations to change into head. The ease with which these changes are effected no doubt depends, in a considerable degree, on the laxity of the uterine parietes and on the greater quantity of amniotic fluid, by both of which the free mobility of the fœtus is favored.

Detection of Fœtal Position by Abdominal Palpation.—The facility with which the position of the fœtus in utero can be ascertained by abdominal palpation has not been generally appreciated in obstetric works, and yet by a little practice it is easy to make it out. Much information of importance can be gained in this way, and it is quite possible, under favorable circumstances, to alter abnormal presentations before labor has begun. For the purpose of making this examination

FIG. 67.



Mode of Ascertaining the Position of the Fœtus by Palpation.

the patient should lie at the edge of the bed, with her shoulders slightly raised and the abdomen uncovered. The first observation to make is to see if the longitudinal axis of the uterine tumor corresponds with that of the mother's abdomen; if it does, the presentation must be either a head or a breech. By spreading the hands over the uterus (Fig. 67) a greater sense of resistance can be felt, in most cases, on one side than on the other, corresponding to the back of the child. By striking the tips of the fingers suddenly inward at the fundus, the hard breech can generally be made out, or the head still more easily if the breech be downward. When the uterine walls are unusually lax, it is often possible to feel the limbs of the child. These observations can be generally corroborated by auscultation, for in head presentations the fœtal heart can usually be heard below the umbilicus, and in breech cases above it. Transverse presentations can even more easily be made

out by abdominal palpation. Here the long axis of the uterine tumor does not correspond with the long axis of the mother's abdomen, but lies obliquely across it. By palpation the rounded mass of the head can be easily felt in one of the mother's flanks, and the breech in the other, while the foetal heart is heard pulsating nearer to the side at which the head is detected.

Explanation of the Position of the Fœtus in Utero.—The reason why the head presents so frequently has been made the subject of much discussion. The oldest theory was, that the head lay over the os uteri as the result of gravitation; and the influence of gravity, although contested by many obstetricians, prominent among whom were Dubois and Simpson, has been insisted upon as the chief cause by others, Dr. Duncan being one of the most strenuous advocates of this view. The objections urged against the gravitation theory were drawn partly from the result of experiments, and partly from the frequency with which abnormal presentations occur in premature labors, when the action of gravity cannot be supposed to be suspended. The experiments made by Dubois went to show that when the fœtus was suspended in water gravitation caused the shoulders, and not the head, to fall lowest. He therefore advanced the hypothesis that the position of the fœtus was due to instinctive movements which it made to adapt itself to the most comfortable position in which it could lie. It need only be remarked that there is not the slightest evidence of the fœtus possessing any such power. Simpson proposed a theory which was much more plausible. He assumed that the foetal position was due to reflex movements produced by physical irritations to which the cutaneous surface of the fœtus is subjected from changes of the mother's position, uterine contractions, and the like. The absence of these movements in the case of the death of the fœtus would readily explain the frequency of mal-presentations under such circumstances. The obvious objection to this theory, complete as it seems to be, is the absence of any proof that such constant extensive reflex movements really do occur in utero. Dr. Duncan has very conclusively disposed of the principal objections which have been raised against the influence of gravitation, and when an obvious explanation of so simple a kind exists it seems useless to seek farther for another. He has shown that Dubois's experiments did not accurately represent the state of the fœtus in utero, and that during the greater part of the day, when the woman is upright or lying on her back, the fœtus lies obliquely to the horizon at an angle of about 30°. The child thus lies, in the former case, on an inclined plane formed by the anterior uterine wall and by the abdominal parietes; in the latter, by the posterior uterine wall and the vertebral column. Down the inclined plane so formed the force of gravity causes the fœtus to slide, and it is only when the woman lies on her side that the fœtus is placed horizontally, and is not subjected in the same degree to the action of gravity (Fig. 68). The frequency of mal-presentations in premature labors is explained by Dr. Duncan partly by the fact that the death of the child (which so frequently precedes such cases) alters its centre of gravity, and partly by the greater mobility of the child and the greater relative amount of liquor amnii (Fig. 69). The influence of gravitation is probably greatly

assisted by the contractions of the uterus which are going on during the greater part of pregnancy. The influence of these was pointed out by Dr. Tyler Smith, who distinctly showed that the contractions of the uterus preceding delivery exerted a moulding or adapting influence on

FIG. 68.





Diagram illustrating the effect of Gravity on the Fœtus. (After Duncan.)

a, b, is parallel to the axis of the pregnant uterus and pelvic brim; *c, d, e,* is a perpendicular line; *e,* the centre of gravity of the fœtus; *d,* the centre of flotation.

the fœtus, and prevented undue alterations of its position. Dr. Hicks proved¹ that these uterine contractions are of constant occurrence from the earliest period of pregnancy, and there can be little doubt that they must have an important influence on the body contained within the

FIG. 69.



Illustrating the greater Mobility of the Fœtus and the larger relative amount of Liquor Amnii in Early Pregnancy. (After Duncan.)

a, b, Axis of pregnant uterus.

b, h. A horizontal line.

uterus. The whole subject has been recently considered by Pinard,² who shows that many factors are in action to produce and maintain the usual position of the fœtus in utero, which may be either of an active or a passive character—the former being chiefly the active movements

¹ *Obst. Trans.*, vol. xiii. p. 216.

² *Annal. de Gyn.*, May and July, 1878.

of the fœtus and the contractions of the uterus and the abdominal muscles; the latter, the form of the uterus and the fœtus, the slippery surface of the amnion, pressure of the amniotic fluid, etc. When any of these factors are at fault, mal-presentation is apt to occur.

Functions of the Fœtus.—The functions of the fœtus are in the main the same, with differences depending on the situation in which it is placed, as those of the separate being. It breathes, it is nourished, it forms secretions, and its nervous system acts. The mode in which some of these functions are carried on in intra-uterine life requires separate consideration.

Nutrition.—During the early part of pregnancy, and before the formation of the umbilical vesicle and the allantois, it is certain that nutritive material must be supplied to the ovum by endosmosis through its external envelope. The precise source, however, from which this is obtained is not positively known. By some it is believed to be derived from the granulations of the discus proligerus which surround it as it escapes from the Graafian follicle, and subsequently from the layer of albuminous matter which surrounds the ovum before it reaches the uterus; while others think it probable that it may come from a special liquid secreted by the interior of the Fallopian tube as the ovum passes along it. As soon as the ovum has reached the uterus there is every reason to believe that the umbilical vesicle is the chief source of nourishment to the embryo, through the channel of the omphalo-mesenteric vessels, which convey matters absorbed from the interior of the vesicle to the intestinal canal of the fœtus. At this time the exterior of the ovum is covered by the numerous fine villousities of the primitive chorion which are imbedded in the mucous membrane of the uterus, and it is thought that they may absorb materials from the maternal system, which may be either directly absorbed by the embryo, or which may serve the purpose of replacing the nutritive matter which has been removed from the umbilical vesicle by the omphalo-mesenteric vessels. This point it is, of course, impossible to decide. Joulin, however, thinks that these villi probably have no direct influence on the nourishment of the fœtus, which is at this time solely effected by the umbilical vesicle, but that they absorb fluid from the maternal system, which passes through the amnion and forms the liquor amnii. As soon as the allantois is developed, vascular communication between the fœtus and the maternal structures is established, and the temporary function of the umbilical vesicle is over: that structure, therefore, rapidly atrophies and disappears, and the nutrition of the fœtus is now solely carried on by means of the chorion villi, lined as they now are by the vascular endo-chorion, and chiefly by those which go to form the substance of the placenta.

This statement is opposed to the views of many physiologists, who believe that a certain amount of nutritive material is conveyed to the fœtus through the channel of the liquor amnii, itself derived from the maternal system, which is supposed either to be absorbed through the cutaneous surface of the fœtus or carried to the intestinal canal by deglutition. The reasons for assigning to the liquor a nutritive function are, however, so slight that it is difficult to believe that it has any appreciable action in this way. They are based on some questionable observations,

such as those of Weydlich, who kept a calf alive for fifteen days by feeding it solely on liquor amnii, and the experiments of Burdach, who found the cutaneous lymphatics engorged in a fœtus removed from the amniotic cavity, while those of the intestine were empty. The deglutition of the liquor amnii for the purposes of nutrition has been assumed from its occasional detection in the stomach of the fœtus, the presence of which may, however, be readily explained by spasmodic efforts at respiration which the fœtus undoubtedly often makes before birth, especially when the placental circulation is in any way interfered with, and during which a certain quantity of fluid would necessarily be swallowed. The quantity of nutritive material, however, in the liquor amnii is so small—not more than 6 to 9 parts of albumen in 1000—that it is impossible to conceive how it could have any appreciable influence in nutrition, even if its absorption, either by the skin or stomach, were susceptible of proof.

That the nutrition of the fœtus is effected through the placenta is proved by the common observation that whenever the placental circulation is arrested, as by disease of its structure, the fœtus atrophies and dies. The precise mode, however, in which nutritive materials are absorbed from the maternal blood is still a matter of doubt, and must remain so until the mooted points as to the minute anatomy of the placenta are settled. The various theories entertained on this subject by the upholders of the Hunterian doctrine of placental anatomy, and by those who deny the existence of a sinus system, have already been referred to in the chapter on the Anatomy of the Placenta, to which the reader is referred (pp. 111–115).

Respiration.—One of the chief functions of the placenta, besides that of nutrition, is the supply of oxygenated blood to the fœtus. That this is essential to the vitality of the fœtus, and that the placenta is the site of oxygenation, is shown by the fact that whenever the placenta is separated, or the access of fœtal blood to it arrested by compression of the cord, instinctive attempts at inspiration are made, and if aerial respiration cannot be performed the fœtus is expelled asphyxiated. Like the other functions of the fœtus during intra-uterine life, that of respiration has been made the subject of numerous more or less ingenious hypotheses. Thus, many have believed that the fœtus absorbed gaseous material from the liquor amnii, which served the purpose of oxygenating its blood, St. Hilaire thinking that this was effected by minute openings in its skin, Beclard and others through the bronchi, to which they believed the liquor amnii gained access. Independently of the entire want of evidence of the absorption of gaseous materials by these channels, the theory is disproved by the fact that the liquor amnii contains no air which is capable of respiration. Serres attributed a similar function to some of the chorion villi, which he believed penetrated the utricular glands of the decidua reflexa and absorbed gas from the hydroperione, or fluid situated between it and the decidua vera, and in this manner he thought the fœtal blood was oxygenated until the fifth month of intra-uterine life, when the placenta was fully formed.

This hypothesis, however, rests on no accurate foundation, for it is certain that the chorion villi do not penetrate the utricular glands in the manner assumed; or, even if they did, the mode in which the oxygen

thus absorbed by the chorion villi reaches the fœtus, which is separated from them by the amnion and its contents, would still remain unexplained.

The mode in which the oxygenation of the fœtal blood is effected before the formation of the placenta remains, therefore, as yet unknown. After the development of that organ, however, it is less difficult to understand, for the fœtal blood is everywhere brought into such close contact with the maternal, in the numerous minute ramifications of the umbilical vessels, that the interchange of gases can readily be effected. The activity of respiration is doubtless much less than in extra-uterine life, for the waste of tissue in the fœtus is necessarily comparatively small, from the fact of its being suspended in a fluid medium of its own temperature, and from the absence of the processes of digestion and of respiratory movements. The quantity of carbonic acid formed would, therefore, be much less than after birth, and there would be a correspondingly small call for oxygenation of venous circulation.

Circulation.—The functions of the lungs being in abeyance, it is necessary that all the fœtal blood should be carried to the placenta to receive oxygen and nutritive materials. To understand the mode in which this is effected we must bear in mind certain peculiarities in the circulatory system which disappear after birth.

1. The two sides of the fœtal heart are not separate, as in the adult. The right ventricle in the adult sends all the venous blood to the lungs through the pulmonary arteries, to be aërated by contact with the atmosphere. In the fœtus, however, only sufficient blood is passed through the pulmonary arteries to ensure their being pervious and ready to carry blood to the lungs immediately after birth.

An aperture of communication, the *foramen ovale*, exists between the two auricles, which is arranged so as to permit the blood reaching the right auricle to pass freely into the left, but not vice versâ. By this means a large portion of the blood reaching the heart through the *venæ cavæ*, instead of passing, as in the adult, into the right ventricle, is directed into the left auricle.

2. Even with this arrangement, however, a larger portion of blood would pass into the pulmonary arteries than is required for transmission to the lungs, and a further provision is made to prevent its going to them by means of a fœtal vessel, the *ductus arteriosus* (Fig. 70), which arises from the point of bifurcation of the pulmonary arteries and opens into the arch of the aorta. In consequence of this arrangement only a very small portion of the blood reaches the lungs at all.

3. The fœtal hypogastric arteries are continued into large arterial trunks, which, passing into the cord, form the *umbilical arteries*, and carry the impure fœtal blood into the placenta.

4. The purified blood is collected into the single *umbilical vein*, through which it is carried to the under surface of the liver, from which point it is conducted, by means of another special fœtal vessel, the *ductus venosus*, into the ascending vena cava and the right auricle.




Diagram of Fœtal Heart.
(After Dalton.)

- 1. Aorta.
- 2. Pulmonary artery.
- 3, 3. Pulmonary branches.
- 4. Ductus arteriosus.

Course of the Fœtal Circulation.—In order to understand the course of the fœtal blood, it may be most conveniently traced from the point where it reaches the under surface of the liver through the umbilical vein. Part of it is distributed to the liver itself, but the greater quantity is carried directly into the inferior vena cava through the ductus venosus. The inferior vena cava also receives the blood from the fœtal veins of the lower extremities and that portion of the blood of the umbilical vein which has passed through the liver. This mixed blood is carried up to the right auricle, from which by far the greater part of it is immediately directed into the left auricle through the foramen ovale. From thence it passes into the left ventricle, which sends the greater part of it into the head and upper extremities through the aorta, a comparatively small quantity being transmitted to the inferior extremities. The blood which is thus sent to the upper part of the body is collected into the vena cava superior, by which it is thrown into the right auricle. Here the mass of it is probably directed into the right ventricle, which expels it into the pulmonary arteries, and from thence, through the ductus arteriosus, into the descending aorta. By this arrangement it will be seen that the descending aorta conveys to the lower part of the body the comparatively impure blood which has already circulated through the head, neck and upper extremities. From the descending aorta a small quantity of blood is conveyed to the lower extremities, the greater part of it being carried for purification to the placenta through the umbilical arteries.

Establishment of Independent Circulation.—As soon as the child is born it generally cries loudly and inflates its lungs, and, in consequence, the pulmonary arteries are dilated, and the greater portion of the blood of the right ventricle is at once sent to the lungs, from whence, after being arterialized, it is returned to the left auricle through the pulmonary veins. The left auricle, therefore, receives more blood than before, the right less, and, the placental circulation being arrested, no more passes through the umbilical vein. In consequence of this, the pressure of the blood in the two auricles is equalized; the mass of the blood in the right auricle no longer passes into the left (the valve of the foramen ovale being closed by the equal pressure on both sides), but directly into the right ventricle, and from thence into the pulmonary arteries, and the ductus arteriosus soon collapses and becomes impervious. The mass of blood in the descending aorta no longer finds its way into the hypogastric arteries, but passes into the lower extremities, and the adult circulation is established.

Changes in Fœtal Circulation after Birth.—The changes which take place in the temporary vascular arrangements of the fœtus prior to their complete disappearance are of some practical interest. The ductus arteriosus, as has been said, collapses, chiefly because the mass of blood is drawn to the lungs, and partly, perhaps, by its own inherent contractility. Its walls are found to be thickened, and its canal closes, first in the centre, and subsequently at its extremities, its aortic end remaining longer pervious on account of the greater pressure of blood from the left side of the heart (Fig. 71). Practical closure occurs within a few days after birth, although Flourens states that it is not completely oblit-

erated until eighteen months or two years have elapsed.¹ According to Schroeder, its walls unite without the formation of any thrombus. The foramen ovale is soon closed by its valve, which contracts adhesion with the edges of the aperture, so as effectually to occlude it. Sometimes, however, a small canal of communication between the two auricles may remain pervious for many months, or even a year and more, without, however, any admixture of blood occurring. A permanently patulous condition of this aperture, however, sometimes exists, giving rise to the disease known as cyanosis.


The umbilical arteries and veins and the ductus venosus soon also become impermeable, in consequence of concentric hypertrophy of their tissue and collapse of their walls. The closure of the former is aided by the formation of coagula in the interior. According to Robin, a longer time than is usually supposed elapses before they become completely closed, the vein remaining pervious until the twentieth or thirtieth day after delivery, the arteries for a month or six weeks. He has also described² a remarkable contraction of the umbilical vessels within their sheaths at the point where they leave the abdominal walls, which takes place within three or four days after birth, and seems to prevent hemorrhage taking place when the cord is detached.

Function of the Liver.—The liver, from its proportionately large size, apparently plays an important part in the foetal economy. It is not until about the fifth month of utero-gestation that it assumes its characteristic structure, and forms bile, previous to that time its texture being soft and undeveloped. According to Claude Bernard, after this period one of its most important offices is the formation of sugar, which is found in much larger amount in the foetus than after birth. Sugar is, however, found in the foetal structures long before the development of the liver, especially in the mucous and cutaneous tissues, and it seems probable that these, as well as the placenta itself, then fulfil the glyco-genic function afterward chiefly performed by the liver. The bile is secreted after the fifth month of pregnancy, and passes into the intestinal canal, and is subsequently collected in the gall-bladder. By some physiologists it has been supposed that the liver during intra-uterine life was the chief seat of depuration of the carbonic acid contained in the venous blood of the foetus. It is, however, more generally believed that this is accomplished solely in the placenta.

The Meconium.—The bile, mixed with the mucous secretion of the intestinal tract, forms the *meconium*, which is contained in the intestines of the foetus, and which collects in them during the whole period of intra-uterine life. It is a thick, tenacious, greenish substance, which is voided soon after birth in considerable quantity.

The Urine.—Urine is certainly formed during intra-uterine life, as is

FIG. 71.

Diagram of Heart of Infant.
(After Dalton.)

1. Aorta. 2. Pulmonary artery.
3. 3. Pulmonary branches.
4. Ductus arteriosus becoming obliterated.

¹ *Acad. des Sciences*, 1854.² *Ibid.*, 1860.

proved by the fact, familiar to all accoucheurs, that the bladder is constantly emptied instantly after birth. It has generally been supposed that the fœtus voids its urine into the cavity of the amnion, and the existence of traces of urea in the liquor amnii, as well as some cases of imperforate urethra, in which the bladder was found to be enormously distended, and some cases of congenital hydronephrosis associated with impervious ureters, have been supposed to corroborate this assumption. The question has been very fully studied by Joulin, who has collected together a large number of instances in which there was imperforate urethra without any undue distension of the bladder. He holds, also, that the amount of urea found in the liquor amnii is far too minute to justify the conclusion that the urine of the fœtus was habitually passed into it, although a small quantity may, he thinks, escape into it from time to time; and he therefore believes that the urine of the fœtus is only secreted regularly and abundantly after birth, and that during intra-uterine life its retention is not likely to give rise to any functional disturbance.¹

Function of the Nervous System.—There is no doubt that the nervous system acts to a considerable extent during intra-uterine life, and some authors have even supposed that the fœtus was endowed with the power of making instinctive or voluntary movements for the purpose of adapting itself to the form of the uterine cavity. Most probably, however, the movements the fœtus performs are purely reflex. That it responds to a stimulus applied to the cutaneous nerves is proved by the experiments of Tyler Smith, who laid bare the amnion in pregnant rabbits, and found that the fœtus moved its limbs when these were irritated through it. Pressure on the mother's abdomen, cold applications, and similar stimuli will also produce energetic fetal movements. The gray matter of the brain in the new-born child is, however, quite rudimentary in its structure, and there is no evidence of intelligent action of the nervous system until some time after birth, and, *a fortiori*, during pregnancy.

CHAPTER III.

PREGNANCY.

As soon as conception has taken place a series of remarkable changes commence in the uterus, which progress until the termination of pregnancy, and are well worthy of careful study. They produce those marvellous modifications which effect the transformation of the small undeveloped uterus of the non-pregnant state into the large and fully-developed uterus of pregnancy, and have no parallel in the whole animal economy.

A knowledge of them is essential for the proper comprehension of the

¹ *Acad. des Sciences*, p. 308.

phenomena of labor and for the diagnosis of pregnancy which the practitioner is so frequently called upon to make. Excluding the varieties of abnormal pregnancy, which will be noticed in another place, we shall here limit ourselves to the consideration of the modifications of the maternal organism which result from simple and natural gestation.

Changes in the Uterus.—The unimpregnated uterus measures $2\frac{1}{2}$ inches in length and weighs about 1 oz., while at the full term of pregnancy it has so immensely grown as to weigh 24 oz. and measure 12 inches. The growth commences as soon as the ovum reaches the uterus, and continues

FIG. 72.



Relations of the Pregnant Uterus at Sixth Month to the Surrounding Parts. (After Martin.)


uninterruptedly until delivery. In the early months the uterus is contained entirely in the cavity of the pelvis, and the increase of size is only apparent on vaginal examination, and that with difficulty. Before the third month the enlargement is chiefly in the lateral direction, so that the whole body of the uterus assumes more of a spherical shape than in the non-pregnant state. If an opportunity of examining the gravid uterus *post-mortem* should occur at this time, it will be found to have the form of a sphere flattened somewhat posteriorly and bulging anteriorly.

After the ascent of the organ into the abdomen it develops more in the vertical direction, so that at term it has the form of an ovoid, with

its large extremity above and its narrow end at the cervix uteri, and its longitudinal axis corresponds to the long diameter of the mother's abdomen, provided the presentation be either of the head or the breech. The anterior surface is now even more distinctly projecting than before—a fact which is explained by the proximity of the posterior surface to the rigid spinal column behind, while the anterior is in relation with the lax abdominal parietes, which yield readily to pressure, and so allow of the more marked prominence of the anterior uterine wall.

Change in Situation.—Before the gravid uterus has risen out of the pelvis no appreciable increase in the size of the abdomen is perceptible. On the contrary, it is an old observation that at this early stage of pregnancy the abdomen is flatter than usual, on account of the partial descent of the uterus in the pelvic cavity as a result of its increased weight. As the growth of the organ advances it soon becomes too large to be contained any longer within the pelvis, and about the middle of the third or the beginning of the fourth month the fundus rises above the pelvic brim—not suddenly, as is often erroneously thought, but slowly and gradually—when it may be felt as a smooth rounded swelling.

FIG. 73.



Size of Uterus at Various Periods of Pregnancy.

Size of Uterine Tumor at Various Periods of Pregnancy.—It is about this time that the movements of the fœtus first become appreciable to the mother, when “*quickening*” is said to have taken place. Toward the end of the fourth month the uterus reaches to about three fingers’ breadth above the symphysis pubis. About the fifth month it occupies the hypogastric region, to which it imparts a marked projection, and the alteration in the figure is now distinctly perceptible to visual examination. About the sixth month it is on a level with, or a little above, the umbilicus. About the seventh month it is about two inches above the umbilicus, which is now projecting and prominent, instead of depressed, as in the non-pregnant state. During the eighth and ninth months it

continues to increase until the summit of the fundus is immediately below the ensiform cartilage (Fig. 73). A knowledge of the size of the uterine tumor at various periods of pregnancy, as thus indicated, is of considerable practical importance, as forming the only guide by which we can estimate the probable period of delivery in certain cases in which the usual data for calculation are absent; as, for example, when the patient has conceived during lactation.

The Uterus Sinks before Delivery.—For about a week or more before labor the uterus generally sinks somewhat into the pelvic cavity, in consequence of the relaxation of the soft parts which precedes delivery, and the patient now feels herself smaller and lighter than before. This

change is familiar to all childbearing women, to whom it is known as "the lightening before labor."

The Direction of the Uterus.—While the uterus remains in the pelvis its longitudinal axis varies in direction, much in the same way as that of the non-pregnant uterus, sometimes being more or less vertical, at others in a state of anteversion or partial retroversion. These variations are probably dependent on the distension or emptiness of the bladder, as its state must necessarily affect the position of the movable organ poised behind it. After the uterus has risen into the abdomen its tendency is to project forward against the abdominal wall, which forms its chief support in front. In the erect position the long axis of the uterine tumor corresponds with the axis of the pelvic brim, forming an angle of about 30° with the horizon. In the semi-recumbent position, on the other hand, as Duncan¹ has pointed out, its direction becomes much more nearly vertical. In women who have borne many children the abdominal parietes no longer afford an efficient support, and the uterus is displaced anteriorly, the fundus in extreme cases even hanging downward.

Lateral Obliquity of the Uterus.—In addition to this anterior obliquity, on account of the projection of the spinal column, the uterus is very generally also displaced laterally, and sometimes to a very marked degree, so that it may be felt entirely in one flank, instead of in the centre of the abdomen. In a large proportion of cases this lateral deviation is to the right side, and many hypotheses have been brought forward to explain this fact, none of them being satisfactory. Thus, it has been supposed to depend on the greater frequency with which women lie on their right side during sleep, on the greater use of the right leg during walking, on the supposed comparative shortness of the right round ligament, which drags the tumor to that side, or on the frequent distension of the rectum on the left side, which prevents the uterus being displaced in that direction. Of these the last is the cause which seems most constantly in operation and most likely to produce the effect.

Changes in the Direction of the Cervix.—The cervix must obviously adapt itself to the situation of the body of the uterus. We find, therefore, that in the early months, when the uterus lies low in the pelvis, it is more readily within reach. After the ascent of the uterus it is drawn up, and frequently so much so as to be reached with difficulty. When the uterus is much anteverted, as is so often the case, the os is displaced backward, so that it cannot be felt at all by the examining finger.

Relation of the Uterus to the Surrounding Parts.—Toward the end of pregnancy the greater part of the anterior surface of the uterus is in contact with the abdominal wall, its lower portion resting on the posterior surface of the symphysis pubis. The posterior surface rests on the spinal column, while the small intestines are pushed to either side, the large intestines surrounding the uterus like an arch.

Changes in the Uterine Parietes.—The great distension of the uterus during pregnancy was formerly supposed to be mainly due to the mechanical pressure of the enlarging ovum within it. If this were so,

¹ *Researches in Obstetrics*, p. 10.

then the uterine walls would be necessarily much thinner than in the non-pregnant state. This is well known not to be the case, and the immense increase in the size of the uterine cavity is to be explained by the hypertrophy of its walls. At the full period of pregnancy the thickness of the uterine parietes is generally about the same as that of the non-pregnant uterus, rather more at the placental site, and less in the neighborhood of the cervix. Their thickness, however, varies in different places, and in some women they are so thin as to admit of the foetal limbs being very readily made out by palpation. Their density is, however, always much diminished, and instead of being hard and inelastic they become soft and yielding to pressure. This change coincides with the commencement of pregnancy, of which it forms, as recognizable in the cervix, one of the earliest diagnostic marks. At a more advanced period it is of value as admitting a certain amount of yielding of the uterine walls to movements of the foetus, thus lessening the chance of their being injured.

Changes in the Cervix during Pregnancy.—Very erroneous views have long been taught, in most of our standard works on midwifery, as to the changes which occur in the cervix uteri during pregnancy. It is generally stated that as pregnancy advances the cervical cavity is greatly diminished in length, in consequence of its being gradually drawn up so as to form part of the general cavity of the uterus, so that in the latter

FIG. 74.



FIG. 76.



FIG. 75.



FIG. 77.




Supposed Shortening of the Cervix at the Third, Sixth, Eighth, and Ninth Months of Pregnancy, as figured in *Obstetric Works*.

months it no longer exists. In almost all midwifery works accurate diagrams are given of this progressive shortening of the cervix (Figs. 74 to 77). The cervix is generally described as having lost one-half of its length at the sixth month, two-thirds at the seventh, and to be entirely obliterated in the eighth and ninth. The correctness of these views was first called in question in recent times by Stoltz in 1826, but Dr. Duncan,¹ in an elaborate historical paper on the subject, has shown

¹ *Researches in Obstetrics.*

that Stoltz was anticipated by Weitbrech in 1750, and to a less degree by Roederer and other writers. This opinion is now pretty generally admitted to be correct, and is upheld by Cazeaux, Arthur Farre, Duncan, and most modern obstetricians. Indeed, various post-mortem examinations in advanced pregnancy have shown that the cavity of the cervix remains in reality of its normal length of one inch, and it can often be measured during life by the examining finger on account of its patulous state (Fig. 78). During the fortnight immediately preceding

FIG. 78.



Cervix from a Woman dying in the Eighth Month of Pregnancy. (After Duncan.)

delivery, however, a real shortening or obliteration of the cervical cavity takes place; but this, as Duncan has pointed out, seems to be due to the incipient uterine contractions which prepare the cervix for labor.

An Apparent Shortening is always Present.—There is no doubt an apparent shortening of the cervix always to be detected during pregnancy, but this is a fallacious and deceptive feeling, due to the softness of the tissue of the cervix, which is exceedingly characteristic of pregnancy, and which to an experienced finger affords one of its best diagnostic marks.

Softening of the Cervix.—In the non-pregnant state the tissue of the cervix is hard, firm, and inelastic. When conception occurs, softening begins at the external os, and proceeds gradually and slowly upward until it involves the whole of the cervix. By the end of the fourth month both lips of the os are thick, softened, and velvety to the touch, giving a sensation likened by Cazeaux to that produced by pressing on a table through a thick, soft cover. By the sixth month at least one-half of the cervix is thus altered, and by the eighth the whole of it, and

so much so that at this time those unaccustomed to vaginal examination experience some difficulty in distinguishing it from the vaginal walls. It is this softening, then, which gives rise to the apparent shortening of the cervix so generally described, and it is an invaluable concomitant of pregnancy, except in some rare cases in which there has been antecedent morbid induration and hypertrophic elongation of the cervix. If, therefore, on examining a woman supposed to be advanced in pregnancy, we find the cervix to be hard and projecting into the vaginal canal, we may safely conclude that pregnancy does not exist. The existence of softening, however, it must be remembered, will not itself justify an opposite conclusion, as it may be produced, to a very considerable extent, by various pathological conditions of the uterus.

The Os Uteri is generally Patulous.—At the same time that the tissue of the cervix is softened its cavity is widened and the external os becomes patulous. This change varies considerably in primiparæ and multiparæ. In the former the external os often remains closed until the end of pregnancy, but even in them it generally becomes more or less patulous after the seventh month, and admits the tip of the examining finger. In women who have borne children this change is much more marked. The lips of the external os are in them generally fissured and irregular, from slight lacerations of its tissue in former labors. It is also sufficiently open to admit the tip of the finger, so that in the latter months of pregnancy it is often quite possible to touch the membranes, and through them to feel the presenting part of the child.

Changes in the Texture of the Uterine Tissues: The Peritoneal Coat.—The remarkable increase in size of the uterus during pregnancy is, as we have seen, chiefly to be explained by the growth of its structures, all of which are modified during gestation. The peritoneal covering is considerably increased, so as still to form a complete covering to the uterus when at its largest size. William Hunter supposed that its extension was effected rather by the unfolding of the layers of the broad ligament than by growth. That the layers of the broad ligament do unfold during gestation, especially in the early months, is probable; but this is not sufficient to account for the complete investment of the uterus, and it is certain that the peritoneum grows *pari passu* with the enlargement of the uterus. In addition, there is a new formation of fibrous tissue between the peritoneal and the muscular coats, which affords strength and diminishes the risk of laceration during labor.

The Muscular Coat.—The hypertrophy of the muscular tissue of the uterus is, however, the most remarkable of the changes produced by pregnancy. Not only do the previously existing rudimentary fibre-cells become enormously increased in size—so as to measure, according to Kölliker, from seven to eleven times their former length and from two to five times their former breadth—but new unstriped fibres are largely developed, especially in the inner layers. These new cells are chiefly found in the first months of pregnancy, and their growth seems to be completed by the sixth month. The connective tissue between the muscular layers is also largely increased in amount. The weight of the muscular tissue of the gravid uterus is therefore much increased, and it has been estimated by Heschl that it weighs at term from 1 to 1.5 lbs.;

that is, about sixteen times more than in the unimpregnated state. This great development of the muscular tissue admits of its dissection in a way which is quite impossible in the unimpregnated state, and the researches of H  lie (p. 61) enable us to understand much better than before how the muscles forming the walls of the gravid uterus act during the expulsion of the child.

The changes in the mucous coat of the uterus which result in the formation of the decidua have already been discussed at length elsewhere (p. 103).

Circulatory Apparatus.—The circulatory apparatus of the uterus during pregnancy has been described when the anatomy of the placenta was under consideration (p. 111).

Lymphatics.—The lymphatics are much increased in size, and recent theories on the production of certain puerperal diseases attribute to them a more important action than has been commonly assigned to them.

Nerves.—The question of the growth of the nerves has been hotly discussed. Robert Lee took the foremost place among those who maintained that the nerves of the uterus share the general growth of its other constituent parts. Dr. Snow Beck, however, believed that they remain of the same size as in the unimpregnated state; and this view is supported by Hirschfeld, Robin, and other recent writers. Robin thought that there is an apparent increase in the size of the nerve-tubes, which, however, is really due to increase in the neurilemma. Kilian describes the nerves as increasing in length, but not in thickness; while Schroeder states that they participate equally with the lymphatics in the enlargement the latter undergo. Whichever of these views may ultimately be found to be correct, it is certain that analogy would lead us to expect an increase of nervous as well as of vascular supply.

General Modifications in the Body produced by Pregnancy.—It is not in the uterus alone that pregnancy is found to produce modifications of importance. There are few of the more important functions of the body which are not, to a greater or less extent, affected: to some of these it is necessary briefly to direct attention, inasmuch as, when carried to excess, they produce those disorders which often complicate gestation and which prove so distressing and even dangerous to the patients. Such of them as are apparent and may aid us in diagnosis are discussed in the chapter which treats of the signs and symptoms of pregnancy: in this place it is only necessary to refer to those which do not properly fall into that category.

Changes in the Blood.—Amongst those which are most constant and important are the alterations in the composition of the blood. The opinion of the profession on this subject has of late years undergone a remarkable change. Formerly it was universally believed that pregnancy was, as the rule, associated with a condition analogous to plethora, and that this explained many characteristic phenomena of common occurrence, such as headache, palpitation, singing in the ears, shortness of breath, and the like. As a consequence, it was the habitual custom, not yet by any means entirely abandoned, to treat pregnant women on an antiphlogistic system—to place them on low diet, to administer low-

ering remedies, and very often to practise venesection, sometimes to a surprising extent. Thus it was by no means rare for women to be bled six or eight times during the latter months, even when no definite symptoms of disease existed; and many of the older authors record cases where depletion was practised every fortnight as a matter of routine, and when the symptoms were well marked even from fifty to ninety times, in the course of a single pregnancy.

Composition of the Blood in Pregnancy.—Numerous careful analyses have conclusively proved that the composition of the blood during pregnancy is very generally—perhaps it would not be too much to say always—profoundly altered. Thus it is found to be more watery, its serum is deficient in albumen, and the amount of colored globules is materially diminished, averaging, according to the analysis of Becquerel and Rodier, 111.8 against 127.2 in the non-gravid state. At the same time the amount of fibrin and of extractive matter is considerably increased. The latter observation is of peculiar importance, as it goes far to explain the frequency of certain thrombotic affections observed in connection with pregnancy and delivery: this hyperinosis of the blood is also considerably increased after labor by the quantity of effete material thrown into the mother's system at that time to be got rid of by her emunctories. The truth is, that the blood of the pregnant woman is generally in a state much more nearly approaching the condition of anemia than of plethora, and it is certain that most of the phenomena attributed to plethora may be explained equally well and better on this view. These changes are much more strongly marked at the latter end of pregnancy than at its commencement; and it is interesting to observe that it is then that the concomitant phenomena alluded to are most frequently met with. Cazeaux, to whom we are chiefly indebted for insisting on the practical bearing of these views, contends that the pregnant state is essentially analogous to chlorosis, and that it should be so treated. More recently the accurate observations of Willcocks¹ have shown that the blood of pregnancy differs from that of chlorosis in the fact that while in both the amount of hæmoglobin is lessened, in pregnancy the individual blood-cells are not impoverished as they are in chlorosis, but simply lessened in comparative number, owing to an increase in the water of the plasma, due to the progressive enlargement of the vascular area during gestation. Objection has not unnaturally been taken to Cazeaux's theory, as implying that a healthy and normal function is associated with a morbid state; and it has been suggested that this deteriorated state of the blood may be a wise provision of nature instituted for a purpose we are not as yet able to understand. It may certainly be admitted that pregnancy, in a perfectly healthy state of the system, should not be associated with phenomena in themselves in any degree morbid. It must not be forgotten, however, that our patients are seldom—we might safely say never—in a state that is physiologically healthy. The influence of civilization, climate, occupation, diet, and a thousand other disturbing causes, that to a greater or less degree are always to be met with, must not be left out of consideration. Making

¹ "Comparative Observations on the Blood in Chlorosis and Pregnancy," by Fred. Willcocks, M. D., *The Lancet*, December 3, 1881.

every allowance, therefore, for the undoubted fact that pregnancy *ought* to be a perfectly healthy condition, it must be conceded, I think, that in the vast majority of cases coming under our notice it is not entirely so; and the deductions drawn by Cazeaux from the numerous analyses of the blood of pregnant women seem to point strongly to the conclusion that the general blood-state is tending to poverty and anæmia, and that a depressing and antiphlogistic treatment is distinctly contraindicated.

Modifications in Certain Viscera.—Closely connected with the altered condition of the blood is the physiological hypertrophy of the heart which is now well known to occur during pregnancy. This was first pointed out by Larcher in 1828, and it has been since verified by numerous observers. It seems to be constant and considerable, and to be a purely physiological alteration, intended to meet the increased exigencies of the circulation which the complex vascular arrangements of the gravid uterus produce. The hypertrophy is limited to the left ventricle, the right ventricle, as well as both auricles, being unaffected. Blot estimates that the whole weight of the heart increases one-fifth during gestation. The more recent researches of Löhlein¹ render it probable that the hypertrophy is less than those authors have supposed. According to Duroziez² the heart remains enlarged during lactation, but diminishes in size immediately after delivery in women who do not suckle, while in women who have borne many children it remains permanently somewhat larger than in nullipare. Similar increase in the size of other organs has been pointed out by various writers; as, for example, in the lymphatics, the spleen, and the liver. Tarnier states that in women who have died after delivery the organs always show signs of fatty degeneration. According to Gassner, the whole body increases in weight during the latter months of pregnancy, and this increase is somewhat beyond that which can be explained by the size of the womb and its contents.

Formation of Osteophytes.—Irregular bony deposits between the skull and the dura mater, in some cases so largely developed as to line the whole cranium, have been so frequently detected in women who have died during parturition that they are believed by some to be a normal production connected with pregnancy. Ducrest found these osteophytes in more than one-third of the cases in which he performed post-mortem examinations during the puerperal period. Rokitansky, who corroborated the observation, believed this peculiar deposit of bony matter to be a physiological and not a pathological condition connected with pregnancy; but whether it be so, or how it is produced, has not yet been satisfactorily determined.

Changes in the Nervous System.—More or less marked changes connected with the nervous system are generally observed in pregnancy, and sometimes to a very great extent. When carried to excess they produce some of the most troublesome disorders which complicate gestation, such as alterations in the intellectual functions, changes in the disposition and character, morbid cravings, dizziness, neuralgia, syncope, and many others. They are purely functional in their cha-

¹ *Zeitschrift für Geburtshilfe, etc.*, 1876.

² *Gaz. des Hôpit.*, 1868.

meter, and disappear rapidly after delivery, and may be best described in connection with the disorders of pregnancy.

Changes in the Respiratory Organs.—Respiration is often interfered with, from the mechanical results of the pressure of the enlarged uterus. The longitudinal dimensions of the thorax are lessened by the upward displacement of the diaphragm, and this necessarily leads to some embarrassment of the respiration, which is, however, compensated, to a great extent, by an increase in breadth of the base of the thoracic cavity.

Changes in the Urine.—Certain changes, which are of very constant occurrence, in the urine of pregnant women have attracted much attention, and have been considered by many writers to be pathognomonic. They consist in the presence of a peculiar deposit, formed when the urine has been allowed to stand for some time, which has received the name of *kiestein*. Its presence was known to the ancients, and it was particularly mentioned by Savonarola in the fifteenth century, but it has more especially been studied within the last thirty years by Eguisier, Golding Bird, and others. If the urine of a pregnant woman be allowed to stand in a cylindrical vessel, exposed to light and air, but protected from dust, in a period varying from two to seven days, a peculiar flocculent sediment, like fine cotton wool, makes its appearance in the centre of the fluid, and soon afterward rises to the surface and forms a pellicle, which has been compared to the fat on cold mutton-broth. In the course of a few days the scum breaks up and falls to the bottom of the vessel. On microscopic examination it is found to be composed of fat particles, with crystals of ammoniaco-magnesium phosphates and phosphate of lime and a large quantity of vibriones. These appearances are generally to be detected after the second month of pregnancy, and up to the seventh or eighth month, after which they are rarely produced. Regnaud explains their absence during the latter months of gestation by the presence in the urine at that time of free lactic acid, which increases its acidity and prevents the decomposition of the urea into carbonate of ammonia. He believes that *kiestein* is produced by the action of free carbonate of ammonia on the phosphate of lime contained in the urine, and that this reaction is prevented by the excess of acid.

Golding Bird believed *kiestein* to be analogous to casein, to the presence of which he referred it, and he states that he has found it in twenty-seven out of thirty cases. Braxton Hicks so far corroborates his view, and states that the deposit of *kiestein* can be much more abundantly produced if one or two teaspoonfuls of rennet be added to the urine, since that substance has the property of coagulating casein. Much less importance, however, is now attached to the presence of *kiestein* than formerly, since a precisely similar substance is sometimes found in the urine of the non-pregnant, especially in anæmic women, and even in the urine of men. Parkes states that it is not of uniform composition—that it is produced by the decomposition of urea, and consists of the free phosphates, bladder mucus, infusoria, and vaginal discharges. Neugebauer and Vogel give a similar account of it, and hold that it is of no diagnostic value. That it is of interest, as indi-

cating the changes going on in connection with pregnancy, is certain ; but inasmuch as it is not of invariable occurrence, and may even exist quite independently of gestation, it is obviously quite undeserving of the extreme importance that has been attached to it.

[Although not a reliable test of pregnancy, it is a remarkable fact that in all the cases of suspected pregnancy in private practice in which I have employed it I never found a woman who proved to be impregnated in whose urine it did not appear. When Dr. Kane was preparing his thesis on kiestein he examined a large number of specimens of urine and considered the test a good one ; but the fact that it may be found in the urine of the non-pregnant destroys its reliability for general use in diagnosis.—ED.]

Glycosuria in Pregnancy.—Toward the end of pregnancy sugar may sometimes be detected in the urine, and after delivery and during lactation it exists in considerable abundance ; thus, out of thirty-five cases tested in the Simpson Memorial Hospital in Edinburgh during the puerperium, it was found in all, the amount varying from 1 to 8 per cent.¹ Kaltenbach has shown that this temporary glycosuria is due to the presence of milk-sugar in the urine, and that it ceases with the disappearance of milk from the breasts.² This physiological glycosuria must be carefully distinguished from true diabetes, which is a grave complication of pregnancy.

CHAPTER IV.

SIGNS AND SYMPTOMS OF PREGNANCY.

Importance of the Subject.—In attempting to ascertain the presence or absence of pregnancy the practitioner has before him a problem which is often beset with great difficulties, and on the proper solution of which the moral character of his patient, as well as his own professional reputation, may depend. The patient and her friends can hardly be expected to appreciate the fact that it is often far from easy to give a positive opinion on the point ; and it is always advisable to use much caution in the examination, and not to commit ourselves to a positive opinion, except on the most certain grounds. This is all the more important because it is just in those cases in which our opinion is most frequently asked that the statements of the patient are of least value, as she is either anxious to conceal the existence of pregnancy, or, if desirous of an affirmative diagnosis, unconsciously colors her statements, so as to bias the judgment of the examiner.

Classification.—Constant attempts have been made to classify the signs of pregnancy : thus, some divide them into the *natural* and *sensible* signs, others into the *presumptive*, the *probable*, and the *certain*. The latter classification, which is that adopted by Montgomery in his classical work

¹ *Edin. Med. Journ.*, Aug., 1881.

² *Zeit. f. Gynäk.*, September 13, 1879.

on the *Signs and Symptoms of Pregnancy*, is no doubt the better of the two, if any be required. The simplest way of studying the subject, however, is the one now generally adopted, of considering the signs of pregnancy in the order in which they occur, and attaching to each an estimate of its diagnostic value.

Signs of a Fruitful Conception.—From the earliest ages authors have thought that the occurrence of conception might be ascertained by certain obscure signs, such as a peculiar appearance of the eyes, swelling of the neck, or by unusual sensations connected with a fruitful intercourse. All of these, it need hardly be said, are far too uncertain to be of the slightest value. The last is a symptom on which many married women profess themselves able to depend, and one to which Cazeaux is inclined to attach some importance.

Cessation of Menstruation.—The first appreciable indication of pregnancy on which any dependence can be placed is the cessation of the customary menstrual discharge, and it is of great importance, as forming the only reliable guide for calculating the probable period of delivery. In women who have been previously perfectly regular, in whom there is no morbid cause which is likely to have produced suppression, the non-appearance of the catamenia may be taken as strong presumptive evidence of the existence of pregnancy; but it can never be more than this, unless verified and strengthened by other signs, inasmuch as there are many conditions besides pregnancy which may lead to its non-appearance. Thus, exposure to cold, mental emotion, general debility, especially when connected with incipient phthisis, may all have this effect. Mental impressions are peculiarly liable to mislead in this respect. It is far from uncommon in newly-married women to find that menstruation ceases for one or more periods, either from the general disturbance of the system connected with the married life or from a desire on the part of the patient to find herself pregnant. Also in unmarried women who have subjected themselves to the risk of impregnation mental emotion and alarm often produce the same result.

Menstruation during Pregnancy.—A further source of uncertainty exists in the fact that in certain cases menstruation may go on for one or more periods after conception, or even during the whole pregnancy. The latter occurrence is certainly of extreme rarity, but one or two instances are recorded by Perfect, Churchill, and other writers of authority, and therefore its possibility must be admitted. The former is much less uncommon, and instances of it have probably come under the observation of most practitioners. The explanation is now well understood. During the early months of gestation, when the ovum is not yet sufficiently advanced in growth to fill the whole uterine cavity, there is a considerable space between the decidua reflexa which surrounds it and the decidua vera lining the uterine cavity. It is from this free surface of the decidua vera that the periodical discharge comes, and there is not only ample surface for it to come from, but a free channel for its escape through the os uteri. After the third month the decidua reflexa and the decidua vera blend together, and the space between them disappears. Menstruation after this time is therefore much more difficult to account for. It is probable that in many supposed cases occasional losses of blood from

other sources, such as placenta prævia, an abraded cervix uteri, or a small polypus, have been mistaken for true menstruation. If the discharge really occurs periodically after the third month, it can only come from the canal of the cervix. The occurrence, however, is so rare that if a woman is menstruating regularly and normally who believes herself to be more than four months advanced in pregnancy, we are justified *ipso facto* in negating her supposition. In an unmarried woman all statements as to regularity of menstruation are absolutely valueless, for in such cases nothing is more common than for the patient to make false statements for the express purpose of deception.

Pregnancy sometimes occurs when Menstruation is normally Absent.—Conception may unquestionably occur when menstruation is normally absent. This is far from uncommon in women during lactation, when the function is in abeyance, and who therefore have no reliable data for calculating the true period of their delivery. Authentic cases are also recorded in which young girls have conceived before menstruation is established, and in which pregnancy has occurred after the change of life.

Estimate of its Diagnostic Value.—Taking all these facts into account, we can only look upon the cessation of menstruation as a fairly presumptive sign of pregnancy in women in whom there is no clear reason to account for it, but one which is undoubtedly of great value in assisting our diagnosis.

Sympathetic Disturbances.—Shortly after conception various sympathetic disturbances of the system occur, and it is only very exceptionally that these are not established. They are generally most developed in women of highly nervous temperament; and they are, therefore, most marked in patients in the upper classes of society, in whom this kind of organization is most common.

Morning Sickness.—Amongst the most frequent of these are various disorders of the gastro-intestinal canal. Nausea or vomiting is very common; and as it is generally felt on first rising from the recumbent position, it is popularly known amongst women as the "morning sickness." It sometimes commences almost immediately after conception, but more frequently not until the second month, and it rarely lasts after the fourth month. Generally there is nausea rather than actual vomiting. The woman feels sick and unable to eat her breakfast, and often brings up some glairy fluid. In other cases she actually vomits; and sometimes the sickness is so excessive as to resist all treatment, seriously to affect the patient's health, and even imperil her life. These grave forms of the affection will require separate consideration.

Cause of the Sickness.—Very different opinions have been held as to the cause of morning sickness. Dr. Henry Bennet believes that when at all severe it is always associated with congestion and inflammation of the cervix uteri. Dr. Graily Hewitt maintains that it depends entirely on flexion of the uterus, producing irritation of the uterine nerves at the seat of the flexion, and consequent sympathetic vomiting. This theory, when broached at the Obstetrical Society, was received with little favor: it seems to me to be sufficiently disproved by the fact, which I believe to be certain, that more or less nausea is a normal and nearly constant phenomenon in pregnancy, for it is difficult to believe that nearly every

pregnant woman has a flexed uterus. The generally received explanation is probably the correct one—viz. that nausea, as well as other forms of sympathetic disturbance, depends on the stretching of the uterine fibres by the growing ovum, and consequent irritation of the uterine nerves. It is, therefore, one, and only one, of the numerous reflex phenomena naturally accompanying pregnancy. It is an old observation that when the sickness of pregnancy is entirely absent, other, and generally more distressing, sympathetic derangements are often met with, such as a tendency to syncope. Dr. Bedford¹ has laid especial stress on this point, and maintains that under such circumstances women are peculiarly apt to miscarry.

Other derangements of the digestive functions, depending on the same cause, are not uncommon, such as excessive or depraved appetite, the patient showing a craving for strange and even disgusting articles of diet. These cravings may be altogether irresistible, and are popularly known as “longings.” Of a similar character is the disturbed condition of the bowels frequently observed, leading to constipation, diarrhœa, and excessive flatulence.

Other Sympathetic Phenomena.—Certain glandular sympathies may be developed, one of the most common being an excessive secretion from the salivary glands. A tendency to syncope is not infrequent, rarely proceeding to actual fainting, but rather to that sort of partial syncope, unattended with complete loss of consciousness, which the older authors used to call “lypothæmia.” This often occurs in women who show no such tendency at other times, and when developed to any extent it forms a very distressing accompaniment of pregnancy. Toothache is common, and is not rarely associated with actual caries of the teeth. When any of these phenomena are carried to excess, it is more than probable that some morbid condition of the uterus exists which increases the local irritation producing them.

Mental Peculiarities.—Mental phenomena are very general. An undue degree of despondency, utterly beyond the patient's control, is far from uncommon; or a change which renders the bright and good-tempered woman fractious and irritable; or even the more fortunate but less common change, by which a disagreeable disposition becomes altered for the better.

The Diagnostic Value of these Sympathetic Disturbances is Small.—All these phenomena of exalted nervous susceptibility are but of slight diagnostic value. They may be taken as corroborating more certain signs, but nothing more, and they are chiefly interesting from their tendency to be carried to excess and to produce serious disorders.


Mammary Changes.—Certain changes in the mammæ are of early occurrence, dependent, no doubt, on the intimate sympathetic relations at all times existing between them and the uterine organs, but chiefly required for the purpose of preparing for the important function of lactation, which on the termination of pregnancy they have to perform.

Changes in the Areolæ.—Generally about the second month of pregnancy the breasts become increased in size and tender. As pregnancy advances they become much larger and firmer, and blue veins may be seen coursing over them. The most characteristic changes are about the

¹ *Diseases of Women and Children*, p. 551.

nipples and areolæ. The nipples become turgid, and are frequently covered with minute branny scales, formed by the desiccation of sero-lactescent fluid oozing from them. The areolæ become greatly enlarged and darkened from the deposit of pigment (Fig. 79). The extent and degree of this discoloration vary much in different women. In fair women it may be so slight as to be hardly appreciable, while in dark women it is

FIG. 79.



Appearance of the Areola in Pregnancy.

generally exceedingly characteristic, sometimes forming a nearly black circle extending over a great part of the breast. The areola becomes moist as well as dark in appearance, and is somewhat swollen, and a number of small tubercles are developed upon it, forming a circle of projections round the nipple. These tubercles are described by Montgomery as being intimately connected with the lactiferous ducts, some of which may occasionally be traced into them and seem to open on their summits. As pregnancy advances they increase in size and number. During the latter months what has been called "the secondary areola" is produced, and when well marked presents a very characteristic appearance. It consists of a number of minute discolored spots all round the outer margin of the areola, where the pigmentation is fainter, and which are generally described as resembling spots from which the color has been discharged by a shower of water-drops. This change, like the darkening of the primary areola, is more marked in brunettes. At this period, especially in women whose skin is of fine texture, whitish silvery streaks are often seen on the breasts. They are produced by the stretching of the cutis vera, and are permanent.

By pressure on the breasts a small drop of serous-looking fluid can very generally be forced out from the nipple, often as early as the third month, and on microscopic examination milk and cholestrum-globules can be seen in it.

Diagnostic Value of Mammary Changes.—The diagnostic value of these mammary changes has been variously estimated. When well marked they are considered by Montgomery to be certain signs of pregnancy. To this statement, however, some important limitations must be made. In women who have never borne children they no doubt are so; for, although various uterine and ovarian diseases produce some darkening of the areola, they certainly never produce the well-marked changes above described. In multiparæ, however, the areolæ often remain permanently darkened, and in them these signs are much less reliable. In first pregnancies the presence of milk in the breasts may be considered an almost certain sign, and it is one which I have rarely failed to detect even from a comparatively early period. It is true that there are authenticated instances of non-pregnant women having an abundant secretion of milk established from mammary irritation. Thus, Baudelocque presented to the Academy of Surgery of Paris a young girl, eight years of age, who had nursed her little brother for more than a month. Dr. Tanner states—I do not know on what authority—that “it is not uncommon in Western Africa for young girls who have never been pregnant to regularly employ themselves in nursing the children of others, the mammæ being excited to action by the application of the juice of one of the Euphorbiaceæ.” Lacteal secretion has even been noticed in the male breast. But these exceptions to the general rule are so uncommon as merely to deserve mention as curiosities; and I have hardly ever been deceived in diagnosing a first pregnancy from the presence of even the minutest quantity of lacteal secretion in the breasts, although even then other corroborative signs should always be sought for. In multiparæ the presence of milk is by no means so valuable, for it is common for milk to remain in the mammæ long after the cessation of lactation, even for several years. Tyler Smith correctly says that “suppression of the milk in persons who are nursing and liable to impregnation is a more valuable sign of pregnancy than the converse condition.” This is an observation I have frequently corroborated.

As a diagnostic sign, therefore, the mammary appearances are of great importance in primiparæ, and when well marked they are seldom likely to deceive. They are specially important when we suspect pregnancy in the unmarried, as we can easily make an excuse to look at the breast without explaining to the patient the reason; and a single glance, especially if the patient be dark-complexioned, may so far strengthen our suspicion as to justify a more thorough examination. In married multiparæ they are less to be depended upon.

Other Pigmentary Changes.—In connection with this subject may be mentioned various irregular deposits of pigment which are frequently observed. The most common is a dark-brownish or yellowish line starting from the pubes and running up to the centre of the abdomen, sometimes as far as the umbilicus only, at others forming an irregular ring round the umbilicus and reaching to the epigastrium. It is, however, of very uncertain occurrence, being well marked in some women, while in others it is entirely absent. Patches of darkened skin are often observed about the face, chiefly on the forehead, and this bronzing sometimes gives a very peculiar appearance. Joulin states that it only occurs

on parts of the face exposed to the sun, and that it is therefore most frequently observed in women of the lower order who are freely exposed to atmospheric influences. These pigmentary changes are of small diagnostic value, and may continue for a considerable time after delivery.

Enlargement of the Abdomen.—The progressive enlargement of the abdomen and the size of the gravid uterus at various periods of pregnancy, as well as the method of examination by means of abdominal palpation, have already been described (pp. 124 and 134).

We will now consider the well-known phenomena, produced by the movements of the fœtus in utero, which are so familiar to all pregnant women. These, no doubt, take place from the earliest period of fœtal life at which the muscular tissue of the fœtus is sufficiently developed to admit of contraction, but they are not felt by the mother until somewhere about the sixteenth week of utero-gestation, the precise period at which they are perceived varying considerably in different cases. The error of the law on this subject, which supposes the child not to be alive, or "quick," until the mother feels its movements, is well known, and has frequently been protested against by the medical profession.

Quickening.—The so-called *quickening*—which certainly is felt very suddenly by some women—is believed to depend on the rising of the uterine tumor sufficiently high to permit of the impulse of the fœtus being transmitted to the abdominal walls of the mother, through the sensory nerves of which, its movements become appreciable. The sensation is generally described as being a feeble fluttering, which when first felt not unfrequently causes unpleasant nervous sensations. As the uterus enlarges the movements become more and more distinct, and generally consist of a series of sharp blows or kicks, sometimes quite appreciable to the naked eye and causing distinct projections of the abdominal walls. Their force and frequency will also vary during pregnancy according to circumstances. At times they are very frequent and distressing; at others the fœtus seems to be comparatively quiet, and they may even not be felt for several days in succession, and thus unnecessary fears as to the death of the fœtus often arise. The state of the mother's health has an undoubted influence upon them. They are said to increase in force after a prolonged abstinence from food or in certain positions of the body. It is certain that causes interfering with the vitality of the fœtus often produce very irregular and tumultuous movements. They can be very readily felt by the accoucheur on palpating the abdomen, and sometimes, in the latter months, so distinctly as to leave no doubt as to the existence of pregnancy. They can also generally be induced by placing one hand on each side of the abdomen and applying gentle pressure, which will induce fœtal motion that can be easily appreciated.

The Diagnostic Value of Fœtal Movements.—As a diagnostic sign the existence of fœtal movements has always held a high place, but care should be taken in relying on it. It is certain that women are themselves very often in error, and fancy they feel the movements of a fœtus when none exists, being probably deceived by irregular contractions of the abdominal muscles or flatus within the bowels. They may even involuntarily produce such intra-abdominal movements as may readily

deceive the practitioner. Of course in advanced pregnancy, when the foetal movements are so marked as to be seen as well as felt, a mistake is hardly possible, and they then constitute a certain sign. But in such cases there is an abundance of other indications and little room for doubt. In questionable cases, and at an early period of pregnancy, the fact that movements are not felt must not be taken as a proof of the non-existence of pregnancy, for they may be so feeble as not to be perceptible, or they may be absent for a considerable period.

Intermittent Uterine Contractions.—Braxton Hicks¹ has directed attention to the value, from a diagnostic point of view, of intermittent contractions of the uterus during pregnancy. After the uterus is sufficiently large to be felt by palpation, if the hand be placed over it and it be grasped for a time without using any friction or pressure, it will be observed to distinctly harden in a manner that is quite characteristic.

This intermittent contraction occurs every five or ten minutes, sometimes oftener, rarely at longer intervals. The fact that the uterus did contract in this way had been previously described, more especially by Tyler Smith, who ascribed it to peristaltic action. But it is certain that no one, before Dr. Hicks, had pointed out the fact that such contractions are constant and normal concomitants of pregnancy, continuing during the whole period of utero-gestation, and forming a ready and reliable means of distinguishing the uterine tumor from other abdominal enlargements. Since reading Dr. Hicks's paper I have paid considerable attention to this sign, which I have never failed to detect, even in the retroverted gravid uterus contained entirely in the pelvic cavity, and I am disposed entirely to agree with him as to its great value in diagnosis. If the hand be kept steadily on the uterus, its alternate hardening and relaxation can be appreciated with the greatest ease. The advantages which this sign has over the foetal movements are that it is constant, that it is not liable to be simulated by anything else, and that it is independent of the life of the child, being equally appreciable when the uterus contains a degenerated ovum or dead foetus. The only condition likely to give rise to error is an enlargement of the uterus in consequence of contents other than the results of conception, such as retained menses or a polypus. The history of such cases—which are, moreover, of extreme rarity—would easily prevent any mistake. As a corroborative sign of pregnancy, therefore, I should give these intermittent contractions a high place. [I once attended the wife of a physician in her second pregnancy, who had lost her first child by abortion, and was supposed to be again threatened with the same misfortune. I found her suffering pain with each intermittent contraction, but beyond this there were no symptoms to indicate an expulsive design on the part of the uterus. These *painful* intermittent contractions persisted for three weeks, and then gradually assumed their normal character under an opiate treatment. The lady went to the full term of gestation and bore a child which lived.—ED.]

Vaginal Signs of Pregnancy.—The vaginal signs of pregnancy are of considerable importance in diagnosis. They are chiefly the changes

¹ *Obst. Trans.*, v. 13.

which may be detected in the cervix, and the so-called *ballottement*, which depends on the mobility of the fœtus in the liquor amnii.

Softening of the Cervix.—The alterations in the density and apparent length of the cervix have been already described (p. 137). When pregnancy has advanced beyond the fifth month the peculiar velvety softness of the cervix is very characteristic, and affords a strong corroborative sign, but one which it would be unsafe to rely on by itself, inasmuch as very similar alterations may be produced by various causes. When, however, in a supposed case of pregnancy advanced beyond the period indicated, the cervix is found to be elongated, dense, and projecting into the vaginal canal, the non-existence of pregnancy may be safely inferred. Therefore the negative value of this sign is of more importance than the positive.

Ballottement.—Ballottement, when distinctly made out, is a very valuable indication of pregnancy. It consists in the displacement, by the examining finger, of the fœtus, which floats up in the liquor amnii, and falls back again on the tip of the finger with a slight tap which is exceedingly characteristic.

Method of Examination.—In order to practise it most easily the patient is placed on a couch or bed in a position midway between sitting and lying, by which the vertical diameter of the uterine cavity is brought into correspondence with that of the pelvis. Two fingers of the right hand are then passed high up into the vagina in front of the cervix. The uterus being now steadied from without by the left hand, the intra-vaginal fingers press the uterine wall suddenly upward, when, if pregnancy exist, the fœtus is displaced, and in a moment falls back again, imparting a distinct impulse to the fingers. When easily appreciable it may be considered as a certain sign, for although an anteflexed fundus or a calculus in the bladder may give rise to somewhat similar sensations, the absence of other indications of pregnancy would readily prevent error. Ballottement is practised between the fourth and seventh months. Before the former time the fœtus is too small, while at a later period it is relatively too large, and can no longer be easily made to rise upward in the surrounding liquor amnii. The absence of ballottement must not be taken as proving the non-existence of pregnancy, for it may be inappreciable from a variety of causes, such as abnormal presentations or the implantation of the placenta upon the cervix uteri.

Vaginal Pulsation.—There are also some other vaginal signs of pregnancy of secondary consequence. Amongst these is the vaginal pulsation pointed out by Oslander, resulting from the enlargement of the vaginal arteries, which may sometimes be felt beating at an early period. Often this pulsation is very distinct, at other times it cannot be felt at all, and it is altogether unreliable, as a similar pulsation may be felt in various uterine diseases.

Uterine Fluctuation.—Dr. Rasch has drawn attention to a previously undescribed sign which he believes to be of importance in the diagnosis of early pregnancy.¹ It consists in the detection of fluctuation through the anterior uterine wall, depending on the presence of the liquor amnii. In order to make this out, two fingers of the right hand must be used,

¹ *Brit. Med. Journ.*, vol. ii., 1873.

as in ballottement, while the uterus is steadied through the abdomen. Dr. Rasch states that by this means the enlarged uterus in pregnancy can easily be distinguished from the enlargement depending on other causes, and that fluctuation can always be felt as early as the second month. If it is associated with suppressed menstruation and darkened areolæ, he considers it a certain sign. In order to detect it, however, considerable experience in making vaginal examinations is essential, and it can hardly be depended on for general use.

Alteration in Color of the Vagina.—A peculiar deep violet hue of the vaginal mucous membrane was relied on by Jacquemier and Klüge as affording a readily observed indication of pregnancy. In most cases it is well marked; sometimes, indeed, the change of color is very intense, and it evidently depends on the congestion produced by pressure of the enlarged uterus. The same effect, however, is constantly seen where similar pressure is effected by large fibroid tumors of the uterus, and, therefore, for diagnostic purposes it is valueless.

Auscultatory Signs of Pregnancy.—By far the most important signs are those which can be detected by abdominal auscultation, and one of these—the hearing of the foetal heart-sounds—forms the only sign which *per se*, and in the absence of all others, is perfectly reliable.

Discovery of Foetal Auscultation.—The fact that the sounds of the foetal heart are audible during advanced pregnancy was first pointed out by Mayor of Geneva in 1818, and the main facts in connection with foetal auscultation were subsequently worked out by Kergaradec, Naegele, Evory Kennedy, and other observers. The pulsations first become audible, as a rule, in the course of the fifth month or about the middle of the fourth month. In exceptional circumstances and by practised observers they have been heard earlier. Depaul believes that he detected them as early as the eleventh week, and Routh has also detected them at an earlier period by vaginal stethoscopy, which, however, for obvious reasons, cannot be ordinarily employed. Naegele never heard them before the eighteenth week, more generally at the end of the twentieth, and for practical purposes the pregnancy must be advanced to the fifth month before we can reasonably expect to detect them. From this period up to term they can almost always be heard—if not at the first attempt, at least afterward, to a certainty, if we have the opportunity of making repeated examinations. Accidental circumstances, such as the presence of an unusual amount of flatus in the intestines, may deaden the sounds for a time, but not permanently. Depaul only failed to hear them in 8 cases out of 906 examined during the last three months of pregnancy; and out of 180 cases which Dr. Anderson of Glasgow carefully examined, he only failed in 12, and in each of these the child was stillborn. They therefore form not only a most certain indication of pregnancy, but of the life of the foetus also.

Description of the Sound.—The sound has always been likened to the double tic-tac of a watch heard through a pillow, which it closely resembles. It consists of two beats, separated by a short interval, the first being the loudest and most distinct, the second being sometimes inaudible. The rapidity of the foetal pulsations forms an important means of distinguishing them from transmitted maternal pulsations, with

which they might be confounded. Their average number is stated by Slater, who made numerous observations on this point, to be 132, but sometimes they reach as high as 140, and sometimes as low as 120. It will thus be seen that the pulsations are always much more rapid than those of the mother's heart, unless, indeed, the latter be unduly accelerated by transient mental emotion or disease. To avoid mistakes, whenever the foetal heart is heard its rate of pulsation should be carefully counted and compared with that of the mother's pulse; if the rate differ, we may be sure that no error has been made. The rapidity of the foetal pulsations remains, as a rule, the same during the whole period of pregnancy, while their intensity gradually increases. They may, however, be temporarily increased or diminished in frequency by disturbing causes, such as the pressure of the stethoscope, which, exciting tumultuous movements of the foetus, may induce greatly increased frequency of its heart-beats. So also during labor, after the escape of the liquor amnii, when the contractions of the uterus have a very distinct influence on the foetus, they may be greatly modified. An acceleration or irregularity of the pulsations, made out in the course of a prolonged labor, may thus be of great practical importance, by indicating the necessity for prompt interference. Similar alterations, associated with tumultuous and unusual fetal movements felt by the mother toward the end of pregnancy, may point to danger to the life of the foetus during the latter months, and may even justify the induction of premature labor. This is especially the case in women who have previously given birth to a succession of dead children owing to disease of the placenta, and in them careful and frequently-repeated auscultations may warn us of the impending danger.

Supposed Difference of Rapidity according to the Sex of the Foetus.—The rapidity of the foetal heart has been supposed by some to afford a means of determining the sex of the child before birth. Frankenhauser, who first directed attention to this point, is of opinion that the average rate of pulsations of the heart is considerably less in male than in female children, averaging 124 in the minute in the former as against 144 in the latter. Steinbach makes the difference somewhat less—viz. 131 for males and 138 for females. He predicted the sex correctly by this means in 45 out of 57 cases, while Frankenhauser was correct in the whole 50 cases which he specially examined with reference to the point. Dr. Hutton of New York¹ was also correct in 7 cases he fixed on for trial. Devilliers found the difference in the sexes to be the same as Steinbach; he attributes it, however, to the size and weight rather than to the sex of the child, and believes the pulsations to be least numerous in large and well-developed children. As male children are usually larger than female, he thus explains the relatively less frequent pulsations of their hearts. Dr. Cumming of Edinburgh also believes that the weight of the child has considerable influence on the frequency of its cardiac pulsations, so that a large female child may have a slower pulse than a small male.² The point, however, is more curious than practical, and the rapidity of the pulsations certainly would not justify any positive prediction on the subject. Circumstances influencing

¹ *New York Med. Journ.*, July, 1872.

² *Edin. Med. Journ.*, 1875.

the maternal circulation seem to have no influence on that of the foetus.

Site at which the Sounds are Heard.—The foetal heart-sounds are generally propagated best by the back of the child, and are therefore most easily audible when this is in contact with the anterior wall of the uterus, as is the case in the large majority of pregnancies. When the child is placed in the dorso-posterior position the sounds have to traverse a larger amount of the liquor amnii, and are further modified by the interposition of the foetal limbs. They are, therefore, less easily heard in such cases, but even in them they can almost always be made out. As the foetus most frequently lies with the occiput over the brim of the pelvis, and the back of the child toward the left side of the mother, the heart-sounds are usually most distinctly audible at a point midway between the umbilicus and the left anterior-superior spine of the ilium. In the next most common position, in which the back of the child lies to the right lumbar region of the mother, they are generally heard at a corresponding point at the right side, but in this case they are frequently more readily made out in the right flank, being then transmitted through the thorax of the child, which is in contact with the side of the uterus. In breech cases, on the other hand, the heart-sounds are generally heard most distinctly *above* the umbilicus, and either to the right or left, according to the side toward which the back of the child is placed. It will thus be seen that the place at which the foetal heart-sounds are heard varies with the position of the foetus; and this, when combined with the information derived from palpation, affords a ready means of ascertaining the presentation of the child before labor. The sounds are only audible over a limited space about two or three inches in diameter; therefore, if we fail to detect them in one place, a careful exploration of the whole uterine tumor is necessary before we are satisfied that they cannot be heard.

Sources of Fallacy.—The only mistake that is likely to be made is taking the maternal pulsations, transmitted through the uterine tumor, for those of the foetal heart. A little care will easily prevent this error, and the frequency of the mother's pulse should always be ascertained before counting the supposed foetal pulsations. If these are found to be 120 or more, while the mother's pulse is only 70 or 80, no mistake is possible. If the latter is abnormally quickened, greater care may be necessary, but even then the rate of pulsation of each will be dissimilar. Braxton Hicks¹ has pointed out that in tedious labor, when the muscular powers of the mother are exhausted, the muscular subsurrus may produce a sound closely resembling the foetal pulsation; but error from this source is obviously very improbable.

Mode of Practising Auscultation.—In listening for the foetal heart-sound the patient should be placed on her back, with the shoulders elevated and the knees flexed. The surface of the abdomen should be uncovered and an ordinary stethoscope employed, the end of which must be pressed firmly on the tumor, so as to depress the abdominal walls. The most absolute stillness is necessary, as it is often far from easy to hear the sounds. Sometimes, after failing with the ordinary

¹ *Obst. Trans.*, vol. xv.

stethoscope, I have succeeded with the bin-aural, which remarkably intensifies them.[¹] When once heard they are most easily counted during a space of five seconds, as, on account of their frequency, it is not always possible to follow them over a longer period.

Value of this Sign of Pregnancy.—When the foetal heart-sounds are heard distinctly, pregnancy may be absolutely and certainly diagnosed. The fact that we do not hear them does not, however, preclude the possibility of gestation, for the foetus may be dead or the sounds temporarily inaudible.

Umbilical Souffle.—There are some other sounds heard in auscultation which are of very secondary diagnostic value. One of these is the so-called *umbilical* or *funic souffle*, which was first pointed out by Every Kennedy. It consists of a single blowing murmur, synchronous with the foetal heart-sounds, and most distinctly heard in the immediate vicinity of the point where these are most audible. Most authors believe it to be produced by pressure on the cord, either when it is placed between a hard part of the foetus and the uterine walls or is twisted round the child's neck. Schroeder and Hecker detected it in 14 or 15 per cent. of all cases, and the latter believed it to be caused by flexure of the first portion of the cord near the umbilicus. For practical purposes it is quite valueless, and need only be mentioned as a phenomenon which an experienced auscultator may occasionally detect.

The Uterine Souffle.—The *uterine souffle* is a peculiar single whizzing murmur which is almost always audible on auscultation. It varies very remarkably in character and position. Sometimes it is a gentle blowing or even musical murmur; at others it is loud, harsh, and scraping; sometimes continuous, sometimes intermittent. It may also be heard at any point of the uterus, but most frequently low down and to one or other side; more rarely above the umbilicus or toward the fundus; and it often changes its position so as to be heard at a subsequent auscultation at a point where it was previously inaudible. It may be heard over a space of an inch or two only, or in some cases over the whole uterine tumor; or, again, it may sometimes be detected simultaneously over two entire distinct portions of the uterus. It is generally to be heard earlier than the foetal heart-sounds, often as soon as the uterus rises above the brim of the pelvis, and it can almost always be detected after the commencement of the fourth month. The sound becomes curiously modified by the uterine contractions during labor, becoming louder and more intense before the pain comes on, disappearing during its acme, and again being heard as it goes off. Hicks attributes to a similar cause—viz. the uterine contractions during pregnancy—the frequent variations in the sound which are characteristic of it.² The uterine souffle is also audible after the death of the foetus, and it is believed by some to be modified and to become more continuously harsh when that event has taken place.

Theories as to its Cause.—Very various explanations have been given

[¹ This instrument was the invention of the late Dr. G. P. Cammann of New York, and thirty years ago was known as the Cammann stethoscope. Dr. C. was an expert in physical explorations of the chest, and devoted his life to the study of its diseases.—Ed.]

² *Op. cit.*, p. 223.

of the causes of this sound. For long it was supposed to be formed in the vessels of the placenta, and hence the name "*placental souffle*" by which it is often talked of; or, if not in the placenta, in the uterine vessels in its immediate neighborhood. The non-placental origin of the sound is sufficiently demonstrated by the fact that it may be heard for a considerable time after the expulsion of the placenta. Some have supposed that it is not formed in the uterus at all, but in the maternal vessels, especially the aorta and the iliac arteries, owing to the pressure to which they are subjected by the gravid uterus. The extreme irregularity of the sound, its occasional disappearance, and its variable site, seem to be conclusive against this view. The theory which refers the sound to the uterine vessels is that which has received most adherents, and which best meets the facts of the case; but it is by no means easy, or even possible, to account for the exact mode of its production in them. Each of the explanations which have been given is open to some objection. It is far from unlikely that the intermittent contractions of the uterine fibres, which are known to occur during the whole course of pregnancy, may have much to do with it, by modifying, at intervals, the rapidity of the circulation in the vessels. Its production in this manner may also be favored by the chlorotic state of the blood, to which Cazeaux and Scanzoni are inclined to attribute an important influence, likening it to the anæmic murmur so frequently heard in the vessels in weakly women.

Its Diagnostic Value.—From a diagnostic point of view the uterine souffle is of very secondary importance, because a similar sound is very generally audible in large fibroid tumors of the uterus, and even in some few ovarian tumors; it is therefore of little or no value in assisting us to decide the character of the abdominal enlargement. The supposed dependence of the sound on the placental circulation has caused its site to be often identified with that of the placenta. It is, however, most frequently heard at the lower part of the uterus, while the placenta is generally attached near the fundus, so that its position cannot be taken as any safe guide in determining the situation of that organ.

Sounds produced by the Movements of the Fœtus.—Occasionally, in practising auscultation, irregular sounds of brief duration may be heard which are not susceptible of accurate description, and which doubtless depend on the sudden movement of the fœtus in the liquor amnii or on the impact of its limbs on the uterine walls. When heard distinctly they are characteristic of pregnancy, and they may be sometimes heard when the other sounds cannot be detected. They are, however, so irregular, and so often entirely absent, that they can hardly be looked upon in any other light than as occasional phenomena.

Sounds referred to Decomposition of the Liquor Amnii and to Separation of the Placenta.—Two other sounds have been described as being sometimes audible which may be mentioned as matters of interest, but which are of no diagnostic value. One is a rustling sound, said by Stoltz to be audible in cases in which the fœtus is dead, and which he refers to gaseous decomposition of the liquor amnii; its existence is, however, extremely problematical. The other is a sound heard after the birth of the child, and referred by Caillaut to the separation of the placental

adhesions. He describes it as a series of rapid, short scratching sounds, similar to those produced by drawing the nails across the seat of a horse-hair sofa. Simpson¹ admitted the existence of the sound, but believed that it is produced by the mere physical crushing of the placenta, and artificially imitated it out of the body by forcing the placenta through an aperture the size of the os uteri.

Relative Value of the Signs and Symptoms of Pregnancy.—It will be seen, then, that although there are numerous signs and symptoms accompanying pregnancy, many of them are unreliable by themselves, and apt to mislead. Those which may be confidently depended on are the pulsations of the fetal heart, which, however, fail us in cases of dead children; the fetal movements when distinctly made out; ballottement; the intermittent contractions of the uterus; and to these we may safely add the presence of milk in the breasts, provided we have to do with a first pregnancy.

The remainder are of importance in leading us to suspect pregnancy, and in corroborating and strengthening other symptoms, but they do not, of themselves, justify a positive diagnosis.

CHAPTER V.

THE DIFFERENTIAL DIAGNOSIS OF PREGNANCY. SPURIOUS PREGNANCY. THE DURATION OF PREGNANCY. SIGNS OF RECENT PREGNANCY.

Importance of the Subject.—The differential diagnosis of pregnancy has of late years assumed much importance on account of the advance of abdominal surgery. The cases are so numerous in which even the most experienced practitioners have fallen into error, and in which the abdomen has been laid open in ignorance of the fact that pregnancy existed, that the subject becomes one of the greatest consequence. Fortunately, it is less so from an obstetrical than from a gynæcological point of view, inasmuch as the converse error, of mistaking some other condition for pregnancy, is of far less consequence, as it is one which time will always rectify. But even in this way carelessness may lead to very serious injury to the character, if not to the health, of the patient; and it will be well to refer briefly to some of the conditions most liable to be mistaken for pregnancy, and to the mode of distinguishing them.

Adipose enlargement of the abdomen may obscure the diagnosis by preventing the detection of the uterus; and if, as is not uncommon with women of great obesity, it is associated with irregular menstruation, the increased size of the abdomen might be supposed to depend on pregnancy. The absence of corroborative signs, such as auscultatory phenomena, mammary changes, and the hardness of the cervix as felt per vaginam, makes it easy to avoid this error.

¹ *Selected Obstet. Works*, p. 151.

Distension of the Uterus by Retained Menses, Hydrometra, etc.—Distension of the uterus by retained menstrual fluid or watery secretion is an occurrence of rarity that could seldom give rise to error. Still, it occasionally happens that the uterus becomes enlarged in this way, sometimes reaching even to the level of the umbilicus, and that the physical character of the tumor is not unlike that of the gravid uterus. The best safeguard against mistakes will be the previous history of the case, which will always be different from that of ordinary pregnancy. Retention of the menses almost always occurs from some physical obstruction to the exit of the fluid, such as imperforate hymen; or if it occur in women who have already menstruated, we may usually trace a history of some cause, such as inflammation following an antecedent labor, which has produced occlusion of some part of the genital tract. The existence of a pelvic tumor in a girl who has never menstruated will of itself give rise to suspicion, as pregnancy under such circumstances is of extreme rarity. It will also be found that general symptoms have existed for a period of time considerably longer than the supposed duration of pregnancy, as judged of by the size of the tumor. The most characteristic of them are periodic attacks of pain due to the addition, at each monthly period, to the quantity of retained menstrual fluid. Whenever, from any of these reasons, suspicion of the true character of the case has arisen, a careful vaginal examination will generally clear it up. In most cases the obstruction will be in the vagina, and is at once detected, the vaginal canal above it, as felt per rectum, being greatly distended by fluid; and we may also find the bulging and imperforate hymen protruding through the vulva. The absence of mammary changes and of ballotement will materially aid us in forming a diagnosis.

Congestive Hypertrophy of the Uterus.—The engorged and enlarged uterus, frequently met with in women suffering from uterine disease, might readily be mistaken for an early pregnancy if it happened to be associated with amenorrhœa. A little time would, of course, soon clear up the point, by showing that progressive increase in size, as in pregnancy, does not take place. This mistake could only be made at an early stage of pregnancy, when a positive diagnosis is never possible. The accompanying symptoms—pain, inability to walk, and tenderness of the uterus on pressure—would prevent such an error.

Ascitic Distension of the Abdomen.—Ascites, per se, could hardly be mistaken for pregnancy, for the uniform distension and evident fluctuation, the absence of any definite tumor, the site of resonance on percussion changing in accordance with alteration of the position of the woman, and the unchanged cervix and uterus, should be sufficient to clear up any doubt. Pregnancy may, however, exist with ascites, and this combination may be difficult to detect, and might easily be mistaken for ovarian disease associated with ascites. The existence of mammary changes, the presence of the softened cervix, ballotement, and amenutation—provided the sounds were not masked by the surrounding fluid—would afford the best means of diagnosing such a case.

Uterine and Ovarian Tumors.—One of the most frequent sources of difficulty is the differential diagnosis of large abdominal tumors, either fibroid or ovarian, or of some enlargements due to malignant disease of

the peritoneum or abdominal viscera. The most experienced have been occasionally deceived under such circumstances. As a rule, the presence of menstruation will prevent error, as this generally continues in ovarian disease, while in fibroids it is often excessive. The character of the tumor—the fluctuation in ovarian disease, the hard nodular masses in fibroid—and the history of the case, especially the length of time the tumor has existed, will aid in diagnosis, while the absence of cervical softening and of auscultatory phenomena will further be of material value in forming a conclusion. Some of the most difficult cases to diagnose are those in which pregnancy complicates ovarian or fibroid disease. Then the tumor may more or less completely obscure the physical signs of pregnancy. The usual shape of the abdomen will generally be altered considerably, and we may be able to distinguish the gravid uterus, separated from the ovarian tumor by a distinct sulcus, or with the fibroid masses cropping out from its surface. Our chief reliance must then be placed in the alteration of the cervix and in the auscultatory signs of pregnancy.

Spurious Pregnancy.—The condition most likely to give rise to errors is that very interesting and peculiar state known as *spurious pregnancy*. In this most of the usual phenomena of pregnancy are so strangely simulated that accurate diagnosis is often far from easy. There are hardly any of the more apparent symptoms of pregnancy which may not be present in marked cases of this kind. The abdomen may become prominent, the areolæ altered, menstruation arrested, and apparent fetal motions felt; and, unless suspicion is aroused and a careful physical examination made, both the patient and the practitioner may easily be deceived.

Cases in which Spurious Pregnancy occurs.—There is no period of the childbearing life in which spurious pregnancy may not be met with; but it is most likely to occur in elderly women about the climacteric period, when it is generally associated with ovarian irritation connected with the change of life; or in younger women, who are either very desirous of finding themselves pregnant, or who, being unmarried, have subjected themselves to the chance of being so. In all cases the mental faculties have much to do with its production, and there is generally either very marked hysteria or even a condition closely allied to insanity. Spurious pregnancy is by no means confined to the human race. It is well known to occur in many of the lower animals. Harvey related instances in bitches, either after unsuccessful intercourse or in connection with their being in heat, even when no intercourse had occurred. In such cases the abdomen swelled and milk appeared in the mammæ. Similar phenomena are also occasionally met with in the cow. In these instances, as in the human female, there is probably some morbid irritation of the ovarian system.

Its Signs and Symptoms.—The physical phenomena are often very well marked. The apparent enlargement is sometimes very great, and it seems to be produced by a projection forward of the abdominal contents due to depression of the diaphragm, together with rigidity of the abdominal muscles, and may even closely simulate the uterine tumor on palpation. After the climacteric it is frequently associated, as Gooch

pointed out, with an undue deposit of fat in the abdominal walls and omentum, so that there may be even some dulness on percussion, instead of resonance of the intestines. The fetal movements are curiously and exactly simulated, either by involuntary contractions of the abdominal walls or by the movement of flatus in the intestines. The patient also generally fancies that she suffers from the usual sympathetic disorders of pregnancy, and thus her account of her symptoms will still further tend to mislead.

Sometimes followed by Spurious Labor.—Not only may the supposed pregnancy continue, but at what would be the natural term of delivery all the phenomena of labor may supervene. Many authentic cases are on record in which regular pains came on, and continued to increase in force and frequency until the actual condition was diagnosed. Such mistakes, however, are only likely to happen when the statements of the patient have been received without further inquiry. When once an accurate examination has been made error is no longer possible.

Methods of Diagnosis.—We shall generally find that some of the phenomena of pregnancy are absent. Possibly menstruation, more or less irregular, may have continued. Examination per vaginam will at once clear up the case, by showing that the uterus is not enlarged and that the cervix is unaltered. It may then be very difficult to convince the patient or her friends that her symptoms have misled her, and for this purpose the inhalation of chloroform is of great value. As consciousness is abolished, the semi-voluntary projection of the abdominal muscles is prevented, the large apparent tumor vanishes, and the bystanders can be readily convinced that none exists. As the patient recovers the tumor again appears.

Duration of Pregnancy.—The duration of pregnancy in the human female has always formed a fruitful theme for discussion amongst obstetricians. The reasons which render the point difficult of decision are obvious. As the large majority of cases occur in married women, in whom intercourse occurs frequently, there is no means of knowing the precise period at which conception took place. The only datum which exists for the calculation of the probable date of delivery is the cessation of menstruation. It is quite possible, however, and indeed probable, that conception occurred, in a considerable number of instances, not immediately after the last period, but immediately before the proper epoch for the occurrence of the next. Hence, as the interval between the end of one menstruation and the commencement of the next averages 25 days, an error to that extent is always possible. Another source of fallacy is the fact, which has generally been overlooked, that even a single coitus does not fix the date of conception, but only that of insemination. It is well known that in many of the lower animals the fertilization of the ovule does not take place until several days after copulation, the spermatozoa remaining in the interval in a state of active vitality within the genital tract. It has been shown by Marion Sims that living spermatozoa exist in the cervical canal in the human female some days after intercourse. It is very probable, therefore, that in the human female, as in the lower animals, a considerable but unknown interval occurs between insemination and actual

impregnation, which may render calculations as to the precise duration of pregnancy altogether unreliable.

Average Time between Cessation of Menstruation and Delivery.—A large mass of statistical observations exists respecting the average duration of gestation which has been drawn up and collated from numerous sources. It would serve no practical purpose to reprint the voluminous tables on this subject that are contained in obstetrical works. They are based on two principal methods of calculation. First, we have the length of time between the cessation of menstruation and delivery. This is found to vary very considerably, but the largest percentage of deliveries occurs between the 274th and 280th day after the cessation of menstruation, the average day being the 278th; but in individual instances very considerable variations both above and below these limits are found to exist. Next we have a series of cases, from various sources, in which only one coitus was believed to have taken place. These are naturally always open to some doubt, but, on the whole, they may be taken as affording tolerably fair grounds for calculation. Here, as in the other mode of calculation, there are marked variations, the average length of time, as estimated from a considerable collection of cases, being 275 days after the single intercourse. It may therefore be taken as certain that there is no definite time which we can calculate on as being the proper duration of pregnancy, and consequently no method of estimating the probable date of delivery on which we can absolutely rely.

Methods of Predicting the Probable Date.—The prediction of the time at which the confinement may be expected is, however, a point of considerable practical importance, and one on which the medical attendant is always consulted. Various methods of making the calculation have been recommended. It has been customary in this country, according to the recommendation of Montgomery, to fix upon ten lunar months, or 280 days, as the probable period of gestation, and, as conception is supposed to occur shortly after the cessation of menstruation, to add this number of days to any day within the first week after the last menstrual period as the most probable period of delivery. As, however, 278 days are found to be the average duration of gestation after the cessation of menstruation, and as this method makes the calculation vary from 281 to 287 days, it is evidently liable to fix too late a date. Naegele's method was to count seven days from the first appearance of the last menstrual period, and then reckon backward three months as the probable date. Thus, if a patient last commenced to menstruate on August 10, counting in this way from August 17 would give May 17 as the probable date of delivery.

Matthews Duncan has paid more attention than any one else to the prediction of the date of delivery. His method of calculating is based on the fact of 278 days being the average time between the cessation of menstruation and parturition; and he claims to have had a greater average of success in his predictions than on any other plan. His rule is as follows: "Find the day on which the female ceased to menstruate, or the first day of being what she calls 'well.' Take that day nine months forward as 275, unless February is included, in which case it is taken as 273 days. To this add three days in the former case, or five if February

is in the count, to make up the 278. This 278th day should then be fixed on as the middle of the week, or, to make the prediction the more accurate, of the fortnight, in which the confinement is likely to occur, by which means allowance is made for the average variation of either excess or deficiency."

Various periodoscopes and tables for facilitating the calculation have been made. The periodoscope of Dr. Tyler Smith (sold by Messrs. John Smith & Co., 52 Long Acre) is very useful for reference in the consulting-room, giving at a glance a variety of information, such as the probable period of quickening, the dates for the induction of premature labor, etc. The following table, prepared by Dr. Protheroe Smith, is also easily read, and is very serviceable:

TABLE FOR CALCULATING THE PERIOD OF UTERO-GESTATION.¹

Nine Calendar Months.			Ten Lunar Months.	
From	To	Days.	To	Days.
January 1	September 30	273	October 7	280
February 1	October 31	273	November 7	280
March 1	November 30	275	December 5	280
April 1	December 31	275	January 5	280
May 1	January 31	276	February 4	280
June 1	February 28	273	March 7	280
July 1	March 31	274	April 6	280
August 1	April 30	273	May 7	280
September 1	May 31	273	June 7	280
October 1	June 30	273	July 7	280
November 1	July 31	273	August 7	280
December 1	August 31	274	September 6	280

Quickening a Fallacious Guide in estimating Date of Delivery.—The date at which the quickening has been perceived is relied on by many practitioners, and still more by patients, in calculating the probable date of delivery, as it is generally supposed to occur at the middle of pregnancy. The great variations, however, at the time at which this phenomenon is first perceived, and the difficulty which is so often experienced of ascertaining its presence with any certainty, render it a very fallacious guide. The only times at which the perception of quickening is likely to prove of any real value are when impregnation has occurred during lactation (when menstruation is normally absent), or when menstruation is so uncertain and irregular that the date of its last appearance cannot be ascertained. As quickening is most commonly felt during the fourth month—more frequently in its first than in its last fortnight—it may thus afford the only guide we can obtain, and that an uncertain one, for predicting the date of delivery.

Is Protraction of Gestation Possible?—From a medico-legal point of

¹ The above obstetric "Ready Reckoner" consists of two columns, one of calendar, the other of lunar, months, and may be read as follows: A patient has ceased to menstruate on July 1: her confinement may be expected at soonest about March 31 (*the end of nine calendar months*), or at latest on April 6 (*the end of ten lunar months*). Another has ceased to menstruate on January 20: her confinement may be expected on September 30, plus 20 days (*the end of nine calendar months*), at soonest, or on October 7, plus 20 days (*the end of ten lunar months*), at latest.

view the question of the possible protraction of pregnancy beyond the average time, and of the limits within which such protraction can be admitted, is of very great importance. The law on this point varies considerably in different countries. Thus, in France it is laid down that legitimacy cannot be contested until 300 days have elapsed from the death of the husband or the latest possible opportunity for sexual intercourse. This limit is also adopted by Austria, while in Prussia it is fixed at 302 days. In England and America no fixed date is admitted, but while 280 days is admitted as the "*legitimum tempus pariendo*," each case in which legitimacy is questioned is to be decided on its own merits. At the early part of the century the question was much discussed by the leading obstetricians in connection with the celebrated Gardner peerage case, and a considerable difference of opinion existed among them. Since that time many apparently perfectly reliable cases have been recorded in which the duration of gestation was obviously much beyond the average, and in which all sources of fallacy were carefully excluded.

Reliable Cases of Protraction.—Not to burden these pages with a number of cases, it may suffice to refer, as examples of protraction, to four well-known instances recorded by Simpson,¹ in which the pregnancy extended respectively to 336, 332, 319, and 324 days after the cessation of the last menstrual period. In these, as in all cases of protracted gestation, there is the possible source of error that impregnation may have occurred just before the expected advent of the next period. Making an allowance of 23 days in each instance for this, we even then have a number of days much above the average—viz. 313, 309, 296, and 301. Numerous instances as curious may be found scattered through obstetric literature. Indeed, the experience of most accoucheurs will parallel such cases, which may be more common than is generally supposed, inasmuch as they are only likely to attract attention when the husband has been separated from the wife beyond the average and expected duration of the pregnancy.

Protraction Common in the Lower Animals.—The evidence in favor of the possible prolongation of gestation is greatly strengthened by what is known to occur in the lower animals. In some of these, as in the cow and the mare, the precise period of insemination is known to a certainty, as only a single coitus is permitted. Many tables of this kind have been constructed, and it has been shown that there is in them a very considerable variation. In some cases in the cow it has been found that delivery took place 45 days, and in the mare 43 days, after the calculated date. Analogy would go strongly to show that what is known to a certainty to occur in the lower animals may also take place in the human female. The fact, indeed, is now very generally admitted; but we are still unable to fix with any degree of precision on the extreme limit to which protraction is possible. Some practitioners have given cases in which, on data which they believe to be satisfactory, pregnancy has been extremely protracted; thus, Meigs and Adler record instances which they believed to have been prolonged to over a year in one case and over fourteen months in the other. These are, however, so problematical that little weight can be attached to them. On the whole, it

¹ *Obstet. Memoirs*, p. 84.

would hardly be safe to conclude that pregnancy can go more than three or four weeks beyond the average time. This conclusion is justified by the cases we possess in which pregnancy followed a single coitus, the longest of which was 295 days.

Evidence from Size of Child.—Dr. Duncan¹ is inclined to refuse credence to every case of supposed protraction unless the size and weight of the child are above the average, believing that lengthened gestation must of necessity cause increased growth of the child. This point requires further investigation, and it cannot be taken as proved that the fœtus necessarily must be large because it has been retained longer than usual in utero; or, even if this be admitted, it may have been originally small, and so at the end of the protracted gestation be little above the average weight. There are, however, many cases which certainly prove that a prolonged pregnancy is at least often associated with an unusually-developed fœtus. Dr. Duncan himself cites several, and a very interesting one is mentioned by Leishman in which delivery took place 295 days after a single coitus, the child weighing 12 lbs. 3 oz.

In some cases Labor may Commence and be Arrested.—It seems possible that, in some cases of protracted pregnancy, labor actually came on at the average time, but, on account of faulty positions of the uterus or other obstructing cause, the pains were ineffective and ultimately died away, not recurring for a considerable time. Joulin relates some instances of this kind. In one of them the labor was expected from the 20th to the 25th of October. He was summoned on the 23d, and found the pains regular and active, but ineffective; after lasting the whole of the 24th and 25th they died away, and delivery did not take place until November 25th, after the lapse of a month. In this instance the apparent cause of difficulty was extreme anterior obliquity of the uterus. A precisely similar case came under my own observation. The lady ceased to menstruate on March 16, 1870. On December 12th—that is, on the 273d day—strong labor-pains came on, the os dilated to the size of a florin, and the membranes became tense and prominent with each pain. After lasting all night they gradually died away, and did not recur until January 12th, 304 days from the cessation of the last period. Here there was no assignable cause of obstruction, and the labor, when it did come on, was natural and easy.

The curious fact that in both these cases, as in others of the same kind that are recorded, labor came on exactly a month after the previous ineffectual attempt at its establishment, affords, so far as it goes, an argument in favor of the view maintained by many that labor is apt to come on at what would have been a menstrual period.

Signs of Recent Delivery.—From a forensic point of view it often becomes of importance to be able to give a reliable opinion as to the fact of delivery having occurred, and a few words may be here said as to the signs of recent delivery. Our opinion is only likely to be sought in cases in which the fact of delivery is denied, and in which we must, therefore, entirely rely on the results of a physical examination. If this be undertaken within the first fortnight after labor, a positive conclusion can be readily arrived at.

¹ *Fecundity and Fertility*, p. 348.

At this time the abdominal walls will still be found loose and flaccid, and bearing very evident marks of extreme distension in the cracks and fissures of the cutis vera. These remain permanent for the rest of the patient's life, and may be safely assumed to be signs of an antecedent pregnancy, provided we can be certain that no other cause of extreme abdominal distension has existed, such as ascites or ovarian tumor.

Within the first few days after delivery the hard round ball formed by the contracted and empty uterus can easily be felt by abdominal palpation, and more certainly by combined external and internal examination. The process of involution, however, by which the uterus is reduced to its normal size, is so rapid that after the first week it can no longer be made out above the brim of the pelvis. In cases in which an accurate diagnosis is of importance the increased length of the uterus can be ascertained by the uterine sound, and its cavity will measure more than the normal $2\frac{1}{2}$ inches for at least a month after delivery. It should not be forgotten that the uterine parietes are now undergoing fatty degeneration, and that they are more than usually soft and friable, so that the sound should be used with great caution and only when a positive opinion is essential. The state of the cervix and of the vagina may afford useful information. Immediately after delivery the cervix hangs loose and patulous in the vagina, but it rapidly contracts, and the internal os is generally entirely closed after the eighth or tenth day. The remainder of the cervix is longer in returning to its normal shape and consistency. It is generally permanently altered after delivery, the external os remaining fissured and transverse, instead of circular with smooth margins, as in virgins. The vagina is at first lax, swollen, and dilated, but these signs rapidly disappear, and cannot be satisfactorily made out after the first few days. The absence of the fourchette may be recognized, and is a persistent sign.

The presence of the lochia affords a valuable sign of recent delivery. For the first few days they are sanguineous, and contain numerous blood-corpuscles, epithelial scales, and the debris of the decidua. After the fifth day they generally change in color and become pale and greenish, and from the eighth or ninth day till about a month after delivery they have the appearance of thick opalescent mucus. They have, however, a peculiar, heavy, sickening odor, which should prevent their being mistaken for either menstruation or leucorrhœal discharge.

The appearance of the breasts will also aid the decision, for it is impossible for the patient to conceal the turgid, swollen condition of the mammæ, with the darkened areolæ, and, above all, the presence of milk. If, on microscopic examination, the milk is found to contain colostrum-corpuscles, the fact of very recent delivery is certain. In women who do not nurse it should be remembered that the secretion of milk often rapidly disappears, so that its absence cannot be taken as a sign that delivery has not taken place. On the whole, there should be no difficulty in deciding that a woman has been delivered, as some of the signs are persistent for the rest of her life; but it is not so easy, unless we see the case within the first eight or ten days, to say how long it is since labor took place.

CHAPTER VI.

ABNORMAL PREGNANCY, INCLUDING MULTIPLE PREGNANCY, SUPERFŒTATION, EXTRA-UTERINE FŒTATION, AND MISSED LABOR.

Plural Births an Abnormal Variety of Pregnancy.—The occurrence of more than one foetus in utero is far from uncommon, but there are circumstances connected with it which justify the conclusion that plural births must not be classified as natural forms of pregnancy. The reasons for this statement have been well collected by Dr. Arthur Mitchell,¹ who conclusively shows that not only is there a direct increase of risk both to the mother and her offspring, but that many abnormalities, such as idiocy, imbecility, and bodily deformity, occur with much greater frequency in twins than in single-born children. He concludes that “the whole history of twin births is exceptional, indicates imperfect development and feeble organization in the product, and leads us to regard twinning in the human species as a departure from the physiological rule, and therefore injurious to all concerned.”

Frequency of Multiple Births.—The frequency of multiple births varies considerably under different circumstances. Taking the average of a large number of cases collected by authors in various countries, we find that twin pregnancies occur about once in 87 labors; triplets, once in 7679. A certain number of quadruple pregnancies, and some cases of early abortion in which there were five foetuses, are recorded, so that there can be no doubt of the possibility of such occurrences; but they are so extremely uncommon that they may be looked upon as rare exceptions, the relative frequency of which can hardly be determined.

The frequency of multiple pregnancy varies remarkably in different races and countries. The following table² will show this at a glance:

RELATIVE FREQUENCY OF MULTIPLE PREGNANCIES IN EUROPE.

Countries.	Proportion of Twin to Single Births.	Proportion of Triplets.	Proportion of Quadruplets.
England	1 : 116	1 : 6,720	
Austria	1 : 94		
Grand Duchy of Baden	1 : 89	1 : 6,575	
Scotland	1 : 95		
France	1 : 99	1 : 8,256	1 : 2,074,306
Ireland	1 : 64	1 : 4,995	1 : 167,226
Mecklenburg-Schwerin	1 : 68.9	1 : 6,436	1 : 183,236
Norway	1 : 81.62	1 : 5,442	
Prussia	1 : 89	1 : 7,820	1 : 394,690
Russia	1 : 50.05	1 : 4,054	
Saxony	1 : 79	1 : 1,000	1 : 400,000
Switzerland	1 : 102		
Württemberg	1 : 862	1 : 6,464	1 : 110,991

¹ *Med. Times and Gaz.*, Nov., 1862.² Puech, *Des Naissances Multiples.*

It will be seen that the largest proportion of multiple births occurs in Russia, and that the number of triple births is greatest where twin pregnancies are most frequent. Puech concludes that the number of multiple pregnancies is in direct proportion to the general fecundity of the inhabitants.

Dr. Duncan has deduced some interesting laws with regard to the production of twins from a large number of statistical observations;¹ especially, that the tendency to the production of twins increases as the age of the woman advances, and is greater in each succeeding pregnancy, exception being made for the first pregnancy, in which it is greater than in any other. Newly-married women appear more likely to have twins the older they are. There can be no doubt that there is often a strong hereditary tendency in individual families to multiple births. A remarkable instance of this kind is recorded by Mr. Curgenvén,² in which a woman had four twin pregnancies, her mother and aunt each one, and her grandmother two. Simpson mentions a case of quadruplets, consisting of three males and one female, who all survived, the female subsequently giving birth to triplets.³

Sex of Children.—In the largest number of cases of twins the children are of opposite sexes, next most frequently there are two females, and twin males are the most uncommon. Thus, out of 59,178 labors, Simpson calculates that twin male and female occurred once in 199 labors, twin females once in 226, and twin males once in 258. The proportion of male to female births is also notably less in twin than in single pregnancies.

Size of Fœtuses.—Twins, and, *a fortiori*, triplets, are almost always smaller and less perfectly developed than single children. Hence the chances of their survival are much less, and Clarke calculates the mortality amongst twin children as one out of thirteen. Of triplets, indeed, it is comparatively rare that all survive; while in quadruplets premature labor and the death of the fœtuses are almost certain. It is a common observation that twins are often unequally developed at birth. By some this difference is attributed to one of them being of a different age to the other. It is probable, however, that in most of these cases the full development of one fœtus has been interfered with by pressure of the other. This is far from uncommonly carried to the extent of destroying one of the twins, which is expelled at term, mummified and flattened between the living child and the uterine wall. In other cases, when one fœtus dies it may be expelled without terminating the pregnancy, the other being retained in utero and born at term; and those who disbelieve in the possibility of superfœtation explain in this way the cases in which it is believed to have occurred.

Causes.—Multiple pregnancies depend on various causes. The most common is probably the simultaneous, or nearly simultaneous, maturation and rupture of two Graafian follicles, the ovules becoming impregnated at or about the same time. It by no means necessarily follows, even if more than one follicle should rupture at once, that both ovules should be impregnated. This is proved by the occurrence of cases in

¹ *On Fecundity, Fertility, and Sterility*, p. 99.

² *Obst. Trans.*, vol. xi.

³ *Obst. Works*, p. 830.

which there are two corpora lutea with only one foetus. There are numerous facts to prove that ovules thrown off within a short time of each other may become separately impregnated, as in cases in which negro women have given birth to twins, one of which was pure negro, the other half-caste.

It may happen, however, that a single Graafian follicle contains more than one ovule, as has actually been observed before its rupture; or, as is not uncommon in the egg of the fowl, an ovule may contain a double germ, each of which may give rise to a separate foetus.

Arrangement of the Fœtal Membranes and Placentæ.—The various modes in which twins may originate explain satisfactorily the variations which are met with in the arrangement of the fœtal membranes and in the form and connections of the placentæ. In a large proportion of cases there are two distinct bags of membranes, the septum between them being composed of four layers—viz. the chorion and amnion of each ovum. The placentæ are also entirely separate. Here it is obvious that each twin is developed from a distinct ovum, having its own chorion and amnion. On arriving in the uterus it is probable that each ovum becomes fixed independently in the mucous membrane and is surrounded by its own decidua reflexa. As growth advances the decidua reflexa generally atrophies from pressure, as it is not usual to find more than four layers of membrane in the septum separating the ova. In other cases there is only one chorion, within which are two distinct amnions, the septum then consisting of two layers only. Then the placentæ are generally in close apposition and become fused into a single mass, the cords, separately attached to each foetus, not infrequently uniting shortly before reaching the placental mass, their vessels anastomosing freely. In other more rare instances both foetuses are contained in a common amniotic sac; but as the amnion is a purely fœtal membrane, it is probable that, when this arrangement is met with, the originally existing septum between the amniotic sacs has been destroyed. In both these latter cases the twins must have been developed from a single ovule containing a double germ, and Schroeder states that they are then always of the same sex. Dr. Brunton¹ has started a precisely opposite theory, and has tried to prove that twins of the same sex are contained in separate bags of membrane, while twins of opposite sexes have a common sac. He says that out of 25 cases coming under his observation, in 15 the children contained in different sacs were of the same sex, but in the remaining 10, in which there was only one sac, they were of opposite sexes. It is difficult to believe that there is not an error in these observations, since twins contained in a single amniotic sac do not occur nearly as often as ten times out of twenty-five cases, and no distinction is made between a common chorion with two amnions and a single chorion and amnion. The facts of double monstrosity also disprove this view, since conjoined twins must of necessity arise from a single ovule with a double germ, and there is no instance on record in which they were of opposite sexes.

Membranes and Placentæ in Triplets.—In triplets the membranes and placentæ may be all separate, or, as is commonly the case, there is one

¹ *Obst. Trans.*, vol. x.

complete bag of membranes, and a second having a common chorion, with a double amnion. It is probable, therefore, that triplets are generally developed from two ovules, one of which contained a double germ.

Diagnosis of Multiple Pregnancy.—It is comparatively seldom that twin pregnancy can be diagnosed before the birth of the first child, and even when suspicion has arisen its indications are very defective. There is generally an unusual size and an irregularity of shape of the uterus, sometimes even a distinct depression or sulcus between the two foetuses. When such a sulcus exists, it may be possible to make out parts of each foetus by palpation on either side of the uterus. The only sign, however, on which the least reliance can be placed is the detection of two foetal hearts. If two distinct pulsations are heard at different parts of the uterus; if, on carrying the stethoscope from one point to another, there is an interspace where pulsations are no longer audible, or when they become feeble and again increase in clearness as the second point is reached; and, above all, if we are able to make out a difference in frequency between them,—the diagnosis is tolerably safe. It must be remembered, however, that the sounds of a single heart may be heard over a larger space than usual, and hence a possible source of error. Twin pregnancy, moreover, may readily exist without the most careful auscultation enabling us to detect a double pulsation, especially if one child lie in the dorso-posterior position, when the body of the other may prevent the transmission of its heart's beat. The so-called placental souffle is generally too diffuse and irregular to be of any use in diagnosis even when it is distinctly heard at separate parts of the uterus.

Superfoetation and Superfecundation.—Closely connected with the subject of multiple pregnancies are the conditions known as *superfecundation* and *superfoetation*, regarding which there has been much controversy and difference of opinion.

By the former is meant the fecundation, at or near the same period of time, of two separate ovules before the decidua lining the uterus has been formed, which by many is supposed to form an insuperable obstacle to subsequent impregnation. The possibility of this occurrence has been incontestably proved by the class of cases already referred to, in which the same woman has given birth to twins bearing evident traces of being the offspring of fathers of different races.

By *superfoetation* is meant the impregnation of a second ovule, when the uterus already contains an ovum which has arrived at a considerable degree of development. The cases which are supposed to prove the possibility of this occurrence are very numerous. They are those in which a woman is delivered simultaneously of foetuses of very different ages, one bearing all the marks of having arrived at term, the other of prematurity; or of those in which a woman is delivered of an apparently mature child, and, after the lapse of a few months, of another equally mature. The possibility of superfoetation is strongly denied by many practitioners of eminence, and explanations are given which doubtless seem to account satisfactorily for a large proportion of the supposed examples. In the former class of cases it is supposed, with much probability, that there is an ordinary twin pregnancy, the development of one foetus being retarded by the presence in utero of another. That this is

not an uncommon occurrence is certain, and the fact has already been alluded to in treating of twin pregnancy. In cases of the latter kind it is possible that some of them may be due to separate impregnation in a bilobed uterus, the contents of one division being thrown off a considerable time before those of the other. Numerous authentic examples of this occurrence are recorded, but by far the most remarkable is that related by Dr. Ross of Brighton, which has been already referred to (p. 67). In this case the patient had previously given birth to many children without any suspicion of her abnormal formation having arisen, and, had it not been detected by Dr. Ross, the case might fairly enough have been claimed as an indubitable example of superfetation.

Making every allowance for these explanations, there remain a considerable number of cases which it is very difficult to account for, except on the supposition that the second child has been conceived a considerable time after the first. Those interested in the subject will find a large number of examples collected in a valuable paper by Dr. Bonnar of Cupar.¹ He has adopted the ingenious plan of consulting the records of the British peerage, where the exact date of the birth of successive children of peers is given, without, of course, any reasonable possibility of error, and he has collected numerous examples of births rapidly succeeding each other which are apparently inexplicable on any other theory. In one case he cites a child was born September 12, 1849, and the mother gave birth to another on January 24, 1850, after an interval of only 127 days. Subtracting from that 14 days, which Dr. Bonnar assumes to be the earliest possible period at which a fresh impregnation can occur after delivery, we reduce the gestation to 113 days—that is, to less than four calendar months. As both these children survived, the second child could not possibly have been the result of a fresh impregnation after the birth of the first; nor could the first child have been a twin prematurely delivered, for if so, it must have only reached rather more than the fifth month, at which time its survival would have been impossible.


Besides the numerous examples of cases of this kind recorded in most obstetric works, there are one or two of miscarriage in the early months, in which, in addition to a fœtus of four or five months' growth, a perfectly fresh ovum of not more than a month's development was thrown off. One such case was shown at the Obstetrical Society in 1862, which was reported on by Drs. Harley and Tanner, who stated that in their opinion it was an example of superfetation. A still more conclusive case is recorded by Tyler Smith:² "A young married woman, pregnant for the first time, miscarried at the end of the fifth month, and some hours afterward a small clot was discharged, enclosing a perfectly healthy ovum of about one month. There were no signs of a double uterus in this case. The patient had menstruated regularly during the time she had been pregnant." This case is of special interest from the fact of the patient having menstruated during pregnancy—a circumstance only explicable on the same anatomical grounds which render superfetation possible. So far as I know, it is the only instance in which the coincidence of superfetation and menstruation during early pregnancy has been observed.

¹ *Edin. Med. Journ.*, 1864-65.

² *Manual of Obstetrics*, p. 112.

Objections to the Admission of Superfetation.—The objections to the possibility of superfetation are based on the assumptions that the decidua so completely fills up the uterine cavity that the passage of the spermatozoa is impossible; that their passage is prevented by the mucous plug which blocks up the cervix; and that when impregnation has taken place ovulation is suspended. It is, however, certain that none of these are insuperable obstacles to a second impregnation. The first was originally based on the older and erroneous view which considered the decidua to be an exudation lining the entire uterine cavity and sealing up the mouths of the Fallopian tubes and the aperture of the internal os uteri. The decidua reflexa, however, does not come into apposition with the decidua vera until about the eighth week of pregnancy, and, therefore, until that time there is a free space between the two membranes through which the spermatozoa might pass to the open mouths of the Fallopian tube, and in which a newly-impregnated ovule might graft itself. A reference to the accompanying figure of a pregnancy in the third month, copied from Coste's work, will readily show that, as far as the decidua is concerned, there is no mechanical obstacle to the descent and lodgment of another impregnated ovule (Fig. 80). Then, as regards

FIG. 80.



Illustrating the Cavity between the Decidua Vera and the Decidua Reflexa during the Early Months of Pregnancy. (After Coste.)

the plug of mucus, it is pretty certain that this is in no way different from the mucus filling the cervix in the non-pregnant state, which offers no obstacle at all to the passage of the spermatozoa. Lastly, respecting the cessation of ovulation during pregnancy, this, no doubt, is the rule, and probably satisfactorily explains the rarity of superfetation. There are, however, a sufficient number of authenticated cases of menstruation during pregnancy to prove that ovulation is not always absolutely in

abeyance; and, as long as it occurs, there is unquestionably no positive mechanical obstruction, at least in the early months of pregnancy, in the way of the impregnation and lodgment of the ovules that are thrown off. The reasonable conclusion, therefore, seems to be that, although a large majority of the supposed cases are explicable in other ways, it cannot be admitted that superfetation is either physiologically or mechanically impossible.

Extra-uterine Pregnancy.—The most important of the abnormal varieties of pregnancy, if we consider the serious and very generally fatal results attending it, is the so-called *extra-uterine fetation*, which consists in the arrest and development of the ovum outside the cavity of the uterus. Of late years this subject has received much well-merited attention, which, it is to be hoped, may lead to the establishment of some definite rules for the management of this most anxious and dangerous class of cases.

Site of Extra-uterine Pregnancy.—The ovum may be arrested and developed in various situations on its way to the uterus, most commonly in some part of the Fallopian tube, or it may be in the cavity of the abdomen, or even quite beyond it, as in a few rare cases in which the ovum has found its way into a hernial sac.

Classification.—Extra-uterine gestation may be subdivided into the following classes: 1st, and most common of all, *Tubal* gestation, and, as varieties of this, although by some made into distinct classes, (a) *interstitial* and (b) *tubo-ovarian* gestation. In the former of these subdivisions the ovum is arrested in the part of the Fallopian tube that is situated in the substance of the uterine parietes; in the latter, at or near the fimbriated extremity of the tube, so that part of its cyst is formed by the tube and part by the ovary. 2d. *Abdominal* gestation, in which an ovum, instead of finding its way into the tube, falls into the peritoneal cavity, and there becomes attached and developed; or the so-called secondary abdominal gestation, in which an extra-uterine pregnancy, originally tubal, becomes ventral through rupture of its cyst and escape of its contents into the abdominal cavity. 3d. *Ovarian* gestation, the existence of which is denied by many writers of eminence, such as Velpéau and Arthur Farre, while it is maintained by others of equal celebrity, such as Kiwisch, Coste, and Hecker. It must be admitted that it is extremely difficult to understand how an ovarian pregnancy, in the strict sense of the word, can occur, for it implies that the ovule has become impregnated before the laceration of the Graafian follicle, through the coats of which the spermatozoa must have passed. Coste, indeed, believes that this frequently happens; but, while spermatozoa have been detected on the surface of the ovary, their penetration into the Graafian follicle has never been demonstrated. Farre has also clearly shown that in many cases of supposed ovarian pregnancy the surrounding structures were so altered that it was impossible to trace their exact origin and to say to a certainty that the fetus was really within the substance of the ovary. Kiwisch gives a reasonable explanation of these cases by supposing that sometimes the Graafian follicle may rupture, but that the ovule may remain within it without being discharged. Through the rent in the walls of the follicle the spermatozoa may reach and impreg-

nate the ovule, which may develop in the situation in which it has been detained. The subject has been recently ably considered by Puech,¹ who admits two varieties of ovarian pregnancy, according as the fœtus has developed in a vesicle which has remained open or in one which has closed immediately after fecundation. He considers that most cases of so-called ovarian pregnancy are either dermoid cysts, ovario-tubal pregnancies, or abdominal pregnancies in which the placenta is attached to the ovary, and that even in the rare cases of true ovarian pregnancies the progress and results do not differ from those of abdominal pregnancy. While, therefore, it is impossible to deny the existence of ovarian pregnancy, it must be considered to be a very rare and exceptional variety, which, as far as treatment and results are concerned, does not differ from tubular or abdominal gestation. 4th. There are two rare varieties in which an ovum is developed either in the supplementary horn of a *bilobed uterus* or in a *hernial sac*.

For the sake of clearness, we may place these varieties of extra-uterine gestation in the following tabular form:

1st. *Tubal*—

(a) Interstitial, (b) Tubo-ovarian.

2d. *Abdominal*—

(a) Primary, (b) Secondary.

3d. *Ovarian*.

4th. In *bilobed uterus*, *hernial*, etc.

Causes.—The etiology of extra-uterine fœtation in any individual case must necessarily be almost always obscure. Broadly speaking, it may be said that extra-uterine fœtation may be produced by any condition which prevents or renders difficult the passage of the ovule to the uterus, while it does not prevent the access of the spermatozoa to the ovule. Thus, inflammatory thickening of the coats of the Fallopian tubes by lessening their calibre, but not sufficiently so to prevent the passage of the spermatozoa, may interfere with the movements of the tube which propel the ovum forward, and so cause its arrest. A similar effect may be produced by various morbid conditions, such as inflammatory adhesions from old-standing peritonitis pressing on the tube, obstruction of its calibre by inspissated mucus or small polypoid growths, the pressure of uterine or other tumors, and the like. The fact that extra-uterine pregnancies occur most frequently in multiparæ, and comparatively rarely in women under thirty years of age, tends to show that these conditions, which are clearly more likely to be met with in such women than in young primiparæ, have considerable influence in its causation. A curiously large proportion of cases occur in women who have either been previously altogether sterile or in whom a long interval of time has elapsed since their last pregnancy. The disturbing effects of fright, either during coition or a few days afterward, have been insisted on by many authors as a possible cause. Numerous cases of this kind are recorded; and, although the influence of emotion in the production of this condition is not susceptible of proof, it is not difficult to imagine that spasms of the Fallopian tubes might be produced in this way which would either interfere with the passage of the ovum or direct

¹ *Annal. de Gynéc.*, July, 1878.

abeyance; and, as long as it occurs, there is unquestionably no positive mechanical obstruction, at least in the early months of pregnancy, in the way of the impregnation and lodgment of the ovules that are thrown off. The reasonable conclusion, therefore, seems to be that, although a large majority of the supposed cases are explicable in other ways, it cannot be admitted that superfetation is either physiologically or mechanically impossible.

Extra-uterine Pregnancy.—The most important of the abnormal varieties of pregnancy, if we consider the serious and very generally fatal results attending it, is the so-called *extra-uterine foetation*, which consists in the arrest and development of the ovum outside the cavity of the uterus. Of late years this subject has received much well-merited attention, which, it is to be hoped, may lead to the establishment of some definite rules for the management of this most anxious and dangerous class of cases.

Site of Extra-uterine Pregnancy.—The ovum may be arrested and developed in various situations on its way to the uterus, most commonly in some part of the Fallopian tube, or it may be in the cavity of the abdomen, or even quite beyond it, as in a few rare cases in which the ovum has found its way into a hernial sac.

Classification.—Extra-uterine gestation may be subdivided into the following classes: 1st, and most common of all, *Tubal* gestation, and, as varieties of this, although by some made into distinct classes, (*a*) *interstitial* and (*b*) *tubo-ovarian* gestation. In the former of these subdivisions the ovum is arrested in the part of the Fallopian tube that is situated in the substance of the uterine parietes; in the latter, at or near the fimbriated extremity of the tube, so that part of its cyst is formed by the tube and part by the ovary. 2d. *Abdominal* gestation, in which an ovum, instead of finding its way into the tube, falls into the peritoneal cavity, and there becomes attached and developed; or the so-called secondary abdominal gestation, in which an extra-uterine pregnancy, originally tubal, becomes ventral through rupture of its cyst and escape of its contents into the abdominal cavity. 3d. *Ovarian* gestation, the existence of which is denied by many writers of eminence, such as Velpeau and Arthur Farre, while it is maintained by others of equal celebrity, such as Kiwisch, Coste, and Hecker. It must be admitted that it is extremely difficult to understand how an ovarian pregnancy, in the strict sense of the word, can occur, for it implies that the ovule has become impregnated before the laceration of the Graafian follicle, through the coats of which the spermatozoa must have passed. Coste, indeed, believes that this frequently happens; but, while spermatozoa have been detected on the surface of the ovary, their penetration into the Graafian follicle has never been demonstrated. Farre has also clearly shown that in many cases of supposed ovarian pregnancy the surrounding structures were so altered that it was impossible to trace their exact origin and to say to a certainty that the foetus was really within the substance of the ovary. Kiwisch gives a reasonable explanation of these cases by supposing that sometimes the Graafian follicle may rupture, but that the ovule may remain within it without being discharged. Through the rent in the walls of the follicle the spermatozoa may reach and impreg-

rially different according to their site, and, for practical purposes, we may consider them as forming two great classes, the tubal (with its varieties) and the abdominal.

Tubal Pregnancies.—When the ovum is arrested in any part of the Fallopian tube, the chorion soon commences to develop villi, just as in ordinary pregnancy, which engraft themselves into the mucous lining of the tube and fix the ovum in its new position. The mucous membrane becomes hypertrophied, much in the same way as that of the uterus under similar circumstances, so that it becomes developed into a sort of pseudo-decidua. Inasmuch, however, as the mucous coat of the tubes is not furnished with tubular glands, a true decidua can scarcely be said to exist, nor is there any growth of membrane around the ovum analogous to the decidua reflexa. The ovum is therefore, comparatively speaking, loosely attached to its abnormal situation, and hence hemorrhage from laceration of the chorion villi can very readily take place.

It is seldom that any development of the chorion villi into distinct placental structure is observed; this is probably owing to the fact that laceration and death generally occur before the period at which the placenta is normally formed. The muscular coat of the tube soon becomes hypertrophied, and as the size of the ovum increases the fibres are separated from each other, so that the ovum protrudes at

FIG. 82.



Tubal Pregnancy. (From a Specimen in the Museum of King's College.)

certain points through them, and at these it is only covered by the stretched and attenuated mucous and peritoneal coats of the tube. At this time the tubal pregnancy forms a smooth oval tumor, which, as a rule, has not formed any adhesions to the surrounding structures (Fig. 82). The part of the tube unoccupied by the ovum may be

found unaltered, and permeable in both directions; or, more frequently, it becomes so stretched and altered that its canal cannot be detected. Most frequently it is that part of the tube nearest the uterus which cannot be made out. The condition of the uterus in this as in other forms of extra-uterine pregnancy has been the subject of considerable discussion. It is now universally admitted that the uterus undergoes a certain amount of sympathetic engorgement, the cervix becomes softened, as in natural pregnancy, and the mucous membrane develops into a true decidua. In many cases the decidua is found on post-mortem examination, in others it is not, and hence the doubts that some have expressed as to its existence. The most reasonable explanation of its absence is that given by Duguet,¹ who has shown that it is far from uncommon for the uterine decidua to be thrown off *en masse* during the hemorrhagic discharges which so frequently precede the fatal issue of extra-uterine gestation.

Interstitial and False Ovarian Pregnancy.—When the ovum is arrested in that portion of the tube passing through the uterus in so-called interstitial pregnancy, the muscular fibres of the uterus become stretched and distended, and form the outer covering of the ovum. When, on the other hand, the site of arrest is in the fimbriated extremity of the tube, the containing cyst is formed partly of the fimbriae of the tube, partly of ovarian tissue; hence it is much more distensible, and the pregnancy may continue without laceration to a more advanced period, or even to term, so that when the ovum is placed in this situation the case much more nearly resembles one of abdominal pregnancy.

Period at which Rupture occurs.—The termination of tubal pregnancy, in the immense majority of cases, is death, produced by laceration giving rise either to internal hemorrhage or to subsequent intense peritonitis. Rupture usually occurs at an early period of pregnancy, most generally from the fourth to the twelfth week, rarely later. However, a few instances are recorded in which it did not take place until the fourth or fifth month, and Saxtorph and Spiegelberg have recorded apparently authentic cases in which the pregnancy advanced to term without laceration. It is generally effected by distension of the tube, which at last yields at the point which is most stretched; and sometimes it seems to be hastened or determined by accidental circumstances, such as a blow or fall or the excitement of sexual intercourse.

Symptoms of Rupture.—The symptoms accompanying rupture are those of intense collapse, often associated with severe abdominal pain, produced by the laceration of the cyst. The patient will be found deadly pale, with a small, thready, and almost imperceptible pulse, perhaps vomiting, but with mental faculties clear. If the hemorrhage be considerable, she may die without any attempt at reaction. Sometimes, however—and this generally occurs in cases in which the tube tears, the ovum remaining intact—the hemorrhage may cease on account of the ovum protruding through the aperture and acting as a plug. The patient may then imperfectly rally, to be again prostrated by a second escape of blood, which proves fatal. If the loss of blood is not of itself sufficient to cause death from shock and anæmia, the fatal issue is generally only

¹ *Annales de Gynécologie*, May, 1874.

postponed, for the effused blood soon sets up a violent general peritonitis, which rapidly carries off the patient. If she should survive the second danger, the case is transformed into one of abdominal pregnancy, the fœtus becoming surrounded by a capsule produced by inflammatory exudation (Fig. 83). The case is then subjected to the rules of treatment

FIG. 83.



Extra-uterine Pregnancy at Term of the Tubo-ovarian Variety. (After a Case of Dr. A. Sibley Campbell's.)

presently to be discussed when considering that variety of extra-uterine gestation.

Diagnosis.—The possibility of diagnosing tubal gestation before rupture occurs is a question of great and increasing interest, from the fact that, could its existence be ascertained, we might very fairly hope to avert the almost certainly fatal issue which is awaiting the patient. Unfortunately, the symptoms of tubal pregnancy are always obscure, and too often death occurs without the slightest suspicion as to the nature of the case having arisen. In the first place, it is to be observed that all the usual sympathetic disturbances of pregnancy exist: the breasts enlarge, the areolæ darken, and morning sickness is present. There is also an arrest of menstruation, but after the absence of one or more periods there is often an irregular hemorrhagic discharge. This is an important symptom, the value of which in indicating the existence of tubal pregnancy has of late years been much dwelt upon by various authors, both

in this country and abroad. Barnes attributes it to partial detachment of the chorion villi, produced by the ovum growing out of proportion to the tube in which it is contained. Whether this is the correct explanation or not, it is a fact that irregular hemorrhage very generally precedes the laceration for several days or more. Accompanying this hemorrhage there is almost always more or less abdominal pain, produced by the stretching of the tissues in which the ovum is placed, and this is sometimes described as being of a very intense and crampy character. If, then, we meet with a case in which the symptoms of early pregnancy exist, in which there are irregular losses of blood, possibly discharge of membranous shreds, and abdominal pain, a careful examination should be insisted on, and then the true nature of the case may possibly be ascertained. Should extra-uterine foetation exist, we should expect to find the uterus somewhat enlarged and the cervix softened, as in early pregnancy, but both these changes are doubtless generally less marked than in normal pregnancy. This fact, of itself, however, is of little diagnostic value, for slight differences of this kind must always be too indefinite to justify a positive opinion.

Presence of a Peri-uterine Tumor.—The existence of a peri-uterine tumor, rounded or oval in outline, and producing more or less displacement of the uterus in the direction opposite to that in which the tumor is situated, may point to the existence of tubular foetation. By bimanual examination, one hand depressing the abdominal wall, while the examining finger of the other acts in concert with it either through the vagina or rectum, the size and relations of the growth may be made out. There are various conditions which give rise to very similar physical signs, such as small ovarian or fibroid growths or the effusion of blood around the uterus; and the differential diagnosis must always be very difficult, and often impossible. A curious example of the difficulty of diagnosis is recorded by Joulin, in which Huguier and six or seven of the most skilled obstetricians of Paris agreed on the existence of extra-uterine pregnancy, and had, in consultation, sanctioned an operation, when the case terminated by abortion, and proved to be a natural pregnancy. The use of the uterine sound, which might aid in clearing up the case, is necessarily contraindicated unless uterine gestation is certainly disproved. Hence it must be admitted that positive diagnosis must always be very difficult. So that the most we can say is, that when the general signs of early pregnancy are present, associated with the other symptoms and signs alluded to, the suspicion of tubal pregnancy may be sufficiently strong to justify us in taking such action as may possibly spare the patient the necessarily fatal consequence of rupture.

Treatment.—If the diagnosis were quite certain, the removal of the entire Fallopian tube and its contents by abdominal section would be quite justifiable, and probably would neither be more difficult nor more dangerous than ovariectomy; for, at this stage of extra-uterine foetation, there are no adhesions to complicate the operation. As yet, however, the uncertainty of the diagnosis has prevented the adoption of the practice.

Opening of the Sac by the Galvano-caustic Knife.—Dr. Thomas of New York¹ has recently recorded a most instructive case, in which he

¹ *New York Med. Journ.*, June, 1875.

saved the life of the patient by a bold and judicious operation. The nature of the case was rendered pretty evident by the signs above described, and Thomas opened the cyst from the vagina by a platinum knife rendered incandescent by a galvano-caustic battery, by which means he hoped to prevent hemorrhage. Through the opening thus made he removed the fœtus. In subsequently attempting to remove the placenta very violent hemorrhage took place, which was only arrested by injecting the cyst with a solution of persulphate of iron. The remains of the placenta subsequently came away piecemeal, after an attack of septicæmia, which was kept within bounds by freely washing out the cyst with antiseptic lotion, the patient eventually recovering. If I might venture to make a criticism on a case followed by so brilliant a success, it would be that in another instance of this kind it would be safer to follow the rule so strictly laid down with regard to gastrotomy in abdominal pregnancies, and leave the placenta untouched, trusting to the injection of antiseptics and the thorough drainage of the cyst to prevent mischief.[¹]

[The advice given by the author in regard to the non-removal of the placenta was *first* urged upon the medical profession in 1791 by Mr. William Trumbull in a paper read before the Medical Society of London, and again in 1795, in a letter² from the late Dr. James Mease of Philadelphia to Dr. Lettsom of London, in which he reported an operation by Dr. Charles McKnight of New York very similar to this of Dr. Thomas, and ending favorably to the woman. The remarks of Dr. Mease on the impropriety of removing the placenta were read before the same society, and concurred in by some of the members present.

It is a little remarkable that the opinion of Dr. Mease originated in an accident which occurred in the operation of Dr. McKnight, by which the funis was ruptured, and in consequence of which the placenta, which was outside of the cyst, could not be found for removal. The value of this discovery appears to have been lost to the profession for a long term of years, as many authors have objected to the operation because of the danger of removing the placenta.

In a second operation, performed on May 10, 1876, in a case of secondary abdominal pregnancy, Dr. Thomas³ operated through the linea alba, and removed a female fœtus weighing 6 lbs. 15 oz. The funis was traced to the left iliac fossa, where it was apparently inserted into the peritoneum, and no placenta was discernible. The cord was cut off at its origin, and the wound closed, except at its lower part, which was kept open by a glass tube. The woman's pulse before the operation was 120, and fell to 107 at the end of the first week; temperature was always 100° and upward, but in the middle of the

[¹ Dr. J. H. Mathieson of St. Mary's, Ontario, performed a similar operation on June 28, 1881, and, as in the King case of Edisto Island in 1816, saved both mother and fœtus. The subject was a woman of 30, and pregnant for the sixth time. After opening the vagina the fœtus, which weighed 8 lbs. 7½ oz., was delivered by the forceps. The placenta, which was three-lobed, was easily peeled off, and a sponge soaked with solution of perchloride of iron was inserted. There was not much hemorrhage, and the cyst was treated with antiseptic syringing. It was three months in closing up (*Trans. Obstetric Soc. London*, May 7, 1884; *Lancet*, May 24, 1884, p. 940).—Ed.]

[² *Memoirs of Med. Soc. London*, vol. iv. p. 342, 1795.]

[³ *Am. Journ. of Obstetrics*, vol. ix. p. 655, 1876.]

fourth week it rose to 103° – 104° , and the pulse to 130. The placenta was found presenting at the opening in the abdomen, and was removed with dressing forceps. It was of the ordinary diameter and had a shrivelled appearance. The removal afforded a decided relief, and the temperature fell within three hours. Antiseptic injections were freely used in the treatment of the case, and the patient made a good recovery.

Prof. T. Gaillard Thomas, above referred to, has reported¹ 27 cases of extra-uterine pregnancy which have come under his personal observation, all but 3 of them having been seen in consultation. Of these, 17 recovered and 10 died. Rupture of the sac occurred in 7 cases, with 6 deaths. Laparotomy was performed in 5 cases, 4 of which recovered. In 2 women the cysts were tapped, and both died. One cyst was incised by the thermo-cautery per vaginam, with recovery, and one was evacuated per rectum, but the patient died. Two recovered after spontaneous evacuation by the rectum, and 3 were cured by destroying the fœtus under galvanism. Prof. Goodell of Philadelphia reports 13 cases, with 3 recoveries.²—ED.]

Means of Destroying the Vitality of the Fœtus.—Another mode of managing these cases is to destroy the fœtus, so as to check its further growth, in the hope that it may remain inert and passive within its sac. Various operations have been suggested and practised for this purpose. Thus, needles have been introduced into the tumor, through which currents of electricity have been passed, either the continuous current, or, as has been suggested by Duchenne, a spark of Franklinic electricity. Hicks, Allen, and others have endeavored to destroy the fœtus by passing an electro-magnetic current through it by means of a needle.^[3] Lusk⁴ relates several successful cases following the use of the faradic current, one pole being passed through the rectum to the site of the ovum, the other being placed on a point in the abdominal wall two or three inches above Poupart's ligament. The current should be passed daily for five or ten minutes, and continued for a week or two until the shrinking of the tumor gives satisfactory evidence of the death of the fœtus. In a case reported by Dr. Bachetti, in which the continuous current was used, the growth of the ovum was arrested and the patient recovered. The same result, however, would probably have followed the simple puncture of the cyst. This has been successfully practised on several occasions, either with a small trocar and canula or with a simple needle. A very interesting case, in which the development of a two months' tubal gestation was arrested in this way, is recorded by Greenhalgh,⁵ and another by Martin of Berlin⁶. Joulin suggested that not only should the cyst be punctured, but that a solution of morphia should be injected into it, which, by its toxic influence, would ensure the destruction of the fœtus; and this is probably one of the best means at our disposal for destroying the fœtus. Other means proposed for effecting the same object, such as pressure or the administration of toxic

[¹ *Trans. Am. Gynecol. Soc.*, vol. vii., for 1882, and vol. ix., for 1884.] [² *Ibid.*]

[³ Dr. Allen did not use needles in applying the current.—ED.]

⁴ *Science and Art of Midwifery*, p. 321.

⁵ *Lancet*, 1867.

⁶ *Monat. f. Geburt*, 1868.

remedies by the mouth, are far too uncertain to be relied on. The simplest and most effectual plan would be to introduce the needle of an aspirator, by which the liquor amnii would be drawn off and the further growth of the fœtus effectually prevented. Parry,¹ indeed, is opposed to this practice, and has collected several cases in which the puncture of the cyst was followed by fatal results, either from hemorrhage or septicæmia. In these, however, an ordinary trocar and canula were probably employed, which would necessarily admit air into the sac. It is difficult to imagine that a fine hair-like aspirating needle, rendered perfectly aseptic by carbolic acid, could have any injurious results; and it could do no harm, even if an error of diagnosis had been made and the suspected extra-uterine fœtation turned out to be some other sort of growth. If the aspirator proves that an extra-uterine fœtation exists, then, if the cyst be of any considerable size, and the pregnancy advanced beyond the second month, we might, if deemed advisable, resort to a more radical operation, such as that so successfully practised by Thomas.

[The safest and most successful of all the methods that have been employed for arresting the development of an extra-uterine fœtus is that of the galvanic current, for the efficient use of which needles and puncturing are not at all necessary. This operation has been more frequently reported in the United States than in any other country, and of 15 cases on record not one was fatal. Dr. Henry J. Garrigues of New York City thus describes his method of operating:² "I used a French one-cell apparatus, composed of two carbon plates and one zinc plate immersed in Bunsen's battery fluid (potass. bichrom. 3ij; acid sulphuric. concentr. f3iss; aquæ fluv. f3xj). The positive electrode, made of a large carbon plate, covered with cloth, was applied on the abdomen over the tumor. The negative electrode, consisting of one insulated brass stem, with knob, was introduced into the vagina and pressed up against the lower part of the tumor. The current was gradually increased to the limit of her endurance, but never enough to cause real pain." Two days later the pulsation of the tumor previously felt in the vagina had disappeared, and the tumor was less in size. The applications were made for ten minutes almost daily for two weeks. In sixteen days the tumor had diminished to the size of an English walnut.—ED.]

Treatment when Rupture has Occurred.—When the chance of arresting the growth of a tubular fœtation has never arisen, and we first recognize its existence after laceration has occurred and the patient is collapsed from hemorrhage, what course are we to pursue? Hitherto, all that ever has been done is to attempt to rally the patient by stimulants, and, in the unlikely event of her surviving the immediate effects of laceration, endeavoring to control the subsequent peritonitis, in the hope that the effused blood may become absorbed, as in pelvic hæmatocele. This is, indeed, a frail reed to rest upon, and when laceration of a tubal gestation, advanced beyond a month, has occurred, death has been the most certain result. It is supposed by Bernutz—and his opinion is shared by Barnes—that rupture which does not prove fatal is probably not very

¹ Parry on *Extra-uterine Pregnancy*, p. 204.

² *Trans. Am. Gynecol. Soc.*, vol. vii., for 1882.]

rare in the first few days of extra-uterine gestation, and that it is not an uncommon cause of certain forms of pelvic hæmatocele. It has more than once being suggested that it would be perfectly justifiable when laceration has occurred to perform gastrotomy, to sponge away the effused blood, and to place a ligature around the lacerated tube and remove it with its contents. This would no doubt be a bold and heroic procedure, but no one who is acquainted with the triumphs of modern abdominal surgery can say that it would be either impossible or hopeless. The sponging out of effused blood from the abdominal cavity is an every-day procedure in ovariectomy, nor is there any apparent difficulty in ligaturing and removing the sac of the extra-uterine pregnancy, for, as a rule, there are no adhesions formed to the surrounding parts. The history of these cases shows that death does not generally follow rupture for some hours, so that there would be usually time for the operation, and the extreme prostration might be, perhaps, temporarily counteracted by transfusion. Pressure on the abdominal aorta, resorted to when the patient is first seen, might possibly be employed with advantage to check further hemorrhage until the question of operation is decided. We must remember that the alternative is death, and hence any operation which would afford the slightest hope of success would be perfectly justifiable. I cannot, therefore, agree with those who hold that because the chances of success are so small the operation should not be tried; and I do not doubt that it will yet fall to the lot of some one by this means to snatch a patient from the jaws of death and still further to extend the successes of abdominal surgery.

[This has already been done upon four occasions, thanks to the boldness and skill of Mr. Lawson Tait of Birmingham, who has operated five times, losing only the first patient. These operations bear the dates of Jan. 17, 1883; March 1, 1884; April 9, 1884; May 25, 1884; and June 5, 1884.¹ There was no later one up to October 1, 1884. Thus, in four consecutive months there were four women saved from death by internal hemorrhage following rupture of Fallopian foetal cysts. In theory, this operation has long been viewed as giving promise of success. Promptness, boldness, and the advantage of an early rupture were the requisites; but permission to operate and agreement in consultation upon the case presented were the obstacles to be overcome. I became fully convinced of the practicability of this measure in making an autopsy of a young married lady in 1856, in whom a cyst had burst at about the fourth week, causing her death in twenty-three hours. In that day no one would have dared to operate in such a case, and, besides, all of the three physicians who saw her under the attack believed she was not pregnant, as she was menstruating. The history of the case, as given to me, indicated to my mind that there had been a rupture of the right Fallopian tube; which proved to be correct. In the last edition of this work I referred to a case upon which Prof. T. Gaillard Thomas of New York would have operated had he not been overruled by several physicians in consultation, who were doubtful of the correctness of his diagnosis and feared to have it tested under the knife. In another case in New York the patient was operated upon by Dr. Charles K. Briddon,

[¹ *British Med. Journ.*, June 23, 1884, p. 1250.]

but the delay in getting a consultation and the final consent of the family was fatal to her: she lived forty-seven hours, but never reacted. At 7 P. M. she was pulseless and thought herself dying; at 9 P. M. a consultation was held, and at 9.30 a second, with a third consultant. Then the family had to consent, and thus hours of vital moment were lost.¹ This success of Mr. Tait will open the way for the performance of the operation by others.—Ed.]

Abdominal Pregnancy.—In the second of the two classes into which, for practical convenience, we have divided extra-uterine gestation, the ovum is developed in the abdominal cavity. It is as yet an open question whether in some cases the pregnancy is primarily abdominal or not. Barnes believes that it probably never is so, on account of the difficulty of admitting that so minute a body as the ovum should be able to fix itself on the smooth peritoneal surface. He therefore thinks that all abdominal pregnancies are primarily either tubal or ovarian, the sac in which they were contained having given way, and the ovum having retained its vitality through partial attachment to the original sac. This theory is opposed to that of the majority of writers, and, although it may perhaps render the facts less difficult to understand, it is purely hypothetical. There is no evidence to show that in most cases there is an early laceration of a tubal or ovarian sac. That the chorion villi do graft themselves upon the surrounding peritoneum is certain, and is observed in all cases of abdominal gestation. It is not more difficult to imagine them doing this from their very first development than a little later; for it must be allowed that if such laceration does occur, in most cases it can only be when pregnancy is very slightly advanced. On the whole, therefore, it seems not unreasonable to admit the usual explanation of these cases, that the ovule, already impregnated, escaped the grasp of the Fallopian tube and fell into the abdominal cavity, where it rooted itself and developed. Some have, indeed, supposed that abdominal pregnancy may occasionally arise in consequence of spermatozoa finding their way into the peritoneal cavity, and there meeting and impregnating an ovule discharged from the Graafian follicle. Such an event one would suppose to be almost impossible, but Koeberle's case, already quoted, proves that it has actually occurred. The probability is that it is by no means rare for impregnated ovules to drop into the peritoneal cavity, and that the majority of those that do so perish without doing any harm. When they do survive, however, the chorion villi sprout, attach themselves to the surrounding structures, and eventually develop into a placenta. The mode in which the chorion villi are attached, and the arrangement of the maternal blood-vessels, have never yet been worked out, and would form a very interesting subject for investigation. The precise seat of attachment varies, and the placenta has been found fixed to most of the abdominal viscera, either those contained in the pelvis proper, or it may be the intestines, or to the iliac fossa; most frequently, apparently, the ovum finds its way into the retro-uterine cul-de-sac.

Formation of a Cyst around the Ovary.—The subsequent changes vary much. In the large majority of cases the ovum produces considerable

[¹ *Medical News*, Phila., Dec. 15, 1883, p. 666.]

irritation, resulting in the exudation of plastic material, which is thrown round it, so as to form a secondary cyst or capsule in which maternal vessels are largely developed, and which stretches, *pari passu*, with the growth of the ovum (Fig. 84). The density and strength of this cyst are found to be very different in different cases; sometimes it forms a complete and strong covering to the ovum, at others it is very thin and

FIG. 84.



Uterus and Fœtus in a Case of Abdominal Pregnancy.

only partially developed, but it is rarely entirely absent. As there is ample space for the development of the ovum, and as the secondary cyst generally stretches and grows along with it, most cases of abdominal pregnancy progress without any very remarkable symptoms, beyond occasional severe attacks of pain, until the full term of pregnancy has been reached. Sometimes, however, the cyst lacerates, and there is an escape of blood into the abdominal cavity, accompanied by more or less prostration and collapse, which may prove fatal, but from which the patient more generally rallies. The fetus, now dead, will remain in the abdomen, and will undergo changes and produce results similar to those which we shall presently describe as occurring in cases progressing to the full period.

Pseudo-labor sometimes Comes on.—In most cases, at the natural termination of pregnancy a strange series of phenomena occur: pseudo-labor comes on, there are more or less frequent and strong uterine contractions, possibly an escape of blood from the vagina, the discharge of the broken-down uterine decidua, and even the establishment of lactation. Sometimes the contractions of the abdominal muscles produced by this ineffective labor have been so strong as to cause the laceration of the adventitious cyst surrounding the fetus, and the escape of blood and liquor amnii into the abdominal cavity, with a rapidly fatal result. More frequently laceration does not occur, and the spurious labor-pains continue at intervals until the fetus dies, possibly from pressure, but more often from effusion of blood into the tissue of the placenta, and consequent asphyxia. Occasionally the fetus has apparently lived a

considerable time, in some cases even for several months, after the natural limit of pregnancy has been reached.

Changes After the Death of the Fœtus.—It is after the death of the fœtus that the dangers of abdominal pregnancy generally commence, and they are numerous and various. The subsequent changes that occur are well worthy of study. Occasionally the fœtus has been retained for a length of time, even until the end of a long life, without producing any serious discomfort, and in many cases of this kind several normal pregnancies and deliveries have subsequently taken place. Even when the extra-uterine gestation appears to be tolerated, and has remained a long time without producing any bad effects, serious symptoms may be suddenly developed, so that no woman, under such circumstances, can be considered safe. The condition of these retained fœtuses varies much. Most commonly the liquor amnii is absorbed, the fœtus shrinks and dies, all its soft structures are changed into adipocere, and the bones only remain unaltered. Sometimes this change occurs with great rapidity. I have elsewhere¹ recorded a case of extra-uterine fœtation in which at the full term of pregnancy the fœtus was alive, and the woman died in less than a year afterward. On post-mortem examination the fœtus was found entirely transformed into a greasy mass of adipocere, studded with fœtal bones, in which not a trace of any of the soft parts could be detected. On the other hand, the fœtus may remain unchanged: in the Museum of the College of Surgeons there is one which was retained in the abdomen for fifty-two years, and which was found to be as fresh and unaltered as a new-born child. In other cases the sac and its contents atrophy and shrink, and calcareous matter is deposited in them, so that the whole becomes converted into a solid mass known as *lithopædion* (Fig. 85). The cases, however, in which the retention of the fœtus gives rise to no mischief are quite exceptional. Generally the fœtus putrefies, and this may either immediately cause fatal peritonitis or septicæmia, or, as more commonly happens, secondary inflammation and suppuration of the sac. Under the influence of the latter the sac opens externally, either directly at some point of the abdominal walls or indirectly through the vagina, the bowels, or even the bladder. Through the aperture or apertures thus formed (for there are often several fistulous openings) pus and the bones and other parts of the broken-down fœtus are discharged; and this may go on for months, and even years, until at last, if the patient's strength does not give way, the whole contents of the cyst are expelled, and recovery takes place. From various statistical observations it appears that the chances of recovery are best when the cyst opens through the abdominal walls, next through the vagina or

FIG. 85.



Lithopædion.

(From a preparation in the Museum of the College of Surgeons.)

¹ *Obst. Trans.*, vol. vii.

bladder, and that the foetus is discharged with most difficulty and danger when the aperture is formed into the bowel. At the best, however, the process is long, tedious, and full of danger; and the patient too often sinks, during the attempt at expulsion, through the irritation and exhaustion produced by the abundant and long-continued discharge.

Diagnosis.—The diagnosis of abdominal gestation is by no means so easy as might be thought, and the most experienced practitioners have been mistaken with regard to it.

The most characteristic symptom, although this is not so common as in tubal gestation, is metrorrhagia combined with the general signs of pregnancy. Very severe and frequently-repeated attacks of abdominal pain are rarely absent, and should at once cause suspicion, especially if associated with hemorrhage and the discharge of a decidual membrane from the uterus. They are supposed by some to depend on intercurrent attacks of peritonitis, by which the foetal cyst is formed. Parry doubts this explanation, and attributes them partly to the distension of the cyst by the growing foetus, and partly to pressure on the surrounding structures. On palpation the form of the abdomen will be observed to differ from that of normal pregnancy, being generally more developed in the transverse direction, and the rounded outline of the gravid uterus cannot be detected. When development has advanced nearly to term, the extreme distinctness with which the foetal limbs can be felt will arouse suspicion. Per vaginam, the os and cervix will be felt softened, as in ordinary pregnancy, but often displaced by the pressure of the cyst, and sometimes fixed by perimetritic adhesions; either of these signs is of great diagnostic value.

By bi-manual examination it may be possible to make out that the uterus is not greatly enlarged, and that it is distinctly separate from the bulk of the tumor; these facts, if recognized, would of themselves disprove the existence of uterine gestation. The diagnosis, if the foetal limbs or heart-sounds could be detected, would be cleared up in any case by the uterine sound, which would show that the uterus was empty and only slightly elongated. But we must be careful not to resort to this test unless the existence of uterine gestation is positively disproved by other means. As, however, it places the diagnosis beyond a doubt, it should always be employed whenever operative procedure is in contemplation. Quite recently I have seen a remarkable case which illustrates the importance of this rule. The case had been diagnosed as abdominal pregnancy by no less than six experienced practitioners, and was actually on the operating-table for the performance of laparotomy. As a precaution, having some doubts of the diagnosis, I suggested the passage of the sound, which entered into a gravid uterus, the case proving to be one of small ovarian tumor jammed down into Douglas's space and displacing the cervix forward. Had it not been for this precaution its true nature would certainly not have been detected.

Treatment.—The treatment of abdominal gestation will always be a subject of anxious consideration, and there is much difference of opinion as to the proper course to pursue. It is pretty generally admitted that it is not advisable to adopt any active measures until the full term of development is reached. Puncturing the cyst, with the view of destroy-

ing the foetus and arresting its further growth, has been practised, but there are good grounds for rejecting it, for there is not the same imminent risk of death from rupture of the cyst as in tubal foetation; and even if the destruction of the foetus could be brought about, there would still be formidable dangers from subsequent attempts at elimination or from internal hemorrhage.

Question as to the Performance of Primary Gastrotomy.—When the full period has arrived, the child being still alive, as proved by auscultation, we have to consider whether it may not be advisable to perform gastrotomy before the foetus perishes, and so at least save the life of the child. There are few questions of greater importance and more difficult to settle. The tendency of medical opinion is rather in favor of immediate operation, which is recommended by Velpeau, Kiwisch, Koeberle, Schroeder, and many other writers, whose opinion necessarily carries great weight. The arguments used in favor of immediate operation are, that while it affords a probability of saving the child, the risks to the mother, great though they undoubtedly are, are not greater than those which may be anticipated by delay. If we put off interference, the cyst may rupture during the ineffectual efforts at labor, and death at once ensue; or, if this does not take place, other risks, which can never be foreseen, are always in store for the patient. She may sink from peritonitis, or from exhaustion, consequent on the efforts at elimination, which in the majority of cases are sooner or later set up, so that, as Barnes properly says, “the patient’s life may be said to be at the mercy of accidents of which we have no sufficient warning.” On the other hand, if we delay, while we sacrifice all hope of saving the child, we at least give the mother the chance of the foetation remaining quiescent for a length of time, as certainly not unfrequently occurs. Thus, Campbell collected 62 cases of ultimate recovery after abdominal gestation, in 21 of which the foetus was retained without injury for a number of years. Then there is the question of secondary gastrotomy, which consists in operating after the death of the foetus when urgent symptoms have arisen—a course which is advocated by Mr. Hutchinson. In favor of this procedure it is urged that by delay the inflammation taking place about the cyst will have greatly increased the chance of adhesions having formed between it and the abdominal parietes, so as to shut off its contents from the cavity of the peritoneum. The more effectually this has been accomplished, the greater are the chances of recovery. When the foetus has been dead for some time, the vascularity of the cyst will also be lessened, and the placental circulation will have ceased, so that the danger of hemorrhage will be much diminished.

It will be seen, therefore, that there are arguments in favor of each of these views. The results of the primary operation are far less favorable than we should have, *a priori*, supposed. Since the first edition of this work appeared the subject has been carefully studied by Dr. Parry in his exhaustive treatise on Extra-uterine Foetation. He has there shown that when the case is left until nature has shown the channel through which elimination is to be effected, the mortality is 17.35 per cent. less than in the cases in which the primary operation was performed. His conclusion is that “the primary operation cannot be too forcibly con-

demned. It is not too much to say that this operation adds only another danger to a life already trembling in the balance, which the delusive hope of saving the uncertain life of a child does not warrant us in assuming." It is only just to remember, as is forcibly pointed out by Keller, that in these days of advanced abdominal surgery a better result might be anticipated than when gastrotomy was performed in the haphazard way which was usual before we had gained experience from ovariectomy. No doubt minute care in the performance of the operation, a due attention to its details, studiously avoiding as much as possible the passage of blood and the contents of the cyst into the peritoneal cavity, and a free use of antiseptics, would materially lessen its peril. This conclusion is well illustrated in a recent interesting paper by Thomas, who relates three successful cases of laparotomy in abdominal pregnancy.¹

Mode of Performing the Operation.—The operation, then, should be performed with all the precautions with which we surround ovariectomy. The incision, best made in the linea alba, should not be greater than is necessary to extract the fetus, and may be lengthened as occasion requires. It has been suggested that should the head be felt presenting above the vagina, the intervening structures should be divided and the fetus withdrawn by the forceps. This procedure was actually adopted with success in 1816 by Dr. John King of Edisto Island, South Carolina. If there are no adhesions, the walls of the cyst should be stitched to the margin of the incision, so as to shut it off as completely as possible from the peritoneal cavity. This has been specially insisted on by Braxton Hicks, and should never be omitted. The special risk is not so much the wounding of the peritoneum as the subsequent entrance of septic matter from the cyst into its cavity. Another cardinal rule, both in primary and secondary gastrotomy, is to make no attempt to remove the placenta. Its attachments are generally so deep-seated and diffused that any endeavor to separate it is likely to be attended with profuse and uncontrollable hemorrhage or with serious injury to the structure to which it is attached. Many of the failures after operating can be traced to a neglect of this rule. The best subsequent course to pursue, after removing the fetus and arresting all hemorrhage either by ligature or the actual cautery, is to sponge out the cyst as gently as possible, sprinkle the cavity with iodoform or with equal parts of tannin and salicylic acid, as recommended by Freund,² and then to bring the upper part of the wound into apposition with sutures, leaving the lower open, with the cord protruding, so as to ensure an outlet for the escape of the placenta as it slips down. The subsequent treatment must be specially directed to favour the escape of the discharge and to prevent the risk of septicæmia. These objects may be much aided by injections of antiseptic fluids, such as solution of carbolic acid or diluted Condy's fluid; and it would probably be advisable to place a drainage-tube in the lower angle of the wound. It may be well to point out that there is no operation in which a scrupulous following of the antiseptic method on Sir Joseph Lister's principles is so likely to be useful.

As long as the placenta is retained the danger is necessarily great, and

¹ *Am. Journ. of Med. Sci.*, Jan., 1873.

² *Edin. Med. Journ.*, Dec., 1883.

it may be many days, or even weeks, before it is discharged. When once this is effected, the sac may be expected to contract, and eventually to close entirely.

[*Relative Risks of Primary and Secondary Laparotomy in Abdominal Pregnancies.*—In view of the improvements in abdominal surgery, especially in its application to women, and the disposition that exists to undertake with little hesitation the performance of operations which a few years ago were regarded as almost necessarily fatal, it becomes a matter of vital importance to consider whether we are justified in attempting to save the life of a fœtus coming to maturity in the abdominal cavity by the use of the knife, either by the past record of such operations or by any reasonable ground of hope that the inherent difficulties of the condition may be overcome through the advantages to be derived from antiseptics, abdominal drainage, irrigation, etc.

We are told by some very prominent gynecologists that the statistics of the past are of little value in this operation; that this is a progressive age in abdominal surgery; and that with the diminution of risks under Listerism we ought to be able to do better than in the past, and to overcome the dangers and difficulties arising from the vascularity of the cyst wall, the activity of the placental circulation, the decomposition and exfoliation of the placenta, with its accompanying hemorrhage or septic poisoning, and the presence of puriform fluid in the cyst or abdominal cavity. It is certainly very tempting to operate in the interest of two lives, and it is true that by postponement we run a risk of the patient being lost before a time considered safer for operating may arrive. This is a delicate question to settle, and one which will be appreciated by some of my contemporaries who have had this unfortunate experience and regretted afterward that they had not operated early. But what is the record of the past? Has not antiseptic surgery had a trial yet? When asked what has been the mortality under the primary operation, a gynecologist will generally refer the applicant to the work of Dr. Parry, already referred to, and in it he finds a table of 20 cases of so-called primary laparotomies "*performed during or at the end of gestation,*" and that 6 of these are stated as having saved the women, with two of their children. This record, if reliable, would give a recovery of 30 per cent.—a far higher one than will be presently shown from a careful re-examination of his statistics. Had Dr. Parry examined the monograph of Dr. Keller of Strasbourg,¹ he would not have fallen into the error I am about to point out, as in this paper some of his primary cases are in the secondary list.

My attention having been directed to some errors in Dr. Parry's statistics, I was led to re-examine his sources of information, and am surprised to find that not one of his six recoveries under primary laparotomy, so called, was entitled to be thus designated.

Case 5 was operated upon by Dr. Schreyer of Hamburg in 1837, and reported in *Casper's Wochenschrift*, No. xlv., 1837, page 726.

Case 6, credited to "Schwanck" (correctly Zwanck), is the same as Case 5, and was taken from a short abstract report in the *Archiv. gen. de Méd.*, June, 1838, p. 227, giving the operation to Dr. Zwanck, who was

[¹ *Des Grossesses Extra-utérines*, Paris, 1872.]

the assistant and reporter. The description of the operation shows clearly that the foetus was taken out of some form of uterine cavity, from which the placenta was readily removed, and that the woman recovered in three weeks. Prof. Litzmann of Kiel, Germany, believes that the foetus was removed from "a very thin-walled uterus."

Case 9. Dr. Découene of Courtrai, France, performed the Cæsarean operation six times in ten years (1841-51), and saved five women and four children. Dr. Parry's reference is incorrect, but these six cases are in the volume named.¹ Prof. Litzmann says that the case (9) was a twin pregnancy, in which laparotomy was performed after uterine rupture.

Case 10. Dr. Frederick A. Stutter's. Pregnancy had existed 45 weeks, and the foetus, which had been dead five weeks, was already in a putrid state. The decomposition hastened the separation of the placenta, which was removed entire in five days.

Case 11. Drs. Ramsbotham and Adams's. Here the foetus died just before full maturity, and the operation was postponed for safety for six months. The woman was then delivered by secondary laparotomy, and recovered: she was pregnant for 15 months.

Case 14. Dr. William D. Hooper's of Liberty, Virginia. Dr. Parry gives the duration of pregnancy as "223 days." Having carefully read the original record, I am inclined to believe that the foetus was carried more than five years. The woman, a two-para, was married a second time when in good health. This soon began to fail under some abdominal malady, but she menstruated regularly from April 1, 1866, to June, 1871, with one exception in the latter year, and did not suppose herself pregnant. Her health began decidedly to fail in January, 1871, and she was a confirmed invalid until operated upon on October 24, 1871. She had passed foetal bones from her rectum four months before, and her abdomen was about to open spontaneously when she was relieved by the knife, and a putrid foetus removed without opening the peritoneal cavity. The foetus was mainly a skeleton, and was computed as one of at least seven months.

In contrast with Dr. Parry's record, Prof. Litzmann² gives one of 10 genuine primary operations. The children were all living, and were delivered alive. The women were pregnant for periods ranging from 29 weeks to full maturity. Nine of the women died—five on the first day, two on the second, one on the third, and one on the sixteenth. Two fully-matured children lived and thrived; a third died in three months, and a fourth on the second day. The remaining six lived from a few hours to fifty.

The only surviving woman was the patient of Mr. Jessop of Leeds, England, operated upon in 1875. This case, thus far, of all the laparotomies undertaken in the hope of saving two lives, is the only one known to me that has not been a failure so far as the mother was concerned; and in this instance the condition of the woman for a long time gave very little encouragement of final recovery, and her child was lost at eleven months with croup and pneumonia. A perusal of Mr. Jessop's

¹ *Gazette des Hôpitaux*, Paris, 1852, p. 221.]

² *Zur Feststellung der Indicationen für die Gastrotomie bei Schwangerschaft ausserhalb der Gebärmutter*, von C. Litzmann, 8vo, pp. 79.]

experience with his patient will certainly not encourage one to try a similar experiment.¹

After an extra-uterine foetus is dead within the abdominal cavity, the risk of its removal, provided time enough shall have passed for the requisite anatomical changes to take place, is very materially diminished, as shown by a number of operations in this country and in Europe performed within a few years. If the woman survives the immediate risk of the false labor that usually takes place when the foetus nears maturity, she usually improves somewhat in health for a time, by reason of the removal of pressure produced by the absorption of much of the liquor amnii. The death of the foetus rendering the placenta no longer of any functional importance, this element of danger in the primary operation ceases in time to be such, by reason of changes in its structure which fit it for undergoing exfoliation without the hemorrhage which follows the primary operation. The question, then, of operating becomes one of time, which itself has no special limit, as the changes in the placental circulation that follow foetal death may be slow or rapid according to circumstances; but experience teaches that from 10 to 12 weeks will be sufficient. Much less than this will be required where decomposition has commenced; and in that event an earlier operation may be demanded on account of the failing health of the woman. Case 10, already noted, was an example of this, the foetus having been dead only five weeks. In one case in which I was consulted a decomposing foetus was removed when four months dead, but for some weeks the patient had been suffering from a slow blood-poisoning, with pulse 105, and attacks of mental distress and nervous excitement at night: this woman is now in excellent health. It is not necessary here to defend the *secondary* operation or to collect its statistics in proof of its comparative safety in skilful hands, as abundant evidence of this is shown in the reports of cases in medical journals. Antiseptic surgery, drainage, and irrigation here avail, as the placenta, if left intact, will usually come away without hemorrhage. Dr. Parry condemned the primary operation in strong terms, although his statistics gave a measure of hope for its success. Later in life he lost a primary case by delay, and regretted not having operated; but he need not have done so, except for the possibility of saving the foetus, and it might be then for a few days only.—ED.]

Treatment when the Foetus is Dead.—When the foetus is dead, or when we have determined not to attempt primary gastrotomy, it is advisable to wait, very carefully watching the patient, until either the gravity of her general symptoms or some positive indication of the channel through which nature is about to attempt to eliminate the foetus shows us that the time for action has arrived. If there be distinct bulging of the cyst in the vagina or in the retro-vaginal cul-de-sac, especially if an opening has formed there, we may properly content ourselves with aiding the passage of the foetus through the channel thus indicated, and removing the parts that present piecemeal as they come within reach, cautiously enlarging the aperture if necessary. If the sac have opened into the intestines, the expulsion of the foetus through this channel is so tedious and difficult,

[¹Dr. Werth of Kiel stated before the Int. Med. Congress of 1884 that he had collected the records of 17 operations, with two recoveries.]

the exhaustion attending it so likely to prove fatal, and the danger from decomposition of the fetus through passage of intestinal gas so great, that it would probably be best to attempt to remove it by gastrotomy, especially if it is only recently dead and the greater portion is still retained.

Mode of Performing Secondary Gastrotomy.—If an opening forms at the abdominal parietes, or if the symptoms determine us to resort to secondary gastrotomy before this occurs, the operation must be performed in the same way and with the same precautions as primary gastrotomy. Here, as before, the safety of the operation must greatly depend on the amount and firmness of the adhesions, for if the cyst be not completely shut off from the peritoneal cavity the risks of the operation will be little less than those of primary gastrotomy. It would obviously materially influence our decision and prognosis if we could determine this point before operating. Unfortunately, it is impossible, as the experience of ovariologists proves, to ascertain the existence of adhesions with any certainty. If, however, we find that the abdominal parietes do not move freely over the cyst, and if the umbilicus be depressed and immovable, the presumption is that considerable adhesions exist. If they are found not to be present, the cyst walls should be stitched to the margin of the incision, in the manner already indicated, before the contents are removed.

If the fetus has been long dead and its tissues greatly altered, its removal may be a matter of difficulty. In the case under my own care, already alluded to, the fetal structures formed a sticky mass of such a nature that I believe it would have been impossible to empty the cyst had an operation been attempted. This would be, to some extent, a further argument in favor of the primary operation.

Opening of Cyst by Caustics.—The importance of adhesions has led some practitioners to recommend the opening of the cyst by potassa fusa or some other caustic, in the hope that it would set up adhesive inflammation around the aperture thus formed. Several successful operations by this method are recorded, and it would be worth trying should the extreme mobility of the cyst lead us to suspect that no adhesions existed. If we have to deal with a case in which fistulous openings leading to the cyst have already formed, it may, perhaps, be advisable to dilate the apertures already existing, rather than make a fresh incision; but in determining this point the surgeon will naturally be guided by the nature of the case and the character and direction of the fistulous openings.

General Treatment.—It is almost needless to say anything of general treatment in these trying cases, but the administration of opiates to allay the sufferings of the patient, and the endeavor to support the severely-taxed vital energies by appropriate food and medication, will form a most important part of the management. Freund specially insists on the importance of careful regulation of the bowels, and on making milk the staple article of diet, as important points in the management of cases prior to operation.

Gestation in a Bilobed Uterus.—A few words may be said as to gestation in the rudimentary horn of a bilobed uterus, to which con-

siderable attention has of late years been directed by the writings of Kussmaul and others. It appears certain that many cases of supposed tubal gestation are really to be referred to this category. Although such cases are of interest pathologically, they scarcely require much discussion from a practical point of view, inasmuch as their history is pretty nearly identical with that of tubal pregnancy. The rudimentary horn is distended by the enlarging ovum, and after a time, when further distension is impossible, laceration takes place. As a matter of fact, all the thirteen cases collected by Kussmaul terminated in this way; and even on post-mortem examination it is often extremely difficult to distinguish them from tubal pregnancies. The best way of doing so is probably by observing the relations of the round ligaments to the tumor, for if the gestation be tubal they will be found attached to the uterus on the inner or uterine side of the cyst, whereas if the pregnancy be in a rudimentary horn of the uterus they will be pushed outward and be external to the sac. In the latter case, moreover, the sac will be probably found to contain a true decidua, which is not the case in tubal pregnancy. The only point in which they differ is that in cornual pregnancy rupture may be delayed to a somewhat later period than in tubal, on account of the greater distensibility of the supplementary horn.

Missed Labor.—The term "*missed labor*" is applied to an exceedingly rare class of cases in which, at the full period of pregnancy, labor has either not come on at all, or, having commenced, the pains have subsequently passed off, and the foetus is retained in utero for a very considerable length of time. Under such circumstances it has usually happened that the membranes have ruptured at or about the proper term, and the access of air to the foetus in utero has been followed by decomposition. A putrid and offensive discharge has then commenced, and eventually portions of the disintegrating foetus have been expelled per vaginam. This discharge may go on until the entire foetus is gradually thrown off, or, more frequently, the patient dies from septicæmia or other secondary result of the presence of the decomposing mass in utero. Thus, McClintock relates one case¹ in which symptoms of labor came on in a woman, 45 years of age, at the expected period of delivery, but passed off without the expulsion of the foetus. For a period of sixty-seven weeks a highly offensive discharge came away, with some few bones, and she eventually died with symptoms of pyæmia. He also cites another case in which the patient died in the same way after the foetus had been retained for eleven years.

Ulceration of the Uterine Walls sometimes Occurs.—Sometimes, when the foetus has been retained for a length of time, a further source of danger has been added by ulceration or destruction of the uterine walls, probably in consequence of an ineffectual attempt at its elimination. This occurred in Dr. Oldham's case (Fig. 86), in which the contained mass is said to have nearly worn through the anterior wall of the uterus; and also in one reported by Sir James Simpson,² in which a patient died three months after term, the foetus having undergone fatty metamorphosis, an opening the size of half a crown having formed between the


¹ *Dublin Quart. Journ.*, Feb. and May, 1864.

² *Edin. Med. Journ.*, 1865.

transverse colon and the uterine cavity. It is also stated that "the uterine walls were as thin as parchment."

In some few cases, however, probably when the entrance of air has been prevented, the fœtus has been retained for a length of time without decomposing and without giving rise to any troublesome symptoms.

FIG. 86.



Contents of the Cyst in Dr. Oldham's Case of Missed Labor.

Such a case is reported by Dr. Cheston,¹ in which the fœtus remained in utero for fifty-two years.

Its Causes are not Properly Understood.—The causes of this strange occurrence are altogether unknown. Generally, the fœtus seems to have died some time before the proper term for labor, and this may have influenced the character of the pains. It is probably also most apt to occur in women of feeble and inert habit of body, possibly where there was some obstacle to the dilatation of the cervix which the pains were unable to overcome. Barnes suggests² that some presumed examples of missed labor "were really cases of interstitial gestation, or gestation in one horn of a two-horned uterus." In several of the cases, however, the details of the post-mortem examination are too minute to admit of the possibility of mistake having been made.

Sometimes Confounded with Extra-uterine Fœtation.—Müller of Nancy has recently attempted to prove, by a critical examination of published cases, that most examples of so-called "missed labor" were in reality cases of extra-uterine fœtation, in which an ineffectual attempt at parturition took place, the fœtus being subsequently retained.

Its Dangers are Serious.—From what has been said, it will be seen

¹ *Med.-Chir. Trans.*, 1814.

² *Diseases of Women*, p. 445.

that the dangers arising from this state are very considerable, and when once the full term has passed beyond doubt, especially if the presence of an offensive discharge shows that decomposition of the fœtus has commenced, it would be proper practice to empty the uterus as soon as possible. The necessary precaution, however, is not to decide too quickly that the term has really passed; and therefore we must either allow sufficient time to elapse to make it quite certain that the case really falls under this category or have unequivocal signs of the death of the fœtus and injury to the mother's health. If we had to deal with the case before any extensive decomposition of the fœtus had occurred, we probably should find little difficulty in its management, for the proper course then would be to dilate the cervix with fluid dilators and remove the fœtus by turning; or, before doing so, we might endeavor to excite uterine action by pressure and ergot. If the case did not come under observation until disintegration of the fœtus had begun, it would be more difficult to deal with. If the fœtus had become so much broken up that it was being discharged in pieces, Dr. McClintock says that "in regard to treatment our measures should consist mainly of palliatives—viz. rest and hip-baths, to subdue uterine irritation; vaginal injections, to secure cleanliness and prevent excoriation; occasional digital examination, so as to detect any fragments of bone that might be presenting at the os and to assist in removing them. These are plain rational measures, and beyond them we shall scarcely, perhaps, be justified in venturing. Nevertheless, under certain circumstances, I would not hesitate to dilate the cervical canal so as to permit of examining the interior of the womb and of extracting any fragments of bone that may be easily accessible; but unless they could thus be easily reached and removed, the safer course would be to defer, for the present, interfering with them."¹

It may be doubted, I think, whether, considering the serious results which are known to have followed so many cases, it would not, on the whole, be safer to make at least one decided effort, under chloroform, to remove as much as possible of the putrefying uterine contents after the os has been fully dilated. Such a procedure would be less irritating than frequently-repeated endeavors to pick away detached portions of the fœtus as they present at the os uteri. When once the os is dilated, antiseptic intra-uterine injections, as of diluted Condyl's fluid, might safely and advantageously be used. Unquestionably, it would be better practice to interfere and empty the uterus as soon as we are quite satisfied of the nature of the case, rather than to delay until the fœtus has been disintegrated.

[From several cases of "missed labor" that have been reported in the United States we find that the failure of the uterus to expel its contents may be due to a variety of causes. If we are certain that the fœtus is actually in utero, that there is no pelvic or vaginal obstruction, and that the uterus is itself of normal form, then we must look for the cause of difficulty in the organ itself. By an examination of our reports of Cæsarean operations we find that there have been several cases in which the power of the uterine contractions was insufficient to overcome the

¹ *Dublin Quart. Journ.*, vol. xxxvii. p. 314.

resistance to expansion in the cervix. This may be due either to a want of contractile force in the muscular coat, to a change in the tissues of the cervix as the result of inflammation, or to both conditions combined. Where the muscular power of the uterus is in its integrity, the resistance in the cervix may be such that the os may remain unchanged after it is slightly opened, and the patient continue in labor until the contractile power of the uterus is exhausted, when all muscular contraction will cease. Efforts at expulsion may recur at intervals covering a period of many months, when they will cease finally. In two Cæsarean cases in the United States, the subjects being black, there was found a calcareous incrustation over and around the internal os uteri. The first operation was performed in Virginia in 1828 upon a multipara of 25.¹ She was taken in labor at term, and had pains for two or three days together, at intervals, for about four weeks, after which pains returned occasionally during fifteen months. The cervix admitted the index finger, and in time the fœtus became putrid. When operated upon she had carried the fœtus two years. There was very little hemorrhage in the operation, although the uterus failed to contract, and for this reason was sutured. The woman died in the second week, of peritonitis, following an attack of indigestion produced by a meal of animal food and cider. The second case, also a multipara, was operated upon in Georgia in 1877, after a labor of four days, by Dr. Theodore Starbuck, who describes the deposit as "ossific." The child was dead, and the woman died of internal hemorrhage very suddenly on the third day.²

In a third case, also black, the cause of retention appears to have been a prevention of the descent of the fœtus, from its arm and leg being secured within the uterus. The woman was 33 years old and the mother of one child, and was operated upon by Dr. J. C. Egan of Shreveport, Louisiana, August 25, 1860.³ On May 4, 1857, while at work in the field, she felt a sudden and violent pain in the left side; fainted, remained insensible so long as to be thought dead, but finally revived, and was pronounced four months pregnant. Labor began in November; the os dilated, head presented, but did not descend; pains continued at intervals for a month. In the fall of 1858 an abscess opened, leaving a fistula $1\frac{1}{4}$ inches below the umbilicus. When operated upon nearly two years later, she was greatly emaciated and affected with hectic fever. The uterus being adherent, the peritoneal cavity was not opened. When the fœtus was extracted, its left foot and hand were wanting, and, search being made, were found in a pouch on the left side of the uterus, enclosed by bands which were cut for their liberation. The uterus was examined bi-manually to make sure that the cervix was sufficiently open for drainage. The decomposed fœtus had been carried 33 months after maturity. Dr. Egan believes that a partial rupture of the uterus took place at the time of her attack in the field, and that the arm and leg were caught in its partial cicatrization. The woman made a good recovery.

[¹ *Am. Journ. Med. Sci.*, vol. xviii. p. 257.]

[² Communicated by the operator, 1880.]

[³ *N. O. Med. and Surg. Journ.*, July, 1877, p. 35; also communicated by operator, 1878.]

Much light is thrown upon a possible way of accounting for some of the mysterious cases of missed labor, which have been claimed to be extra-uterine in order to account for them, by a case recently operated upon in Portland, Maine, by Dr. Stanley P. Warren, and kindly reported to me by letter. The woman was a native, of Scotch-Irish descent, aged 32, and mother of a child of 13. She last menstruated in January, 1884. Supposed accidental abortion in May, as there was hemorrhage; the physician said he had removed the placenta, and there was a thick "molasses-like" discharge afterward. Dr. Warren was called in a week later; found metro-peritonitis and a tumor of about four inches in diameter in the right groin. The peritonitis became general, and Dr. W. was in attendance for 15 days. On July 1st the tumor was in the median line, and foetal movements and heart-sounds distinct. Labor expected about October 28th; subsequent gestation normal. Was called October 28th, at 11 P. M.; found no true pains; pain apparently abdominal, rather than uterine, and continuous in the back and over the sides of the uterus. Foetus transverse, with head to right; pulse 152. No change for several days. Second week in November found child dead. Next four weeks slight occasional chills, and temperature 102° for two or three nights, but usually normal. Absolutely no expulsive pains. Cervix reached with difficulty, and finger passed through a long tubular neck, but foetus not reached. Cervix absolutely closed from December 21st to 29th; pulse 120, temperature 100° to 102° . Attempted to dilate with sponge tent, but could not pass it into the uterine cavity. December 30th attempted to open cervix by digital dilatation, and succeeded finally in passing a cranioclast, but the parts closed as soon as the dilators were removed. Patient in a profound shock. After stimulating for an hour performed Cæsarean section; hemorrhage slight; peritoneum adherent everywhere to uterus; uterine wall $\frac{1}{2}$ inch thick; child presented by right arm and side; placenta thin and far advanced in fatty degeneration; no hemorrhage on its removal; uterus did not contract; sutured by continuous stitch with cat-gut. Child 8½ lbs. Woman rallied slightly, but died of shock in 28 hours. Drs. T. A. Foster and S. C. Gordon were associated with Dr. Warren in the management of the case.

It would appear in this instance of missed labor that the changes produced by metro-peritonitis prevented the natural dilatation of the cervix and the contractile action of the muscular coat of the uterus. Possibly, fatty degeneration of the muscular fibres had taken place, but this could not be ascertained, as there was no autopsy.

The Cæsarean case of Dr. Brodie S. Herndon of Fredericksburg, Virginia, operated upon with success in 1845, bears a close resemblance in many of its features to that of Dr. Warren. The subject was a white multipara of 30, whose pains of labor gave place to the continuous pain and other characteristic symptoms of peritonitis. This disease lasted a month, during which time the fluid contents of the uterus escaped, and the vaginal discharge became very offensive. Five weeks after the peritonitis commenced the os uteri admitted two fingers, and attempts at dilatation were made, but failed. Under ergot an offensive placenta was expelled, but the foetus could not be removed. The woman being

greatly wasted and her room filled with stench, the Cæsarean operation was performed on November 16th, 46 days after the first signs of labor appeared. The uterus being adherent, the peritoneal cavity was not exposed; the uterus was sponged out, but did not contract; it was closed in the suturing of the abdomen. The patient made a good recovery. As in the Warren case, the uterus became unsuited for performing the functions of labor by reason of changes in its tissues effected by inflammatory action.—Ed.]

Nov 15.
Nov. 8.

CHAPTER VII.

DISEASES OF PREGNANCY.

Diseases of Pregnancy.—The diseases of pregnancy form a subject so extensive that they might well of themselves furnish ample material for a separate treatise. The pregnant woman is of course liable to the same diseases as the non-pregnant, but it is only necessary to allude to those whose course and effects are essentially modified by the existence of pregnancy, or which have some peculiar effect on the patient in consequence of her condition. There are, moreover, many disorders which can be distinctly traced to the existence of pregnancy. Some of them are the direct results of the sympathetic irritations which are then so commonly observed; and of these several are only exaggerations of irritations which may be said to be normal accompaniments of gestation. These functional derangements may be classed under the head of neuroses, and they are sometimes so slight as merely to cause temporary inconvenience, at others so grave as seriously to imperil the life of the patient. Another class of disorders is to be traced to local causes in connection with the gravid uterus, and are either the mechanical results of pressure or of some displacement or morbid state of the uterus; while the origin of others may be said to be complex, being partly due to sympathetic irritation, partly to pressure, and partly to obscure nutritive changes produced by the pregnant state.

Derangements of the Digestive System.—Among the sympathetic derangements there are none which are more common, and none which more frequently produce distress and even danger, than those which affect the digestive system. Under the heading of "The Signs of Pregnancy" the frequent occurrence of nausea and vomiting has already been discussed, and its most probable causes considered (p. 145). A certain amount of nausea is, indeed, so common an accompaniment of pregnancy that its consideration as one of the normal symptoms of that state is fully justified. We need here only discuss those cases in which the nausea is excessive and long continued, and leads to serious results from inanition and from the constant distress it occasions. Fortunately, a pregnant woman may bear a surprising amount of nausea and sick-

ness without constitutional injury, so that apparently almost all aliments may be rejected without the nutrition of the body very materially suffering. At times the vomiting is limited to the early part of the day, when all food is rejected, and when there is a frequent retching of glairy transparent fluid, in several cases mixed with bile, while at the latter part of the day the stomach may be able to retain a sufficient quantity of food and the nausea disappears. In other cases the nausea and vomiting are almost incessant. The patient feels constantly sick, and the mere taste or sight of food may bring on excessive and painful vomiting. The duration of this distressing accompaniment of pregnancy is also variable. Generally it commences between the second and third months, and disappears after the woman has quickened. Sometimes, however, it begins with conception, and continues unabated until the pregnancy is over.

Symptoms of the Graver Cases.—In the worst class of cases, when all nourishment is rejected and when the retching is continuous and painful, symptoms of very great gravity, which may even prove fatal, develop themselves. The countenance becomes haggard from suffering, the tongue dry and coated, the epigastrium tender on pressure, and a state of extreme nervous irritability, attended with restlessness and loss of sleep, becomes established. In a still more aggravated degree there is general feverishness, with a rapid, small, and thready pulse. Extreme emaciation supervenes, the result of wasting from lack of nourishment. The breath is intensely fetid and the tongue dry and black. The vomited matters are sometimes mixed with blood. The patient becomes profoundly exhausted, a low form of delirium ensues, and death may follow if relief is not obtained.

Prognosis.—Symptoms of such gravity are fortunately of extreme rarity, but they do from time to time arise, and cause much anxiety. Gueniot collected 118 cases of this form of the disease, out of which 46 died; and out of the 72 that recovered, in 42 the symptoms only ceased when abortion, either spontaneous or artificially produced, had occurred. When pregnancy is over the symptoms occasionally cease with marvellous rapidity. The power of retaining and assimilating food is rapidly regained, and all the threatening symptoms disappear.

Treatment.—In the milder forms of obstinate vomiting one of the first indications will be to remedy any morbid state of the primæ viæ. The bowels will not unfrequently be found to be obstinately constipated, the tongue loaded, and the breath offensive; and when attention has been paid to the general state of the digestive organs by general aperient medicines and antacid remedies, such as bismuth and soda and liquor pepticus after meals, the tendency to vomiting may abate without further treatment.

Regulation of Diet.—The careful regulation of the diet is very important. Great benefit is often derived from recommending the patient not to rise from the recumbent position in the morning until she has taken something. Half a cup of milk and lime-water, or a cup of strong coffee, or a little rum and milk, or cocoa and milk, a glass of sparkling kouniss, or even a morsel of biscuit, taken on waking, often has a remarkable effect in diminishing the nausea. When any attempt at

allowing solid food brings on vomiting, it is better to give up all pretence of keeping to regular meals and to order such light and easily-digested food, at short intervals, as can be retained. Iced milk, with lime or soda-water, given frequently and not more than a mouthful at a time, will frequently be retained when nothing else will. Cold beef-jelly, a spoonful at a time, will also be often kept down. Sparkling lemonade has been strongly recommended as very useful in such cases, and is worthy of trial. It is well, however, to bear in mind, in regulating the diet, that the stomach is fickle and capricious, and that the patient may be able to retain strange and apparently unlikely articles of food, and that, if she express a desire for such, the experiment of feeding her have them should certainly be tried.

Medicinal Treatment.—The medicines that have been recommended are innumerable, and the practitioner will often have to try one after the other unsuccessfully, or may find, in an individual case, that a remedy will prove valuable which in another may be altogether powerless. Amongst those most generally useful are effervescent draughts containing from three to five minims of dilute hydrocyanic acid; the creasote mixture of the Pharmacopœia; tincture of *nux vomica*, in doses of five or ten minims; single minim-doses of *vinum ipecacuanhæ*, every hour in severe cases, three or four times daily in those which are less urgent; *salicin*, in doses of three to five grains three times a day, recommended by Tyler Smith; oxalate of cerium in the form of pill, of which three to five grains may be given three times a day—a remedy strongly advocated by Sir James Simpson, and which occasionally is of undoubted service, but more often fails; the compound pyroxylic spirit of the London Pharmacopœia, in doses of five minims every four hours, with a little compound tincture of cardamom—a drug which is comparatively little known, but which occasionally has a very marked and beneficial effect in checking vomiting; opiates in various forms—which sometimes prove useful, more often not—may be administered either by the mouth, in pills containing from half a grain to a grain of opium, or in small doses of the solution of the bimeconate of morphia or of Battley's sedative solution, or subcutaneously—a mode of administration which is much more often successful. If there is much tenderness about the epigastrium, one or two leeches may be advantageously applied, or one-third of a grain of morphia may be sprinkled on the surface of a small blister, or cloths saturated in landanum may be kept over the pit of the stomach. The administration per rectum of twenty grains of chloral, combined with the same amount of bromide of potassium, in a small enema, is said to be very useful. In many cases I have found that the application of a spinal ice-bag to the cervical vertebræ, in the manner recommended by Dr. Chapman, has checked the vomiting when all drugs have failed. The ice may be placed in one of Chapman's spinal ice-bags, and applied for half an hour or an hour twice or three times a day. It inevitably produces a comforting sensation of warmth, which is always agreeable to the patient. Ice may be given to suck *ad libitum*, and is very useful; while, if there be much exhaustion, small quantities of good champagne may also be given from time to time.

Local Treatment.—Inasmuch as the vomiting unquestionably has its

origin in the uterus, it is only natural that practitioners should endeavor to check it by remedies calculated to relieve the irritability of that organ. Thus, morphia in the form of pessaries per vaginam, or belladonna applied to the cervix, has been recommended; the former especially are often of undoubted service. A pessary containing one-third to half a grain of morphia may be introduced night and morning without interfering with other methods of treatment. Dr. Henry Bennet directs especial attention to the cervix, which, he says, is almost always congested and inflamed and covered with granular erosions. This condition he recommends to be treated by the application of nitrate of silver through the speculum. Dr. Clay of Manchester corroborates this view, and strongly advocates, especially when vomiting continues in the latter months, that one or two leeches should be applied to the cervix. Exception may fairly be taken to both these methods of treatment as being somewhat hazardous, unless other means have been tried and failed. I have little doubt, however, that in many cases a state of uterine congestion is an important factor in keeping up the unduly irritable condition of the uterine fibres, and an endeavor should always be made to lessen it by insisting on absolute rest in the recumbent posture. Of the importance of this precaution in obstinate cases there can be no question. Dr. Chapman of Norwich strongly recommended dilatation of the cervix by the finger, and stated that he found it very serviceable in checking nausea. It is obvious that this treatment must be adopted with great caution, as, roughly performed, it might lead to the production of abortion. Dr. Hewitt's views as to the dependence of sickness on flexions of the uterus have already been adverted to, and reasons have been given for doubting the general correctness of his theory. It is quite likely, however, that well-marked displacements of the uterus, either forward or backward, may serve to intensify the irritability of the organ. Cazeaux mentions an obstinate case immediately cured by replacing a retroverted uterus. A careful vaginal examination should therefore be instituted in all intractable cases, and if distinct displacement be detected an endeavor should be made to support the uterus in its normal axis. If retroverted, a Hodge's pessary may be safely employed; if anteverted, a small air-ball pessary, as recommended by Hewitt, should be inserted. I believe, however, that such displacements are the exception, rather than the rule, in cases of severe sickness.

Importance of Promoting the Nutrition of the Patient.—The importance of promoting nutrition by every means in our power should always be borne in mind. The effervescing koumiss, which can now be readily obtained, I have found of great value, as it can often be retained when all other aliment is rejected. The exhaustion produced by want of food soon increases the irritable state of the nervous system, and if the stomach will not retain anything, we can only combat it by occasional nutritive enemata of strong beef-tea, yolk of egg, and the like.

The Production of Artificial Abortion.—Finally, in the worst class of cases, when all treatment has failed and when the patient has fallen into the condition of extreme prostration already described, we may be driven to consider the necessity of producing abortion. Fortunately, cases justifying this extreme resource are of great rarity, but nevertheless

there is abundant evidence that every now and then women do die from uncontrollable vomiting whose lives might have been saved had the pregnancy been brought to an end. The value of artificial abortion has been abundantly proved. Indeed, it is remarkable how rapidly the serious symptoms disappear when the uterus is emptied and the tension of the uterine fibres lessened. It has fortunately but rarely fallen to my lot to have to perform this operation for intractable vomiting. In one such case the patient was reduced to a state of the utmost prostration, having kept hardly any food on her stomach for many weeks, and when I first saw her she was lying in a state of low muttering delirium. Within a few hours after abortion was induced all the threatening symptoms had disappeared, the vomiting had entirely ceased, and she was next day able to retain and absorb all that was given to her. The value of the operation, therefore, I believe to be undoubted. Where it has failed it seems to have been on account of undue delay. Owing to the natural repugnance which all must feel toward this plan, it has generally been postponed until the patient has been too exhausted to rally. If, therefore, it is done at all, it should be before prostration has advanced so far as to render the operation useless. In these cases the obvious indication is to lessen the tension of the uterus at once, and therefore the membranes should be punctured by the uterine sound, so as to let the liquor amnii drain away; and this may of itself be sufficient to accomplish the desired effect. It is almost needless to add that no one would be justified in resorting to this expedient without having his opinion fortified by consultation with a fellow-practitioner.

Other Disorders of the Digestive System.—Other disorders of the digestive system may give rise to considerable discomfort, but not to the serious peril attending obstinate vomiting. Amongst them are a loss of appetite, acidity and heartburn, flatulent distension, and sometimes a capricious appetite, which assumes the form of longing for strange and even disgusting articles of diet. Associated with these conditions there is generally derangement of the whole intestinal tract, indicated by furred tongue and sluggish bowels, and they are best treated by remedies calculated to restore a healthy condition of the digestive organs, such as a light, easily-digested diet, mineral acids, vegetable bitters, occasional aperients, bismuth and soda, and pepsin. The indications for treatment are not different from those which accompany the same symptoms in the non-pregnant state.

Diarrhœa.—Diarrhœa is an occasional accompaniment of pregnancy, often depending on errors of diet. When excessive and continuous it has a decided tendency to induce uterine contractions, and I have frequently observed premature labor to follow a sharp attack of diarrhœa. It should, therefore, not be neglected, and if at all excessive should be checked by the usual means, such as chalk mixture with aromatic confection and small doses of laudanum or chlorodyne. The possibility of apparent diarrhœa being associated with actual constipation, the fluid matter finding its way past the solid materials blocking up the intestines, should be borne in mind.

Constipation.—Constipation is much more common, and is indeed a very general accompaniment of pregnancy, even in women who do not

suffer from it at other times. It partly depends on the mechanical interference of the gravid uterus with the proper movements of the intestines, and partly on defective innervation of the bowels resulting from the altered state of the blood. The first indication will be to remedy this defect by appropriate diet, such as fresh fruits, brown bread, oatmeal porridge, etc. Some medicinal treatment will also be necessary, and in selecting the drugs to be used care should be taken to choose such as are mild and unirritating in their action and tend to improve the tone of the muscular coat of the intestine. A small quantity of aperient mineral water in the early morning, such as the Hunyadi, Friedrichshall, or Pullna water, often answers very well; or an occasional dose of the confection of sulphur; or a pill containing three or four grains of the extract of colocynth, with a quarter of a grain of the extract of nux vomica and a grain of extract of hyoscyamus at bedtime; or a teaspoonful of the compound liquorice powder in milk at bedtime. Constipation is also sometimes effectually combated by administering, twice daily, a pill containing a couple of grains of the inspissated ox-gall with a quarter of a grain of extract of belladonna. Enemata of soap and water are often very useful, and have the advantage of not disturbing the digestion. In the latter months of pregnancy, especially in the few weeks preceding delivery, the irritation produced by the collection of hardened feces in the bowel is a not infrequent cause of the annoying false pains which then so commonly trouble the patient. In order to relieve them it will be necessary to empty the bowels thoroughly by an aperient, such as a good dose of castor oil, to which fifteen or twenty minims of laudanum may be advantageously added. Should the rectum become loaded with scybalous masses, it may be necessary to break down and remove them by mechanical means, provided we are unable to effect this by copious enemata.

Hemorrhoids.—The loaded state of the rectum so common in pregnancy, combined with the mechanical effect of the pressure of the gravid uterus on the hemorrhoidal veins, often produces very troublesome symptoms from piles. In such cases a regular and gentle evacuation of the bowels should be secured daily, so as to lessen as much as possible the congestion of the veins. Any of the aperients already mentioned, especially the sulphur electuary, may be used. Dr. Fordyce Barker¹ insists that, contrary to the usual impression, one of the best remedies for this purpose is a pill containing a grain or a grain and a half of powdered aloes, with a quarter of a grain of extract of nux vomica, and that castor oil is distinctly prejudicial and apt to increase the symptoms. I have certainly found it answer well in several cases. When the piles are tender and swollen, they should be freely covered with an ointment consisting of four grains of muriate of morphia to an ounce of simple ointment, or with the ung. gallæ c. opio of the Pharmacopœia; and, if protruded, an attempt should be made to push them gently above the sphincter, by which they are often unduly constricted. Relief may also be obtained by frequent hot fomentations, and sometimes, when the piles are much swollen, it will be found useful to puncture them, so as to lessen the congestion before any attempt at reduction is made.

¹ *The Puerperal Diseases*, p. 33.

Ptyalism.—A profuse discharge from the salivary glands is an occasional distressing accompaniment of pregnancy. It is generally confined to the early months, but it occasionally continues during the whole period of gestation, and resists all treatment, only ceasing when delivery is over. Under such circumstances the discharge of saliva is sometimes enormous, amounting to several quarts a day, and the distress and annoyance to the patient are very great. In one case under my care the saliva poured from the mouth all day long, and for several months the patient sat with a basin constantly by her side, incessantly emptying her mouth, until she was reduced to a condition giving rise to really serious anxiety. This profuse salivation is no doubt a purely nervous disorder, and not readily controlled by remedies. Astringent gargles, containing tannin and chlorate of potass, frequent sucking of ice or of tannin lozenges, inhalation of turpentine and creasote, counter-irritation over the salivary glands by blisters or iodine, the bromides, opium internally, small doses of belladonna or atropine, may all be tried in turn, but none of them can be depended on with any degree of confidence.

Toothache and Caries of the Teeth.—Severe dental neuralgia is also a frequent accompaniment of pregnancy, especially in the early months. When purely neuralgic, quinine in tolerably large doses is the best remedy at our disposal; but not unfrequently it depends on actual caries of the teeth, and attention should always be paid to the condition of the teeth when facial neuralgia exists. There is no doubt that pregnancy predisposes to caries, and the observation of this fact has given rise to the old proverb, "For every child a tooth." Mr. Oakley Coles, in an interesting paper¹ on the condition of the mouth and teeth during pregnancy, refers the prevalence of caries to the coexistence of acid dyspepsia, causing acidity of the oral secretions. There is much unreasonable dread amongst practitioners as to interfering with the teeth during pregnancy, and some recommend that all operations, even stopping, should be postponed until after delivery. It seems to me certain that the suffering of severe toothache is likely to give rise to far more severe irritation than the operation required for its relief, and I have frequently seen badly-decayed teeth extracted during pregnancy, and with only a beneficial result.

Affections of the Respiratory Organs.—Amongst the derangements of the respiratory organs, one of the most common is spasmodic cough, which is often excessively troublesome. Like many other of the sympathetic derangements accompanying gestation, it is purely nervous in character, and is unaccompanied by elevated temperature, quickened pulse, or any distinct auscultatory phenomena. In character it is not unlike whooping cough. The treatment must obviously be guided by the character of the cough. Expectorants are not likely to be of service, while benefit may be derived from some of the antispasmodic class of drugs, such as belladonna, hydrocyanic acid, opiates, or bromide of potassium. Such remedies may be tried in succession, but will often be found to be of little value in arresting the cough. Dyspnoea may also be nervous in character, and sometimes symptoms not unlike those of

¹ *Trans. of the Odontological Society.*

spasmodic asthma are produced. Like the other sympathetic disorders, it, as well as nervous cough, is most frequently observed during the early months. There is another form of dyspnoea, not uncommonly met with, which is the mechanical result of the interference with the action of the diaphragm and lungs by the pressure of the enlarged uterus. Hence this is most generally troublesome in the latter months, and continues unrelieved until delivery or until the sinking of the uterine tumor which immediately precedes it. Beyond taking care that the pressure is not increased by tight-lacing or injudicious arrangement of the clothes, there is little that can be done to relieve this form of breathlessness.

[In some instances the difficulty of respiration is particularly distressing when the patient attempts to lie down in bed, and sleep is rendered broken and unrefreshing. In such cases two points are indicated: we must elevate the chest, and at the same time relieve the tension of the abdomen. This is best accomplished by the use of an inclined plane, in the form of a wide board padded with pillows, resting on the head and middle of the bed at its two ends. The patient is to rest her back upon this in a half-reclining position, and have her knees elevated with a pillow under them, on which she virtually, as it were, sits. This I have found to give great relief, especially to primiparæ, who are apt to suffer from diaphragmatic pressure and abdominal resistance. Inunction of the abdomen will also be found of value.—ED.]

Palpitation.—Palpitation, like dyspnoea, may be due either to sympathetic disturbance or to mechanical interference with the proper action of the heart. When occurring in weakly women, it may be referred to the functional derangements which accompany the chlorotic condition of the blood often associated with pregnancy, and is then best remedied by a general tonic regimen and the administration of ferruginous preparations. At other times antispasmodic remedies may be indicated, and it is seldom sufficiently serious to call for much special treatment.

Syncope.—Attacks of fainting are not rare, especially in delicate women of highly-developed nervous temperament, and are, perhaps, most common at or about the period of quickening. In most cases these attacks cannot be classed as cardiac, but are more probably nervous in character, and they are rarely associated with complete abolition of consciousness. They rather, therefore, resemble the condition described by the older authors as *lypothæmia*. The patient lies in a semi-unconscious condition with a feeble pulse and widely-dilated pupils, and this state lasts for varying periods, from a few minutes to half an hour or more. In one very troublesome case under my care they often recurred as frequently as three or four times a day. I have observed that they rarely occur when the more common sympathetic phenomena of pregnancy, especially vomiting, are present. Sometimes they terminate with the ordinary symptoms of hysteria, such as sobbing. The treatment should consist during the attack in the administration of diffusible stimulants, such as ether, sal-volatile, and valerian, the patient being placed in the recumbent position, with the head low. If frequently repeated it is unadvisable to attempt to rally the patient by the too free administration of stimulants. In the intervals a generally tonic regimen

and the administration of ferruginous remedies are indicated. If they recur with great frequency the daily application of the spinal ice-bag has proved of much service.

Extreme Anæmia and Chlorosis.—In connection with disorders of the circulatory system may be noticed those which depend on the state of the blood. The altered condition of the blood, which has already been described as a physiological accompaniment of pregnancy (p. 139), is sometimes carried to an extent which may fairly be called morbid; and either on account of the deficiency of blood-corpuscles or from the increase in its watery constituents a state of extreme anæmia and chlorosis may be developed. This may be sometimes carried to a very serious extent. Thus, Gusserow¹ records five cases in which nothing but excessive anæmia could be detected, all of which ended fatally. Generally, when such symptoms have been carried to an extreme extent, the patient has been in a state of chlorosis before pregnancy. The treatment must, of course, be calculated to improve the general nutrition and enrich the impoverished blood; a light and easily-assimilated diet, milk, eggs, beef-tea, and animal food—if it can be taken—attention to the proper action of the bowels, a due amount of stimulants, and abundance of fresh air, will be the chief indications in the general management of the case. Medicinally, ferruginous preparations will be required. Some practitioners object, apparently without sufficient reason, to the administration of iron during pregnancy as liable to promote abortion. This unfounded prejudice may probably be traced to the supposed emmenagogue properties of the preparations of iron; but if the general condition of the patient indicate such medication they may be administered without any fear. Preparations of phosphorus, such as the phosphide of zinc, or free phosphorus, also promise favorably, and are worthy of trial.

Edema associated with Hydræmia.—Some of the more aggravated cases are associated with a considerable amount of serous effusion into the cellular tissue, generally limited to the lower extremities, but occasionally extending to the arms, face, and neck, and even producing ascites and pleuritic effusion. Under the latter circumstances this complication is, of course, of great gravity, and it is said that after delivery the disappearance of the serous effusion may be accompanied by metastasis of a fatal character to the lungs or the nervous centres. This form of œdema must be distinguished from the slight œdematous swelling of the feet and legs so commonly observed as a mechanical result of the pressure of the gravid uterus, and also from those cases of œdema associated with albuminuria. The treatment must be directed to the cause, while the disappearance of the effusion may be promoted by the administration of diuretic drinks, the occasional use of saline aperients, and rest in the horizontal position.

Albuminuria.—The existence of albumen in the urine of pregnant women has for many years attracted the attention of obstetricians, and it is now well known to be associated, in ways still imperfectly understood, with many important puerperal diseases. Its presence in most cases of puerperal eclampsia was long ago pointed out by Lever in this country

¹ *Arch. f. Gyn.*, ii. 2, 1871.

and Rayer in France, and its association with this disease gave rise to the theory of the dependence of the convulsion on uræmia which is generally still entertained. It has been shown of late years, especially by Braxton Hicks, that this association is by no means so universal as was supposed ; or, rather, that in some cases the albuminuria follows and does not precede the convulsions, of which it might therefore be supposed to be the consequence rather than the cause ; so that further investigations as to these particular points are still required. Modern researches have shown that there is an intimate connection between many other affections and albuminuria ; as, for example, certain forms of paralysis, either of special nerves, as puerperal amaurosis, or of the spinal system ; cephalalgia and dizziness ; puerperal mania ; and possibly hemorrhage. It cannot, therefore, be doubted that albuminuria in the pregnant woman is liable, at any rate, to be associated with grave disease, although the present state of our knowledge does not enable us to define very distinctly its precise mode of action.

Causes of Puerperal Albuminuria.—The presence of albumen in the urine of pregnant women is far from a rare phenomenon. Blot and Litzman met with albuminuria in 20 per cent. of pregnant women, which is, however, far above the estimate of other authors ; Fordyce Barker¹ thinks it occurs in about 1 out of 25 cases, or 4 per cent. ; while Hofmier² found it in 137 out of 5000 deliveries in the Berlin Gynæcological Institution, or 2.74 per cent. As in the large majority of these cases it rapidly disappears after delivery, it is obvious that its presence must, in a large portion of cases, depend on temporary causes, and has not always the same serious importance as in the non-pregnant state. This is further proved by the undoubted fact that albumen, rapidly disappearing after delivery, is often found in urine of pregnant women who go to term and pass through labor without any unfavorable symptoms.

Pressure by the Gravid Uterus.—The obvious facts that in pregnancy the vessels supplying the kidneys are subjected to mechanical pressure from the gravid uterus, and that congestion of the venous circulation of those viscera must necessarily exist to a greater or less degree, suggest that here we may find an explanation of the frequent occurrence of albuminuria. This view is further strengthened by the fact that the albumen rarely appears until after the fifth month, and therefore not until the uterus has attained a considerable size ; and also that it is comparatively more frequently met with in primiparæ, in whom the resistance of the abdominal parietes, and consequent pressure, must be greater than in women who have already borne children. It is indeed probable that pressure and consequent venous congestion of the kidneys have an important influence in its production ; but there must be, as a rule, some other factors in operation, since an equal or even greater amount of pressure is often exerted by ovarian and fibroid tumors without any such consequences. They are probably complex. One important condition is doubtless the increased amount of work the kidneys have to do in excreting the waste products of the fœtus, as well as those of the mother. The increased arterial tension throughout the body, associated with hypertrophy of the heart, known to exist in pregnancy, also operates in

¹ *American Journal of Obstetrics*, July, 1878.

² *Berlin klin. Woch.*, Sept., 1878.

and the administration of ferruginous remedies are indicated. If they recur with great frequency the daily application of the spinal ice-bag has proved of much service.

Extreme Anæmia and Chlorosis.—In connection with disorders of the circulatory system may be noticed those which depend on the state of the blood. The altered condition of the blood, which has already been described as a physiological accompaniment of pregnancy (p. 139), is sometimes carried to an extent which may fairly be called morbid; and either on account of the deficiency of blood-corpuscles or from the increase in its watery constituents a state of extreme anæmia and chlorosis may be developed. This may be sometimes carried to a very serious extent. Thus, Gusserow¹ records five cases in which nothing but excessive anæmia could be detected, all of which ended fatally. Generally, when such symptoms have been carried to an extreme extent, the patient has been in a state of chlorosis before pregnancy. The treatment must, of course, be calculated to improve the general nutrition and enrich the impoverished blood; a light and easily-assimilated diet, milk, eggs, beef-tea, and animal food—if it can be taken—attention to the proper action of the bowels, a due amount of stimulants, and abundance of fresh air, will be the chief indications in the general management of the case. Medicinally, ferruginous preparations will be required. Some practitioners object, apparently without sufficient reason, to the administration of iron during pregnancy as liable to promote abortion. This unfounded prejudice may probably be traced to the supposed emmenagogue properties of the preparations of iron; but if the general condition of the patient indicate such medication they may be administered without any fear. Preparations of phosphorus, such as the phosphide of zinc, or free phosphorus, also promise favorably, and are worthy of trial.

Edema associated with Hydræmia.—Some of the more aggravated cases are associated with a considerable amount of serous effusion into the cellular tissue, generally limited to the lower extremities, but occasionally extending to the arms, face, and neck, and even producing ascites and pleuritic effusion. Under the latter circumstances this complication is, of course, of great gravity, and it is said that after delivery the disappearance of the serous effusion may be accompanied by metastasis of a fatal character to the lungs or the nervous centres. This form of œdema must be distinguished from the slight œdematous swelling of the feet and legs so commonly observed as a mechanical result of the pressure of the gravid uterus, and also from those cases of œdema associated with albuminuria. The treatment must be directed to the cause, while the disappearance of the effusion may be promoted by the administration of diuretic drinks, the occasional use of saline aperients, and rest in the horizontal position.

Albuminuria.—The existence of albumen in the urine of pregnant women has for many years attracted the attention of obstetricians, and it is now well known to be associated, in ways still imperfectly understood, with many important puerperal diseases. Its presence in most cases of puerperal eclampsia was long ago pointed out by Lever in this country

¹ *Arch. f. Gyn.*, ii. 2, 1871.

by the drain of albuminous materials through the kidneys. This fact has been observed by many writers. A good illustration of it is given by Tanner,¹ who states that four out of seven women he attended suffering from Bright's disease during pregnancy aborted, one of them three times in succession.

Symptoms.—The symptoms accompanying albuminuria in pregnancy are by no means uniform or constantly present. That which most frequently causes suspicion is the anasarca—not only the cedematous swelling of the lower limbs which is so common a consequence of the pressure of the gravid uterus, but also of the face and upper extremities. Any puffiness or infiltration about the face, or any cedema about the hands or arms, should always give rise to suspicion and lead to a careful examination of the urine. Sometimes this is carried to an exaggerated degree, so that there is anasarca of the whole body.

Nervous Phenomena.—Anomalous nervous symptoms—such as headache, transient dizziness, dimness of vision, spots before the eyes, inability to see objects distinctly, sickness in women not at other times suffering from nausea, sleeplessness, irritability of temper—are also often met with, sometimes to a slight degree, at others very strongly developed, and should always arouse suspicion. Indeed, knowing as we do that many morbid states may be associated with albuminuria, we should make a point of carefully examining the urine of all patients in whom any unusually morbid phenomena show themselves during pregnancy.

Character of the Urine.—The condition of the urine varies considerably, but it is generally scanty and highly colored, and, in addition to the albumen, especially in cases in which the albuminuria has existed for some time, we may find epithelium-cells, tube-casts, and occasionally blood-corpuscles.

Treatment.—The treatment must be based on what has been said as to the causes of the albuminuria. Of course it is out of our power to remove the pressure of the gravid uterus, except by inducing labor; but its effects may at least be lessened by remedies tending to promote an increased secretion of urine, and thus diminishing the congestion of the renal vessels. The administration of saline diuretics, such as the acetate of potash or bitartrate of potash, the latter being given in the form of the well-known imperial drink, will best answer this indication. The action of the bowels may be solicited by purgatives producing watery motions, such as occasional doses of the compound jalap powder. Dry cupping over the loins, frequently repeated, has a beneficial effect in lessening the renal hyperæmia. The action of the skin should also be promoted by the use of the vapor bath, and with this view the Turkish bath may be employed with great benefit and perfect safety. Jaborandi and pilocarpin have been given for this purpose, but have been found by Fordyce Barker to produce a dangerous degree of depression. The next indication is to improve the condition of the blood by appropriate diet and medication. A very light and easily-assimilated diet should be ordered, of which milk should form the staple. Tarnier² has recorded several cases in which a purely milk diet was very successful

¹ *Signs and Diseases of Pregnancy*, p. 428.

² *Annul. de Gynéc.*, Jan., 1876.

the same direction. But in the large majority of cases, although these conditions are present, no albuminuria exists, and they must therefore be looked upon as predisposing causes, to which some other is added before the albumen escapes from the vessels. What this is, generally escapes our observation, but probably any condition producing sudden hyperæmia of the kidneys and giving rise to a state analogous to the first stage of Bright's disease—such, for example, as sudden exposure to cold and impeded cutaneous action—may be sufficient to set a light to the match already prepared by the existence of pregnancy. It has more recently been pointed out that a transient albuminuria, disappearing in a few days, is very common after delivery, and probably depends on a catarrhal condition of the urinary tract. Ingersten observed this in 50 out of 153 deliveries, and in 15 only had any albumen existed before the confinement.¹ In addition to these temporary causes it must not be forgotten that pregnancy may supervene in a patient already suffering from Bright's disease, when, of course, the albumen will exist in the urine from the commencement of gestation.

The Effects of Puerperal Albuminuria.—The various diseases associated with the presence of albumen in the urine will require separate consideration. Some of these, especially puerperal eclampsia, are amongst the most dangerous complications of pregnancy. Others, such as paralysis, cephalalgia, dizziness, may also be of considerable gravity. The precise mode of their production, and whether they can be traced, as is generally believed, to the retention of urinary elements in the blood, either urea or free carbonate of ammonia produced by its decomposition, or whether the two are only common results of some undetermined cause, will be considered when we come to discuss puerperal convulsions. Whatever view may ultimately be taken on these points, it is sufficiently obvious that albuminuria in a pregnant woman must constantly be a source of much anxiety, and must induce us to look forward with considerable apprehension to the termination of the case.

Prognosis.—We are scarcely in possession of a sufficiently large number of observations to justify any very accurate conclusions as to the risk attending albuminuria during pregnancy, but it is certainly by no means slight. Hofmïer believes that albuminuria is a most severe complication both for woman and child, even when uncomplicated with eclampsia. The prognosis, he thinks, depends on whether it is acute in its onset—that is, coming on within a few days of labor—or is extended over several weeks. The former is more likely to pass entirely away after delivery, while in the latter there is more risk of the morbid state of the kidneys becoming permanent and leading to the establishment of Bright's disease after the pregnancy is over. Goubeyre estimated that 49 per cent. of primiparæ who have albuminuria, and who escape eclampsia, die from morbid conditions traceable to the albuminuria. This conclusion is probably much exaggerated, but if it even approximate to the truth the danger must be very great.

Tendency to Produce Abortion.—Besides the ultimate risk to the mother, albuminuria strongly predisposes to abortion, no doubt on account of the imperfect nutrition of the fœtus by blood impoverished

¹ *Zeitschrift f. Geburt*, Band v. Heft 2.

CHAPTER VIII.

DISEASES OF PREGNANCY (CONTINUED).

Disorders of the Nervous System.—There are many disorders of the nervous system met with during the course of pregnancy. Among the most common are morbid irritability of temper or a state of mental despondency and dread of the results of the labor, sometimes almost amounting to insanity or even progressing to actual mania. These are but exaggerations of the highly susceptible state of the nervous system generally associated with gestation. Want of sleep is not uncommon, and, if carried to any great extent, may cause serious trouble from the irritability and exhaustion it produces. In such cases we should endeavor to lessen the excitable state of the nerves by insisting on the avoidance of late hours, overmuch society, exciting amusements, and the like; while it may be essential to promote sleep by the administration of sedatives, none answering so well as the chloral hydrate, in combination with large doses of the bromide of potassium or sodium, which greatly intensify its hypnotic effects.

Headaches and Neuralgicæ.—Severe headaches and various intense neuralgicæ are common. Amongst the latter the most frequently met with are pain in the breasts, due to the intimate sympathetic connection of the mammae with the gravid uterus, and intense intercostal neuralgia, which a careless observer might mistake for pleuritic or inflammatory pain. The thermometer, by showing that there is no elevation of temperature, would prevent such a mistake. Neuralgia of the uterus itself or severe pains in the groins or thighs—the latter being probably the mechanical results of dragging on the attachments of the abdominal muscles—are also far from uncommon. In the treatment of such neuralgic affections attention to the state of the general health, and large doses of quinine and ferruginous preparations whenever there is much debility, will be indicated. Locally sedative applications, such as belladonna and chloroform liniments, friction with aconite liniment when the pain is limited to a small space, and, in the worst cases, the subcutaneous injection of morphia, will be called for. Those pains which apparently depend on mechanical causes may often be best relieved by lessening the traction on the muscles by wearing a well-made elastic belt to support the uterus.

Paralysis depending on Pregnancy.—Among the most interesting of the nervous diseases are various paralytic affections. Almost all varieties of paralysis have been observed, such as paraplegia, hemiplegia (complete or incomplete), facial paralysis, and paralysis of the nerves of special sense, giving rise to amaurosis, deafness, and loss of taste. Churchill records 22 cases of paralysis during pregnancy, collected by him from various sources. A large number have also been brought together by Imbert Goubeyre in an interesting memoir on the subject,

and others are recorded by Fordyce Barker, Joulin, and other authors, so that there can be no doubt of the fact that paralytic affections are common during gestation. In a large proportion of the cases recorded the paralyzes have been associated with albuminuria, and are doubtless uræmic in origin. Thus in 19 cases related by Goubeyre albuminuria was present in all; Darcy,¹ however, found no albuminuria in 5 out of 14 cases. The dependency of the paralysis on a transient cause explains the fact that in the large majority of these cases the paralysis was not permanent, but disappeared shortly after labor. In every case of paralysis, whatever be its nature, special attention should be directed to the state of the urine, and, should it be found to be albuminous, labor should be at once induced. This is clearly the proper course to pursue, and we should certainly not be justified in running the risk that must attend the progress of a case in which so formidable a symptom has already developed itself. When the cause has been removed the effect will also generally rapidly disappear, and the prognosis is therefore, on the whole, favorable. Should the paralysis continue after delivery, the treatment must be such as we would adopt in the non-pregnant state, and small doses of strychnia, along with faradization of the affected limbs, would be the best remedies at our disposal.

Paralyses which are not Uræmic in their Origin.—There are, however, unquestionably some cases of puerperal paralysis which are not uræmic in their origin and the nature of which is somewhat obscure. Hemiplegia may doubtless be occasioned by cerebral hemorrhage, as in the non-pregnant state. Other organic causes of paralysis, such as cerebral congestion or embolism, may now and again be met with during pregnancy, but cases of this kind must be of comparative rarity. Other cases are functional in their origin. Tarnier relates a case of hemiplegia which he could only refer to extreme anæmia. Some, again, may be hysterical. Paraplegia is apparently more frequently unconnected with albuminuria than the other forms of paralysis, and it may either depend on pressure of the gravid uterus on the nerves as they pass through the pelvis or on reflex action, as is sometimes observed in connection with uterine disease. When in such cases the absence of albuminuria is ascertained by frequent examination of the urine, there is obviously not the same risk to the patient as in cases depending on uræmia, and therefore it may be justifiable to allow pregnancy to go on to term, trusting to subsequent general treatment to remove the paralytic symptoms. As the loss of power here depends on a transient cause, a favorable prognosis is quite justifiable. Partial paralysis of one lower extremity, generally the left, sometimes occurs from pressure of the foetal occiput, and may continue for days or weeks, with a gradual improvement, after parturition.

Chorea.—Chorea is not unfrequently observed, and forms a serious complication. It is generally met with in young women of delicate health and in the first pregnancy. In a large proportion of the cases the patient has already suffered from the disease before marriage. On the occurrence of pregnancy the disposition to the disease again becomes evoked and choreic movements are re-established. This fact may be

¹ *Thèse de Paris*, 1877.

explained partly by the susceptible state of the nervous system, partly by the impoverished condition of the blood.

Prognosis.—That chorea is a dangerous complication of pregnancy is apparent by the fact that out of 56 cases collected by Dr. Barnes¹ no less than 17, or 1 to 3, proved fatal. Nor is it danger to life alone that is to be feared, for it appears certain that chorea is more apt to leave permanent mental disturbance when it occurs during pregnancy than at other times. It has also an unquestionable tendency to bring on abortion or premature labor, and in most cases the life of the child is sacrificed.

Treatment.—The treatment of chorea during pregnancy does not differ from that of the disease under more ordinary circumstances, and our chief reliance will be placed on such drugs as the liquor arsenicalis, bromide of potassium, and iron. In the severe form of the disease the incessant movements and the weariness and loss of sleep may very seriously imperil the life of the patient, and more prompt and radical measures will be indicated. If, in spite of our remedies, the paroxysms go on increasing in severity and the patient's strength appears to be exhausted, our only resource is to remove the most evident cause by inducing labor. Generally, the symptoms lessen and disappear soon after this is done. There can be no question that the operation is perfectly justifiable, and may even be essential under such circumstances. It should be borne in mind that the chorea often recurs in a subsequent pregnancy, and extra care should then always be taken to prevent its development.

Disorders of the Urinary Organs.—Disorders of the urinary organs are of frequent occurrence. Retention of urine may be met with, and this is often the result of a retroverted uterus. The treatment, therefore, must then be directed to the removal of the cause. This subject will be more particularly considered when we come to discuss that form of displacement (p. 216); but we may here point out that retention of urine, if long continued, may not only lead to much distress, but to actual disease of the coats of the bladder. Several cases have been recorded in which cystitis, resulting from urinary retention in pregnancy, eventually caused the exfoliation of the entire mucous membrane of the bladder,² which was cast off, sometimes entire, sometimes in shreds, and occasionally with portions of the muscular coat attached to it. The possibility of this formidable accident should teach us to be careful not to allow any undue retention of urine, but by a timely use of the catheter to relieve the symptoms, while we at the same time endeavor to remove the cause.

Irritability of the Bladder.—Irritability of the bladder is of frequent occurrence. In the early months it seems to be the consequence of sympathetic irritation of the neck of the bladder, combined with pressure, while in the later months it is probably solely produced by mechanical causes. When severe it leads to much distress, the patient's rest being broken and disturbed by incessant calls to micturate, and the suffering induced may produce serious constitutional disturbances. I have elsewhere pointed out³ that irritability of the bladder in the later months of pregnancy is frequently associated with an abnormal position of the

¹ *Obst. Trans.*, vol. x.

² *Ibid.*, vol. xi.

³ *Ibid.*, vol. xiii.

fœtus, which is placed transversely or obliquely. The result is either that undue pressure is applied to the bladder or that it is drawn out of its proper position. The abnormal position of the fœtus can readily be detected by palpation, and is readily altered by external manipulation. In some of the cases I have recorded altering the position of the fœtus was immediately followed by relief, the symptoms recurring after a time when the fœtus had again assumed an oblique position. Should the fœtus frequently become displaced, an endeavor may be made to retain it in the longitudinal axis of the uterus by a proper adaptation of bandages or pads. In cases not referable to this cause we should attempt to relieve the bladder symptoms by appropriate medication, such as small doses of liquor potassæ if the urine be very acid; tincture of belladonna; the decoction of triticum repens, an old but very serviceable remedy; and vaginal sedative pessaries containing morphia or atropine.

[In one case where a lady had borne two children with very little inconvenience I found great suffering from the pressure of the fœtus on the bladder, commencing as early as the fifth month. This continued for a period of two months, when she very fortunately miscarried. In making a digital exploration I recognized that the fœtus was anencephalous, and for this reason descended too low in the pelvis.—ED.]

Incontinence of Urine.—Women who have borne many children are often troubled with incontinence of urine during pregnancy, the water dribbling away on the slightest movement. Through this much irritation of the skin surrounding the genitals is produced, attended with troublesome excoriations and eruptions. Relief may be partially obtained by lessening the pressure on the bladder by an abdominal belt, while the skin is protected by applications of simple ointment or glycerin.

Phosphatic Deposit.—Dr. Tyler Smith has directed attention to a phosphatic condition of the urine occurring in delicate women, whose constitutions are severely tried by gestation. This condition can easily be altered by rest, nutritious diet, and a course of restorative medicines, such as steel, mineral acids, and the like.

Leucorrhœa.—A profuse whitish leucorrhœal discharge is very common during pregnancy, especially in its latter half. The discharge frequently alarms the patient, but, unless it is attended with disagreeable symptoms, it does not call for special treatment. When at all excessive it may lead to much irritation of the vagina and external generative organs. The labia may become excoriated and covered with small aphthous patches, and the whole vulva may be hot, swollen, and tender. Warty growths, similar in appearance to syphilitic condylomata, are occasionally developed in pregnant women, unconnected with any specific taint and associated with the presence of an irritating leucorrhœal discharge. According to Thibierge,¹ these resist local applications, such as sulphate of copper or nitrate of silver, but spontaneously disappear after delivery. Inasmuch as the leucorrhœal discharge is dependent on the congested condition of the generative organs accompanying pregnancy, we can hope to do little more than alleviate it. In the severer forms, as has been pointed out by Henry Bennet, the cervix will be found to be

¹ *Arch. gén. de Méd.*, 1856.

abraded or covered with granular erosion, and it may be from time to time cautiously touched with the nitrate of silver or a solution of carbolic acid. Generally speaking, we must content ourselves with recommending the patient to wash the vagina out gently with diluted Condyl's fluid, or with a solution of the sulpho-carbolate of zinc of the strength of four grains to the ounce of water, or with plain tepid water. For obvious reasons, frequent and strong vaginal douches are to be avoided, but a daily gentle injection, for the purpose of ablution, can do no harm.

Pruritus.—A very distressing pruritus of the vulva is frequently met with along with leucorrhœa, especially when the discharge is of an acrid character, which in some cases leads to intense and protracted suffering, forcing the patient to resort to incessant friction of the parts. Pruritus, however, may exist without leucorrhœa, being apparently sometimes of a neuralgic character, at others associated with aphthous patches on the mucous membrane, ascarides in the rectum, or pediculi in the hairs of the mons veneris and labia. Cases are even recorded in which the pruritic irritation extended over the whole body. The treatment is difficult and unsatisfactory. Various sedative applications may be tried, such as weak solutions of Goulard's lotion, or a lotion composed of an ounce of the solution of the muriate of morphia, with a drachm and a half of hydrocyanic acid, in six ounces of water, or one formed by mixing one part of chloroform with six of almond oil. A very useful form of medication consists in the insertion into the vagina of a pledget of cotton wool, soaked in equal parts of the glycerin of borax and sulphurous acid; this may be inserted at bedtime, and withdrawn in the morning by means of a string attached to it. Smearing the parts with an ointment consisting of boracic acid and vaseline often answers admirably. In the more obstinate cases the solid nitrate of silver may be lightly brushed over the vulva, or, as recommended by Tarnier, a solution of bichloride of mercury, of about the strength of two grains to the ounce, may be applied night and morning. The state of the digestive organs should always be attended to, and aperient mineral water may be usefully administered. When the pruritus extends beyond the vulva, or even in severe local cases, large doses of bromide of potassium may perhaps be useful in lessening the general hyperæsthetic state of the nerves.

Œdema of the Lower Limbs.—Some of the disorders of pregnancy are the direct results of the mechanical pressure of the gravid uterus. The most common of these are œdema and a varicose state of the veins of the lower extremities, or even of the vulva. The former is of little consequence, provided we have assured ourselves that it is really the result of pressure, and not of albuminuria, and it can generally be relieved by rest in the horizontal position. A varicose state of the veins of the lower limbs is very common, especially in multiparæ, in whom it is apt to continue after delivery. Occasionally the veins of the vulva, and even of the vagina, are also enlarged and varicose, producing considerable swelling of the external genitals. Rest in the recumbent position and the use of an abdominal belt, so as to take the pressure off the veins as much as possible, are all that can be done to relieve this trouble-

some complication. If the veins of the legs are much swollen, some benefit may be derived from an elastic stocking or a carefully applied bandage.

Occasional Serious Results from Laceration of the Veins.—Serious and even fatal consequences have followed the accidental laceration of the swollen veins. When laceration occurs during or immediately after delivery—a not uncommon result of the pressure of the head—it gives rise to the formation of a vaginal thrombus. It has occasionally happened from an accidental injury during pregnancy, as in the cases recorded by Simpson, in which death followed a kick on the pudenda, producing laceration of a varicose vein, or in one mentioned by Tarnier, where the patient fell on the edge of a chair. Severe hemorrhage has followed the accidental rupture of a vein in the leg. The only satisfactory treatment is pressure, applied directly to the bleeding parts by means of the finger or by compresses saturated in a solution of the perchloride of iron. The treatment of vaginal thrombus following labor must be considered elsewhere. Occasionally the varicose veins inflame, become very tender and painful, and coagula form in their canals. In such cases absolute rest should be insisted on, while sedative lotions, such as the chloroform and belladonna liniments, should be applied to relieve the pain.

Displacements of the Gravid Uterus.—Certain displacements of the gravid uterus are met with which may give rise to symptoms of great gravity.

Prolapse, which is rare, is almost always the result of pregnancy occurring in a uterus which had been previously more or less procident. Under such circumstances the increasing weight of the uterus will at first necessarily augment the previously existing tendency to prolapse of the womb, which may come to protrude partially or entirely beyond the vulva. In the great majority of cases, as pregnancy advances, the prolapse cures itself, for at about the fourth or fifth month the uterus will rise above the pelvic brim. It has been said that in some cases of complete procidentia pregnancy has gone even to term with the uterus lying entirely outside the vulva. Most probably these cases were imperfectly observed, the greater part of the uterus being in reality above the pelvic brim, a portion only of its lower segment protruding externally; or, as has sometimes been the case, the protruding portion has been an old-standing hypertrophic elongation of the cervix, the internal os uteri and fundus being normally situated. Should a prolapsed uterus not rise into the abdominal cavity as pregnancy advances, serious symptoms will be apt to develop themselves; for, unless the pelvis be unusually capacious, the enlarging uterus will get jammed within its bony walls, the rectum and urethra will be pressed upon, defecation and micturition will be consequently impeded, and severe pain and much irritation will result. In all probability such a state of things would lead to abortion. The possibility of these consequences should therefore teach us to be careful in the management of every case of prolapse, however slight, in which pregnancy occurs. Absolute rest in the horizontal position should be insisted on, while the uterus should be supported in the pelvis by a full-sized Hodge's pessary, which should be worn until at least the

sixth month, when the uterus would be fully within the abdominal cavity. After delivery prolonged rest should be recommended, in the hope that the process of involution may be accompanied by a cure of the prolapse. There can be no doubt that pregnancy carried to term affords an opportunity of curing even old-standing displacements which should not be neglected.

Anteversión of the gravid uterus seldom produces symptoms of consequence. In all probability it is common enough when pregnancy occurs in a uterus which is more than usually anteverted or is anteflexed. Under such circumstances there is not the same risk of incarceration in the pelvic cavity as in cases in which pregnancy exists in a retroflexed uterus, for as the uterus increases in size it rises without difficulty into the abdominal cavity. In the early months the pressure of the fundus on the bladder may account for the irritability of that viscus then so commonly observed. It will be remembered that Graily Hewitt attributes great importance to this condition as explaining the sickness of pregnancy—a theory, however, which has not met with general acceptance.

Extreme anteversion of the uterus at an advanced period of pregnancy is sometimes observed in multiparæ with very lax abdominal walls, occasionally to such an extent that the uterus falls completely forward and downward, so that the fundus is almost on a level with the patient's knees. This form of pendulous belly may be associated with a separation of the recti muscles, between which the womb forms a ventral hernia, covered only by the cutaneous textures. When labor comes on, this variety of displacement may give rise to trouble by destroying the proper relation of the uterine and pelvic axes. The treatment is purely mechanical, keeping the patient lying on her back as much as possible and supporting the pendulous abdomen by a properly adjusted bandage. A similar forward displacement is observed in cases of pelvic deformity, and in the worst forms, in rachitic and dwarfed women, it exists to a very exaggerated degree.

Retroversion.—The most important of the displacements, in consequence of its occasional very serious results, is retroversion of the gravid uterus. It was formerly generally believed that this was most commonly produced by some accident, such as a fall, which dislocated a uterus previously in a normal position. Undue distension of the bladder was also considered to have an important influence in its production by pressing the uterus backward and downward.

Its Causes.—It is now almost universally admitted that, although the above-named causes may possibly sometimes produce it, in the very large proportion of cases it depends on pregnancy having occurred in a uterus previously retroverted or retroflexed. The merit of pointing out this fact unquestionably belongs to the late Dr. Tyler Smith, and further observations have fully corroborated the correctness of his views.

In the large majority of cases in which pregnancy occurs in a uterus so displaced, as the womb enlarges it straightens itself and rises into the abdominal cavity, without giving any particular trouble; or, as not unfrequently happens, the abnormal position of the organ interferes so much with its enlargement as to produce abortion. Sometimes, however,

some complication. If the veins of the legs are much swollen, some benefit may be derived from an elastic stocking or a carefully applied bandage.

Occasional Serious Results from Laceration of the Veins.—Serious and even fatal consequences have followed the accidental laceration of the swollen veins. When laceration occurs during or immediately after delivery—a not uncommon result of the pressure of the head—it gives rise to the formation of a vaginal thrombus. It has occasionally happened from an accidental injury during pregnancy, as in the cases recorded by Simpson, in which death followed a kick on the pudenda, producing laceration of a varicose vein, or in one mentioned by Tarnier, where the patient fell on the edge of a chair. Severe hemorrhage has followed the accidental rupture of a vein in the leg. The only satisfactory treatment is pressure, applied directly to the bleeding parts by means of the finger or by compresses saturated in a solution of the perchloride of iron. The treatment of vaginal thrombus following labor must be considered elsewhere. Occasionally the varicose veins inflame, become very tender and painful, and coagula form in their canals. In such cases absolute rest should be insisted on, while sedative lotions, such as the chloroform and belladonna liniments, should be applied to relieve the pain.

Displacements of the Gravid Uterus.—Certain displacements of the gravid uterus are met with which may give rise to symptoms of great gravity.

Prolapse, which is rare, is almost always the result of pregnancy occurring in a uterus which had been previously more or less procident. Under such circumstances the increasing weight of the uterus will at first necessarily augment the previously existing tendency to prolapse of the womb, which may come to protrude partially or entirely beyond the vulva. In the great majority of cases, as pregnancy advances, the prolapse cures itself, for at about the fourth or fifth month the uterus will rise above the pelvic brim. It has been said that in some cases of complete procidentia pregnancy has gone even to term with the uterus lying entirely outside the vulva. Most probably these cases were imperfectly observed, the greater part of the uterus being in reality above the pelvic brim, a portion only of its lower segment protruding externally; or, as has sometimes been the case, the protruding portion has been an old-standing hypertrophic elongation of the cervix, the internal os uteri and fundus being normally situated. Should a prolapsed uterus not rise into the abdominal cavity as pregnancy advances, serious symptoms will be apt to develop themselves; for, unless the pelvis be unusually capacious, the enlarging uterus will get jammed within its bony walls, the rectum and urethra will be pressed upon, defecation and micturition will be consequently impeded, and severe pain and much irritation will result. In all probability such a state of things would lead to abortion. The possibility of these consequences should therefore teach us to be careful in the management of every case of prolapse, however slight, in which pregnancy occurs. Absolute rest in the horizontal position should be insisted on, while the uterus should be supported in the pelvis by a full-sized Hodge's pessary, which should be worn until at least the

should, under natural circumstances, always be possible. By bi-manual examination we can make out, with due care, the alternate relaxation and contraction of the uterine parietes characteristic of the gravid uterus, and so differentiate the swelling from any other in the same situation. The accompanying phenomena of pregnancy will also prevent any mistake of this kind.

Retroversion going on to Term.—In some few cases retroversion has been supposed to go on to term. Strictly speaking, this is impossible; but in the supposed examples, such as the well-known case recorded by Oldham, part of a retroflexed uterus remained in the pelvic cavity, while the greater part developed in the abdominal cavity. The uterus is therefore divided, as it were, into two portions—one, which is the flexed fundus, remaining in the pelvis; the other, containing the greater part of the fetus, rising above it. Under these circumstances a tumor in the vagina would exist in combination with an abdominal tumor, and pregnancy might go on to term. Considerable difficulty may even arise in labor, but the malposition generally rectifies itself before it gives rise to any serious results.

Treatment.—The treatment of retroversion of the gravid uterus should be taken in hand as soon as possible, for every day's delay involves an increase in the size of the uterus, and therefore greater difficulty in reposition. Our object is to restore the natural direction of the uterus by lifting the fundus above the promontory of the sacrum. The first thing to be done is to relieve the patient by emptying the bladder, the retention of urine having probably originally called attention to the case. For this purpose it is essential to use a long elastic male catheter of small size, as the urethra is too elongated and compressed to admit of the passage of the ordinary silver instrument. Even then it may be extremely difficult to introduce the catheter, and sometimes it has been found to be quite impossible. Under such circumstances, provided reposition cannot be effected without it, the bladder may be punctured an inch or two above the pubes by means of the fine needle of an aspirator, and the urine drawn off. Dieulafoy's work on aspiration proves conclusively that this may be done without risk, and the operation has been successfully performed by Schatz and others. It very rarely happens, however, and in long-neglected cases only, that the withdrawal of the urine is found to be impossible.

Mode of Effecting Reduction.—The bladder being emptied, and the bowels being also opened, if possible, by copious enemata, we proceed to attempt reduction. For this purpose various procedures are adopted. If the case is not of very long standing, I am inclined to think that the gentlest and safest plan is the continuous pressure of a caoutchouc bag, filled with water, placed in the vagina. The good effect of steady and long-continued pressure of this kind was proved by Tyler Smith, who effected in this way the reduction of an inverted uterus of long standing, and it is not difficult to understand that it may succeed when a more sudden and violent effort fails. I have tried this plan successfully in two cases, a pyriform india-rubber bag being inserted into the vagina and distended as far as the patient could bear by means of a syringe. The water must be let out occasionally to allow the patient to empty

the bladder, and the bag immediately refilled. In both my cases reposition occurred within twenty-four hours. Barnes has failed with this method, but it succeeded so well in my cases, and is so obviously less likely to prove hurtful than forcible reposition with the hand, that I am inclined to consider it the preferable procedure, and one that should be tried first. Failing with the fluid pressure, we should endeavor to replace the uterus in the following way: The patient should be placed at the edge of the bed, in the ordinary obstetric position, and thoroughly anæsthetized. This is of importance, as it relaxes all the parts and admits of much freer manipulation than is otherwise possible. One or more fingers of the left hand are then inserted into the rectum—if the patient be deeply chloroformed, it is quite possible, with due care, even to pass the whole hand—and an attempt is then made to lift or push the fundus above the promontory of the sacrum. At the same time reposition is aided by drawing down the cervix with the fingers of the right hand per vaginam. It has been insisted that the pressure should be made in the direction of one or other sacro-iliac synchondrosis rather than directly upward, so that the uterus may not be jammed against the projection of the promontory of the sacrum. Failing reposition through the rectum, an attempt may be made per vaginam, and for this some have advised the upward pressure of the closed fist passed into the canal. Others recommend the hand-and-knee position as facilitating reposition, but this prevents the administration of chloroform, which is of more assistance than any change of position can possibly be. Various complex instruments have been invented to facilitate the operation, but they are all more or less dangerous, and are unlikely to succeed when manual pressure has failed.

As soon as the reduction is accomplished, subsequent descent of the uterus should be prevented by a large-sized Hodge's pessary, and the patient should be kept at rest for some days, the state of the bladder and bowels being particularly attended to. When reposition has been fairly effected a relapse is unlikely to occur.

Treatment when Reduction is found Impossible.—In cases in which reduction is found to be impossible our only resource is the artificial induction of abortion. Under such circumstances this is imperatively called for. It is best effected by puncturing the membranes, the discharge of the liquor amnii of itself lessening the size of the uterus, and thus diminishing the pressure to which the neighboring parts are subjected. After this reposition may be possible, or we may wait until the fœtus is spontaneously expelled. It is not always easy to reach the os uteri, although we can generally do so with a curved uterine sound. If we cannot puncture the membranes, the liquor amnii may be drawn off through the uterine walls by means of the aspirator, inserted through either the rectum or vagina. The injury to the uterine walls thus inflicted is not likely to be hurtful, and the risk is certainly far less than leaving the case alone. Naturally, so extreme a measure would not be adopted until all the simpler means indicated have been tried and failed.

Diseases Coexisting with Pregnancy.—The pregnant woman is of course liable to contract the same diseases as in the non-pregnant state,

and pregnancy may occur in women already the subject of some constitutional disease. There is no doubt yet much to be learned as to the influence of coexisting disease on pregnancy. It is certain that some diseases are but little modified by pregnancy, and that others are so to a considerable extent, and that the influence of the disease on the foetus varies much. The subject is too extensive to be entered into at any length, but a few words may be said as to some of the more important affections that are likely to be met with.

Eruptive Fevers: Small-pox.—The eruptive fevers have often very serious consequences, proportionate to the intensity of the attack. Of these variola has the most disastrous results, which are related in the writings of the older authors, but which are, fortunately, rarely seen in these days of vaccination. The severe and confluent forms of the disease are almost certainly fatal to both the mother and child. In the discrete form and in modified small-pox after vaccination the patient generally has the disease favorably, and although abortion frequently results, it does not necessarily do so.

Scarlet Fever.—If scarlet fever of an intense character attacks a pregnant woman, abortion is likely to occur and the risks to the mother are very great. The milder cases run their course without the production of any untoward symptoms. Should abortion occur, the well-known dangerous effect of this zymotic disease after delivery will gravely influence the prognosis. Cazeaux was of opinion that pregnant women are not apt to contract the disease, while Montgomery thought that the poison when absorbed during pregnancy might remain latent until delivery, when its characteristic effects were produced.

Measles.—Measles, unless very severe, often runs its course without seriously affecting the mother or child. I have myself seen several examples of this. De Tourcoing, however, states that out of 15 cases the mother aborted in 7, these being all very severe attacks. Some cases are recorded in which the child was born with the rubeolous eruption upon it.

Continued Fevers.—The pregnant woman may be attacked with any of the continued fevers, and, if they are at all severe, they are apt to produce abortion. Out of 22 cases of typhoid, 16 aborted, and the remaining 6, who had slight attacks, went on to term; out of 63 cases of relapsing fever, abortion or premature labor occurred in 23. According to Schweden, the main cause of danger to the foetus in continued fevers is the hyperpyrexia, especially when the maternal temperature reaches 104° or upward. The fevers do not appear to be aggravated as regards the mother, and the same observation has been made by Cazeaux with regard to this class of disease occurring after delivery.

Pneumonia.—Pneumonia seems to be specially dangerous, for of 15 cases collected by Grisolle,¹ 11 died—a mortality immensely greater than that of the disease in general. The larger proportion also aborted, the children being generally dead, and the fatal result is probably due, as in the severe continued fevers, to hyperpyrexia. The cause of the maternal mortality does not seem quite apparent, since the same danger does not appear to exist in severe bronchitis or other inflammatory affections.

¹ *Arch. gén. de Méd.*, vol. xii. p. 291.

Phthisis.—Contrary to the usually-received opinion, it appears certain that pregnancy has no retarding influence on coexisting phthisis, nor does the disease necessarily advance with greater rapidity after delivery. Out of 27 cases of phthisis collected by Grisolle,¹ 24 showed the first symptoms of the disease after pregnancy had commenced. Phthisical women are not apt to conceive—a fact which may probably be explained by the frequent coexistence in such cases of uterine disease, especially severe leucorrhœa. The entire duration of the phthisis seems to be shortened, as it averaged only nine and a half months in the 27 cases collected—a fact which proves, at least, that pregnancy has no material influence in arresting its progress. If we consider the tax on the vital powers which pregnancy naturally involves, we must admit that this view is more physiologically probable than the one generally received, and apparently adopted without any due grounds.

Heart Disease.—The evil effects of pregnancy and parturition on chronic heart disease have of late received much attention from Spiegelberg, Fritsch, Peter, and other writers. The subject has been ably discussed² in a series of elaborate papers by Dr. Angus McDonald, which are well worthy of study. Out of 28 cases collected by him, 17, or 60 per cent., proved fatal. This, no doubt, is not altogether a reliable estimate of the probable risk of the complication, but, at any rate, it shows the serious anxiety which the occurrence of pregnancy in a patient suffering from chronic heart disease must cause. Dr. McDonald refers the evils resulting from pregnancy in connection with cardiac lesions to two causes: first, destruction of that equilibrium of the circulation which has been established by compensatory arrangements; secondly, the occurrence of fresh inflammatory lesions upon the valves of the heart already diseased.

The dangerous symptoms do not usually appear until after the first half of the pregnancy has passed, and the pregnancy seldom advances to term. The pathological phenomena generally met with in fatal cases are pulmonary congestion, especially of the bronchial mucous membrane, and pulmonary œdema, with occasional pneumonia and pleurisy. Mitral stenosis seems to be the form of cardiac lesion most likely to prove serious, and next to this aortic incompetency. The obvious deduction from these facts is, that heart disease, especially when associated with serious symptoms, such as dyspnoea, palpitation, and the like, should be considered a strong contraindication of marriage. When pregnancy has actually occurred, all that can be done is to enjoin the careful regulation of the life of the patient, so as to avoid exposure to cold and all forms of severe exertion.

Syphilis.—The important influence of syphilis on the ovum is fully considered elsewhere. As regards the mother, its effects are not different from those at other times. It need only, therefore, be said that whenever indications of syphilis in a pregnant woman exist the appropriate treatment should be at once instituted and carried on during her gestation, not only with the view of checking the progress of the disease, but in the hope of preventing or lessening the risk of abortion or of the birth of an infected infant. So far from pregnancy contraindicating

¹ *Arch. gén. de Méd.*, vol. xxii.

² *Obst. Journ.*, 1877.

mercurial treatment, there rather is a reason for insisting on it more strongly. As to the precise medication, it is advisable to choose a form that can be exhibited continuously for a length of time without producing serious constitutional results. Small doses of the bichloride of mercury, such as one-sixteenth of a grain, thrice daily, or of the iodide of mercury, or of the hydrargyrum cum creta, in combination with reduced iron, answer this purpose well; or in the early stages of pregnancy the mercurial vapor bath or cutaneous inunction may be employed.

Dr. Weber of St. Petersburg¹ has made some observations showing the superiority of the latter methods, which he found did not interfere with the course of pregnancy; the contrary was the case when the mercury was administered by the mouth, probably, as he supposes, from disturbance of the digestive system. It must be borne in mind that in married women it may sometimes be expedient to prescribe an anti-syphilitic course without their knowledge of its nature, so that inunction is not always feasible.

Epilepsy.—The influence of pregnancy on epilepsy does not appear to be as uniform as might perhaps be expected. In some cases the number and intensity of the fits have been lessened; in others the disease becomes aggravated. Some cases are even recorded in which epilepsy appeared for the first time during gestation. On account of the resemblance between epilepsy and eclampsia there is a natural apprehension that a pregnant epileptic may suffer from convulsions during delivery. Fortunately, this is by no means necessarily the case, and labor often goes on satisfactorily without any attack.

Jaundice, the result of acute yellow atrophy of the liver, is occasionally observed, and is said to have been sometimes epidemic. Independently of the grave risks to the mother, it is most likely to produce abortion or the death of the fœtus. According to Davidson,² it originates in catarrhal icterus, the excretion of the bile-products being impeded in consequence of pregnancy, and their retention giving rise to the fatal blood-poisoning which accompanies the severer forms of the disease. Slight and transient attacks of jaundice may occur without being accompanied by any bad consequences. Their production is probably favored by the mechanical pressure of the gravid uterus on the intestines and the bile-ducts.

Carcinoma.—The occurrence of pregnancy in a woman suffering from malignant disease of the uterus is by no means so rare as might be supposed, and must naturally give rise to much anxiety as to the result. The obstetrical treatment of these cases will be discussed elsewhere. Should we be aware of the existence of the disease during gestation, the question will arise whether we should not attempt to lessen the risks of delivery by bringing on abortion or premature labor. The question is one which is by no means easy to settle. We have to deal with a disease which is certain to prove fatal to the mother before long, and the progress of which is probably accelerated after labor, while the manipulations necessary to induce delivery may very unfavorably influence the diseased structures. Again, by such a measure we necessarily sacrifice the child,

¹ *Allgem. Med. Cent. Zeit.*, Feb., 1875.

² *Monat. f. Geburt*, 1867.

while we are by no means certain that we materially lessen the danger to the mother. The question cannot be settled except on a consideration of each particular case. If we see the patient early in pregnancy, by inducing abortion we may save her the dangers of labor at term—possibly of the Cæsarean section—if the obstruction be great. Under such circumstances the operation would be justifiable. If the pregnancy has advanced beyond the sixth or seventh month, unless the amount of malignant deposit be very small indeed, it is probable that the risks of labor would be as great to the mother as at term, and it would then be advisable to give her the advantage of the few months' delay.

Ovarian Tumor.—Cases are occasionally met with in which pregnancy occurs in women who are suffering from ovarian tumor, and their proper management has given rise to considerable discussion. There can be no doubt that such cases are attended with very dangerous and often fatal consequences, for the abdomen cannot well accommodate the gravid uterus and the ovarian tumor, both increasing simultaneously. The result is that the tumor is subject to much contusion and pressure, which have sometimes led to the rupture of the cyst and the escape of its contents into the peritoneal cavity; at others to a low form of inflammation, attended with much exhaustion, the death of the patient supervening either before or shortly after delivery. The danger during delivery from the same cause in the cases which go on to term is also very great. Of 13 cases of delivery by the natural powers, which I collected in a paper on "Labor Complicated with Ovarian Tumor,"¹ far more than one-half proved fatal. [In one instance in this city a lady well known to the editor gave birth to three of her four children during the existence of an ovarian tumor. The children all lived to grow up, and their mother died of her disease at the age of 75, after being repeatedly tapped during fifty years. The ovarian tumor was discovered by Dr. Benjamin Rush soon after her first child was born in 1809, and she was first tapped by Dr. Physick in 1811. In 1812, 1815, and 1818 she gave birth to the children mentioned, the third being delicate, sickly, and weighing six pounds. This last died of phthisis when 45; one still lives.² According to the teaching of Mr. Lawson Tait, this may have been a parovarian cyst, and not an ovarian cystoma.—Ed.] Another source of danger is twisting of the pedicle, and consequent strangulation of the cyst, of which several instances are recorded. It is obvious, then, that the risks are so manifold that in every case it is advisable to consider whether they can be lessened by surgical treatment.

Methods of Treatment.—The means at our disposal are either to induce labor prematurely, to treat the tumor by tapping, or to perform ovariectomy. The question has been particularly discussed by Spencer Wells in his works on *Ovariectomy*, and by Barnes in his *Obstetric Operations*. The former holds that the proper course to pursue is to tap the tumor when there is any chance of its being materially lessened in size by that procedure, but that when it is multilocular or when its contents are solid ovariectomy should be performed at as early a period of pregnancy as possible. Barnes, on the other hand, maintains that the safer course is

¹ *Obst. Trans.*, vol. ix.

[² *Trans. Phila. Obstet. Soc.*, vol. i., 1873, p. 64, reported by Ed.]

to imitate the means by which nature often meets this complication, and bring on premature labor without interfering with the tumor. He thinks that ovariectomy is out of the question, and that tapping may be insufficient and leave enough of the tumor to interfere seriously with labor. So far as recorded cases go, they unquestionably seem to show that tapping is not more dangerous than at other times, and that ovariectomy may be practised during pregnancy with a fair amount of success. Wells records 10 cases which were surgically interfered with. In 1 tapping was performed, and in 9 ovariectomy; and of these 8 recovered, the pregnancy going on to term in 5. On the other hand, 5 cases were left alone, and either went to term or spontaneous premature labor supervened; and of these 3 died. The cases are not sufficiently numerous to settle the question, but they certainly favor the view taken by Wells rather than that by Barnes. It is to be observed that unless we give up all hope of saving the child and induce abortion, the risk of induced premature labor, when the pregnancy is sufficiently advanced to hope for a viable child, would almost be as great as that of labor at term; for the question of interference will only have to be considered with regard to large tumors, which would be nearly as much affected by the pressure of a gravid uterus at seven or eight months as by one at term. Small tumors generally escape attention, and are more apt to be impacted before the presenting part in delivery. The success of ovariectomy during pregnancy has certainly been great, and we have to bear in mind that the woman must necessarily be subjected to the risk of the operation sooner or later, so that we cannot judge of the case as one in which abortion terminates the risk. Even if the operation should put an end to the pregnancy—and there is at least a fair chance that it will not do so—there is no certainty that that would increase the risk of the operation to the mother, while as regards the child we should only have the same result as if we intentionally produced abortion. On the whole, then, it seems that the best chance to the mother, and certainly the best to the child, is to resort to the apparently heroic practice recommended by Wells. The determination must, however, be to some extent influenced by the skill and experience of the operator. If the medical attendant has not gained that experience which is so essential for a successful ovariectomist, the interests of the mother would be best consulted by the induction of abortion at as early a period as possible. One or other procedure is essential; for, in spite of a few cases in which several successive pregnancies have occurred in women who have had ovarian tumors, the risks are such as not to justify an expectant practice. Should rupture of the cyst occur, there can be no doubt that ovariectomy should at once be resorted to, with the view of removing the lacerated cyst and its extravasated contents.

Fibroid Tumors.—Pregnancy may occur in a uterus in which there are one or more fibroid tumors. If these are situated low down and in a position likely to obstruct the passage of the fetus, they may very seriously complicate delivery. When they are situated in the fundus or body of the uterus they may give rise to risk from hemorrhage or from inflammation of their own structure. Inasmuch as they are structurally similar to the uterine walls, they partake of the growth of the uterus

during pregnancy, and frequently increase remarkably in size. Cazeaux says: "I have known them in several instances to acquire a size in three or four months which they would not have done in several years in the non-pregnant condition." Conversely, they share in the involution of the uterus after delivery, and often lessen greatly in size, or even entirely disappear. Of this fact I have elsewhere recorded several curious examples;¹ and many other instances of the complete disappearance of even large tumors have been described by authors whose accuracy of observation cannot be questioned.

Treatment.—The treatment will vary with the position of the tumor. If it is such as to be certain to obstruct the passage of the child, abortion should be induced as soon as possible. If the tumor is well out of the way, this is not so urgently called for. The principal danger, then, is that the tumor will impede the post-partum contraction of the uterus and favor hemorrhage. Even if this should happen, the flooding could be controlled by the usual means, especially by the injection of the perchloride of iron. I have seen several cases in which delivery has taken place under such circumstances without any untoward accident. The danger from inflammation and subsequent extrusion of the fibroid masses would probably be as great after abortion or premature labor as after delivery at term. It seems, therefore, to be the proper rule to interfere when the tumors are likely to impede delivery, and in other cases to allow the pregnancy to go on, and be prepared to cope with any complications as they arise. The risks of pregnancy should be avoided in every case in which uterine fibroids of any size exist, the patients being advised to lead a celibate life.

[Fibroid tumors may so obstruct the pelvis as to make delivery *per vias naturales* impossible. If the obstacle cannot be forced up out of the pelvis with the hand, delivery by the abdomen will be required if the child is to be saved. This form of obstruction makes the Cæsarean operation more than usually hazardous, and likewise its modification by Porro. Ten Cæsarean operations have been performed in consequence of obstruction by uterine fibroids in the United States, with the saving of four women and five children. Two fatal Porro operations have also been performed.—Ed.]

CHAPTER IX.

PATHOLOGY OF THE DECIDUA AND OVUM.

Pathology of the Decidua.—Comparatively little is, unfortunately, known of the pathological changes which occur in the mucous membrane of the uterus during pregnancy. It is probable that they are of much more consequence than is generally believed to be the case, and it is certain that they are a frequent cause of abortion.

¹ *Obst. Trans.*, vols. v., xiii., and xix.

Endometritis.—One of the most generally observed probably depends on endometritis antecedent to conception. When the impregnated ovule reached the uterus it engrafted itself on the inflamed mucous membrane, which was in an unfit condition for its reception and growth. A not uncommon result, under such circumstances, is the laceration of some of the decidual vessels, extravasation of the blood between the decidua and the uterine walls, and consequent abortion at an early stage of pregnancy. As this morbid state of the uterine mucous membrane is likely to continue after abortion is completed, the same history repeats itself on each impregnation, and thus we may have constant early miscarriages produced. It does not necessarily follow, however, that the pregnancy is immediately terminated when this state of things is present. Some-

FIG. 87.



Hypertrophied Decidua laid open, with the Ovum attached to its Fundal Portion.
(After Duncan.)

times a condition of hyperplasia of the decidua is produced, the membrane becomes much thickened and hypertrophied in consequence of proliferation of its interstitial connective tissue, and the decidual cells are greatly increased in size (Fig. 87). In other instances the internal surface of the decidua becomes studded with rough polypoid growths,¹ depending on proliferation of its interstitial tissue. Duncan has found that the hypertrophied decidua is always in a state of fatty degeneration, more advanced in some places than in others.² The result of these alter-

¹ *Virchow's Archiv. für Path.*, 1861, 1st ed.

² *Researches in Obstetrics*, p. 293.

ations is frequently to produce dwindling or death of the ovum, which, however, retains its connection with the decidua, until, after a lapse of time, the decidua is expelled in the form of a thick triangular fleshy substance, with the atrophied ovum attached to some part of its inner surface. In other cases, in which the hyperplasia has advanced to a less extent, the nutrition of the fetus is not interfered with, and pregnancy may continue to term, the changes in the decidua being recognizable after delivery. Other diseases besides endometritis may give rise to similar alterations in the decidua, one of these being, as Virchow maintains, syphilis. The converse condition, an imperfect development of the decidua, especially of the decidua reflexa, has also been noted as a cause of abortion. The ovum will then hang loosely in the uterine cavity, without the support which the growth of the decidua reflexa around it ought to afford, and its premature expulsion readily follows (Fig. 88).

FIG 88.



Imperfectly developed Decidua Vera, with the Ovum. (After Duncan.)

Hydrorrhœa Gravidarum.—The peculiar condition known as *hydrorrhœa gravidarum* most probably depends on some obscure morbid state of the uterine mucous membrane. By it is meant a discharge of clear watery fluid at intervals during pregnancy. It may happen at any period of gestation, but is most commonly met with in the latter months. It may commence with a mere dribbling, or there may be a sudden and copious discharge of fluid. Afterward the watery fluid, which is generally of a pale yellowish color and transparent like the liquor amnii, may continue to escape at intervals for many weeks, and sometimes in very great abundance, so as to saturate the patient's clothes. Very frequently it is expelled in gushes, and at night when the patient is lying quietly in bed; its escape is then probably due to uterine contraction.

Many theories have been held as to its cause. By some it is attributed to the rupture of a cyst placed between the ovum and the uterine walls; Baudelocque referred it to a transudation of the liquor amnii through the membranes; while Burgess and Dubois believed it to depend on a

laceration of the membranes at a distance from the os uteri. Mattei more recently has attributed it to the existence of a sac between the chorion and the amnion. It may be that in some instances a single discharge of fluid may come from one of the two last-mentioned causes. But if it be continuous or repeated another source must be sought for. Hegar¹ maintains that it is the result of abundant secretion from the glands of the mucous membrane, which accumulates between the decidua and chorion and escapes through the os uteri. If this occur, the decidua is probably in a hypertrophied and otherwise morbid state. Hydrorrhœa is chiefly of interest from the error of diagnosis it is likely to give rise to, for on being summoned to a case in which watery discharge has occurred for the first time, we are naturally apt to suppose that the membranes have ruptured and that labor is imminent. Nor is there any very certain means of deciding if this be so. In hydrorrhœa we find that pains are absent, the os uteri unopened, and ballottement may be made out. Even if the membranes be ruptured, there will be no indication for interference unless labor has actually commenced; and the repetition of the discharge and the continuance of the pregnancy will soon clear up the diagnosis. Hydrorrhœa, although apt to alarm the patient, need not give rise to any anxiety. The pregnancy generally progresses favorably to the full period, although in exceptional cases premature labor may supervene. No treatment is necessary, nor is there any that could have the least effect in controlling the discharge.

Pathology of the Chorion.—The only important disease of the chorion with which we are acquainted is the well-known condition which is variously described as *uterine hydatids*, *cystic disease of the ovum*, *hydatidiform degeneration of the chorion*, or *vesicular mole*. The name of uterine hydatids was long given to it on the supposition that the grape-like vesicles which characterize the disease were true hydatids, similar to those which develop in the liver and other structures. This idea has long been exploded, and it is now known as a certainty that the disease originates in the villi of the chorion. The precise mode and the causes of its production are, however, not yet satisfactorily settled. The disease is characterized by the existence in the cavity of the uterus of a large number of translucent vesicles containing a clear limpid fluid, which has been found on analysis to bear close resemblance to the liquor amnii. These small bladder-like bodies, which vary in size from that of a millet-seed to an acorn, are often described as resembling a bunch of grapes or currants. On more minute examination they are found not to be each attached to independent pedicles, as is the case in a bunch of grapes, but some of them grow from other vesicles, while others have distinct pedicles attached to the chorion, the pedicles themselves sometimes being distended by fluid (Fig. 89). This peculiar arrangement of the vesicles is explained by their mode of growth.

Causes of Cystic Degeneration.—There has been considerable discussion as to the etiology of this disease. By some it is supposed always to follow death of the fetus, and, the whole developmental energy being expended on the chorion, which retains its attachment to the decidua, the result is its abnormal growth and cystic degeneration. This is the

¹ *Monat. f. Geburt*, 1863.

view maintained by Gierse and Graily Hewitt, and it is favored by the undoubted fact that in almost all cases the foetus has entirely disappeared, and by the occasional occurrence of cases of twin conceptions in which one chorion has degenerated, the other remaining healthy until term. On the other hand, it is maintained

that the starting-point is connected with the maternal organism. Virchow thinks it originates in a morbid state of the decidua, while others have attributed it to some blood-dyscrasia on the part of the mother, such as syphilis. There are many reasons for believing that causes of this nature may originate the affection. Thus it is often found to occur more than once in the same person, and alterations of a similar kind, although limited in extent, are not unfrequently found in connection with the placenta and membranes of living children. On this theory the death of the foetus is secondary, the consequence of impaired nutrition from the morbid state of the chorion. The probability is that both views may be right, the disease sometimes following the death of the embryo, and at others being the result of obscure maternal causes.



FIG. 89.
Hydatidiform Degeneration of the Chorion.

commenced to form. In that case the entire superficies of the chorion becomes affected. The disease, however, may not begin until after the greater part of the chorion villi have atrophied, and then it is limited to the placenta. The epithelium of the villi appears to be the part first affected, and the whole interior of the diseased villus becomes filled with cells. The connective tissue of the villus undergoes a remarkable proliferation, and collects in masses at individual spots, the remainder of the villus being unaffected. By the growth of these elements the villus becomes distended and many of the cells liquefy, the intercellular fluid thus produced widely separating the connective tissue, so as to form a network in the interior of the villus.¹ Thus are formed the peculiar grape-like bodies which characterize the disease. When once the degeneration has commenced, the diseased tissue has a remarkable power of increase, so that it sometimes forms a mass as large as a child's head and several pounds in weight.

The nutrition of the altered chorion is maintained by its connections with the decidua, which is also generally diseased and hypertrophied. Sometimes the adhesion of the mass to the uterine walls is very firm, and may interfere with its expulsion, while in a few rare cases it has been found that the villi have forced their way into the substance of the uterus, chiefly through the uterine sinuses, and thus caused atrophy and

¹ Braxton Hicks, *Guy's Hospital Reports*, vol. ii., 3d Series, p. 380.

thinning of its muscular structure. Cases of this kind are related by Volkmann, Waldeyer,¹ and Barnes, and it is obvious that the intimate adhesion thus effected must seriously add to the gravity of the prognosis.

Taking this view of the etiology of this disease, it is obvious that it is essentially connected with pregnancy, and that there is no valid ground for maintaining, as has sometimes been done, that it may occur independently of conception. It is just possible, however, that true entozoa may form in the substance of the uterus, which, being expelled per vaginam, might be taken for the results of cystic disease, and thus give rise to groundless suspicions as to the patient's chastity. Hewitt has related one case in which true hydatids, originally formed in the liver, had extended to the peritoneum, and were about to burst through the vagina at the time of death. This occurred in an unmarried woman. One or two other examples of true hydatids forming in the substance of the uterus are also recorded. A very interesting case is also related by Hewitt² in which undoubted accephalocysts were expelled from the uterus of a patient who ultimately recovered. A careful examination of the cyst and its contents would show their true nature, as the echinococci heads, with their characteristic hooklets, would be discoverable by the microscope.

It is also possible that unfounded suspicions might arise from the fact of a patient expelling a mass of hydatids long after impregnation. In the case of a widow or woman living apart from her husband serious mistakes might thus be made. This has been specially pointed out by McClintock,³ who says: "Hydatids may be retained in utero for many months or years, or a portion only may be expelled, and the residue may throw out a fresh crop of vesicles, to be discharged on a future occasion."

Symptoms and Progress of the Disease.—The symptoms of cystic disease of the ovum are by no means well marked. At first there is nothing to point to the existence of any morbid condition, but as pregnancy advances its ordinary course is interfered with. There is more general disturbance of the health than there ought to be, and the reflex irritations, such as vomiting, may be unusually developed. The first physical sign remarked is rapid increase of the uterine tumor, which soon does not correspond in size to the supposed period of pregnancy. Thus, at the third month the uterus may be found to reach up to or beyond the umbilicus. About this time there generally are more or less profuse watery and sanguineous discharges, which have been described as resembling currant-juice. They no doubt depend on the breaking down and expulsion of the cysts, caused by painless uterine contractions. They are sometimes excessive in amount, recur with great frequency, and often reduce the patient extremely. Portions of cysts may now generally be found mingled with the discharge, and sometimes large masses of them are expelled from time to time. Indeed, the discovery of portions of cysts is the only certain diagnostic sign. Vaginal examination, before the os has dilated, will give no information except the absence of ballottement. An unusual hardness or density of the uterus—described by Leishman, who attributes much importance to it, as "a peculiar doughy,

¹ *Virchow's Archiv*, vol. xliv. p. 86.

² McClintock's *Diseases of Women*, p. 398.

³ *Obs. Trans.*, vol. xii.

boggy feeling"—has been pointed out by several writers. The contour of the uterine tumor, moreover, is often irregular. In addition, we of course fail to discover the usual auscultatory signs of pregnancy. All this may aid in diagnosis, but nothing except the presence of cysts in the watery bloody discharge will enable us to pronounce with certainty as to the nature of the disease.

Treatment.—As soon as the diagnosis is established the indications for treatment are obvious. The sooner the uterus is cleared of its contents the better. Ergot may be given with advantage to favor uterine contraction and the expulsion of the diseased ovum. Should this fail, more especially if the hemorrhage be great, the fingers or the whole hand must be introduced into the uterus, and as much as possible of the mass removed. As the os is likely to be closed, its preliminary dilatation by sponge or laminaria tents, or by a Barnes's bag if it be already opened to some extent, will in most cases be required. If chloroform be then administered, the remaining steps of the operation will be easy. On account of the occasional firm adhesion of the cystic mass to the uterus, too energetic attempts at complete separation should be avoided. Any severe hemorrhage after the operation can be controlled by swabbing out the uterine cavity with the perchloride of iron solution.

Myxoma Fibrosum.—Under the name of *Myxoma fibrosum* a more rare degeneration of the chorion has been described by Virchow and Hildebrandt,¹ characterized not by vesicular but by fibroid degeneration of the connective tissue of the chorion. This is, however, too little understood to require further observation.

Pathology of the Placenta.—The pathology of the placenta has of late years attracted much attention, and it has an important practical bearing, in consequence of its effect on the child.


Placentæ vary considerably in shape. They may be crescentic, or spread over a considerable surface, in consequence of the chorion villi entering into communication with a larger portion of the decidua than usual (*placenta membranacea*). Such forms, however, are merely of scientific interest. The only anomaly of shape of any practical importance is the formation of what have been called *placentæ succenturiæ*. These consist of one or more separate masses of placental tissue, produced by the development of isolated patches of chorion villi. Hohl believes that they always form exactly at the junction of the anterior and posterior walls of the uterus, which in early pregnancy is a mere line. As the uterus expands, the portions of placenta on each side of this become separated from each other. They are only of consequence from the possibility of their remaining unnoticed in the uterus after delivery and giving rise to secondary post-partum hemorrhage. The rare form of double placenta with a single cord, figured in the accompanying woodcut (Fig. 90), was probably formed in this way, and the supplementary portion in such a case might readily escape notice.

The placenta may also vary in dimensions. Sometimes it is of excessive size, generally when the child is unusually big, but not unfrequently in connection with hydramnios, the child being dead and shrivelled. In other cases it is remarkably small, or at least appears to be

¹ *Monat. f. Geburt*, May, 1865.

so. If the child be healthy, this is probably of no pathological importance, as its smallness may be more apparent than real, depending on its vessels not being distended with blood. When true atrophy of the placenta exists, the vitality of the foetus may be seriously interfered with. This condition may depend either on a diseased state of the chorion villi

FIG. 90.



Double Placenta, with Single Cord.

or of the decidua in which they are implanted.¹ The latter is the more common of the two; and it generally consists in hyperplasia of the connective tissue of the decidua, which presses on the villi and vessels and gives rise to general or local atrophy. This change is similar in its nature to that observed in cirrhosis of the liver and certain forms of Bright's disease. It has generally been ascribed to inflammatory changes, and under the name of *placentitis* has been described by many authors, and has been considered to be a common disease. To it are attributed many of the morbid alterations which are commonly observed in placentæ, such as hepatizations, circumscribed purulent deposits, and adhesions to the uterine walls. Many modern pathologists have doubted whether these changes are in any proper sense inflammatory. Whittaker observes on this point: "The disposition to reject placentitis altogether increases in modern times. Indeed, it is impossible to conceive of inflammation on the modern theory (Cohnheim) of that process, since there are no capillaries, in the maternal portion at least, through whose walls a 'migration' might occur, and there are no nerves to regulate the contractility of the vessel walls in the entire structure." Robin thus

¹ Whittaker, *Amer. Journ. of Obst.*, vol. iii. p. 229.

explains the various pathological changes above alluded to: "What has been taken for inflammation of the placenta is nothing else than a condition of transformation of blood-clots at various periods. What has been regarded as pus is only fibrin in the course of disorganization, and in those cases where true pus has been found the pus did not come from the placenta, but from an inflammation of the tissue of the uterine vessels and an accidental disposition in the tissue of the placenta." The extravasations of blood here alluded to are of very common occurrence, and they are found in all parts of the organ—in its substance, on its decidual surface, or immediately below the amnion, where they serve as points of origin for the cysts that are there often observed. The fibrin thus deposited undergoes retrograde metamorphosis, as in other parts of the body; it becomes decolorized, it undergoes fatty degeneration, or it becomes changed into calcareous masses; and in this way, it is supposed, may be explained the various pathological changes which are so commonly observed. The amount of retrograde metamorphosis, and the precise appearance presented, will of course depend on the time that has elapsed since the blood-extravasations took place.

Fatty Degeneration.—Fatty degeneration of the placenta, and its

FIG. 91.



Fatty Degeneration of the Placenta.

influence on the nutrition of the fœtus, have been specially studied in this country by Barnes and Druitt. Yellowish masses of varying sizes are very commonly met with in placentæ, and these are found to consist, in great part, of molecular fat, mixed with a fine network of fibrous tissue. The true fatty degeneration, however, specially affects the chorion

villi (Fig. 91). On microscopic examination they are found to be altered and misshaped in their contour and to be loaded with fine granular fat-globules. Similar changes are observed in the cells of the decidua. The influence on the fœtus will, of course, depend on the extent to which the functions of the villi are interfered with. The probable cause of this degeneration is no doubt some obscure alteration in the nutrition of the tissue, depending on the state of the mother's health. Barnes believes that syphilis has much influence in its production. Drutt has pointed out that some amount of fatty degeneration is always present in a mature placenta, and is probably connected with the physiological separation of the organ; and Goodell has more recently suggested that an unusual amount of this change may be merely an anticipation of the natural termination of the life of the placenta.¹

Other Morbid States.—Other morbid states of the placenta, of greater rarity, are occasionally met with, as an œdematous infiltration of its tissue, always occurring, according to Lange, in cases of hydramnios, pigmentary and calcareous deposits, and tumors of various kinds; but these require only a passing mention.

Pathology of the Umbilical Cord.—The umbilical cord may be of excessive length, varying from 18 to 20 inches, which is its average measurement, up to 50 or 60 inches, and a case is recorded in which it even reached the extraordinary length of 9 feet. If unusually long, it may be twisted round the limbs or neck of the child, and the latter position may, in exceptional instances, prove injurious during labor.

Some authors refer cases of spontaneous amputation of fœtal limbs in utero to constrictions by the umbilical cord, but this accident is more probably produced by filamentous adnexa of the amnion. Knots in the cord are not uncommon, and they result from the fœtus, in its movements, passing through a loop of the cord (Fig. 92). If there is an average amount of Wharton's jelly in the cord, the vessels are protected from pressure, and no bad effects follow. Géry, in a recent paper on this subject,² attempts to show that such knots are more important than is generally believed, and relates two cases in which he believes them to have caused the death of the fœtus.

Extreme torsion of the cord, an exaggeration of the spiral twists generally observed, may prove injurious, and even fatal, to the child, by obstructing the circulation in the vessels. Spæth mentions three cases in which this caused the death of the fœtus, the cord being twisted until it was reduced to the thickness of a thread.

Anomalies in the distribution of the vessels of the cord are of common occurrence.




FIG. 92.

Knots of the Umbilical Cord.

The cord may be attached

¹ *American Journal of Obstetrics*, vol. ii. p. 535.

² *L'Union médicale*, Oct., 1876.

to the edge, instead of to the centre, of the placenta (*battledore placenta*). It may break up into its component parts before reaching the placenta, the vessels running through the membranes; and if, in such a case, traction on the cord be made, the separate vessels may lacerate and the cord become detached. There may be two veins and one artery, or only one vein and one artery, or there may be two separate cords to one placenta. These and other anomalies that might be mentioned are of little practical importance.

Pathology of the Amnion.—The principal pathological condition of the amnion with which we are acquainted is that which is associated with excessive secretion of the liquor amnii, and is generally known under the name of *hydramnios*, which term Kidd¹ limits to cases in which more than two quarts of amniotic fluid exist. Its precise cause is still a matter of doubt. By some it is referred to inflammation of the amnion itself; at other times it is apparently connected with some morbid state of the decidua, which may be found diseased and hypertrophied. The fœtus is very often dead and shrivelled and the placenta enlarged and oedematous. It does not necessarily follow, however, that hydramnios causes the death of the child. Out of 33 cases, McClintock found that 9 children were born dead,² and of the 24 born alive, 10 died within a few hours; the remainder survived. There does not appear to be any marked relation between the state of the mother's health and the occurrence of this disease; and it is certainly not necessarily present when the mother is suffering from dropsical effusions in other parts of the body. The theory that the disease is of purely local origin is favored by the fact that when hydramnios occurs in twin pregnancy one ovum only is generally affected. Its effects, as regards the mother, are chiefly mechanical. It rarely begins to show itself before the fifth or sixth month of pregnancy, but when once it has commenced it rapidly produces a feeling of discomfort and enlargement altogether beyond that which should exist at the period of pregnancy which has been reached. In advanced stages the distress produced is often very great, the enlarged uterus pressing upon the diaphragm and producing much embarrassment of respiration. Premature expulsion of the fœtus very often supervenes. Four out of McClintock's patients died after labor, showing that the maternal mortality is high—a result which he refers to the debilitated state of the women who were the subjects of the disease.

Its Diagnosis.—The diagnosis is not, as a rule, difficult. It has to be distinguished from ascitic distension of the abdomen, from enlargement of the uterus from twin pregnancy, and from ovarian tumor or pregnancy complicated with ovarian tumor. The first will be recognized by the superficial position of the fluid; by the difficulty of feeling the contour of the uterus, which is obscured by the surrounding fluid, and the results of percussion, which show that the fluid is free in the peritoneal cavity; and by the coexistence of dropsical effusions in other parts of the body. The second may be difficult, and even impossible, to diagnose from it: generally, however, in hydramnios the uterine

¹ "On the Diagnosis of Dropsy of the Amnion," *Proceedings of the Obstetrical Society of Dublin*, May 11, 1878.

² *Diseases of Women*, p. 383.

tumor is more distinctly tense or fluctuating, the foetal limbs cannot be felt on palpation, and the lower segment of the uterus, as felt per vaginam, is unusually distended, the presenting part not being appreciable. Ovarian tumors alone or complicating pregnancy may also be difficult to distinguish from dropsy of the amnion. The general history of the case, and the presence or absence of signs of pregnancy, may enable us to arrive at a diagnosis; and Kidd points out that the position of the uterus, whether gravid or not, is usually low down in the pelvis in ovarian dropsy, while in dropsy of the amnion it is drawn high up and reached with difficulty on vaginal examination.

Its Effect on Labor.—During labor an excessive amount of liquor amnii is often a cause of deficient uterine action and delay, the pains being feeble and ineffective. This, of course, tells chiefly in the first stage, which is often much prolonged, unless the membranes are punctured early and the superabundant fluid allowed to escape.

Treatment.—No treatment is known to have any effect on the disease. If the discomfort and distension are very great, it may be absolutely necessary to puncture the membranes and allow the water to escape. This inevitably brings on labor. If the pregnancy be not sufficiently advanced to give hope for the birth of a living child, we would not of course resort to this expedient unless the mother's health was seriously imperilled. It is possible that in such cases the patient might be relieved by inserting the minute needle of an aspirator through the os and removing a certain quantity of the liquor amnii by aspiration, without inducing the labor. I have never had an opportunity of trying this expedient; but it seems a possibility.

Deficiency of Liquor Amnii.—A defective amount of liquor amnii is said to favor certain malformations by allowing the uterus to compress the foetus unduly. It certainly occasionally gives rise to adhesion between the foetus and the membranes, and to the formation of amniotic bands which are capable of producing certain fetal deformities (pp. 235 and 240).

Appearance of the Liquor Amnii.—The liquor amnii itself varies much in appearance. It is sometimes thick and treacly, instead of limpid, and it may be offensive in odor. The cause of these variations is not well understood.

Pathology of the Foetus.—There is abundant evidence that the foetus in utero is subject to many diseases, some of which cause its death, and others leave distinct traces of their existence, although not proving fatal. The subject is of great importance, and is well worthy of study. There is still much to be done in this direction which may lead to important practical results. I can, however, do little more than enumerate some of the principal affections which have been observed.

Blood Diseases transmitted through the Mother: Small-pox.—It is a well-established fact that the various eruptive fevers from which the mother may suffer may be communicated to the foetus in utero. When the mother is attacked with confluent small-pox she almost always aborts, but not necessarily so when it is discrete or modified. In such cases it has often happened that the foetus has been born with evident marks of small-pox. Cases are on record which prove that the foetus

was attacked subsequently to the mother. Thus a mother attacked with small-pox has miscarried, and has given birth to a living child showing no trace of the disease, which, however, showed itself in two or three days—proving that it had been contracted, and had run through its usual period of incubation, when the fœtus was still in utero. It does not follow, however, that the fœtus is affected, as Serres has collected 22 cases in which women suffering from small-pox gave birth to children who had not contracted the disease. It has been supposed that in such cases the child is protected from small-pox, though it has shown no symptom of having had the disease. Tarnier, however, cites two instances in which such children had small-pox two years after birth. Madge and Simpson record cases in which vaccination performed on the mother during pregnancy protected the fœtus, on whom all subsequent attempts at vaccination failed. There is evidence also to prove that the disease may be transmitted to the fœtus through a mother who is herself unsusceptible of contagion, the child having been covered with small-pox eruption, the mother being quite free from it. It is probable that the same facts which have been observed with regard to small-pox hold true with reference to other zymotic diseases, such as scarlet fever and measles, although there is not sufficient evidence to justify a positive assertion to that effect.

Malaria and Lead-Poisoning.—Amongst other maternal diseases, malaria and lead-poisoning are known to affect the fœtus in utero. Dr. Stokes relates cases in which the mother suffered from tertian ague, the child having also attacks, as evidenced by its convulsive movements, appreciable by the mother, which took place at the regular intervals, but at a different time from the mother's paroxysms. In other cases the febrile paroxysm comes on at the same time in the fœtus as in the mother; and the fact has been verified by the observation that the paroxysms continued to recur simultaneously after delivery. The fœtus has also been born with distinct malarious enlargement of the spleen. From the frequency with which largely hypertrophied spleens are seen in mere infants in malarious districts, I imagine that the intra-uterine disease must be common. I have frequently observed this fact in India, although, of course, without any possibility of ascertaining if the mothers had suffered from intermittent fever during pregnancy. Lead-poisoning is also known to have a most prejudicial effect on the fœtus, and frequently to lead to abortion. M. Paul has collected 81 cases¹ in which it caused the death of the fœtus, in some not until after birth; and occasionally it seems to have affected the fœtus even when the mother escaped.

Syphilis.—Of all blood-dyscrasie transmitted to the fœtus, the most important is syphilis. Its influence in producing repeated abortion will be elsewhere described (p. 247). It may unquestionably be transmitted to the fœtus without producing abortion, and at term the mother may be either delivered of a living child bearing evident traces of the disease, of a dead child similarly affected, or of an apparently healthy child in whom the disease develops itself after a lapse of a month or two. These varying effects probably depend on the intensity of the poison; and the longer the time that has elapsed since the origin of the

¹ *Arch. gén. de Méd.*, 1860.

disease in the affected parents, the better will be the chance for the child. The disease is no doubt generally transmitted through the mother, and if she be affected at the time of conception, the infection of the fœtus seems certain. If, however, she contracts the disease at an advanced period of pregnancy, the child may entirely escape. Ricord even believes that syphilis, contracted after the sixth month of pregnancy, never affects the child. The father alone may transmit the disease to the ovum; and Hutchinson has recorded cases to show that the mother may become secondarily affected through the diseased fœtus. The evidences of syphilitic taint in a living or dead child are sufficiently characteristic. The child is generally puny and ill-developed. An eruption of pemphigus is common, either fully developed bullæ or their early stage, when they form circular copper-colored patches. This eruption is always most marked on the hands and feet, and a child born with such an eruption may be certainly considered syphilitic. On post-mortem examination the most usual signs are small patches of suppuration in the thymus, similar localized suppurations in the tissues of the lungs, indurated yellowish patches in the liver, and peritonitis, the importance of which in causing the death of syphilitic children has been specially dwelt on by Simpson.¹

Inflammatory Diseases.—The most important of the inflammatory diseases affecting the fœtus is peritonitis. Simpson has shown that traces of it are very frequently met with, and that it is not always syphilitic. Sometimes it has been observed when the mother has been in bad health during pregnancy, and at others it seems to have resulted from some morbid condition of the fetal viscera. Pleurisy with effusion is another inflammatory affection which has been noticed.

Dropsies.—The dropsical affections most generally met with are ascites and hydrocephalus, which may both have the effect of impeding delivery. Of these, hydrocephalus is the more common, and may give rise to much difficulty in labor. Its causes are uncertain, but it probably depends on some altered state of the mother's health, as it is apt to recur in several successive pregnancies, and is not infrequently associated with an imperfectly developed vertebral column and spina bifida. The fluid collects in the ventricles, which it greatly distends, and these then produce expansion and thinning of the cranium, the bones of which are widely separated from each other at the sutures, which are prominent and fluctuating. In a few cases internal hydrocephalus may be complicated, and the diagnosis in labor consequently obscured by the coexistence of what has been called "external hydrocephalus." This consists of a collection of fluid between the skull and the scalp, which may be either formed there originally or may collect from a rupture of one of the sutures or fontanelles during labor, through which the intracranial fluid escapes.

Ascites is generally associated with hydramnios, and sometimes with hydrothorax or other dropsical effusions. It is a rare affection, and according to Depaul² extreme distension of the bladder is not infrequently mistaken for it.

Tumors.—Tumors of different kinds may be met with in various parts of the child's body, which sometimes grow to a great size and impede

¹ *Obst. Works*, vol. i. p. 117.

² *Tarnier's Cazeaux*, p. 855.

delivery. Tarnier records cases of meningocele larger than a child's head, and large cystic growths have been observed attached to the nates, pectoral region, or other parts of the body. Cancerous tumors of considerable size, either external or of the viscera, have also been met with. Other fetal tumors may be produced by congenital deformities, such as projection of the liver or other abdominal viscera through a deficiency of the abdominal wall; or spina bifida from imperfectly developed vertebræ. The amount of dystocia produced by such causes will, of course, vary much in proportion to the size, consistency, and accessibility of the tumor.

Wounds and Injuries of the Fœtus.—Accidents of serious gravity to the fœtus may happen from violence to which the mother has been subjected, such as falls or blows, without necessarily interfering with gestation. Many curious examples of this kind are on record. Thus, a child has been born presenting a severe lacerated wound extending the whole length of the spine, where both the skin and the muscles had been torn, and which seems to have resulted from the mother having fallen in the last month of pregnancy. Similar lacerations and contusions have been observed in other parts of the body, the wounds being in various stages of cicatrization corresponding to the lapse of time since the accident had occurred. Intra-uterine fractures are not rare, apparently arising from similar causes. In some of these cases the broken ends of the bones had united, but, from want of accurate apposition, at an acute angle, so as to give rise to much subsequent deformity. Chaussier records two cases in which there were many fractures in the same child, in one 113 and in another 42, which were in different stages of repair. He attributes this curious occurrence to some congenital defect in the nutrition of the bones, possibly allied to mollities ossium.¹

Intra-uterine Amputations of Fœtal Limbs.—Intra-uterine amputations of fetal limbs have not unfrequently been observed. Children are occasionally born with one extremity more or less completely absent, and cases are known in which the whole four extremities were wanting (Fig. 93). The mode in which these malformations are produced has given rise to much discussion. At one time it was supposed that the deficiency of the limb was due to gangrene of the extremity and subsequent separation of the sphacelated parts. Reuss, who has studied the whole subject very minutely,² considers gangrene in the unruptured ovum to be an impossibility, for that change cannot occur unless there is access of oxygen, and when portions of the separated extremity are found in utero, as is often the case, they show evidences of maceration, but not of decomposition. The general belief is that these intra-uterine amputations depend on constriction of the limb by folds or bands of the amnion—



Intra-uterine Amputation of both Arms and Legs.

most often met with when the liquor amnii is deficient in quantity—

¹ *Gazette hebdom.*, 1860.

² Scanzoni's *Beiträge*, 1869.

which obstruct the circulation and thus give rise to atrophy of the part below the constriction. It has been supposed that the umbilical cord might, by encircling the limb, produce a like result. It appears doubtful, however, whether this cause is sufficient to produce complete separation of the limb, as any great amount of constriction would interfere with the circulation through the cord. Sometimes, when intra-uterine amputation occurs, the separated portion of the limb is found lying loose in the amniotic cavity, and is expelled after the child. Cases of this kind have been recorded by Martin, Chaussier, and Watkinson. More often no trace of the separated extremity can be found. The explanation probably depends upon the period of utero-gestation at which amputation took place. If it occurred at a very early period of pregnancy, before the third month, the detached portion would be minute and soft, and would easily disappear by solution. If at a later period, this could hardly happen, and the detached portion would remain in utero. In cases of the latter kind cicatrization of the stump has often been observed to be incomplete. Simpson pointed out the occasional existence of rudimentary fingers or toes on the stump of an amputated limb, such as are seen on the thighs in Fig. 93. These he attributed to an abortive reproduction of the separated extremity, analogous to what is observed in some of the lower animals. This explanation has been contested with much show of reason. Martin believes that the reproduction is only apparent, and that the rudimentary extremities are, in reality, instances of arrested development. The constricting agents interfered with the circulation sufficiently to arrest the growth of the limb below the site of constriction, but not sufficiently to effect complete separation. If constriction occurred at a very early stage of development, an appearance similar to that observed by Simpson would be produced. It does not follow, however, that all cases of absence of limbs depend on intra-uterine amputations. In some cases they would appear to be the result of a spontaneous arrest of development or of congenital monstrosity. Mr. Scott¹ relates a case in which a distinct hereditary tendency was evident, and here the deformity certainly could not have resulted from the constriction of amniotic bands. In this family the grandfather had both forearms wanting, with rudimentary fingers attached; the next generation escaped, but the grandchild had a deformity precisely similar to the grandfather.

Death of Fœtus.—When from any cause the fœtus has died during pregnancy, it may be either soon expelled or it may be retained in utero for a longer or shorter time, or even to the full period. The changes observed in such fœtuses vary considerably according to the age of the fœtus at the time of death or the time that it has been retained in utero. If it die at an early period, when the tissues are very soft, it may entirely dissolve in the liquor amnii, and no trace of it may be found when the membranes are expelled. Or it may shrivel or mummify; and if this happen in a twin pregnancy, as sometimes occurs, the growing fœtus may compress and flatten the dead one against the uterine wall.

Appearance of a Putrid Fœtus.—At a later period of pregnancy a dead fœtus undergoes changes ascribed to putrefaction, but which pro-

¹ *Obst. Trans.*, vol. xiii. p. 94.

duce appearances different from those of decomposition in animal textures exposed to the atmosphere. There is no offensive smell, as in ordinary decay. The tissues are all softened and flaccid. The more manifest changes are in the skin, the epidermis of which is separated from the cutis vera, which has a deep reddish color. This is especially apparent on the abdomen, which is flaccid and hollow in the centre. The internal organs are much altered. The brain is diffuent and pulpy, and the cranial bones loose within the scalp. The structures of the muscles and viscera are in various stages of transformation, many having undergone fatty changes, and contain crystals of margaric and cholesterin. The extent to which these changes occur depends, in a great measure, on the length of time the fœtus has been dead, but they do not admit of our estimating with any degree of accuracy what that time has been.

Symptoms and Diagnosis of the Death of the Fœtus.—The symptoms and diagnosis of the death of the fœtus may here be considered. They are, unfortunately, not very reliable. The cessation of the fœtal movements cannot be depended on, as they are frequently unfelt for days or weeks when the child is alive and well. Sometimes the death of the fœtus is preceded by its irregular and tumultuous movements, and in women who have been delivered of several dead children in succession this sensation may guide us in our diagnosis. This suspicion may be confirmed by auscultation. The mere fact that we are unable, at any given time, to hear the fœtal heart will not justify an opinion that the fœtus is dead. If, however, the fœtal heart has been distinctly heard, and after one or two careful examinations, repeated at separate times, it cannot again be made out, the probability of the child being dead may be assumed. Certain changes in the mother's health have been noted in connection with the death of the fœtus, such as depression and lowness of spirits, a feeling of coldness and weight about the lower parts of the abdomen, paleness of the face, a livid circle round the eyes, irregular shiverings and feverishness, shrinking of the breasts, and diminution in the size of the abdominal tumor. All these, however, are too indefinite to justify a positive diagnosis, and they are not infrequently altogether absent. At most they can do no more than cause a suspicion as to what has happened.

CHAPTER X.

ABORTION AND PREMATURE LABOR.

Importance and Frequency of Abortion.—The premature expulsion of the fœtus is an event of great frequency. The number of fœtal lives thus lost is enormous. There are few multiparæ who have not aborted at one time or other of their lives. Hegar estimates that about 1 abortion occurs to every 8 or 10 deliveries at term. Whitehead has calcu-

lated that at least 90 per cent. of married women who lived to the change of life had aborted. The influence of this incident on the future health of the mother is also of great importance. It rarely, indeed, proves directly fatal, but it often produces great debility from the profuse loss of blood accompanying it; and it is one of the most prolific causes of uterine disease in after-life, possibly because women are apt to be more careless during convalescence than after delivery, and the proper involution of the uterus is thus more frequently interfered with.

Definition.—A not uncommon division of the subject is into *abortion*, *miscarriage*, and *premature labor*, the first name being applied to expulsion of the ovum before the end of the fourth month of utero-gestation; *miscarriage*, to expulsion from the end of the fourth to the end of the sixth month; and *premature labor*, to expulsion from the end of the sixth month to the term of pregnancy. This is, however, a needless and confusing subdivision, which leads to no practical result. It suffices to apply the term abortion or miscarriage indiscriminately to all cases in which pregnancy is terminated before the fœtus has arrived at a viable age, and premature labor to those in which there is a possibility of its survival. There is little or no hope of a fetus living before the 28th week or seventh lunar month, and this period is therefore generally fixed on as the limit between premature labor and abortion. The rule is, however, not without an occasional, although very rare, exception. Dr. Keiller of Edinburgh has recorded an instance in which a fœtus was born alive at the fourth month, nine days after the mother had experienced the sensation of quickening. I myself recently attended a lady who miscarried in the fifth month of pregnancy, the child being born alive and living for three hours. Several cases are on record in which after delivery in the sixth month the child survived and was reared. The possibility of the birth of a living child under such circumstances should be recognized, as it may give rise to legal questions of importance; but the exceptions to the ordinary rule are so rare that they need not interfere with the division of the subject usually made.

Abortion is most Common in Multiparæ.—Multiparæ abort far more frequently than primiparæ. This is contrary to the statement in many obstetrical works. Thus, Tyler Smith says "there seems to be a greater danger of this accident in the first pregnancy." Schroeder,¹ however, states that 23 multiparæ abort to 3 primiparæ; and Dr. Whitehead of Manchester, who has particularly studied the subject, believes that abortion is more apt to occur after the third and fourth pregnancies, especially when these take place toward the time for the cessation of menstruation.

Liability to a Recurrence of Abortion.—There can be no doubt that women who have aborted more than once are peculiarly liable to a recurrence of the accident. This can generally be traced to the existence of some predisposing cause which persists through several pregnancies, as, for example, a syphilitic taint, a uterine flexion, or a morbid state of the lining membrane of the uterus. It is probable that in many women a recurrence of the accident induces a habit of abortion, or perhaps it might be more accurate to say a peculiar irritable condition of

¹ Schroeder, *Manual of Midwifery*, p. 149.

the uterus, which renders the continuance of pregnancy a matter of difficulty, independently of any recognizable organic cause.

Very Early Abortions are often Unrecognized.—The frequency of abortion varies much at different periods of pregnancy; and it occurs much more often in the early months, because of the comparatively slight connection then existing between the chorion and the decidua. At a very early period of pregnancy the ovum is cast off with such facility, and is of such minute size, that the fact of abortion having occurred passes unrecognized. Very many cases in which the patient goes one or two weeks over her time, and then has what is supposed to be merely a more than usually profuse period, are probably instances of such early miscarriages. Velpeau detected an ovum of about fourteen days which was not larger than an ordinary pea; and it is easy to understand how so small a body should pass unnoticed in the blood which escapes along with it.

Before the End of the Third Month the Ovum is generally Expelled Entire.—Up to the end of the third month, when miscarriage occurs, the ovum is generally cast off *en masse*, the decidua subsequently coming away in shreds or as an entire membrane. The abortion is then comparatively easy. From the third to the sixth month, after the placenta is formed, the amnion is, as a rule, first ruptured by the uterine contractions, and the foetus is expelled by itself. The placenta and membranes may then be shed as in ordinary labor. It often happens, however, that on account of the firmness of the placental adhesion at this period the secundines are retained for a greater or less length of time. This subjects the patient to many risks, especially to those of profuse hemorrhage and of septicæmia. For this reason, premature termination of the pregnancy is attended by much greater danger to the mother between the third and sixth months than at an earlier or later date. After the sixth month the course of events is not different from that attending ordinary labor. The prognosis to the child is more unfavorable in proportion to the distance from the full period of gestation at which premature labor takes place.

Causes.—The causes of abortion may conveniently be subdivided into the *predisposing* and *exciting*, the latter being often slight, and such as would have no effect in inducing uterine contractions in women unless associated with one or more of the former class of causes. The predisposition to abortion may depend on some condition interfering with the vitality of the ovum or its relation to the maternal structures, or on certain conditions directly affecting the mother's health.

Causes Referable to the Fœtus.—One of the most common antecedents of abortion is the death of the foetus, which leads to secondary changes, and ultimately produces the uterine contractions which end in its expulsion. The precise causes of death in any given case cannot always be accurately ascertained, as they sometimes depend on conditions which are traceable to the maternal structures, at others to the ovular, or it may be to a combination of the two. Nor does it by any means follow that the death of the ovum immediately results in its expulsion. The mode in which death of the ovum produces abortion is not difficult to understand, for it necessarily leads to changes in the relations between

the ovular and maternal structures; these changes cause hemorrhages—partly external and partly into the membranes—which in their turn excite uterine contraction. Extravasations of blood may take place in various positions. One of the most common is into the decidual cavity, between the decidua vera and the decidua reflexa, or between the decidua vera and the uterine walls. If the hemorrhage is only slight, and especially if it comes from that portion of the decidua near the internal os and at a distance from the ovum, there need be no material separation, and pregnancy may continue. This explains the cases occasionally met with in which there is more or less hemorrhage without subsequent abortion. When the amount of extravasated blood is at all great, separation and abortion necessarily result, and the decidua will be found on expulsion to have coagula on its surface and between its various layers which are found to project into the cavity of the amnion (Fig. 94). In

FIG. 94.



An Apoplectic Ovum, with Blood effused in masses under the Fetal Surface of the Membranes.

other cases hemorrhage is still more extensive, and, after breaking through the decidua reflexa, it forms clots between it and the chorion, and even in the cavity of the amnion. Supposing expulsion to take place shortly after coagula are deposited among the membranes, the blood is little altered, and we have an ordinary abortion. If, however, the ovum is retained, the coagulated fibrin and the placenta or membranes undergo secondary changes which lead to the formation of moles. The so-called *fleshy mole* (Fig. 95) is often retained for many weeks or months after the death of the fetus, and during this time there may be but little modification of the usual symptoms of pregnancy; or, as is frequently the case, it gives rise to occasional hemorrhage, until at last uterine contractions come on, and it is cast off in the form of a thick fleshy mass, having but little resemblance to the ordinary products of

conception. The most probable explanation of its formation is, that when hemorrhage originally took place the effusion of blood was not sufficient to effect the entire separation and expulsion of the ovum. Part of the membranes or of the placenta—if that organ had commenced to form—retained its organic connection with the uterus, while the fœtus

FIG. 95.



Blighted Ovum, with Fleshy Degeneration of the Membranes.

perished. The attached portion of the placenta or membranes continues to be nourished, although abnormally. The fœtus generally entirely disappears, especially if it has perished at an early period of utero-gestation, when it becomes dissolved in the liquor amnii. Or it may become macerated, shrivelled, and greatly altered in appearance. The effused blood becomes decolorized from the absorption of the corpuscles, and, according to Scanzoni, fresh vessels are developed in the fibrin, which increase the vascular attachment of the mole to the uterine walls. The placenta and membranes may go on increasing in thickness until they form a mass of considerable size. Careful microscopic examination will almost always enable us to discover the villi of the chorion, altered in appearance, often loaded with granular fatty molecules, but sufficiently distinct to be readily recognizable.

Causes depending on the Maternal State.—Important as are the causes of abortion arising from some morbid condition of the ovum, they are not more so than those which depend on the maternal state, and it is to be observed that the former are often indirect causes, produced by primary maternal changes. Many of these maternal causes act by causing hyperæmia of the uterus, which leads to extravasation of blood. Thus, abortion is apt to occur in women who lead unhealthy lives, such as those who occupy overheated and ill-ventilated rooms or indulge to excess in the fatigues and pleasures of society, in the use of alcoholic drinks, and the like. Over-frequent coitus has been, for the same reason, observed to produce a remarkable tendency to abortion, and Parent-

Duchatelet has noted that it is of very frequent occurrence amongst women of loose life. Many diseases strongly predispose to it, such as fevers, zymotic diseases of all kinds, measles, scarlet fever, small-pox; and diseases of the respiratory organs, such as bronchitis and pneumonia. *Syphilis* is well known to be one of the most frequent causes, and one that is likely to act in successive pregnancies. It may act so that the pregnancy is brought to a premature termination, time after time, until the constitutional disease is eradicated by appropriate treatment. It acts in some cases through the influence of the father in producing a diseased ovum, and it is the only cause which can with certainty be traced to the state of the father's health. Many other morbid conditions of the blood also dispose to abortion. It has been observed to be a frequent result of lead-poisoning; also of the presence of noxious gases in the atmosphere, such as an excess of carbonic acid.

Causes acting through the Nervous System.—Many causes act through the nervous system, such as fright, anxiety, sudden shock, and the like. Thus, there are numerous instances on record in which women aborted suddenly after the receipt of some bad news, and it is said to have been of frequent occurrence in women immediately before execution. The influence of irritation propagated through the nervous system from a distance, tending to produce uterine contraction and abortion through the agency of reflex action, has been specially dwelt upon by Tyler Smith. Thus he points out that abortion not unfrequently occurs from the irritation of constant suckling in women who become pregnant during lactation. The effect of suckling in producing uterine contraction is indeed well known, and the application of the child to the breast for this purpose has long been recognized as a method of treatment in post-partum hemorrhage. The irritation of the trifacial in severe toothache, of the renal nerves in cases of gravel, in albuminuria, etc., of the intestinal nerves in excessive vomiting, in diarrhoea, obstinate constipation, ascariides, etc., all act in the same way. We may perhaps also explain by this hypothesis the fact that women are more apt to abort at what would have been the menstrual epoch than at other times, as the ovarian nerves may then be subject to undue excitement. It is probable, however, that there may be also at these times more or less active congestion of the decidua, which may predispose to laceration of its capillaries and blood-extravasation. Such congestion exists in those exceptional cases in which menstruation continues for one or more periods after conception, the blood probably escaping from the space between the decidua vera and reflexa; and therefore there is no reason to question its also happening even when such abnormal menstruation is not present.

Physical Causes.—Certain physical causes may produce abortion by separating the ovum. Thus it may follow a fall, a blow, or other accidents of a trivial character. On the other hand, women may be subjected to injuries of the severest kind without aborting. The probability, therefore, is that these apparently trivial causes only operate in women who for some other reason are predisposed to the accident. This is borne out by the fact—which is well known in these days, when the artificial production of abortion is, unhappily, far from a very rare

event—that it is by no means easy to destroy the vitality of the fœtus. I myself know of a case in which the uterine sound was passed several times into a pregnant uterus without producing abortion, the pregnancy proceeding to term. Oldham has related a similar case in which he in vain attempted to produce abortion by the sound in a case of contracted pelvis; and Duncan has mentioned an instance in which an intra-uterine stem pessary was unwittingly introduced, and worn for some time by a pregnant woman, without any bad effect. The fact that pregnancy is with difficulty interfered with when there is a healthy relation between the ovum and the uterus no doubt explains the disastrous effects of criminal abortion which have been especially insisted on by many of our American brethren.

Cases depending on Morbid States of the Uterus.—Morbid states of the uterus have an important influence in the production of abortion. Any condition which mechanically interferes with the proper development of the uterus is apt to operate in this way. Amongst these may be mentioned fibroid tumors; the presence of old peritoneal adhesions, rendering the womb a more or less fixed organ; but, above all, flexion and displacement of the uterus. Retroflexion of the uterus is unquestionably one of the most frequent factors in its production, not only on account of the irritation which the abnormal position sets up, but from interference with the uterine circulation, which leads to the effusion of blood and the death of the ovum. An inflamed condition of the cervical and uterine mucous membranes will act in the same way should pregnancy have occurred, although such a condition more often prevents conception taking place.

Symptoms.—One of the earliest indications of impending abortion is more or less hemorrhage. This may at first be slight, and may last for a short time only, recurring after an interval of time, or it may commence with a sudden and profuse discharge. Occasionally it is very abundant, and its continuance and amount form one of the gravest symptoms of the accident. After the loss of blood has continued for a greater or less length of time—it may be even for some days—uterine contractions come on, recurring at regular intervals, and eventually lead to the expulsion of the ovum. More rarely the impending miscarriage commences with pains which lead to laceration of vessels and hemorrhage.

When Pain and Hemorrhage Coexist.—As long as one or other of these symptoms exist alone we may hope to avert the threatened miscarriage, but when both occur together there is little or no chance of its being arrested. Certain premonitory symptoms are described by authors as common in abortion, such as feverishness, shivering, a sensation of coldness; all of which are obscure and unreliable, and are certainly much more frequently absent than present.

If the pregnancy be early, it is probable that the entire ovum will be shed with little trouble, and it often passes unperceived in the clots which surround it. It is therefore of importance that all the discharges should be very carefully examined. After the second month the rigid and undilated cervix presents a formidable obstacle to the escape of the ovum, and it may be a considerable time before there is sufficient dilata-

tion to admit of its passage. This is gradually effected by the continuance of pains, but not without a severe loss of blood. It may be that the amnion is ruptured and the fœtus expelled first. After a lapse of time the secundines are also shed, but there may be a considerable delay, amounting even to days, before this is effected. As long as any portions of the membranes are retained in utero the patient is necessarily subjected to considerable risk, not only from the continuance of hemorrhage, but also from septicæmia. Hence it may be laid down as a rule that we can never consider our patient out of danger until we have satisfied ourselves that the whole of the uterine contents have been expelled.

Treatment.—Our first endeavor in any case of impending miscarriage will be, of course, to avert the threatened accident. If hemorrhage has not been excessive, and if on vaginal examination—which should always be practised—we find no dilatation of the os, we may entertain a reasonable hope of success. If, on the contrary, we find the os beginning to open, if we are able to insert the finger through it so as to touch the ovum, especially if pains also exist, we are justified in considering abortion to be inevitable, and the indication will then be to have the ovum expelled and the case terminated as soon as possible. In the former case the most absolute rest is the first thing to insist on. The patient should be placed in bed, not overburdened with clothes, in a cool temperature, and she should have a light and easily-assimilated diet. All movements, even rising out of bed to empty the bladder or bowels, should be absolutely prohibited. To avert the tendency to the commencement of uterine contraction there is no remedy so useful as opium, which must be given freely and frequently repeated. It may be administered either in the form of laudanum or of Battley's sedative solution, which has the advantage of producing less general disturbance. It may be advantageously exhibited in doses of from 20 to 30 minims, and repeated after a few hours. A still better preparation is chlorodyne, which I have found of extreme value in arresting impending miscarriage, in doses of 10 minims repeated every third or fourth hour. If, from any other cause, it is considered unadvisable to give the sedative by the mouth, it may be administered in a small starch enema *per rectum*. In all cases it will be necessary to keep the patient more or less under the influence of the drug for several days and until all symptoms of miscarriage have passed away. Care should be taken that the bowels do not become locked up by the action of the opiates—as this might of itself be a cause of irritation—and their constipating effects ought to be obviated by small doses of castor oil or other gentle aperient. Various subsidiary methods of treatment have been recommended, such as bleeding from the arm or the local application of leeches in supposed plethoric states of the system; revulsives, such as dry cupping to the loins; the application of ice, to check hemorrhage; astringents, such as acetate of lead or gallic acid, for the same purpose. Most of these, if not hurtful, will be at least useless. The cases in which venesection would be beneficial are extremely rare, and the local applications, especially cold, are much more apt to favor than to prevent uterine action.

Prophylactic Treatment.—In cases of repeated miscarriage in successive pregnancies a special course of prophylactic treatment is indicated, and

is often attended with much success. In cases of this kind the first indication, and one which ought to be carefully attended to, is to seek for and, if possible, to remove or mitigate the cause which has given rise to the former abortions. Those causes which depend on constitutional states must first be carefully investigated, and treated according to the indications present. These may be obscure and not easily discovered; but it is certainly unwise to assume too readily the existence of what has been called "a habit of abortion," which further inquiry may prove to be only an indication of constitutional debility, degeneracy of the placental structures, or a latent and unsuspected syphilitic taint. If constitutional debility be present to a marked extent, a generous diet and a restorative course of treatment (preparations of iron, quinine, and other suitable tonics) may effect the desired object.

[As an evidence of the efficiency of opium, I once succeeded in arresting a labor at four and a half months by repeated doses of sulphate of morphia, in the case of a lady who was in a decidedly parturient state for ten hours, with recurrent uterine contractions, accompanied by pain and a considerable loss of blood. Under the narcotic the pains became more and more infrequent until they finally ceased, and the patient carried the fetus to the full period. It was a small female child, and lived some months, its delicacy being largely due to the fact that its mother was phthisical. In another case labor was arrested at eight months, and the fetus was carried to the full period. In the hands of some obstetricians the fluid extract of *Viburnum prunifolium* would appear to act efficiently as a preventive of abortion in cases where the habit is known to exist. In one case of this habit, after repeated failures, a residence in a mountainous region carried the patient through the usual period of danger, and the fetus is now a young lady. The mother was of very full habit, asthmatic, and at times rheumatic, but otherwise in excellent health. She usually aborted at six weeks. After three months she had occasional threatenings, but not any during the last two months. Rest in bed checked any apparent tendency to miscarry.—ED.]

Treatment in Cases depending on Local Causes.—Local congestion of the uterus or a general plethoric state of the patient has often been supposed to be an efficient cause of recurring abortion. Dr. Henry Bennet has especially dwelt on the influence of congestion and abrasions of the cervix in causing premature expulsion of the fetus,¹ and recommends the topical application of nitrate of silver or other caustics to the inflammatory abrasions existing on the neck of the womb. Formerly venesection was a favorite remedy; and many authors have recommended the local abstraction of blood by leeches applied to the groin or round the anus, or even to the cervix. The influence of general plethora is more than doubtful; and although local congestions are, probably, much more effective causes, still it would seem more judicious to treat them by rest and local sedatives rather than by topical applications, which, injudiciously applied, might produce the very accident they were intended to prevent.

The position of the uterus should be carefully investigated. If it be found to be retroflexed, a well-fitting Hodge's pessary should be ap-

¹ *On Inflammation of the Uterus*, p. 432.

plied, so as to support it until it has completely risen out of the pelvis.

Treatment in Cases depending on Syphilis.—The possibility of syphilitic infection should always be inquired into, for this poison may act on the product of conception long after all appreciable traces of it have disappeared from the infected parent. Should there be recurrent abortions in a patient who had formerly suffered from syphilis or whose husband had at any time contracted the disease, no time should be lost in using appropriate antisyphilitic remedies, which should invariably be administered both to the husband and wife. Diday especially insists that in such cases it is not sufficient to submit the father and mother to a mercurial course in the absence of pregnancy, but that, as each successive impregnation occurs, the mother should again commence antisyphilitic treatment, even though she has no visible traces of the disease.¹ In this way there is reasonable ground for hoping that infection of the ovum may be prevented. I think, too, that we may be the more encouraged to persevere in the treatment of these unfortunate cases from the fact that the syphilitic poison tends to wear itself out. I have seen several cases in which this taint at first produced early abortion, then each successive pregnancy was of longer duration, until eventually a living child was born.

In fatty degeneration of the chorion villi and in other morbid states of the placenta, which act by preventing the proper nutrition of the fœtus and the due aëration of its blood, there is no reliable means of treatment except the general improvement of the mother's health. Simpson strongly recommended the administration of chlorate of potash in cases in which the child habitually dies in the latter months of pregnancy, on the supposition that it supplied to the blood a large amount of oxygen, and thus made up for any deficiency in the supply of that element through the placental tufts. The theory is at best a doubtful one, although I believe the drug to be unquestionably beneficial in cases of the kind.^[2] It probably acts by its tonic properties rather than in the manner Simpson supposed. It may be given in doses of 15 to 20 grains three times a day, and may be advantageously combined with small doses of dilute hydrochloric acid. In frequently-recurring premature labors with dead children Simpson strongly recommended the induction of premature labor a little before the time at which we had reason to believe that the fœtus has usually perished; or, in other words, before the placental disease had advanced sufficiently far to interfere with its nutrition. The practice has constantly been adopted with success, and is perfectly legitimate, but the difficulty, of course, is to fix on the right time. Careful auscultation of the fetal heart may be of some use in guiding us to a decision, as the death of the fœtus is generally preceded for some days by irregular, tumultuous, and intermittent action of the heart.

Treatment where no Cause can be Discovered.—There will always remain a certain number of cases in which no appreciable cause can be

¹ Diday. *Infantile Syphilis*, Syd. Soc. Trans., p. 207.

^[2] Chlorate of potash will not part with its oxygen except at a red heat. In the system it remains unchanged, as proved by Prof. Leeds of Hoboken.—Ed.]

discovered. Under such circumstances prolonged rest, at least until the time has passed at which abortion formerly took place, will afford the best chance of avoiding a recurrence of the accident. There must always be some difficulty in carrying out this indication, inasmuch as the patient's health is apt to suffer in other ways from the confinement and the want of fresh air and exercise which it entails. The strictness with which rest should be insisted on must vary in different cases, but it should be specially attended to at what would have been the menstrual periods. At these times the patient should remain in bed altogether; at others she may lie on a sofa, and, if circumstances permit, spend part of the day at least in the open air. Sexual intercourse should be prohibited. Should actual symptoms of abortion come on, the preventive treatment, already indicated, may be resorted to. Great care, however, should be used in prescribing opiates as preventives, and they should be given for a specified time only. I have seen, more than once, an incurable habit of opium-eating originate from the incautious and too long-continued exhibition of the drug in such cases.

When we have satisfied ourselves that abortion is inevitable, we must proceed to employ treatment that favors the expulsion of the ovum.

Removal of the Ovum when within Reach.—If the os be sufficiently dilated and the pains strong, we may find the ovum separated and protruding from the os. We may then be able to detach it by the finger. For this purpose the uterus is depressed from without by the left hand, while an endeavor is made to scoop out the ovum with the examining finger. If it be out of reach, and yet appears detached, chloroform should be administered, the whole hand introduced into the vagina, and the finger into the uterine cavity. The complete detachment of the ovum can in this way be far more readily and safely effected than by using any of the many ovum-forceps which have been invented for the purpose.

Plugging of the Vagina.—If the ovum be not sufficiently separated or the os be undilated, means must be taken to control the hemorrhage until the former can be removed or expelled. It is here that plugging of the vagina finds its most useful application. This may be done in various ways. That most usually employed is filling the vagina with a tolerably large sponge, in the interstices of which the blood coagulates. A better plan is to soak a number of pledgets of cotton wool in carbolized water and tie a string round each. The vagina can be completely and effectively packed with these, and this is best done through a speculum. Each pledget should be covered with glycerin, which completely prevents the offensive odor which otherwise arises. The pledgets can be removed by traction on the strings, but if these are not used much pain is caused in getting them out of the vagina. The plug should never be left in for more than six or eight hours, after which a fresh one may be inserted if necessary. Two or three full doses of the liquid extract of ergot, of fʒss to fʒj each, or a subcutaneous injection of ergotin, may be given while the plug is in position. The plug itself is a strong excitant of uterine action, and the two combined often effect complete detachment, so that on the removal of the tampon the ovum may be found lying loose in the os uteri. If the os be undilated and the

ovum entirely out of reach, the former may be opened by means of sponge or laminaria tents. I think a well-prepared sponge tent the most effectual, and it can be maintained *in situ* by a vaginal plug below it. It also acts as a most efficient plug, effectually controlling all hemorrhage. In a few hours it opens up the os sufficiently to admit the finger.

Retention of the Membranes.—The most troublesome cases are those in which the fetus is first expelled and the placenta and membranes remain in utero. As long as this is the case the patient can never be considered safe from the occurrence of septicæmia. Dr. Priestley has strongly insisted on the importance of removing the secundines as soon as possible. There can be no doubt that this should be done whenever it is feasible. Cases, however, are frequently met with in which any forcible attempt at removal would be likely to prove very hurtful, and in which it is better practice to control hemorrhage by the plug or sponge tent, and wait until the placenta is detached, which it will generally be in a day or two at most. Under such circumstances fetor and decomposition of the secundines may be prevented by intra-uterine injections of diluted Condyl's fluid. Provided the os be sufficiently patulous to prevent the collection of the fluid in the uterine cavity, and not more than a drachm or two of fluid be injected at a time, so as simply to wash away and disinfect decomposing detritus, they can be used with perfect safety. Sometimes cases are met with in which the os has entirely closed, and in which we can only suspect the retention of the placenta by the history of the case, the continuance of hemorrhage, or the presence of a fetid discharge. Should we see reason to suspect this, the os must be dilated with sponge or laminaria tents and the uterine cavity thoroughly explored under chloroform. This condition of things is far from uncommon in women who have not had medical assistance from the first, and it often gives rise to very troublesome and anxious symptoms. It has been said that placenta thus retained have been completely absorbed, and cases of the kind have been related by Nægele and Oslander. The spontaneous absorption, however, of so highly organized a body as the placenta would be a phenomenon of the most remarkable character; and it seems more natural to suppose that in most cases of the kind the placenta has been cast off without the knowledge of the patient. Sometimes the placenta never entirely becomes detached, and, retaining organic connection with the uterine walls, forms what has been called a "*placental polypus*." This may produce secondary hemorrhages, in the same way as an ordinary fibroid polypus. Barnes recommends the removal of these masses by means of the wire *écraseur*. Before their detection the os uteri must be opened up.

Retention in Utero of a Blighted Ovum.—The cases previously alluded to, in which an ovum has perished in early pregnancy and is retained in utero, are often puzzling, and may give rise to serious moral and medico-legal questions. The blighted ovum may be retained for many months, the outside limit, according to McClintock,¹ by whom the subject has been ably discussed, being nine months. The appearance of the ovum when thrown off will give no reliable clue to the length of time which has elapsed since it perished. The symptoms are often very obscure.

¹ Sydenham's Society's ed. of *Smellie's Midwifery*, vol. i. p. 169.

Generally there have been the usual indications of pregnancy, which, with or without signs of impending miscarriage, disappear or are modified, and then follows a period of ill-health, with pelvic uneasiness and irregular metrorrhagia, which may be mistaken for menstruation. Occasionally, but by no means necessarily, there is a fetid discharge, and this probably exists only when the membranes have broken and air has access to the ovum. In some cases obscure septicæmic symptoms have been observed. Such symptoms are obviously too indefinite to lead to an accurate diagnosis. In the course of time the ovum is generally thrown off, with more or less hemorrhage. If the nature of the case is detected, ergot may be given to promote the expulsion of the uterine contents, and it may even be advisable to dilate the cervix with sponge or laminaria tents and remove them artificially.

Subsequent Management.—The frequency with which abortion leads to chronic uterine disease should lead us to attach much more importance to the subsequent management of the patient than has been customary. The usual practice is to confine the patient to bed for two or three days only, and then to allow her to resume her ordinary avocations, on the supposition that a miscarriage requires less subsequent care than a confinement. The contrary of this is, however, most probably the case, for the uterus has been emptied when it is unprepared for involution, and that process is often very imperfectly performed. We should therefore insist on at least as much attention being paid to rest as after labor at term. [A common cause of *uterine hyperplasia* is the imperfect involution which is apt to follow an abortion at an early period of gestation, especially when the patient is not properly treated by rest in a recumbent position. Displacements of the uterus are apt to be produced by the erect position and exercise while the organ is heavy and its ligaments relaxed.—ED.]

PART III.

LABOR.

CHAPTER I.

THE PHENOMENA OF LABOR.

Delivery at Term.—In considering delivery at term we have to discuss two distinct classes of events.

One of these is the series of vital actions brought into play in order to effect the expulsion of the child ; and the other consists of the movements imparted to the child, the body to be expelled—in other words, the mechanism of delivery.

Causes of Labor.—Before proceeding to the consideration of these important topics a few words may be said as to the determining causes of labor. This subject has been from the earliest times a *questio vexata* among physiologists, and many and various are the theories which have been broached to explain the curious fact that labor spontaneously commences, if not at a fixed epoch, at any rate approximately so. It must be admitted that even yet there is no explanation which can be implicitly accepted.

Fœtal or Maternal Causes.—The explanations which have been given may be divided into two classes—those which attribute the advent of labor to the fœtus, and those which refer it to some change connected with the maternal generative organs.

The former is the opinion which was held by the older accoucheurs, who assigned to the fœtus some active influence in effecting its own expulsion. It need hardly be said that such fanciful views have no kind of physiological basis. Others have supposed that there might be some change in the placental circulation or in the vascular system of the fœtus which might solve the mystery. The latest hypothesis of this kind—which, however, is not fortified by any evidence—is by Barnes, who says: “I rather incline to the opinion that when the fœtus has attained its full development, when its organs are prepared for external life, some change takes place in its circulation which involves a correlative disturbance in the maternal circulation, which excites the attempt at labor.”¹

The majority of obstetricians, however, refer the advent of labor to purely maternal causes. Among the more favorite theories is one which

¹ *Diseases of Women*, p. 434.

was originally started in this country by Dr. Power and adopted and illustrated by Depaul, Dubois, and other writers. It is based on the assumption that there is a sphincter action of the fibres of the cervix analogous to that of the sphincters of the bladder and rectum, and that when the cervix is taken up into the general uterine cavity as pregnancy advances, the ovum presses upon it, irritates its nerves, and so sets up reflex action, which ends in the establishment of uterine contraction. This theory was founded on erroneous conceptions of the changes that occurred in the neck of the uterus; and, as it is certain that obliteration of the cervix does not really take place in the manner that Power believed when his theory was broached, it is obvious that its supposed result cannot follow.

Distension of the Uterus.—Extreme distension of the uterus has been held to be the determining cause of labor—a view lately revived by Dr. King of Washington,¹ who believes that contractions are induced because the uterus ceases to augment in capacity, while its contents still continue to increase. This hypothesis is sufficiently disproved by a number of clinical facts which show that the uterus may be subject to excessive and even rapid distension—as in cases of hydramnios, multiple pregnancy, and hydatidiform degeneration of the ovum—without the supervention of uterine contractions.

Fatty Degeneration of the Decidua.—Another inciter of uterine action has been supposed to be the separation of the ovum from its connections to the uterine parietes in consequence of fatty degeneration of the decidua occurring at the end of pregnancy. The supposed result of this change, which undoubtedly occurs, is that the ovum becomes so detached from its organic adhesions as to be somewhat in the position of a foreign body, and thus incites the nerves so largely distributed over the interior of the uterus. This theory, which has been widely accepted, was originally started by Sir James Simpson, who pointed out that some of the most efficient means of inducing labor (such, for example, as the insertion of a gum-elastic catheter between the ovum and the uterine walls) probably act in the same way—viz. by effective separation of the membranes and detachment of the ovum.

Barnes instances, in opposition to this idea, the fact that ineffectual attempts at labor come on at the natural term of gestation in cases of extra-uterine pregnancy, when the fœtus is altogether independent of the uterus, and therefore, he argues, the cause cannot be situated in the uterus itself. A fair answer to this argument would be that although, in such cases, the womb does not contain the ovum, it does contain a decidua, the degeneration and separation of which might suffice to induce the abortive and partial attempts at labor then witnessed.

Objections to these Theories.—A serious objection to all these theories, which are based on the assumption that some local irritation brings on contraction, is the fact, which has not been generally appreciated, that uterine contractions are always present during pregnancy as a normal occurrence, and that they may be, and often are, readily intensified at any time, so as to result in premature delivery.

It is, indeed, most likely that, at or about the full term, the nervous

¹ *American Journal of Obstetrics*, vol. iii.

supply of the uterus is so highly developed, and in so advanced a state of irritability, that it more readily responds to stimuli than at other times. If by separation of the decidua, or in some other way, stimulation of the excitor nerves is then effected, more frequent and forcible contractions than usual may result, and, as they become stronger and more regular, terminate in labor. But, allowing this, it still remains quite unexplained why this should occur with such regularity at a definite time.

Tyler Smith's Ovarian Theory.—Tyler Smith tried, indeed, to prove that labor came on naturally at what would have been a menstrual epoch, the congestion attending the menstrual nixus acting as the exciter of uterine contraction. He therefore refers the onset of labor to ovarian rather than to uterine causes. Although this view is upheld with all its author's great talent, there are several objections to it difficult to overcome. Thus, it assumes that the periodic changes in the ovary continue during pregnancy, of which there is no proof. Indeed, there is good reason to believe that ovulation is suspended during gestation, and with it, of course, the menstrual nixus. Besides, as has been well objected by Cazeaux, even if this theory were admitted, it would still leave the mystery unsolved, for it would not explain why the menstrual nixus should act in this way at the tenth menstrual epoch rather than at the ninth or eleventh.

In spite, then, of many theories at our disposal, it is to be feared that we must admit ourselves to be still in entire ignorance of the reason why labor should come on at a fixed epoch.

Mode in which the Expulsion of the Child is Effected.—The expulsion of the child is effected by the contractions of the muscular fibres of the uterus, aided by those of some of the abdominal muscles. These efforts are in the main entirely independent of volition. So far as regards the uterine contractions, this is absolutely true, for the mother has no power of originating, lessening, or increasing the action of the uterus. As regards the abdominal muscles, however, the mother is certainly able to bring them into action and to increase their power by voluntary efforts; but, as labor advances and as the head passes into the vagina and irritates the nerves supplying it, the abdominal muscles are often stimulated to contract, through the influence of reflex action, independently of volition on the part of the mother.

The Chief Factor in Expulsion.—There can be little doubt that the chief agent in the expulsion of the child is the contraction of the uterus itself. This opinion is almost unanimously held by accoucheurs, and the influence of the abdominal muscles is believed to be purely accessory. Dr. Haughton, however, maintains a view which is directly contrary to this. From an examination of the force of the uterine contractions, arrived at by measuring the amount of muscular fibre contained in the walls of the uterus, he arrives at the conclusion that the uterine contractions are chiefly influential in rupturing the membranes and dilating the os uteri, bringing into action, if needful, a force equivalent to 54 lbs.; but when this is effected, and the second stage of labor has commenced, he thinks the remainder of the labor is mainly completed by the contractions of the abdominal muscles, to which he attributes enormous powers

—equivalent, if needful, to a pressure of 523.65 lbs. on the area of the pelvic canal.

These views bear on a topic of primary consequence in the physiology of labor. They have been fully criticised by Duncan, who has devoted much experimental research to the study of the powers brought into action in the expulsion of the child. His conclusions are that, so far from the enormous force being employed that Haughton estimated, in the large majority of cases the effective force brought to bear on the child by the combined action of both the uterine and abdominal muscles is less than 50 lbs. ; that is, less than the force which Haughton attributed to the uterus alone. In extremely severe labors, when the resistance is excessive, he thinks that extra power may be employed, but he estimates the maximum as not above 80 lbs., including in this total the action of both the uterine and abdominal muscles. Joulin arrived at the conclusion that the uterine contractions were capable of resisting a maximum force of about one hundredweight. Both these estimates, it will be observed, are much under that of Haughton, which Duncan describes as representing "a strain to which the maternal machinery could not be subjected without instantaneous and utter destruction."

Reasons on which this Conclusion is Based.—There are many facts in the history of parturition which make it certain that the chief factor in the expulsion of the child is the uterus. Among these may be mentioned occasional cases in which the action of the abdominal muscles is materially lessened, if not annulled—as in profound anæsthesia and in some cases of paraplegia—in which, nevertheless, uterine contractions suffice to effect delivery. The most familiar example of its influence, however, and one that is a matter of every-day observation in practice, is when inertia of the uterus exists. In such cases no effort on the part of the mother, no amount of voluntary action that she can bring to bear on the child, has any appreciable influence on the progress of the labor, which remains in abeyance until the defective uterine action is re-established or until artificial aid is given.

The contraction of the uterus, then, being the main agent in delivery, it is important for us to appreciate its mode of action and its effect on the ovum.

Uterine Contractions at the Commencement of Labor.—We have seen that intermittent and generally painless uterine contractions exist during pregnancy. As the period for delivery approaches these become more frequent and intense, until labor actually commences, when they begin to be sufficiently developed to effect the opening up of the os uteri with a view to the passage of the child. They are now accompanied by pain, which increases as labor advances, and is so characteristic that "pains" are universally used as a descriptive term for the contractions themselves. It does not necessarily follow that uterine contractions are painless until they commence to effect dilatation of the os uteri. On the contrary, during the last days or even weeks of pregnancy women constantly have irregular contractions, accompanied by severe suffering, which, however, pass off without producing any marked effect on the cervix. When labor has actually begun, if the hand is placed on the uterus when a pain commences the contraction of its muscular tissue is very apparent,

and the whole organ is observed to become tense and hard, the rigidity increasing until the pain has reached its acme, the uterine walls then relaxing and remaining soft until the next pain comes on. At the commencement of labor these pains are few, separated from each other by a considerable interval, and of short duration. In a perfectly typical labor the interval between the pains becomes shorter and shorter, while, at the same time, the duration of each pain is increased. At first they may occur only once in an hour or more, while eventually there may not be more than a few minutes' interval between them.

Mode in which Dilatation of the Cervix is Effected.—If, when the pains are fairly established, a vaginal examination be made, the os uteri will be found to be thinned and dilated in proportion to the progress of the labor. During the contraction the bag of membranes will be felt to bulge, to become tense from the downward pressure of the liquor amnii within it, and to protrude through the os if it be sufficiently open. The membranes, with the contained liquor amnii, thus form a fluid wedge, which has a most important influence in dilating the os uteri (see Frontispiece). This does not, however, form the sole mechanism by which the os uteri is dilated, for it is also acted upon by the contractions of the muscular fibres of the uterus, which tend to pull it open. It is probable that the muscular dilatation of the os is effected chiefly by the longitudinal fibres, which, as they shorten, act upon the os uteri, the part where there is least resistance.

Partly, then, by muscular contraction, partly by mechanical pressure, the cervical canal is dilated, and as it opens up it becomes thinner and thinner, until it is entirely taken up into the uterine cavity.

Rupture of the Membranes.—There is no longer any obstacle to the passage of the presenting part of the child into the cavity of the pelvis, and the force of the pains now generally effects the rupture of the membranes and the escape of the liquor amnii. There is often observed, at this time, a temporary relaxation in the frequency of the pains, which had been steadily increasing; but they soon recommence with increased vigor. If the abdomen be now examined, it will be observed to be much diminished in size, partly in consequence of the escape of the liquor amnii, partly from the descent of the fœtus into the pelvic cavity.

Change in the Character of the Pains.—The character of the pains soon changes. They become stronger, longer in duration, separated by a shorter interval, and accompanied by a distinct forcing effort, being generally described as “the bearing-down” pains. Now is the time at which the accessory muscles of parturition come into operation. The patient brings them into play in the manner which will be subsequently described, and the combined action of the uterine and abdominal muscles continues until the expulsion of the child is effected.

Mode of Action of the Uterus.—The precise mode of uterine contraction is still somewhat a matter of dispute. It is generally described as commencing in the cervix, passing gradually upward by peristaltic action, the wave then returning downward toward the os uteri. This view was maintained by Wigand, and has been endorsed by Rigby, Tyler Smith, and many other writers. In support of it they instance the fact that on the accession of a pain the presenting part first recedes;

the bag of membranes then becomes tense and protrudes through the os, and it is not until some time that the presenting part of the child itself is pushed down. It is very doubtful if this view is correct; and a careful examination of the course of the pains would rather lead to the belief that the contractions commence at the fundus, where the muscular tissue is most largely developed, and gradually proceed downward to the cervix, the waves of contraction being, however, so rapid that the whole organ seems to harden *en masse*. The apparent recession of the presenting part and the bulging of the bag of membranes are certainly no proof that the contractions begin at the cervix; for the commencing contraction would necessarily push down the fluid in front of the head, and cause the membranes to bulge and the os to become tense before its force was brought to bear on the fœtus itself. Indeed, did the contraction commence at the lower part of the uterus, we should expect the opposite of what takes place to occur, and the waters to be pushed upward and away from the cervix. The fundal origin of the contraction is further illustrated by what is observed when the hand of the accoucheur is placed in the uterine cavity, as often happens in certain cases of hemorrhage or turning; for if a pain then comes on, it will be felt to start at the fundus, and gradually compress the hand from above downward.

Value of the Intermittent Character of the Pains.—The intermittent character of the contractions is of great practical importance. Were they continuous, not only would the muscular powers of the patient be rapidly exhausted, but by the obliteration of the vessels produced by the muscular contraction the circulation through the placenta would be interfered with and the life of the child imperilled. Hence, one of the chief dangers of protracted labor, especially after the escape of the liquor amnii, is that the uterine fibres may enter into a state of tonic rigidity—a condition that cannot be long continued without serious risks both to the mother and child.

The Contractions are Incited through the Sympathetic Nerves.—The fact that the uterine contractions are altogether involuntary proves them to be excited—as indeed we would *a priori* infer from our knowledge of the anatomical arrangement of the nerves of the uterus—solely by the sympathetic system. Still, it is a fact of every-day observation that they can be largely influenced by emotions. Various stimuli applied to the spinal system of nerves (as, for example, when the mammæ are irritated) have also a marked effect in inducing uterine contraction. The precise mode in which such influence is conveyed to the uterus, in spite of the numerous experiments which have been made for the purpose of determining how far labor is affected by destruction of the spinal cord, is still a matter of doubt. After the fœtus has passed through the cervix the spinal nerves distributed to the vagina and perineum are excited by the pressure of the presenting part, and through them the accessory powers of parturition are chiefly brought into play. The contraction of the muscles of the vagina itself is supposed to have some influence in favoring the expulsion of the fœtus after the birth of part of the body, and also in promoting the expulsion of the placenta. In the lower animals the vagina has a very marked contractile property, and is, in some of them,

the main agent by which the young are expelled. In the human subject this influence is certainly of very secondary importance.

Character and Source of Pains during Labor.—The amount of suffering experienced during labor varies much in different cases, and is in direct proportion to the nervous susceptibility of the patient. There are some women who go through labor with little or no pain at all. This is proved by the cases (of which there are numerous authentic instances recorded) in which labor has commenced during sleep, and the child has been actually born without the mother awaking. I am acquainted with a lady who has had a large family, who assures me that, though labor is accompanied by a sense of pressure and discomfort, she experiences nothing which can be called actual pain. Such a happy state of affairs is, however, extremely exceptional, and in the vast majority of cases parturition is accompanied by intense suffering during its whole course, in some cases amounting to anguish which has probably no parallel under any other condition.

The precise cause of the pain has been much discussed, and is no doubt complex.

In the First Stage.—In the early stage of labor, and before the dilatation of the os, it is chiefly seated in the back, from whence it shoots round the loins and down the thighs. It is then probably produced partly by pressure on the nerve-filaments, caused by contraction of the muscular fibres to which they are distributed, and partly by stretching and dilatation of the muscular tissue of the cervix. M. Beau believes that in this stage the pain is not produced, strictly speaking, in the uterus itself, but is rather a neuralgia of the lumbo-abdominal nerves. The pains at this time are generally described as “acute” and “grinding”—terms which sufficiently well express their nature. In highly-nervous women these pains are often much less well borne, than those of a later stage, and the suffering they undergo is indicated by their extreme restlessness and loud cries as each contraction supervenes. As the os dilates and the labor advances into the expulsive stage other sources of suffering are added.

In the Second Stage.—The presenting part now passes into the vagina and presses on the vaginal nerves, as well as on the large nervous plexuses lying in the pelvis. As it descends lower it stretches the perineum and vulva and presses on the bladder and rectum. Hence cramps are produced in the muscles supplied by the nerve-plexuses, as well as an intolerable sense of tearing and stretching in the vulva and perineum, and often a distressing feeling of tenesmus in the bowels. By this time the accessory muscles of parturition are brought into action, and they, as well as the uterine muscles, are thrown into frequent and violent contractions, which, independently of the other causes mentioned, are sufficient of themselves to produce great pain, likened to that of colic produced by involuntary and repeated contraction of the muscles of the intestines.

Taking all these causes into consideration, there is no lack of sufficient explanation of the intolerable suffering which is so constant an accompaniment of childbirth.

Effect of the Pains on the Mother and Fœtus.—The effect of the pains

on the mother's circulation is well marked. The rapidity of the pulse increases distinctly with each contraction, and as the pain passes off it again declines to its former state. A similar observation has been made with regard to the sounds of the foetal heart, especially after the expulsion of the liquor amnii. Hicks has pointed out that during a pain the muscular vibrations give rise to a sound which often resembles that of the foetal heart, and which completely disappears when the muscular tissue relaxes. The effect of the pain in intensifying the uterine souffle has been already mentioned. The strong muscular efforts would naturally lead us to expect a marked elevation of temperature during labor. Further observations on this point are required, but Squire asserts that there is generally only a very slight increase in temperature during delivery, rapidly passing off as soon as labor is over.

Division of Labor into Stages.—Such being the physiological facts in connection with the labor-pains, we may now describe the ordinary progress of a natural labor—that is, one terminated by the natural powers and with a head presenting.

For facility of description obstetricians have long been in the habit of dividing the course of labor into *stages*, which correspond pretty accurately with the natural sequence of events. For this purpose we generally talk of three stages—viz. : 1, from the commencement of regular pains until the complete dilatation of the cervix ; 2, from the complete dilatation of the cervix until the expulsion of the child ; 3, the concluding stage, comprising the permanent contraction of the uterus and the separation and expulsion of the placenta. To these we may conveniently add a preparatory stage, antecedent to the regular commencement of the labor.

Preparatory Stage.—For a short time before delivery, varying from a few days to a week or two, certain premonitory symptoms generally exist which indicate the approaching advent of labor. Sometimes they are well marked, and cannot be mistaken ; at others they are so slight as to escape observation. Amongst the most common is a sinking of the uterus into the pelvic cavity, resulting from the relaxation of the soft parts preceding delivery. The result is, that the upper edge of the uterine tumor is less high than before, and, in consequence, the pressure on the respiratory organs is diminished, and the woman often feels lighter, and altogether less unwieldy, than in the previous weeks. If a vaginal examination be made at this time, the lower segment of the uterus will be found to have sunk lower into the pelvic cavity ; and the consequence of this is that, while the respiration is less embarrassed and the patient feels less bulky, other accompaniments of pregnancy, such as hemorrhoids, irritability of the bladder and bowels, and cedema of the limbs, become aggravated. The increased pressure on the bowels often induces a sort of temporary diarrhœa, which is so far advantageous that it empties the bowels of feces which may have collected within them. As has already been pointed out, the contractions which have been going on at intervals during the latter months of pregnancy now get more and more marked, and they have the effect of producing a real shortening of the cervix, which is of great value preparatory to its dilatation. More marked mucous discharge from the cavity of the cervix also generally

occurs a short time before labor, and it is not unfrequently tinged with blood from the laceration of minute capillary vessels. This discharge, popularly known as the "*shows*," is a pretty sure sign that labor is not far off. It may, however, be entirely absent, even until the birth of the child. When copious it serves to lubricate the passages, and is generally coincident with rapid dilatation of the parts and a speedy labor.

False Pains.—During this time (premonitory stage) painful uterine contractions are often present, which, however, have no effect in dilating the cervix. In some cases they are frequent and severe, and are very apt to be mistaken for the commencement of real labor. Such "*false pains*," as they are termed, are often excited and kept up by local irritations, such as a loaded or disordered state of the intestinal canal; and they frequently give rise to considerable distress and much inconvenience both to the patient and practitioner. They are, it should be remembered, only the normal contractions of the uterus, intensified and accompanied with pain.

First Stage, or Dilatation.—As labor actually commences the uterine contractions become stronger, and the fact that they are "*true*" pains can be ascertained by their effect on the cervix. If a vaginal examination be made during one of these, the membranes will be felt to become tense and bulging during the pain, and the os uteri will be found partially dilated and thinned at its edges. As labor advances this effect on the os becomes more and more marked. At first the dilatation is very slight, perhaps not more than enough to admit the tip of the examining finger, and both the upper and lower orifices of the cervix can be made out. As the pains get stronger and more frequent, dilatation proceeds in the way already described, and the cervix gets more thin and tense, until we can feel a thin circular ring (which is lax between the pains, but becomes rigid and tense during the contraction, when the bag of waters bulges through it), without any distinction between the upper and lower orifices. During this time the patient, although she may be suffering acutely, is generally able to sit up and walk about. The amount of pain experienced varies much according to the character of the patient. In emotional women of highly-developed nervous susceptibilities it is generally very great. They are restless, irritable, and desponding, and when the pain comes on cry out loudly. The character of the cry is peculiar and well marked during the first stage, and has constantly been described by obstetric writers as characteristic. It is acute and high, and is certainly very different from the deep groans of the second stage, when the breath is involuntarily retained to assist the parturient effort. When dilatation is nearly completed various reflex nervous phenomena often show themselves. One of these is nausea and vomiting; another is uncontrollable shivering, which is not accompanied by a sense of coldness, the patient being often hot and perspiring. Both these symptoms indicate that the propulsive stage will shortly commence, and they may be regarded as favorable rather than otherwise, although they are apt to alarm the patient and her friends. By this time the os is fully dilated, the membranes generally rupture spontaneously, and a considerable portion of the liquor amnii flows away. The head, if presenting, often acts as a sort of ball-valve, and, falling down on the aperture of the cervix, pre-

LABOR.

the complete evacuation of the liquor amnii, which escapes by the vagina during the rest of the labor or may be retained in considerable quantity until the birth of the child.

It frequently happens, if the membranes are somewhat tougher than usual, the pains frequent and strong, that the foetus is pushed through the pelvis, and even expelled, surrounded by the membranes. When this occurs the child is said to be born with a "caul;" and this would doubtless happen more frequently than it does were it not the custom of the accoucheur to rupture the membranes artificially as soon as the os is completely opened up, after which time their integrity is no longer of any value.

Second Stage, or Propulsion.—The os is now entirely retracted over the presenting part, and is no longer to be felt, the vagina and the uterine cavity forming a single canal. Now the mucous discharge is generally abundant, so that the examining finger brings away long strings of glairy, transparent mucus, tinged with blood. The pains, after a short interval of rest, become entirely altered in character. The uterus contracts tightly round the foetus, the presenting part descends into the pelvis, and the true propulsive pains commence. The accessory muscles of parturition now come into play. With each pain the patient takes a deep inspiration, and thus fills the chest, so as to give a *point d'appui* to the abdominal muscles. For the same reason she involuntarily seizes hold of some point of support, as the hand of a bystander or a towel tied to the bed, and at the same time pushes with her feet against the end of the bed, and so is able to bear down to advantage. The cries are no longer sharp and loud, but consist of a series of deep suppressed groans, which correspond to a succession of short expirations made during the straining effort. In this way the abdominal muscles contract forcibly on the uterus, which they further stimulate to action by pressing upon it. It is to be observed that these straining efforts are, to a considerable extent, under the control of the patient. By encouraging her to hold her breath and bear down they can be intensified, while if we wish to lessen them we can advise her to call out, and when she does so the abdominal muscles have no longer a fixed point of action. Although the patient may thus lessen the effect of these accessory muscles, it is entirely out of her power to stop their action altogether. As labor advances the head descends lower and lower, receding somewhat in the intervals between the pains, until eventually it comes down on the perineum, which it soon distends.

Distension of the Perineum and Birth of the Child.—The pains now get stronger and more frequent, often with scarcely a perceptible interval between them, until the perineum gets stretched by the advancing head. In the interval between the pains the elasticity of the perineal structures pushes the head upward, so as to diminish the tension to which the perineum is subjected, the next pain again putting it on the stretch and protruding the head a little farther than before. By this alternate advance and recession the gradual yielding of the structures is favored and risk of laceration greatly diminished. During this time the pressure of the head mechanically empties the bowel of its contents. During the last pains, when the perineum is stretched to the utmost, the

anal aperture is dilated, sometimes to the size of a five-shilling piece; and in this way the perineum is relaxed, just as the distension, and consequent risk of laceration, are at their maximum. The apex of the head now protrudes more and more through the vulva, surrounded by the orifice of the vagina, and eventually it glides over the perineum and is expelled. The intensity of the suffering at this moment generally causes the patient to call out loudly. The force of the abdominal muscles is thus lessened at the last moment; and this, in combination with the relaxation of the sphincter ani, forms an admirable contrivance for lessening the risk of perineal injury. The rest of the body is generally expelled immediately by a single pain, and with it are discharged the remains of the liquor amnii and some blood-clots from separation of the placenta; and so the second stage of labor terminates.

The Third Stage: Its Importance.—The third stage commences after the expulsion of the child. It is of paramount importance to the safety of the mother that it should be conducted in a natural and efficient manner, for it is now that the uterine sinuses are closed; and the frail barrier by which nature effects this may be very readily interfered with, and serious and even fatal loss of blood ensue. Unfortunately, it is too often the case that the practitioner's entire attention is fixed on the expulsion of the child, so that the natural history of the rest of delivery is generally imperfectly studied and understood.


Contraction of the Uterus and Detachment of the Placenta.—As soon as the child is expelled the uterine fibres contract in all directions, and the hand, following the uterus down, will find that it forms a firm rounded mass lying in the lower part of the abdominal cavity. By retraction of its internal surface the placental attachments are generally separated, and the after-birth remains in the cavity of the uterus as a foreign body.

Mode in which Hemorrhage is Prevented.—The escape of blood from the open mouths of the uterine sinuses is now prevented in two ways—viz. (1) by the contractions of the uterine walls, and the more firm, persistent, and tonic this is the more certain is the immunity from hemorrhage; (2) by the formation of coagula in the mouths of the vessels. Any undue haste in promoting the expulsion of the placenta tends to prevent the latter of these two hæmostatic safeguards, and is apt to be followed by loss of blood. After a certain time, averaging from a quarter to half an hour, the uterus will be felt to harden, and, if the case be solely left to nature, what has been aptly called a miniature labor occurs. Pains come on, and the placenta is spontaneously expelled from the uterus, either into the canal of the vagina or even externally. In most obstetric works it is stated that the after-birth may be separated either from its centre or edge, and that it is very generally expelled through the os in an inverted form, with its fetal surface downward, and folded transversely on itself. That this is the mode in which the placenta is often expelled when traction on the cord is practised is a matter of certainty. It then passes through the os very much in the shape of an inverted umbrella. It is certain, however, that this is not the natural mechanism of its delivery. What this is has been well illustrated by Duncan,¹ who has very clearly shown that when this stage

¹ *Edin. Med. Journ.*, April, 1871.

of labor is left entirely to nature the separated placenta is expelled edge-ways, its uterine and detached surface gliding along the inner surface of the uterus, the foldings of its structure being parallel to the long diameter of the uterine cavity (Fig. 96). In this way it is expelled into the vagina, and during the process little or no hemorrhage occurs. When the placenta is drawn out in the way too generally practised, it obstructs the aperture of the os, and, acting like the piston of a pump, tends to promote hemorrhage. The corollaries as to treatment drawn from these facts will be subsequently considered.

FIG. 96.



Mode in which the Placenta is Naturally Expelled. (After Duncan.)

I am anxious, however, here to direct attention to Nature's mechanism, because I believe there is no part of labor about the management of which erroneous views are more prevalent than that of this stage, and none in which they are more apt to lead to serious consequences; and unless the mode in which Nature effects the expulsion of the placenta and prevents hemorrhage is thoroughly understood, we shall certainly fail in assisting her in a proper manner. In the large proportion of cases, when left entirely to themselves, the placenta would be retained, if not in the uterus, at any rate in the vagina, for a considerable time—possibly for several hours; and such delay would very unnecessarily tire the patience of the practitioner and be prejudicial to the patient. It is therefore our duty in the majority of cases to promote the expulsion of the after-birth; and when this is properly and scientifically done we increase rather than diminish the patient's safety and comfort. But in order to do this we must assist Nature, and not act in opposition to her method, as is so often the case.

After-Pains.—When once the placenta is expelled, the uterus contracts still more firmly, and in a typical case is felt just within the pelvic brim, hard and firm and about the size of a cricket-ball. Generally for several hours, or even for one or two days, it occasionally relaxes and contracts, and these contractions give rise to the "*after-pains*" from which women often suffer much. The object of these pains is, no doubt, to expel any coagula that may remain in the uterus; and therefore, however unpleasant they may be to the patient, they must be considered, unless very excessive, to be salutary rather than otherwise.

Duration of Labor.—The length of labor varies extremely in different cases, and it is quite impossible to lay down any definite rules with regard to it. Subject to exceptions, labor is longer in primiparæ than in multiparæ, on account of the greater resistance of the soft parts in the former, especially of the structures about the vagina and vulva. It is also generally stated that the difficulty of labor increases with the age of the patient, and that in elderly primiparæ it is likely to be unusually tedious from rigidity of the soft parts. It is very doubtful if this opinion has any real basis, and in such cases the practitioner often finds himself agreeably disappointed in the result. Mr. Roper,¹ indeed, argues

¹ *Obst. Trans.*, v. 7.

that the wasting of the tissues which occurs after forty years of age diminishes their resistance, and that first labors after that age are easier, as a rule, than in early life. The habits and mode of life of patients have no doubt a considerable influence on the duration of labor, but we are not in possession of any very reliable facts with regard to this subject. It is reasonable to suppose that the tissues of large, muscular, strongly-developed women will offer more resistance than those of slighter build. On the other hand, women of the latter class, especially in the upper ranks of life, more often develop nervous susceptibilities which may be expected to influence the length of their labors. The average duration of labor, calculated from a large number of cases, is from eight to ten hours; even in primipare, however, it is constantly terminated in one or two hours from its commencement, and may be extended to twenty-four hours without any symptoms of urgency arising. In multipare it is frequently over in even a shorter time. Indications calling for interference may arise at any time during the progress of labor independently of its length. The proportion between the length of the first and second stages also varies considerably. The first stage is generally the longest; and it is stated by Cazeaux to be normally about twice the length of the second. This is probably under the mark, and I believe Joulin to be nearer the truth in stating that the first stage should be to the second as four or five to one, rather than as two to one. Often when the first stage has been very prolonged the second is terminated rapidly.

Necessity of Caution in Expressing an Opinion as to the Possible Duration of Labor.—The practitioner is constantly asked as to the probable length of labor, and the uncertainty of this should always lead him to give a most guarded opinion. Even when labor is progressing apparently in the most satisfactory manner, the pains frequently die away and delivery may be delayed for many hours. In the first stage a cervix that is apparently rigid and unyielding may rapidly and unexpectedly dilate and delivery soon follow. In either case, if the practitioner has committed himself to a positive opinion, he is apt to incur blame, and it is far better always to be extremely cautious in our predictions on this point.

Period of the Day at which Labor Occurs.—A somewhat larger proportion of deliveries occur in the early hours of the morning than at other times. Thus, West¹ found that out of 2019 deliveries, 780 took place from 11 P. M. to 7 A. M., 662 from 7 A. M. to 3 P. M., and 577 from 3 P. M. to 11 P. M.

¹ *Amer. Med. Journ.*, 1854.

CHAPTER II.

MECHANISM OF DELIVERY IN HEAD PRESENTATIONS.

Importance of the Subject.—It is quite impossible to over-estimate the importance of thoroughly understanding the mechanism of the passage of the fœtus through the pelvis. This dominates the whole scientific practice of midwifery, and the practitioner cannot acquire more than a merely empirical knowledge, such as may be possessed by an uneducated midwife, or conduct the more difficult cases requiring operative interference with safety to the patient or satisfaction to himself, unless he thoroughly masters the subject.

In treating the physiological phenomena of labor it was assumed that we had to do with an ordinary case of head presentation, the description being applicable, with slight variations, to presentations of other parts of the fœtus. So in discussing the mechanical phenomena of delivery I shall describe more in detail the mechanism of head presentations, reserving any account of the mechanism of other presentations until they are separately studied.

Frequency of Head Presentations.—Head presentation is so much more frequent than that of any other part—amounting to 95 per cent. of all cases—that this mode of studying the subject is fully justified; and when once the student has mastered the phenomena of delivery in head presentations, he will have little difficulty in understanding the mechanism when other parts of the fœtus present, based, as it always is, on the same general plan.

Mode of Recognizing the Position of the Head by its Sutures and Fontanelles.—In entering on this study we come to appreciate the importance of the sutures and fontanelles in enabling us to detect the position of the fetal head and to watch its progress through the pelvis; and unless the “tactus eruditus” by which these can be distinguished from each other has been acquired, the practitioner will be unable to satisfy himself of the exact progress of the labor. Nor is this always easy. Indeed, it requires considerable experience and practice before it is possible to make out the position of the head with absolute certainty; but this knowledge should always be aimed at, and the student will never regret the time and trouble he spends in acquiring it.

Position of the Head at the Commencement of Labor.—At the commencement of labor the long diameter of the head lies in almost any diameter of the pelvic brim except in the antero-posterior, where there is not space for it. In the large majority of cases, however, it enters the pelvis in one or other of the oblique diameters, or in one between the oblique and transverse; but until it has fairly passed through the brim it more frequently lies directly in the transverse diameter than has been generally supposed. Hence obstetricians are in the habit of describing the head as lying in four positions, according to the parts of the pelvis

to which the occiput points; the first and third positions being those in which the long diameter of the head occupies the right oblique diameter of the pelvis, the second and fourth those in which it lies in the left oblique. Many subdivisions of these positions have been made, which only complicate the subject and render it more difficult to understand.

The positions, then, of the foetal head after it has entered the brim, which it is of importance to be able to distinguish in practice, are:

First (or left occipito-cotyloid).—The occiput points to the left foramen ovale, the sinciput to the right sacro-iliac synchondrosis, and the long diameter of the head lies in the right oblique diameter of the pelvis.

Second (or right occipito-cotyloid).—The occiput points to the right foramen ovale, the forehead to the left sacro-iliac synchondrosis, and the long diameter of the head lies in the left oblique diameter of the pelvis.

Third (or right occipito-sacro-iliac).—The occiput points to the right sacro-iliac synchondrosis, the forehead to the left foramen ovale, and the long diameter of the head lies in the right oblique diameter of the pelvis. This position is the reverse of the first.

Fourth (or left occipito-sacro-iliac).—The occiput points to the left sacro-iliac synchondrosis, the forehead to the right foramen ovale, and the long diameter of the head lies in the left oblique diameter of the pelvis. This position is the reverse of the second.

The Relative Frequency of these Positions.—The relative frequency of these positions has long been, and still is, a matter of discussion among obstetricians. According to Naegele, to whose classical essay we owe the greater part of our knowledge of the subject, the head lies in the right oblique diameter in 99 per cent. of all cases. More recent researches have thrown some doubt on the accuracy of these figures, and many modern obstetricians believe that the second position, which Naegele believed only to be observed as a transitional stage in the natural progress of the third position, is much more common than he supposed. This question will be more fully discussed when we treat of the mechanism of occipito-posterior delivery, and, in the mean time, it may serve to show the discrepancy which exists in the opinions of modern writers if we append the following table of the relative frequency of the various positions,¹ copied from Leishman's work:

	First Position.	Second Position.	Third Position.	Fourth Position.	Not Classified.
Naegele	70.	29.	1.
Naegele, Jr.	64.64	32.88	2.47
Simpson and Barry	76.45	.29	22.68	.58	
Dubois	70.83	2.87	25.66	.62	
Murphy	63.23	16.18	16.18	4.42	
Swayne	86.36	9.79	1.04	2.8	

Here it will be seen that all obstetricians are agreed as to the immensely greater frequency of the first position, the only point at issue being the relative frequency of the second and third.

¹ Leishman's *System of Midwifery*, p. 341.

CHAPTER II.

MECHANISM OF DELIVERY IN HEAD PRESENTATIONS.

Importance of the Subject.—It is quite impossible to over-estimate the importance of thoroughly understanding the mechanism of the passage of the fœtus through the pelvis. This dominates the whole scientific practice of midwifery, and the practitioner cannot acquire more than a merely empirical knowledge, such as may be possessed by an uneducated midwife, or conduct the more difficult cases requiring operative interference with safety to the patient or satisfaction to himself, unless he thoroughly masters the subject.

In treating the physiological phenomena of labor it was assumed that we had to do with an ordinary case of head presentation, the description being applicable, with slight variations, to presentations of other parts of the fœtus. So in discussing the mechanical phenomena of delivery I shall describe more in detail the mechanism of head presentations, reserving any account of the mechanism of other presentations until they are separately studied.

Frequency of Head Presentations.—Head presentation is so much more frequent than that of any other part—amounting to 95 per cent. of all cases—that this mode of studying the subject is fully justified; and when once the student has mastered the phenomena of delivery in head presentations, he will have little difficulty in understanding the mechanism when other parts of the fœtus present, based, as it always is, on the same general plan.


Mode of Recognizing the Position of the Head by its Sutures and Fontanelles.—In entering on this study we come to appreciate the importance of the sutures and fontanelles in enabling us to detect the position of the fœtal head and to watch its progress through the pelvis; and unless the “tactus eruditus” by which these can be distinguished from each other has been acquired, the practitioner will be unable to satisfy himself of the exact progress of the labor. Nor is this always easy. Indeed, it requires considerable experience and practice before it is possible to make out the position of the head with absolute certainty; but this knowledge should always be aimed at, and the student will never regret the time and trouble he spends in acquiring it.

Position of the Head at the Commencement of Labor.—At the commencement of labor the long diameter of the head lies in almost any diameter of the pelvic brim except in the antero-posterior, where there is not space for it. In the large majority of cases, however, it enters the pelvis in one or other of the oblique diameters, or in one between the oblique and transverse; but until it has fairly passed through the brim it more frequently lies directly in the transverse diameter than has been generally supposed. Hence obstetricians are in the habit of describing the head as lying in four positions, according to the parts of the pelvis

term which has received various definitions, the best of which is probably that adopted by Tyler Smith : viz. "that portion of the foetal head felt most prominently within the circle of the os uteri, the vagina, and the os tincæ in the successive stages of labor." If the tip of the examining finger be passed slightly upward, it will feel the sagittal suture running obliquely across the pelvis, and if this be traced downward and to the left, it will come upon the triangular posterior fontanelle, with the lambdoidal sutures diverging from it. If the finger could be passed sufficiently high in the opposite direction, upward and to the right, it would come upon the large anterior fontanelle, but at this time that is too high up to be within reach. The chin is slightly flexed upon the sternum, this flexion, as we shall presently see, being greatly increased as the head begins to descend.

The head, at the commencement of labor, generally lies within the pelvic brim, especially in primiparæ. In multiparæ, owing to the

FIG. 98.



First Position : Movement of Flexion.

relaxation of the abdominal parietes, the uterus is apt to fall somewhat forward, and the head consequently is more entirely above the brim, but is pushed within it as soon as labor actually commences.

Nægele's Views.—Nægele—and his description has been adopted by most subsequent writers—describes the head at this period as lying obliquely in relation to the brim, the right parietal bone, on which the examining finger impinges, being supposed by him to be much lower than the left. The accuracy of this view has of late years been contested, and it is now pretty generally admitted that this obliquity does not exist, and that the head enters the brim of the pelvis with both parietal bones on the same level, and with its bi-parietal diameter parallel to the plane of the inlet (Fig. 98). Nægele's view was adopted, partly because the finger always felt the right parietal protuberance lowest, and partly because it was at that point that the "*caput succedaneum*," or swelling observed on the head after delivery, was always

Explanation.—Various explanations have been given of the greater frequency with which the head lies in the right oblique diameter. By some it is referred to the natural tendency of the back of the foetus, as shown by the experimental researches of Höning and other writers, to be directed, in consequence of gravitation, forward and to the left side of the mother in the erect attitude, and backward and to her right side in the recumbent. The explanation given by Simpson was that the head lay in the right oblique diameter in consequence of the measurement of the left oblique being more or less lessened by the presence of the rectum. When the rectum is collapsed, indeed, the narrowing of the diameter is slight; but it is so often distended by fecal matter—sometimes, when constipation exists, to a very great extent—that it may really have a very important influence in determining the position of the fetal head.

In describing the mechanism of delivery it will be well for us to concentrate our attention on the first or most common position, dwelling subsequently more briefly on the differences between it and the less common ones.

Description of the First Position.—In this position, when the head commences to descend, the occiput lies in the brim pointing to the left ileo-pectineal eminence, the forehead is directed to the right sacro-iliac synchondrosis, and the sagittal suture runs obliquely across the pelvis in the right-oblique diameter. The back of the child is turning toward

FIG. 97.



Attitude of Child in First Position. (After Hodge.)

the left side of the mother's abdomen, the right shoulder to her right side, the left to her left side (Fig. 97). If a vaginal examination be (the patient lying in the ordinary obstetric position), and the ^{lv} open, the finger will impinge upon the protuberance of bone, which is described as the "presenting part"—a

and the chin is no longer so much flexed on the sternum. This change is due to the fact that the anterior end of the ovoid experiences greater resistance than the posterior, and as soon as this resistance counterbalances and exceeds that applied to the latter, the sinciput must descend. The right side of the head also descends more than the left from a similar cause, so that the head becomes, as it were, slightly flexed on the right shoulder. This obliquity of the head on its transverse diameter

FIG. 99.



First Position: Occiput in the Cavity of the Pelvis. (After Hodge.)

in the lower part of the pelvis has been denied by Küncke,¹ who maintains that the head passes through the entire pelvis in the same position as it enters the brim—that is, with both parietal bones on a level—so that the point of intersection of the transverse and antero-posterior diameters of the pelvis would correspond with the sagittal suture. There is, however, good reason to believe that in the lower half of the pelvic cavity the head is not truly syncletic, as Küncke describes, but that the right parietal bone is on a somewhat lower level than the left.

4. *Rotation*.—The movement of *rotation* is very important. By it the long diameter of the head is changed from the oblique diameter of the pelvic cavity to the antero-posterior diameter of the outlet (Fig. 100),

FIG. 100.



First Position: Occiput at Outlet of the Pelvis. (After Hodge.)

or to a diameter nearly corresponding to it, so that the long diameter of the head is brought into relation with the longest diameter of the pelvic outlet. This alteration almost always takes place, and may be readily observed by the accoucheur who carefully watches the progress of labor. Various explanations have been given of its causes. The one most generally adopted is, that it is due to the projection inward of the ischial spines, which narrow the transverse diameter of the pelvic outlet. As

¹ *Die Vier Factoren der Geburt*, Berlin, 1869.

the pains force the occiput downward, its rotation backward is prevented by the projection of the left ischial spine, while its rotation forward is favored by the smooth bevelled surface of the ascending ramus of the ischium. Similarly, the ischial spine on the opposite side prevents the rotation forward of the forehead, which is guided backward to the cavity of the sacrum by the smooth surface of the sacro-ischiatic ligaments. These arrangements, therefore, give a screw-like form to the interior of the pelvis; and as the pains force the head downward, they are effectual in imparting to it the rotatory movement which is of such importance in adapting it to the longest measurement of the outlet.

By most of the German obstetricians the influence of the ischial spines and of the smooth pelvic planes in producing rotation is not admitted. They rather refer the change of direction to the increased resistance the head meets from the posterior wall of the pelvis and from the perineal structures. Whichever part of the head first meets this resistance, which is much greater than that of the interior part of the pelvis, must necessarily be pressed forward; and as, in the large majority of cases, the posterior fontanelle descends first, it is thus pressed forward until rotation is effected. This view has the advantage of accounting equally well for the rotation in occipito-posterior as in occipito-anterior positions, the former of which, on the more ordinarily received theory, are not quite satisfactorily explicable. It does not follow that the smooth surfaces of the pelvic planes are without influence in favoring the rotation—on the contrary, they probably greatly facilitate it—but it is more simply and effectually explained by the latter theory than by that which attributes so important an action to the ischial spines.

In some rare cases the head escapes rotation and reaches the perineum still lying in the oblique diameter. Even here, however, rotation is generally effected, often suddenly, just as the head is about to pass the vulva, and it is very rarely expelled in the oblique position. The movement at this stage may be explained by the perineum, which is attached at its sides and grooved in its centre: to the hollow so formed the long diameter of the head accommodates itself, and is thus rotated into the antero-posterior diameter of the outlet.

5. *Extension*.—By the process just described the face is turned back into the hollow of the sacrum; but the head does not lie absolutely in the antero-posterior diameter of the pelvic outlet, but rather in one between it and the oblique. The occiput is still forced down by the pains, and, in consequence of its altered position, is enabled to pass between the rami of the pubes, and advances until its further descent is checked by the nape of the neck, which is pressed under and against the arch of the pubes. By this means the occiput is fixed, and, the pains continuing, the uterine force no longer acts on the occiput, but on the anterior part of the head, which is now pushed down and separated from the sternum. This constitutes *extension*. As the head descends the soft structures of the perineum are stretched and the coccyx pushed back so as to enlarge the outlet. The pains continue to distend the perineum more and more, the head advancing and receding with each pain. As the forehead descends, the sub-occipito-bregmatic, the sub-occipito-frontal, and the sub-occipito-mental diameters successively present; the

occiput turns more and more upward in front of the pubes (Fig. 101), and at last the face sweeps over the perineum and is born.

The mechanical cause of this movement may be readily explained. As soon as the occiput has passed under the arch of the pubis, and is

FIG. 101.



First Position : Head Delivered. (After Hodge.)

no longer resisted by the anterior pelvic walls, the head is subjected to the action of two forces—that of the uterine pressure, acting downward and backward, and that of the resistance of the posterior walls of the pelvis and the soft parts, acting almost directly forward. The necessary result is that the head is pushed in a direction intermediate between these two opposing forces—that is, downward and forward in the axis of the pelvic outlet.

In addition to the slight obliquity which exists as regards the direct relation of the long diameter of the head to the antero-posterior diameter of the outlet at the moment of its expulsion, the head also lies somewhat obliquely in relation to its own transverse diameter, so that, in the majority of cases, the right parietal bone is expelled before the left.

FIG. 102.



External Rotation of Head in First Position. (After Hodge.)

6. *External Rotation.*—Shortly after the head is expelled, as soon as renewed uterine action commences, it may be observed to make a distinct rotatory movement, the occiput turning to the left thigh of the mother, and the face turning upward to the right thigh (Fig. 102). The

the pains force the occiput downward, its rotation backward is prevented by the projection of the left ischial spine, while its rotation forward is favored by the smooth bevelled surface of the ascending ramus of the ischium. Similarly, the ischial spine on the opposite side prevents the rotation forward of the forehead, which is guided backward to the cavity of the sacrum by the smooth surface of the sacro-ischiatic ligaments. These arrangements, therefore, give a screw-like form to the interior of the pelvis; and as the pains force the head downward, they are effectual in imparting to it the rotatory movement which is of such importance in adapting it to the longest measurement of the outlet.

By most of the German obstetricians the influence of the ischial spines and of the smooth pelvic planes in producing rotation is not admitted. They rather refer the change of direction to the increased resistance the head meets from the posterior wall of the pelvis and from the perineal structures. Whichever part of the head first meets this resistance, which is much greater than that of the interior part of the pelvis, must necessarily be pressed forward; and as, in the large majority of cases, the posterior fontanelle descends first, it is thus pressed forward until rotation is effected. This view has the advantage of accounting equally well for the rotation in occipito-posterior as in occipito-anterior positions, the former of which, on the more ordinarily received theory, are not quite satisfactorily explicable. It does not follow that the smooth surfaces of the pelvic planes are without influence in favoring the rotation—on the contrary, they probably greatly facilitate it—but it is more simply and effectually explained by the latter theory than by that which attributes so important an action to the ischial spines.

In some rare cases the head escapes rotation and reaches the perineum still lying in the oblique diameter. Even here, however, rotation is generally effected, often suddenly, just as the head is about to pass the vulva, and it is very rarely expelled in the oblique position. The movement at this stage may be explained by the perineum, which is attached at its sides and grooved in its centre: to the hollow so formed the long diameter of the head accommodates itself, and is thus rotated into the antero-posterior diameter of the outlet.

5. *Extension*.—By the process just described the face is turned back into the hollow of the sacrum; but the head does not lie absolutely in the antero-posterior diameter of the pelvic outlet, but rather in one between it and the oblique. The occiput is still forced down by the pains, and, in consequence of its altered position, is enabled to pass between the rami of the pubes, and advances until its further descent is checked by the nape of the neck, which is pressed under and against the arch of the pubes. By this means the occiput is fixed, and, the pains continuing, the uterine force no longer acts on the occiput, but on the anterior part of the head, which is now pushed down and separated from the sternum. This constitutes *extension*. As the head descends the soft structures of the perineum are stretched and the coccyx pushed back so as to enlarge the outlet. The pains continue to distend the perineum more and more, the head advancing and receding with each pain. As the forehead descends, the sub-occipito-bregmatic, the sub-occipito-frontal, and the sub-occipito-mental diameters successively present; the

ward, the anterior fontanelle forward, while the examining finger impinges on the left parietal bone. The mechanism of delivery in these cases is of much interest. In the large majority of cases, during the progress of delivery the occiput rotates forward along the right side of the pelvis until it comes to lie almost in the antero-posterior diameter of the outlet and passes under the pubic arch, the forehead passing over the perineum. It will be seen that during part of this extensive rotation the head must lie in the second position, and the case terminates just as if it had been in the second position from the commencement of labor.

Manner in which the Occiput is Rotated Forward.—How is it that this rotation is effected, and that the sinciput, occupying the position of the occiput in the first position, should not be rotated forward to the pubes as that is? This, no doubt, may be explained by the fact that the uterine force transmitted through the vertebral column causes the occiput to descend lower than the sinciput, so that in most cases, in making a vaginal examination, the posterior fontanelle can be readily felt, while the anterior is high up and out of reach. The head is therefore extremely flexed, and so descends into the pelvic cavity, until the occiput, being now below the right ischial spine, experiences the resistance of the pelvic floor opposite the right sacro-ischiatic ligament, by which it is directed forward. The forehead is at this time, supposing flexion to be marked, too high to be influenced by the anterior pelvic plane. Pressure continuing, the occiput rotates forward, the forehead passes round the left side of the pelvis, and labor is terminated as in the second position.


The period of labor at which rotation takes place varies. In the majority of cases it does not occur until the head is on the floor of the pelvis, for it is then that resistance is most felt; but the greater the resistance, the sooner will rotation be produced. Hence it is more likely to occur early when the head is large and the pelvis comparatively small.

The facility with which this movement is effected obviously depends upon the complete flexion of the chin on the sternum, by which the anterior fontanelle is so elevated that its rotation backward is not resisted by the inward projection of the left ischial spine, and the occiput is correspondingly depressed. If, however, this flexion is not complete, and the anterior fontanelle is so low as to be readily within reach of the finger, considerable difficulty is likely to be experienced. In many such cases rotation is still eventually effected, but in others it is not; and the labor is then terminated with the face to the pubes, but at the expense of considerable delay and difficulty. According to Dr. Uvedale West of Alford, who devoted much careful study to the subject, this termination occurs in about 4 per cent. of occipito-posterior positions. When it is about to happen the anterior fontanelle may be felt very low down, and sometimes even the forehead and superciliary ridges. The uterine force pushes down the occiput, the sinciput being fixed behind the pubes, which it obviously cannot pass under, as does the occiput in the first position. The sinciput therefore becomes more flexed and pushed upward, while the resistance of the pelvic floor directs the occiput forward. The perineum now becomes enormously distended by the back part of the head, and is in great danger of laceration. The occiput is eventually, but not without much difficulty, expelled. A process

reason of this is evident. When the head descends in the right oblique diameter, the shoulders lie in the opposite or left oblique diameter, and as the head rotates into the antero-posterior diameter they are necessarily placed more nearly in the transverse. As soon as the head is expelled the shoulders are subjected to the same uterine force and pelvic resistance as the head has just been, and they are acted on in precisely the same way. Consequently they too rotate, but in the opposite direction—into the antero-posterior diameter of the outlet, or nearly so, just as the head did—and as they do so they necessarily carry the head with them and cause its external rotation.

The two shoulders are soon expelled, the left shoulder generally the first, sweeping over the perineum in the same manner as the face. This is, however, not always the case, and they are often expelled simulta-

FIG. 103.



Third Position of Occiput at Brim of Pelvis.

neously, or the right shoulder may come first. The body soon follows; and the second stage of labor is completed.

Second Position.—In the second position (right occipito-cotyloid) the long diameter of the head lies in the left oblique diameter of the pelvis. On making a vaginal examination in the ordinary obstetric position, the finger, passing upward and to the right, feels the small posterior fontanelle; downward and to the left, it feels the anterior. The sagittal suture lies obliquely across the pelvis in the left oblique diameter. The description of the mechanism of delivery is precisely the same as in the first position, substituting the word left for right. Thus the finger impinges on the left parietal bone; the occiput turns from right to left during rotation. After the birth of the head the occiput turns to the right thigh of the mother, the face to the left thigh.

Third or Right Occipito-sacro-iliac Position.—In the third position the head enters the pelvic brim with the occiput directed backward to the right sacro-iliac synchondrosis, and the sinciput forward to the left foramen ovale (Fig. 103). The posterior fontanelle is directed back-

Formation of the Caput Succedaneum.—The formation of the caput succedaneum has been already alluded to. This term is applied to the œdematous swelling which forms on the head, and is produced by effusion from the obstruction of the venous circulation caused by the pressure to which the head is subjected. It follows that the size of the swelling is in direct proportion to the length of the labor. In rapid deliveries, in which the head is forced through the pelvis quickly, it is scarcely, if at all, developed, while after protracted labors it is large and distinct, and may obscure the diagnosis of the position by preventing the sutures and fontanelles being felt. Its situation varies according to the position of the head; thus, in the first and fourth positions it forms on the right parietal bone, in the second and third on the left; and we may therefore verify, by inspection of its site, the accuracy of our diagnosis.

An ordinary mistake which has been made by obstetricians is to regard the caput succedaneum as formed at the point where the head has been most subjected to pressure; while, in fact, it forms on that part which is most unsupported by the maternal structures, and where the swelling may consequently most readily occur. Therefore, in the early stages of the labor it always forms on the part of the head which lies in the circle of the os uteri; while in subsequent stages it forms on that which lies in the axis of the vaginal canal, and eventually is most prominent on the part that is first expelled from the vulva.

Alteration in the Shape of the Head from Moulding.—A few words may be said as to the alteration in the form of the fetal head which occurs in tedious labors, and results from the moulding which it has undergone in its passage through the pelvis. The smaller the pelvis and the greater the pressure applied to the head during delivery, the more marked this is. The result is, that in vertex presentations the occipito-mental and occipito-frontal diameters are elongated to the extent of an inch or even more, while the transverse diameters are lessened, from compression of the parietal bones. This moulding is of unquestionable value in facilitating the birth of the child. The amount of apparent deformity is very considerable, and may even give rise to some anxiety. It is well to remember, therefore, that it is always transient, and that in a few hours, or days at most, the elasticity of the soft cranial bones causes them to resume their natural form. The caput succedaneum also disappears rapidly; therefore, no amount of deformity from either of these causes need give rise to anxiety or call for any treatment.

of extension now occurs, the nape of the neck being fixed, as it were, against the centre of the perineum, the expelling force now acting on the forehead and producing rotation of the head on its transverse axis. The forehead and face are thus protruded, and the body follows without difficulty.

It is said that in a few exceptional cases, where the anterior fontanelle is much depressed, the labor may terminate by the conversion of the presentation into one of the face, the head rotating on its transverse axis, the forehead passing to the posterior part of the pelvis, and the chin emerging under the perineum. It is obvious, however, that this change can only occur when the head is unusually small, and it must of necessity be extremely rare.

Relative Frequency of Second and Third Positions.—Reference has already been made to Naegele's views as to the rarity of the second position, and to his opinion that cases in which the occiput was found to point to the right foramen ovale were only transitional stages in the rotation of occipito-posterior positions. Such an assumption, however, is unwarrantable unless the case has been watched from the very commencement of labor. Many perfectly-qualified observers have arrived at the conclusion that second positions are far more common than Naegele supposed; and in the table already quoted it will be seen that while Murphy estimates the second and third as being equally frequent, Swayne believes the second to be much more common than the third. It is probable that the weight of Naegele's authority has induced many observers to classify second positions as third positions in which partial rotation has already been accomplished. My own experience would certainly lead me to think that second positions are very far from uncommon. The question, however, must be considered to be in abeyance

FIG. 104.



Fourth Position of Occiput at Pelvic Brim.

until further observations by competent authorities enable us to decide conclusively.

Fourth, or Left Occipito-sacro-iliac.—The fourth position is just as much the reverse of the second as the third is of the first. The occiput points to the left (Fig. 104) sacro-iliac synchondrosis, and the finger impinges on the right parietal bone. The mechanism is precisely the same as in the third position, the rotation taking place from left to right.

Failing this, it is better to avoid the use of the corset altogether, and to have as little pressure on the uterus as possible, although many women cannot do without the support to which they are accustomed. To multiparæ, especially if there be much laxity of the abdominal parietes, a well-fitting elastic abdominal belt is often a great comfort. This is constructed so that it can be tightened when the patient is walking and in the erect position, when such support is most required, and readily loosened when desired.

Necessity of Attending to the First Summons.—It is hardly necessary to insist on the necessity of the practitioner attending immediately to the first summons to the patient. It is true that he may very often be sent for long before he is actually required. But, on the other hand, it is quite impossible to foresee what may be the state of any individual case. By prompt attention he may be able to rectify a malposition or prevent some impending catastrophe, and thus save his patient from consequences of the utmost gravity.

Articles to be Taken by the Accoucheur.—The practitioner should always be provided with the articles which he may require. The ordinary obstetric cases, containing one or two bottles and a catheter, such as are sold by most instrument-makers, are cumbrous and useless, while "obstetric bags" are expensive luxuries not within the reach of all. Every one can manufacture an excellent obstetric bag for himself, at a small expense, by having compartments for holding bottles stitched on to the sides of an ordinary leather bag, such as is sold for a few shillings at any portmanteau-maker's. It is a great comfort to have at hand all that may be required, and the bag should contain chloroform or other anæsthetic, chloral, laudanum, the liquor ferri perchloridi of the Pharmacopœia, the liquid extract of ergot, and a hypodermic syringe, with bottles containing ether and a solution of ergotin for subcutaneous injection. If it also contain a Higginson's syringe, a small elastic catheter, a good pair of forceps, and one or two suture-needles, with some silver wire or carbolized catgut, the practitioner is provided against any ordinary contingency. Other articles that may be required, such as thread, scissors, and the like, are generally provided by the nurse or patient.

Duties on First Visiting the Patient.—On arriving at the house the practitioner should have his visit announced to the patient; and he will very often find that the first effect of his presence is to arrest the pains that have been hitherto progressing rapidly, thereby affording a very conclusive proof of the influence of mental impressions on the progress of labor. If the pains be not already propulsive, it is well that he should occupy himself at first in general inquiries from the attendants as to the progress of the labor, and in seeing that all the necessary arrangements are satisfactorily carried out, so as to allow the patient time to get accustomed to his presence. If he have any choice in the matter, he should endeavor to secure a large, airy, and well-ventilated apartment for the lying-in room, as far removed as possible from without. He may also see to the bed, which should be without curtains and prepared for the labor by having a water-proof sheeting laid under a folded blanket or sheet, on which the patient lies. These

CHAPTER III.

MANAGEMENT OF NATURAL LABOR.

ALTHOUGH labor is a strictly physiological function, and in a large majority of cases might, no doubt, be safely accomplished without assistance from the accoucheur, still, medical aid, properly given, is always of value in facilitating the process, and is often absolutely essential for the safety of the mother and child.

Preparatory Treatment.—The management of the pregnant woman before delivery is a point which should always receive the attention of the medical attendant, since it is of consequence that the labor should come on when she is in as good a state of health as possible. For this purpose ordinary hygienic precautions should never be neglected in the latter months of gestation. The patient should take regular and gentle exercise, short of fatigue, and, if the weather permit, should spend as much of her time as possible in the open air. Hot rooms, late hours, and excitement of all kinds should be strictly avoided. The diet should be simple, nutritious, and unstimulating. The state of the bowels should be particularly attended to. During the few days preceding labor the descent of the uterus often causes pressure on the rectum and prevents its evacuation. Hence it is customary to prescribe occasional gentle aperients, such as small doses of castor oil, for a few days before the expected period of delivery. Some caution, however, is necessary, as it is certainly not very uncommon for labor to be determined rather sooner than was anticipated in consequence of the irritation of too large a purgative dose. The state of the bowels should always be inquired into at the commencement of labor, and if there be any reason to suspect that they are loaded a copious enema should be administered. This is always a proper precaution to take, for a loaded rectum is a common cause of irregular and ineffective uterine action; and even when it does not produce this result, the escape of the feces, in consequence of pressure on the bowel during the propulsive stage, is always disagreeable both to the patient and practitioner.

Dress of Patient during Pregnancy.—The dress of the patient during pregnancy may be here adverted to, for much discomfort may arise, and the satisfactory progress of labor may even be interfered with, from errors in this respect.

After the uterus has risen out of the pelvis the ordinary corset which most women wear is apt to produce very injurious pressure; still more so when attempts are made to conceal the increased size by tight-lacing. After the fourth or fifth month, therefore, the comfort of the patient is much increased by wearing a specially-constructed pair of stays with elastic let into the sides and front, so that they accommodate themselves to the gradual increase of the figure. Such are made by all stay-makers, and should be worn whenever the circumstances of the patient permit.

at first slight and short, and gradually recur with increased force and regularity. The situation of the two kinds of pains also varies, the false pains being chiefly situated in front, while the true pains are felt most in the back and gradually shoot round toward the abdomen. Nothing short of a vaginal examination will enable us to clear up the diagnosis satisfactorily. If the labor have actually commenced, the os will be more or less dilated and its edges thinned, while with each pain the cervix will become rigid and the membranes tense and prominent. The false pains, on the contrary, have no effect on the cervix, which remains flaccid and undilated, or, if the os be sufficiently open to admit the tip of the finger, the membranes will not become prominent during the contraction. Under such circumstances we may confidently assure the patient that the pains are false, and measures should be taken to remove the irritation which produces them. In the large majority of cases the cause of the spurious pains will be found to be some disordered state of the intestinal tract; and they will be best remedied by a gentle aperient—such as castor oil or the compound colocynth pill with hyoscyamus—followed by, or combined with, a sedative, such as twenty minims of laudanum or chlorodyne. Shortly after this has been administered the false pains will die away, and not recur until true labor commences.

Mode of Conducting a Vaginal Examination.—For a vaginal examination the patient is placed by the nurse on her left side, close to the edge of the bed, with the legs flexed on the abdomen. The practitioner, being seated by the edge of the bed, passes the index finger of the right hand, previously lubricated with carbolized oil or cold cream, up to the vulva, and gently insinuates it into the orifice of the vagina, then pushes it backward in the axis of the vaginal outlet, and finally turns it upward and forward so as to more readily reach the cervix. This it may not always be easy to do, for at the commencement of labor the cervix may be so high as to be reached with difficulty, or it may be directed backward so as to point toward the cavity of the sacrum. The exploration is often much facilitated by depressing the uterus from without by the left hand placed on the abdomen. Our object is not only to ascertain the state of the cervix as to softness and dilatation, but also the presentation, the condition of the vagina, and the capacity of the pelvis. The examination is generally commenced during a pain, at which time it is less depressing to the patient; but in order to be satisfactory the finger must remain in the vagina until the pain is over, the examination being concluded in the interval between this pain and the next.

In head presentations the round mass of the cranium is generally at once felt through the lower part of the uterus, and then we have the satisfaction of being able to assure the patient that all is right. If the os be sufficiently dilated, we can also feel through it the occiput covered by the membranes. It is impossible at this time to make out the exact position of the head by means of the sutures and fontanelles, which are too high up to be within reach. Nor should any attempt be made to do so, for fear of prematurely rupturing the membranes. The fact that the head is presenting is all that we require to know at this stage of the labor.

The Condition of the Os as Indicating the Progress of Labor.—The

condition of the os itself, as to rigidity and dilatation, will materially assist us in forming an opinion as to the progress and probable duration of the labor; but, although the friends will certainly press for an opinion on this point, the cautious practitioner will be careful not to commit himself to a positive statement, which may so easily be falsified. It will suffice to assure the friends that everything is satisfactory, but that it is impossible to say with any certainty how rapidly, or the reverse, the case may progress.

If the pains be not very frequent or strong, and the os not dilated to more than the size of a shilling, a considerable delay may be anticipated, and the presence of the medical attendant is useless. He may, therefore, safely leave the patient for an hour or more, provided he be within easy reach. It is needless to say that this should never be done unless the exact presentation be made out. If some part other than the head be presenting, it will probably be impossible to make it out until dilatation has progressed further; and the practitioner must be incessantly on the watch until the nature of the case be made out, so as to be able to seize the most favorable moment for interference, should that be necessary.

Position of Patient during First Stage.—The position of the patient is a matter of some moment in the first stage. It is a decided advantage that she should not be then in a recumbent position on her side, as is usual in the second stage; for it is of importance that the expulsive force should act in such a way as to favor the descent of the head into the pelvis—i. e. perpendicularly to the plane of its brim—and also that the weight of the child should operate in the same way. Therefore, the ordinary custom of allowing the patient to walk about or to recline in a chair is decidedly advantageous; and it will often be observed that the pains are more lingering and ineffective if she lie in bed. If the patient be a multipara, or if the abdomen be somewhat pendulous, an abdominal bandage, by supporting the uterus, will greatly favor the progress of this stage. Keeping the patient out of bed has the further advantage of preventing her being unduly anxious for the termination of the labor; and a little cheerful conversation will keep up her spirits and obviate the mental depression which is so common. Good beef-tea may be freely administered, with a little brandy and water occasionally if the patient be weak, and will be useful in supporting her strength.

Vaginal Examinations.—Over-frequent vaginal examinations at this period should be avoided, for they serve no useful purpose and are apt to irritate the cervix. It will be necessary, however, to ascertain the progress of the dilatation at intervals.

Artificial Rupture of the Membranes.—When once the os is fully dilated, the membranes may be artificially ruptured if they have not broken spontaneously, for they no longer serve any useful purpose, and only retard the advent of the propulsive stage. This can be easily done by pressing on them, when they are rendered tense during a pain, by some pointed instrument, such as the end of a hairpin, which is always at hand. In some cases, indeed, it is even expedient to rupture the membranes before the os is fully dilated. Thus, it not unfrequently happens, when the amount of liquor amnii is at all excessive, that the os dilates to the size of a five-shilling piece or more; but, although it is perfectly

soft and flaccid, it opens up no farther until the liquor amnii is evacuated, when the propulsive pains rapidly complete its dilatation. Some experience and judgment is required in the detection of such cases, for if we evacuate the liquor amnii prematurely the pressure of the head on the cervix may produce irritation and seriously prolong the labor. This manœuvre is most likely to be useful when the pains are strong and the os perfectly flaccid, and when the membranes do not protrude through the os and effect further dilatation.

It is sometimes not easy to ascertain whether the membranes are ruptured or not. This is most likely to be the case when the head is low down and the amount of liquor amnii is so small that the pouch does not become prominent during the pains. A little care, however, will enable us, if the membranes be ruptured, to feel the rugosities of the scalp covered with hair, and to distinguish it from the smooth polished surface of the membranes.

Treatment of the Propulsive Stage.—After the evacuation of the liquor amnii there is generally a lull in the progress of the labor, the pains, however, soon recurring with increased force and frequency, and propelling the head through the pelvic cavity. The change in the character of the pains is soon appreciated by the bearing-down efforts by which they are accompanied, as well as by their increased length and intensity.

Position of the Patient during the Second Stage.—It is now advisable that the patient be placed in bed; and in this country it is usual for her to lie on her left side, with her nates parallel to the edge of the bed and her body lying across it. This is the established obstetric position in England, and it would be useless to attempt to insist on any other, even if it were advisable. Although the dorsal position is preferred on the Continent, it is difficult to see wherein its advantages consist. It certainly leads to unnecessary exposure of the person, and it is, on the whole, less easy to reach the patient, so placed, for the necessary manipulations. Moreover, the dorsal position increases the risk of laceration of the perineum, by bringing the weight of the child's head to bear more directly upon it. Thus, Schroeder found that lacerations occurred in 37.6 per cent. of cases delivered on the back, as against 24.4 per cent. in other positions.

The patient usually remains in bed during the whole of this stage, and it is customary for the nurse to tie to the foot of the bed a jack-towel, which is laid hold of and used as a support in making bearing-down efforts. If the pains be few and far between, and the patient finds it more comfortable to get up occasionally, there is no reason why she should not do so. On the contrary, as we shall subsequently see in treating of lingering labor, the pains under such circumstances are often increased in the sitting posture, in consequence of the weight of the child producing increased pressure on the nerves of the vagina.

Detection of the Position of Head.—At this time vaginal examination, which should be more frequently repeated than in the first stage, enables us to ascertain precisely the position of the head by means of the sutures and fontanelles, as well as to watch its progress.

Management of the Anterior Lip of Cervix when Impacted between the Head and Pelvis.—It not unfrequently happens that the head descends

condition of the os itself, as to rigidity and dilatation, will materially assist us in forming an opinion as to the progress and probable duration of the labor; but, although the friends will certainly press for an opinion on this point, the cautious practitioner will be careful not to commit himself to a positive statement, which may so easily be falsified. It will suffice to assure the friends that everything is satisfactory, but that it is impossible to say with any certainty how rapidly, or the reverse, the case may progress.

If the pains be not very frequent or strong, and the os not dilated to more than the size of a shilling, a considerable delay may be anticipated, and the presence of the medical attendant is useless. He may, therefore, safely leave the patient for an hour or more, provided he be within easy reach. It is needless to say that this should never be done unless the exact presentation be made out. If some part other than the head be presenting, it will probably be impossible to make it out until dilatation has progressed further; and the practitioner must be incessantly on the watch until the nature of the case be made out, so as to be able to seize the most favorable moment for interference, should that be necessary.

Position of Patient during First Stage.—The position of the patient is a matter of some moment in the first stage. It is a decided advantage that she should not be then in a recumbent position on her side, as is usual in the second stage; for it is of importance that the expulsive force should act in such a way as to favor the descent of the head into the pelvis—*i. e.* perpendicularly to the plane of its brim—and also that the weight of the child should operate in the same way. Therefore, the ordinary custom of allowing the patient to walk about or to recline in a chair is decidedly advantageous; and it will often be observed that the pains are more lingering and ineffective if she lie in bed. If the patient be a multipara, or if the abdomen be somewhat pendulous, an abdominal bandage, by supporting the uterus, will greatly favor the progress of this stage. Keeping the patient out of bed has the further advantage of preventing her being unduly anxious for the termination of the labor; and a little cheerful conversation will keep up her spirits and obviate the mental depression which is so common. Good beef-tea may be freely administered, with a little brandy and water occasionally if the patient be weak, and will be useful in supporting her strength.

Vaginal Examinations.—Over-frequent vaginal examinations at this period should be avoided, for they serve no useful purpose and are apt to irritate the cervix. It will be necessary, however, to ascertain the progress of the dilatation at intervals.

Artificial Rupture of the Membranes.—When once the os is fully dilated, the membranes may be artificially ruptured if they have not broken spontaneously, for they no longer serve any useful purpose, and only retard the advent of the propulsive stage. This can be easily done by pressing on them, when they are rendered tense during a pain, by some pointed instrument, such as the end of a hairpin, which is always at hand. In some cases, indeed, it is even expedient to rupture the membranes before the os is fully dilated. Thus, it not unfrequently happens, when the amount of liquor amnii is at all excessive, that the os dilates to the size of a five-shilling piece or more; but, although it is perfectly

favor, laceration, in consequence of the pressure producing increased uterine action just at the time when forcible distension of the perineum is likely to be hurtful. Therefore, some hold that the perineum ought to be left entirely alone, and that the head should be allowed gradually to distend it, without any assistance on the part of the practitioner.

Much error may be traced to a misconception of what is required. The term "supporting the perineum" conveys an unquestionably erroneous idea, and it is certain that no one can prevent laceration by mechanical support. If the term "relaxation of the perineum" was employed, we should have a far more accurate idea of what should be aimed at, and if this be borne in mind I think it cannot be questioned that nature may be most usefully assisted at this stage.

Dr. Goodell's Method.—Dr. Goodell of Philadelphia has specially studied this subject, and has recommended a method the object of which is to relax the perineum. His advice is, that one or two fingers of the left hand should be inserted into the rectum, by which the perineum should be hooked up and pulled forward over the head, toward the pubes, the thumb of the same hand being placed on the advancing head, so as to restrain its progress if needful. I have adopted this plan frequently, and believe that it admirably answers its purpose, especially when the perineum is greatly distended and laceration is threatened. It must be admitted that the insertion of the fingers into the anal orifice, in the manner recommended, is repugnant both to the practitioner and patient, and the same result can be obtained in a less unpleasant way. I mention it, however, to show what it is that the practitioner must aim at. If, when the head is distending the perineum greatly, the thumb and fore finger of the right hand are placed along its sides, it can be pushed gently forward over the head at the height of the pain, while the tips of the fingers may at the same time press upon the advancing vertex, so as to retard its progress if advisable (Fig. 106). By this means the sudden and forcible stretching of the perineal structures is prevented, and the chance of laceration reduced to a minimum, while nature's mode of relaxing the tissues, by dilatation of the anal orifice, is favored. This is very different from the mechanical support that is usually recommended, and the less pressure that is applied directly to the perineum the better. Nor is it either needful or advisable to sit by the patient with the hand applied to the perineum for hours, as is so often practised. Time should be given for the gradual distension of the tissues by the alternate advance and recession of the head, and we need only intervene to assist relaxation when the stretching has reached its height and the head is about to be expelled. A napkin may be interposed between the hand and the skin for the purpose of cleanliness. Should the perineum be excessively tough and resistant, assiduous fomentation with a hot sponge may be resorted to, and will be of some service in promoting relaxation.

Incision of the Perineum.—When the tension is so great that laceration seems inevitable, it is generally recommended that a slight incision should be made on each side of the central raphé, with the view of preventing spontaneous laceration. This may no doubt be done with perfect safety, but I question if it is likely to be of use. The idea is that

an incised wound is likely to heal more readily than a lacerated one. When, however, a distended perineum ruptures, its structures are so thinned that the tear is always linear, and, as a matter of fact, the edges of the tear are always as clean and as closely in apposition as if the cut had been made with a knife. Moreover, the laceration invariably heals

FIG. 106.



Mode of Effecting Relaxation of the Perineum.

perfectly if only the edges be brought into contact at once with one or two metallic sutures. I believe, therefore, that Goodell is right in stating that incision of the perineum is rarely, if ever, necessary, unless it is hardened by previous cicatrization. In almost all first labors the fourchette is torn, but requires no treatment of any kind. In some cases, do what we will, more or less laceration occurs, and the perineum should always be examined after the expulsion of the child to see if any tear has taken place.

Treatment of Lacerations.—If it has given way to any extent, I believe that it is good practice to insert one or two interrupted sutures of silver wire or carbolized gut at once. Immediately after delivery the sensibility of the tissues is deadened by the distension to which they have been subjected, and the sutures can be inserted with little or no pain. It is quite true that lacerations of an inch or less will generally heal perfectly well of themselves, but this is not invariably the case, while healing almost certainly follows if the edges be brought together at once. In the severer forms of laceration, extending back to, or even through, the sphincter, the precaution is all the more necessary, and a subsequent operation of gravity may in this way be avoided. The sutures can be removed without difficulty in a week or so, when complete adhesion has taken place.

Expulsion of the Child.—The head, when expelled, should be received in the palm of the right hand, while the left hand is placed upon the abdomen to follow down the uterus as it contracts and expels the body.

There is generally some little delay after the expulsion of the head, and we should now see if the cord surround the neck, and if it does so it should be drawn over the head; and, if this is not possible, it may be tied and divided between the ligatures. The expulsion of the body should be left entirely to the uterine contractions. If there be undue delay, we may endeavor to excite uterine action by friction on the fundus, and it will rarely happen that sufficient contraction does not now come on. If we display undue haste in withdrawing the body, we run the risk of emptying the uterus while its tissues are relaxed, and so favor hemorrhage. If, however, there seem serious danger of the child being asphyxiated, its expulsion may be favored by gently passing the fore finger of each hand within the axillæ and using traction; but it is only very exceptionally that such interference is required.

Promotion of Uterine Contraction after the Birth of the Child.—As the uterus contracts it should be carefully followed down through the abdominal parietes by the left hand, which should grasp it as the body is expelled, with the view of seeing that it is efficiently contracted. This is a point of vital importance in preventing hemorrhage, which will presently be more especially considered.

Ligature of the Cord.—As soon as the child cries we may proceed to tie and separate the cord. For this purpose the nurse usually provides ligatures composed of several strands of whitey-brown thread, but tape or any other suitable material may be employed. It is important, especially if the cord be very thick and gelatinous, to see that it is thoroughly compressed, so that the vessels are obliterated, otherwise secondary hemorrhage might occur. The cord is tied about an inch and a half from the child, and it is usual, though of course not essential, to place a second ligature about two inches nearer the placental extremity of the cord. The latter is perhaps of some use by retaining the blood, and thus increasing the size of the placenta and favoring its more ready expulsion by uterine contraction. The cord is then divided with scissors between the ligatures, the child wrapped up in flannel, and given to the nurse or a bystander to hold, while the attention of the practitioner is concentrated on the mother with a view to the proper management of the third stage of labor. The researches of Budin,¹ Ribemont,² and others show that there is a distinct advantage in not tying the cord until the child has cried lustily, as the act of respiration tends to withdraw the placental blood, and thus increases the entire amount of blood in the fœtus. It is said that after late ligature of the cord the child is more vigorous and active than when it is tied too early.

Treatment of the Cord by Laceration.—The cord may, if preferred, be treated with perfect safety by laceration. This method was first brought under my notice by my friend Dr. Stephen, who has employed it for many years and in several hundred cases. The cord is twisted round the index fingers of both hands and torn through, the lacerated vessels retracting without any hemorrhage. It is a close imitation of the method instinctively adopted by the lower animals, who gnaw the cord asunder, and has the advantage of dispensing with ligatures altogether.

¹ Budin, *Progress médicale*, 1876.

² *Archiv. de Toxolog.*, Oct., 1879.

I have used it myself in a large number of cases, but prefer, on the whole, the plan usually adopted.

Importance of Proper Management of Third Stage.—There is unquestionably no period of labor where skilled management is more important and none in which mistakes are more frequently made. By proper care at this time the risk of post-partum hemorrhage is reduced to a minimum, the efficient contraction of the uterus is secured, the amount and intensity of after-pains are lessened, and the safety and comfort of the patient greatly promoted. Moreover, the general practice as to the management of this stage is opposed to the natural mechanism of placental expulsion, and is far from being well adapted to secure the important objects which we ought to have in view. Let us see what is the practice usually recommended and followed, and then we shall be in a position to understand in what respects it is erroneous. For this purpose I cannot do better than copy the directions contained in one of our most deservedly popular obstetric text-books, which undoubtedly expresses the usual practice in the management of this stage: "When the binder is applied the patient may be allowed to rest a while if there is no flooding; after which, *when the uterus contracts*, gentle traction may be made by the funis to ascertain if the placenta be detached. If so, and especially if it be in the vagina, it may be removed by continuing the traction steadily in the axis of the upper outlet at first, at the same time making pressure on the uterus."¹

[In this country, for many years, the uniform teaching has been that the binder should not be applied until the uterus has expelled the placenta and become firmly contracted. Although the plan of expression was not carried out as completely as is now taught under the Credé method, that of stimulating the contractions of the uterus by manipulation and pressure was certainly in use forty years ago. When the size and solidity of the uterus, as ascertained by the compressing hand, indicate that the placenta has been expelled into the vagina, it is a question whether we shall cause it to be forced through the vulva by pressing down the uterus upon it, or make traction upon it by the finger hooking down its edge. Occasionally, we find a patient who is very sensitive to pressure made upon her uterus after it has become firmly contracted; and in such a case it may be well to depend partly upon traction for completing the delivery of the secundines. That it is possible for the uterus to expel the placenta suddenly from the vagina where no pressure has been made is evident from the fact that a physician of this city, who was making traction upon the cord under the old method some years ago, was surprised to find the placenta shoot out from the vulva and dangle by the funis as he held it in his hand. In such a case the uterus must have been aided during a contraction by voluntary abdominal pressure, causing the os to descend nearly to the vulva. It is very evident that the uterus is subject to muscular fatigue and to the exhaustion of its contractile power when long in action; hence, there is a greater risk of uterine atony and hemorrhage after a long labor than a short one, and we may expect a more complete expulsion of the placenta in the latter. It is also clear, from cases in my own experience, that the

¹ Churchill's *Theory and Practice of Midwifery*, p. 162.


muscular power of the uterus is by no means in proportion to the general strength of the woman. The power to assist by bearing down no doubt is, but the independent power of the organ itself does not appear to be. Certainly some of the most perfect in parturient power that have come under my care were small women, with little general muscular force. One little woman of 86 pounds weight appeared almost to have escaped the curse pronounced upon Eve; and another, still smaller, expelled a placenta from her vagina almost without any loss of blood.—ED.]

Objections to Ordinary Practice.—This may fairly be taken as a sufficiently accurate description of the practice usually followed.¹ The objections I have to make are: (1) That it inculcates the common error of relying on the binder as a means of promoting uterine contraction, advising its application before the expulsion of the placenta; while I hold that the binder should never be applied until after the placenta is expelled, and not even then unless the uterus is perfectly and permanently contracted. (2) That it teaches that traction on the cord should be used as a means of withdrawing the placenta, whereas the uterus itself should be made to expel the after-birth, and in nineteen cases out of twenty the finger need never be introduced into the vagina after the birth of the child, nor the cord touched. This may seem an exaggerated statement to those who have accustomed themselves to the usual method of dealing with the placenta; but I feel confident that all who have learnt the method of expression of the placenta would testify to its accuracy.

Expression of the Placenta.—The cardinal point to bear in mind is,

¹This practice is further illustrated by the annexed diagram, contained in most obstetric works, which represents the accoucheur as withdrawing the placenta by traction, and which I insert as an illustration of what ought *not* to be done (Fig. 107):

FIG. 107.



Usual Method of Removing the Placenta by Traction on the Cord.

that the placenta should be expelled from the uterus by a *vis a tergo*, not drawn out by a *vis a fronte*. That uterine pressure after the birth of the child has been recommended by many English writers is certain, and the Dublin school especially have dwelt on its importance as a preventive of post-partum hemorrhage; but the distinct enunciation of the doctrine that the placenta should be pressed, and not drawn, out of the uterus, we owe to Credé and other German writers; and it is only of late years that this practice has become at all common. Those who have not seen placental expression practised find it difficult to understand that in the large majority of cases the uterus may be made to expel the placenta out of the vagina; but such is unquestionably the fact. A little practice is no doubt necessary to effect this satisfactorily; but when once the knack has been learnt, there is little difficulty likely to be experienced.

Importance of Not Removing the Placenta Hurriedly.—Before describing the method of placental expression a word of caution may be said against undue haste in attempting expression of the placenta—a mistake that is often made, and which, I believe, tends to increase the risk of post-partum hemorrhage. So long as we satisfy ourselves that the uterus is fairly contracted, so as to avoid the possibility of its distension with blood, a certain delay after the birth of the child is useful, from its giving time for coagula to form within the uterine sinuses, by which their open mouths are closed up. The importance of this point has been specially dwelt upon by McClintock, who lays down the rule that fifteen or twenty minutes should be allowed to elapse after the birth of the child before any attempt to remove the after-birth is made. This is a good and safe practical rule, as it gives ample time for the complete detachment of the placenta and the coagulation of the blood in the uterine sinuses.

Mode of effecting Expression of the Placenta.—During this interval the practitioner or nurse should sit by the bedside, with the hand on the uterus to secure contraction and prevent distension, but not kneading or forcibly compressing it. When we judge that a sufficient time has elapsed we may proceed to effect expulsion. For this purpose the fundus should be grasped in the hollow of the left hand, the ulnar edge of the hand being well pressed down behind the fundus, and *when the uterus is felt to harden* strong and firm pressure should be made downward and backward in the axis of the pelvic brim. If this manœuvre be properly carried out and sufficiently firm pressure made, in almost every case the uterus may be made to expel the placenta into the bed, along with any coagula that may be in its cavity (Fig. 108). The uterine surface of the placenta is generally expelled first, as is represented in the diagram, the cord being within the membranes; whereas the foetal surface and root of the cord are the parts which appear first when the placenta is removed by traction (Fig. 107). If we do not succeed at the first effort, which is rarely the case if extrusion be not attempted too soon after the birth of the child, we may wait until another contraction takes place, and then reapply the pressure. I repeat that, after a little practice, the placenta may be entirely expelled in this way, in nineteen cases out of twenty, without even touching the cord,

and the bugbear of retained placenta will cease to be a source of dread.

Should we fail in causing the uterus to expel the placenta, a vaginal examination may be made, and, if the placenta be found lying entirely in the vagina, it may be carefully withdrawn. If, however, the cord can be traced up through the os, showing that the placenta is still within the uterine cavity, we must again resort to pressure to effect its expulsion, and not attempt to withdraw it by traction. Such cases may fairly be classed as retained placenta, but they should be very rarely met with, and are discussed elsewhere. When they do occur often in the hands of the same practitioner, it is fair to conclude that he has not properly acquired the art of managing this stage of labor. Generally speaking, the placenta should be expelled within twenty minutes

FIG. 108.



Illustrating Expression of the Placenta.

after the birth of the child, but no doubt in the large majority of cases expulsion might be effected sooner were it advisable to attempt it.

Management of the Membranes.—When the mass of the placenta is expelled, the membranes generally still remain in the vagina, and they should be twisted into a rope and very gently withdrawn, so as not to leave any portion behind. This is a precaution the importance of which I would strongly urge, for I believe that the chance of part of the membranes being torn off and left in utero is the one objection to the method recommended. With due care, however, this accident may be avoided; and the risk will be lessened if the placenta is received into the palm of the right hand, on expression, so as to avoid any strain on the membranes.

Compression of the Uterus after the Expulsion of the Placenta.—The duties of the medical attendant are not even now over. For at least ten minutes after the extrusion of the placenta he should keep his hand on the firmly-contracted uterus, gently kneading it, without any force, for the purpose of promoting firm and equable contraction and causing it to throw off the coagula that may form in its cavity.

Administration of Ergot of Rye.—The subsequent comfort and safety of the patient may be promoted by administering at this time a full dose of ergot of rye, such as a drachm or more of the liquid extract. The property possessed by this drug of producing tonic and persistent contraction of the uterine fibres, which renders it of doubtful utility as an oxytocic during labor, is of special value after delivery, when such contraction is precisely what we desire. I have long been in the habit of administering the drug at this period, and believe it to be of great value, not only as a prophylactic against hemorrhage, but as a means of lessening after-pains.

Application of the Binder.—When we are satisfied that the uterus is permanently contracted we may apply the binder, but this should rarely be done until at least half an hour after the birth of the child. The soiled clothes should be gently withdrawn from under the patient, moving her as little as possible, and the binder should be, at the same time, slipped under the body, taking care that it is passed well below the hips, so as to secure a firm hold. No kind of bandage is better than a piece of stout jean, of sufficient breadth to extend from the trochanters to the ensiform cartilage: a jack-towel or bolster slip answers the purpose very well. These are preferable, at any rate at first, to the shaped binders that are often used. One or two folded napkins are generally placed over the uterus, so as to form a pad to keep up pressure. Once in position, the binder is pulled tight and fastened by pins. The utility of careful bandaging after delivery can scarcely be doubted, although some years ago it became the fashion to dispense with it. It gives a comfortable support to the lax abdominal walls, keeps up a certain amount of pressure on the uterus, and tends to restore the figure of the patient. After the bandage is applied a warm napkin should be placed on the vulva as a means of estimating the quantity of the discharge, and the patient may be allowed to rest.

After-Treatment.—Unless the labor have been very long and fatiguing, an opiate, often exhibited as a matter of routine, is unadvisable, although it may be well to leave one with the nurse, to be given if the patient cannot sleep or if the after-pains be very troublesome. The practitioner may now leave the room, but not the house, and at least an hour should elapse after delivery before he takes his departure. Before doing so he should visit the patient, inspect the napkin to see that there is not too much discharge, and satisfy himself that the uterus is contracted and not distended with coagula. He should also count the pulse, which, if the patient be progressing satisfactorily, will be found at its normal average. If, however, it be beating over 100 per minute, he should on no account leave, for such a rapidity of the circulation renders it extremely probable that hemorrhage is impending. This is a good practical rule, laid down by McClintock in his excellent paper "On the Pulse in Childbed," attention to which may often save the patient from disastrous consequences.

Before leaving, the practitioner should see that the room is darkened, all bystanders excluded, and the patient left as quiet as possible to recover from the shock of labor.

CHAPTER IV.

ANÆSTHESIA IN LABOR.

A FEW words may be said as to the use of anæsthetics during labor—a practice which has become so universal that no argument is required to establish its being a perfectly legitimate means of assuaging the sufferings of childbirth. Indeed, the tendency in the present day is in the opposite direction, and a common error is the administration of chloroform to an extent which materially interferes with the uterine contractions and predisposes to subsequent post-partum hemorrhage.

Agents Employed.—Practically speaking, the only agent hitherto employed in this country is chloroform, although the bichloride of methylene and ether have been occasionally tried. Of late years, chloral has been extensively used by some, and, as I believe it to be an agent of very great value, I shall first indicate the circumstances under which it may be employed.

Chloral.—The peculiar value of chloral in labor is that it may be safely administered at a time when chloroform cannot be generally employed. The latter, while it annuls suffering, very frequently tends in a marked degree to diminish uterine action. This is a familiar observation to all who have employed it much during labor, as the diminution of the force and intensity of the pains, and the consequent retardation of the labor, often oblige us to suspend its inhalation, at least temporarily. Indeed, this very property of annulling uterine action is one of its most valuable qualities in obstetrics, as in certain cases of turning. For such purposes it is necessary to give it to the surgical extent, which we endeavor to avoid when it is used simply to lessen the suffering of ordinary labor. Still, it is not always easy to limit its action in this way, and thus it very frequently does more than we wish. Such diminution in the intensity of uterine contraction is comparatively of less consequence in the propulsive stage, and it is generally more than counterbalanced by the relief it affords. In the first stage it is otherwise, and, practically speaking, chloroform is generally not admissible until the head is in the pelvic cavity.

Chloral is especially the Anæsthetic of the First Stage.—Chloral, on the other hand, has no such relaxing effects on uterine contraction. It cannot, it is true, compete with chloroform in its power of relieving pain, but it produces a drowsy state in which the pain is not felt nearly so acutely as before. It is therefore in the first stage of labor, while the pains are cutting and grinding, and during the dilatation of the cervix, that it finds its most useful application. It is especially valuable in those cases, so frequently met with in the upper classes, in which the pains produce intolerably acute suffering, but with little effect on the progress of the labor. In them the os is often thin and rigid and the pains very frequent and acute, but little or no dilatation is effected. When the patient

is brought under the influence of chloral, however, the pains become less frequent but stronger, nervous excitement is calmed, and the dilatation of the cervix often proceeds rapidly and satisfactorily. Indeed, I know of nothing which answers so well in cases of rigid, undilatable cervix, and I believe its administration to be far more effective, under such circumstances, than any of the remedies usually employed.

Object and Mode of Administration.—The object is to produce a somnolent condition which shall be protracted as long as possible. For this purpose 15 grains of chloral may be administered every twenty minutes until three doses are given. This generally suffices to produce the desired effect. The patient becomes very drowsy, dozes between the pains, and wakes up as each contraction commences. It may be necessary to give a fourth dose at a longer interval, say an hour after the third dose, to keep up and prolong the soporific action; but this is seldom necessary, and I have rarely given more than a drachm of chloral during the entire progress of labor. Another advantage of this treatment is that, while it does not interfere with the use of chloroform in the second stage, it renders it necessary to give less than otherwise would be called for, and thus its action can be more easily kept within bounds. On the whole, therefore, I am inclined to consider chloral a very valuable aid in the management of labor, and believe that it is destined to be much more extensively used than is at present the case. So far as my experience has yet gone, I have not met with any symptoms which have led me to think that it has produced bad effects; and I have known many patients sleep quietly through labor, without expressing any excessive suffering or asking for chloroform, who under ordinary circumstances would have been most urgently calling for relief. It occasionally happens that the patient cannot retain the chloral from its tendency to produce sickness; it may then be readily given per rectum in the form of enema.

Chloroform.—Generally speaking, we do not think of giving chloroform until the os is fully dilated, the head descending, and the pains becoming propulsive. It has often, indeed, been administered earlier, for the purpose of aiding the dilatation of a rigid cervix; and there is no doubt that it often succeeds well when employed in this way, but I have already stated my belief that chloral answers this purpose better.

Only to be Given during the Pains.—There is one cardinal rule to be remembered in giving chloroform during the propulsive stage, and that is that it should be administered intermittently, and never continuously. When the pain comes on a few drops may be scattered over a Skinner's inhaler, which affords one of the best means of administering it in labor, or placed within the folds of a handkerchief twisted into the form of a cone. * During the acme of the pain the patient inhales it freely, and at once experiences a sense of great relief; and as soon as the pain dies away the inhaler should be removed. In the interval between the pains the effect of the drug passes off, so that the higher degree of anæsthesia should never be produced. Indeed, when properly given consciousness should not be entirely abolished, and the patient, between the pains, should be able to speak and understand what is said to her. This intermittent administration constitutes the peculiar safety of chloroform administered in labor; and it is a fortunate circumstance that, as yet,

there is, I believe, no case on record of death during the inhalation of chloroform for obstetric purposes.[¹] This is obviously due to the effect of each inhalation passing off before a fresh dose is administered.

The effect on the pains should be carefully watched. If they become very materially lessened in force and frequency, it may be necessary to stop the inhalation for a short time, commencing again when the pains get stronger; which effect may be often completely and easily prevented by mixing the chloroform with about one-third of absolute alcohol, which, originally recommended, I believe, by Dr. Sansom, increases the stimulating effects of chloroform, and thus diminishes its tendency to produce undue relaxation. The amount administered must vary, of course, with the peculiarities of each individual case and the effect produced, but it need never be large. As the head distends the perineum and the pains get very strong and forcing, it may be given more freely, and to the extent of inducing even complete insensibility just before the child is born.

Ether as a Substitute for Chloroform.—In cases in which chloroform has lessened the force of the pains ether may be given instead with great advantage. It certainly often acts well when chloroform is inadmissible on account of its effects on the pains, and, so far as my experience goes, it has not the property of relaxing the uterus, but, on the contrary, has sometimes seemed to me distinctly to intensify the pains. Of late I have used a mixture of one part of absolute alcohol, two of chloroform, and three of ether. This is less disagreeable than ether, and has not the over-relaxing effects of chloroform.

Precautions.—Bearing in mind the tendency of chloroform to produce uterine relaxation, more than ordinary precautions should always be taken against post-partum hemorrhage in all cases in which it has been freely administered.

In cases of operative midwifery it is often given to the extent of producing complete anæsthesia. In all such cases it should be administered, when possible, by another medical man, and not by the operator, because the giving of chloroform to the surgical degree requires the undivided attention of the administrator, and no man can do this and operate at the same time. I once learnt an important lesson on this point. I had occasion to apply the forceps in the case of a lady who insisted on having chloroform. When commencing the operation I noticed some suspicious appearances about the patient, who was a large stout woman with a feeble circulation. I therefore stopped, allowed her to regain consciousness, and delivered her without anæsthesia, much to her own annoyance. Just one month after labor she went to a dentist to have a tooth extracted, and took chloroform, during the inhalation of which she died. This impressed on my mind the lesson that no man can do two things at the same time. The partial unconsciousness of incomplete anæsthesia, in which the patient is restless and tossing about, renders the application of forceps, as well as all other operations, very difficult.

[¹ In the *Transactions of the American Gynecological Society* for the year 1877 are five cases of chloroform-poisoning occurring in obstetrical cases, reported by Dr. W. T. Lusk of New York. In three, restoration was effected by artificial respiration, but in two death resulted absolutely.—Ed.]

Therefore, unless the patient can be completely and fully anæsthetized, it is better to operate without chloroform being given at all.

[In the United States the dangers attending the use of chloroform in obstetric practice have, in large measure, banished it from the lying-in chamber. Some obstetricians in our chief cities still resort to it with little hesitation, believing that by great carefulness in its administration, and by the substitution of ether in exceptional cases, all danger may be avoided. Others have a very great fear of it, and universally trust to the safer anæsthetic. It is an error to suppose that the parturient state robs chloroform of much of its danger, the apparent immunity being due to its intermittent and incomplete administration; complete anæsthesia being but a fraction less dangerous than in surgical operations upon women who are not pregnant. Dr. Lusk, already quoted, after a large experience with the use of chloroform, says: "*Patients in labor do not enjoy any absolute immunity from the pernicious effects of chloroform.*"¹ It is much to be regretted that this more pleasant anæsthetic is so much more dangerous than ether as an inhalant; but in consideration of the difference of risk, that of their relative effects upon the nose and trachea is scarcely to be considered. Chloroform acts upon the respiratory centres just as ether does; and this is an element of danger in each, but is capable of being counteracted by artificial respiration. But, beyond this, chloroform is far more dangerous, in acting upon the motor ganglia of the heart and producing sudden death. According to the experiments of Vulpian upon animals, not more than one case of cardiac failure in forty can be restored by artificial respiration. He affirms that there is danger at the commencement, during the course, and at the close of chloroformization, and even some hours or days subsequent to it. Nélaton made the important discovery that the cerebral anæmia produced by chloroform, with its accompanying death-like condition, might be remedied by long perseverance in artificial respiration with the patient turned head downward.

Anæsthesia in labor is much less popular, both with obstetricians and patients in this country, than it was soon after its introduction. Improvements in the purity of sulphuric ether have made the narcosis more reliable, but the general effect upon patients varies very decidedly, being all that can be desired in some, and just the reverse in others. Some of the undesirable effects I have witnessed are intoxication, with cessation of labor, hysterical excitement, nightmare, and post-partum inertia and hemorrhage. I have also witnessed the most delightful results from ether that could be desired. In a small, delicate multipara, whose mother died of phthisis, and to whom I had been obliged to administer stimulants in the first and much of the second stage of labor, the use of ether had the effect to revolutionize her condition. Her pulse became strong; her expulsive power increased; she had no suffering; her placenta was expelled without accompanying blood; and there was no subsequent uterine relaxation. But such cases are, unfortunately, exceptional.—ED.]

[¹ *Opus cit.*]

CHAPTER V.

PELVIC PRESENTATIONS.

UNDER the head of *pelvic* presentations it is customary to include all cases in which any part of the lower extremities of the child presents. By some these are further subdivided into *breech*, *footling*, and *knee* presentations; but, although it is of consequence to be able to recognize the feet and the knee when they present, so far as the mechanism and management of delivery are concerned the cases are identical, and therefore may be most conveniently considered together.

Frequency.—Presentations coming under this head are far from uncommon; those in which the breech alone occupies the pelvis are met with, according to Churchill, once in 52 labors, while Ramsbotham estimates that it presents more frequently—viz. once in 38.8 labors. Footling presentations occur only once in 92 cases. They are probably often the mere conversion of original breech presentations, the feet having come down during the labor, either in consequence of the sudden escape of the liquor amnii when the breech was still freely movable above the brim, or from some other cause. Knee presentations are extremely rare, as may be readily understood if it be borne in mind that to admit them the thighs must be extended, hence the vertical measurement of the child must be greatly increased, and therefore it could not be readily accommodated within the uterine cavity unless of unusually small size. As a matter of fact, Mme. La Chapelle found only one knee presentation in upward of 3000 cases.

Causes.—The causes of pelvic presentations are not known. They are probably the same as those which produce other varieties of malpresentations; and it is not unlikely that in certain women there may be some peculiarity in the shape of the uterine cavity which favors their production. It would be difficult otherwise to explain such a case as that mentioned by Velpeau in which the breech presented in six labors.

Prognosis.—The results, as regards the mother, are in no way more unfavorable than in vertex presentation. The first stage of the labor is generally tedious, since the large rounded mass of the breech does not adapt itself so well as the head to the lower segment of the uterus, and dilatation of the cervix is consequently apt to be retarded. The second stage is, however, if anything, more rapid than in vertex cases; and even when it is protracted the soft breech does not produce such injurious pressure on the maternal structures as the hard and unyielding head.

The Infantile Mortality in Pelvic Presentations.—The result is very different as regards the child. Dubois calculated that 1 out of 11 children was stillborn. Churchill estimates the mortality as much higher—viz. 1 in 3½. The latter certainly indicates a larger number of stillbirths than is consistent with the experience of most practitioners, and more than should occur if the cases be properly managed; but there can be

no doubt that the risk to the child is, even under the most favorable circumstances, very great. Even when the child is not lost it may be seriously injured. Dr. Ruge has tabulated a series of 29 cases in which there were found to be fractures of bones or other injuries.¹

Causes of Fœtal Mortality.—The chief source of danger is pressure on the umbilical cord in the interval elapsing between the birth of the body and the head. At this time the cord is very generally compressed between the head of the child and the pelvic walls, so that circulation in its vessels is arrested. Hence the aëration of the fœtal blood cannot take place, and, pulmonary respiration not having been yet established, the child dies asphyxiated. There are other conditions present which tend, although in a minor degree, to produce the same result. One of these is that the placenta is probably often separated by the uterine contractions when the bulk of the body is being expelled, as, indeed, takes place, under analogous circumstances, when the vertex presents, the necessary result being the arrest of placental respiration. Joulin thinks that the same effect may be produced by the compression of the placenta between the contracted uterus and the hard mass of the fœtal skull. Probably all these causes combine to arrest the functions of the placenta, and if the delivery of the head, and consequently the establishment of pulmonary respiration, be delayed, the death of the child is almost inevitable. The corollary is, that the danger to the child is in direct proportion to the length of time that elapses between the birth of the body and that of the head.

The risk to the child is greater in footling than in breech cases, because in the former the maternal structures are less perfectly dilated in consequence of the small size of the feet and thighs, and therefore the birth of the head is more apt to be delayed.

Diagnosis.—Inasmuch as the long axis of the child corresponds with the long axis of the uterus in pelvic as in vertex presentations, there is nothing in the shape of the uterus to arouse suspicion as to the character of the case. Still, it is often sufficiently easy to recognize a pelvic presentation by abdominal examination if we have occasion to make one. The facility with which it may be done depends a good deal on the individual patient. If she be not very stout, and if the abdominal parietes be lax and non-resistant, we shall generally be able to feel the round head at the upper part of the uterus, much firmer and more defined in outline than the breech. The conclusion will be fortified if we hear the fœtal heart beating on a level with or above the umbilicus. The greater resistance on one side of the abdomen will also enable us to decide with tolerable accuracy to which side the back of the child is placed. Information thus acquired is, at the best, uncertain, and we can never be quite sure of the existence of a pelvic presentation until we can corroborate the diagnosis by vaginal examination.

Results of Vaginal Examination.—The first circumstance to excite suspicion on examination *per vaginam*, even when the os is undilated, is the absence of the hard globular mass felt through the lower segment of the uterus, so characteristic of vertex presentations. When the os is sufficiently open to allow the membranes to protrude, although the pre-

¹ *Bull. gén. de Thérap.*, August, 1875.

senting part is too high up to be within reach, we may be struck with the peculiar shape of the bag of membranes, which, instead of being rounded, projects a considerable distance through the os, like the finger of a glove. This is a peculiarity met with in all mal-presentations alike, and is, indeed, much less distinct in breech than in footling presentations, because in the former the membranes are more stretched, just as they are in vertex cases. When the membranes rupture, instead of the waters dribbling away by degrees, they often escape with a rush, in consequence of the pelvic extremity not filling up the lower part of the uterus so accurately as the head, which acts as a sort of ball-valve and prevents the sudden and complete discharge of the waters.

Diagnosis of the Breech.—Often, on first examining, even when the membranes are ruptured, the presentation is too high up to be made out accurately. All that we can be certain of is that it is not the head; and the case must be carefully watched and examinations frequently repeated until the precise nature of the presentation can be established. If the breech present, the finger first impinges on a round, soft prominence, on depressing which a bony protuberance, the trochanter major, can be felt. On passing the finger upward it reaches a groove, beyond which a similar fleshy mass, the other buttock, can be felt. In this groove various characteristic points, diagnostic of the presentation, can be made out. Toward one end we can feel the movable tip of the coccyx, and above it the hard sacrum with its rough projecting prominences. These points, if accurately made out, are quite characteristic, and resemble nothing in any other presentation. In front there is the anus, in which it is sometimes, but by no means always, possible to insert the tip of the finger. If this can be done, it is easy to distinguish it from the mouth, with which it might be confounded, by observing that the hard alveolar ridges are not contained within it. Still more in front we may find the genital organs, the scrotum in male children being often much swollen if the labor has been protracted. Thus it is often possible to recognize the sex of the child before birth.

Differential Diagnosis.—The breech might be mistaken for the face, especially if the latter be much swollen; but this mistake can readily be avoided by feeling the spinous processes of the sacrum.

The knee is recognized by its having two tuberosities with a depression between them. It might be confounded with the heel, the elbow, or the shoulder. From the heel it is distinguished by having two tuberosities instead of one; from the elbow, by the latter having one sharp tuberosity, with a depression on one side, instead of a central depression and two lateral prominences; and from the shoulder, by the latter being more rounded, having only one prominence, running from which the acromion and clavicle can be traced.

Diagnosis of the Foot.—The foot may be mistaken for the hand. This error will be avoided by remembering that all the toes are in the same line, and that the great toe cannot be brought into apposition with the others, as the thumb can with the fingers. The internal border of the foot is much thicker than the external, whereas the two borders of the hand are of the same thickness. Moreover, the foot is articulated at right angles to the leg, and cannot be brought into a line with it, as

the hand can with the arm. Finally, the projection of the calcaneum is characteristic, and resembles nothing in the hand.

Mechanism.—As is the case in other presentations, obstetricians have very variously subdivided breech presentations, with the effect of needlessly complicating the subject. The simplest division, and that which will most readily impress itself on the memory of the student, is to describe the breech as presenting in four positions, analogous to those of the vertex, the sacrum being taken as representing the occiput, and the positions being numbered according to the part of the pelvis to which it points. Thus we have—

First, or left sacro-anterior (corresponding to the first position of the vertex). The sacrum of the child points to the left foramen ovale of the mother.

Second, or right sacro-anterior (corresponding to the second vertex position). The sacrum of the child points to the right foramen ovale of the mother.

Third, or right sacro-posterior (corresponding to the third vertex position). The sacrum of the child points to the right sacro-iliac synchondrosis of the mother.

Fourth, or left sacro-posterior (corresponding to the fourth vertex position). The sacrum of the child points to the left sacro-iliac synchondrosis of the mother,

Of these, as with the corresponding vertex positions, the first and third are the most common, their comparative frequency, no doubt, depending on the same causes. The mechanical conditions to which the presenting part is subjected are also identical, but the alterations of position of the breech in its progress are by no means so uniform as those of the head, on account of its less perfect adaptation to the pelvic cavity. The mechanism of the delivery of the shoulders and head in breech presentations, moreover, is of much greater practical importance than that of the body in vertex presentations, inasmuch as the safety of the child depends on its speedy and satisfactory accomplishment. Bearing these facts in mind, it will suffice to describe briefly the phenomena of delivery in the first and third breech positions.

Position of the Child at Brim.—In the first position (Fig. 109) the sacrum of the child points to the left foramen ovale; its back is consequently placed to the left side of the uterus and anteriorly, and its abdomen looks to the right side of the uterus and posteriorly. The sulcus between the buttocks lies in the right oblique diameter of the pelvis, while the transverse diameter of the buttocks lies in the left oblique diameter, the left buttock being most easily within reach. As in vertex presentations, the hips of the child lie on the same level at the pelvic brim, although Naegle describes the left hip as placed lower than the right.

Descent.—As the pains act on the body of the child the breech is gradually forced through the pelvic cavity, retaining the same relations as at the brim, its progress being generally more slow than that of the head, until it reaches the lower pelvic strait, when the same mechanism which produces rotation of the occiput comes to operate upon it. The result is a rotation of the child's pelvis, so that its transverse diameter

comes to lie approximately in the antero-posterior diameter of the outlet; its antero-posterior diameter corresponds to the transverse diameter of the mother's pelvis, the left hip lies behind the pubes, and the right toward the sacrum. This rotation, which is admitted by the majority of obstetricians, is altogether denied by Naegele. There can be no doubt, however, that it does generally take place, but by no means so constantly as the corresponding rotation of the vertex; and it is not uncommon for it to be entirely absent and for the hips to be born in the oblique diameter of the outlet. The body of the child is said frequently not to follow the movement imparted to the hips, so that there is more or less of a twist in the vertebral column.

Expulsion of the Hips and Body.—The left hip now becomes firmly fixed behind the pubes, and a movement of extension, analogous to that

FIG. 109.



First, or Left Sacro-anterior, Position of the Breech.

of the head in vertex presentations, takes place. The right, or posterior, hip revolves round the fixed one, gradually distends the perineum, and is expelled first, the left hip rapidly following. As soon as both hips are born the feet slip out, unless the legs are completely extended upon the child's abdomen. The shoulders soon follow, lying in the left oblique diameter of the pelvis (Fig. 110). The left shoulder rotates forward behind the pubes, where it becomes fixed, the right shoulder sweeping over the perineum and being born first. The arms of the child are generally found placed upon its thorax, and are born before the shoulders. Sometimes they are extended over the child's head, thus causing considerable delay and greatly increasing the risk to the child. It is now generally admitted that such extension is most apt to occur when traction has been made on the child's body with the view of hastening delivery, and that it is rarely met with when the expulsion of the body is left entirely to the natural powers.

Delivery of the Head.—When the shoulders are expelled the head enters the pelvis in the opposite, or right oblique, diameter, the face looking to the right sacro-iliac synchondrosis. As the greater part of

FIG. 110.



Passage of the Shoulders and Partial Rotation of the Thorax.

the child is now expelled, and as the head has entered the vagina, the uterus, having a comparatively small mass to contract upon, must obviously act at a mechanical disadvantage. Still, the pressure of the head on the vagina is a powerful inciter, the accessory muscles of parturition are brought into strong action, and there may be sufficient force to ensure expulsion of the head without artificial aid. On account of the great resistance to the descent of the occiput from its articulation with the spinal column, the pains have the effect of forcing down the anterior portion of the head, and this ensures the complete flexion of the chin upon the sternum (Fig. 111). This is a great advantage from a mechanical point of view, as it causes the short occipito-mental diameter of the

FIG. 111.



Descent of the Head.

head to enter the pelvis in the axis of the uterus and the brim. If the head should be in a state of partial extension, as sometimes happens when the pelvis is unusually roomy, the occipito-frontal diameter is

placed in a similar relation to the brim—a position certainly less favorable to the easy birth of the head. As the head descends it experiences a movement of rotation, the occiput passing forward and to the right, behind the pubic arch, the face turning backward into the hollow of the sacrum. The body of the child will be observed to follow this movement, so that its back is turned toward the mother's abdomen, its anterior surface to the perineum. The nape of the neck now becomes firmly fixed under the arch of the pubes; the pains act chiefly on the anterior portion of the head, and cause it to sweep over the perineum, the chin being first born, then the mouth and forehead, and lastly the occiput.

Sacro-posterior Positions.—It is needless to describe the differences between the mechanism of the second and first positions, which the student, who has mastered the subject of vertex presentations, will readily understand. It is necessary, however, to say a few words as to sacro-posterior positions, choosing for that purpose the third, which is the more common of the two. This is exactly the opposite of the first position. The sacrum of the child points to the right sacro-iliac synchondrosis; its abdomen looks forward and to the left side of the mother. The transverse diameter of the child's pelvis lies in the left oblique diameter, the right hip being anterior. The birth of the body generally takes place exactly in the way that has been already described, the right hip being toward the pubes.

As the head descends into the pelvis the occiput most usually rotates along its right side—the rotation having been often already partially effected when that of the hips had been made—until it comes to rest behind the pubes, the face passing backward along the left side of the pelvis into the hollow of the sacrum. This change corresponds exactly to the anterior rotation of the occiput in occipito-posterior positions, and is the natural and favorable termination.

Sometimes, forward rotation does not take place, and the occiput then turns backward into the hollow of the sacrum. What then generally occurs is, that the pains continue, for the reason already mentioned, to depress the chin and produce strong flexion of the face on the sternum, the occiput becoming fixed on the anterior border of the perineum. The pains continuing to act chiefly on the anterior part of the head, the face is born first behind the pubes, the occiput only slipping over the perineum after the forehead has been expelled.

Second Mode in which such Cases occasionally End.—A second mode of termination of such positions is mentioned in most works on the authority of one or two recorded cases; but, although mechanically possible, it is certainly an event of extreme rarity. The chin, instead of being flexed on the sternum, is greatly extended, so that the face of the child looks upward toward the pelvic brim. The child then hitches over the upper edge of the pubes, and becomes fixed there, while the force of the uterine contractions is expended on the posterior part of the head, which descends through the pelvis, distending the perineum, and is born first, the face subsequently following.

Mechanism of Feet Presentations.—The mechanism of the delivery of the body and head in cases in which the feet originally present does

not differ in any important respect from that which has been already described, and requires no separate notice.

Treatment.—From what has been said of the natural mechanism it is evident that one of the most fruitful causes of difficulty and delay is undue interference on the part of the practitioner. It is to be avoided, tempting to use traction on the partially-born trunk in the hope of expediting delivery, but when it is remembered that this is almost certain to produce extension of the arms above the head, and consequently extension of the osipex on the spine, both of which seriously increase the difficulty of delivery, the necessity of leaving the case as much as possible to nature will be apparent.

Having once, therefore, determined the existence of a pelvic presentation, nothing more should be done until the birth of the breech. The membranes should be even more carefully prevented from prematurely rupturing than in vertex presentations, since they serve to dilate the genital passages better than the presenting part. Hence they should be preserved intact, if possible, until they reach the floor of the pelvis, instead of being punctured as soon as the os is fully dilated. The breech when born should be received and supported in the palm of the hand.

Danger to Child.—When the body is expelled as far as the umbilicus the dangers to the child commence, for now the cord is apt to be pressed between the body of the child and the pelvic walls. To obviate this risk as much as possible, a loop of the cord should be pulled down and carried to that part of the pelvis where there is most room, which will generally be opposite one or the other sacro-iliac synchondrosis. As long as the cord is freely pulsating we may be satisfied that the life of the child is not gravely imperilled, although delay is fraught with danger from other sources which have been already indicated. In most cases the arms now slip out, but it may happen, even without any fault on the part of the accoucheur, that they are extended above the head; and it is of great importance that we should be thoroughly acquainted with the best means of liberating them from their abnormal position.

Management when the Arms are Extended above the Head.—They must, of course, never be drawn directly downward, or the almost certain result would be fracture of the fragile bones. We should endeavor to make the arm sweep over the face and chest of the child, so that the natural movements of its joints should not be opposed. If the shoulders be within easy reach, the finger of the accoucheur should be slipped over that which is posterior—because there is likely to be more space for this manœuvre toward the sacrum—and gently carried downward toward the elbow, which is drawn over the face, and then onward, so as to liberate the forearm. The same manœuvre should then be applied to the opposite arm. It may be that the shoulders are not easily reached, and then they may be depressed by altering the position of the child's body. If this be carried well up to the mother's abdomen, the posterior shoulder will be brought lower down; and, by reversing this procedure and carrying the body back over the perineum, the anterior shoulder may be similarly depressed. It is only very exceptionally, however, that these expedients are required.

Birth of the Head.—The arms being extracted, some degree of artificial assistance is at this time almost always required. If there be much delay the child will almost certainly perish. Attempts have been made, in cases in which delivery of the head could not be rapidly effected, to establish pulmonary respiration by passing one or two fingers into the vagina, so as to press it back and admit air to the child's mouth, or by passing a catheter or tube into the mouth. Neither of these expedients is reliable, and we should rather seek to aid nature in completing the birth of the head as rapidly as possible. The first thing to do, supposing the face to have rotated into the cavity of the sacrum, is to carry the body of the child well up toward the pubes and abdomen of the mother without applying any traction, for fear of interfering with the all-important flexion of the chin on the sternum. If now the patient bear down strongly, the natural powers may be sufficient to complete delivery. If there be any delay, traction must be resorted to, and we must endeavor to apply it in such way as to ensure flexion. For this purpose, while the body of the child is grasped by the left hand and drawn upward toward the mother's abdomen, the index and middle fingers of the right hand are placed on the back of the child's neck, so that their tips press on either side of the base of the occiput and push the head into a state of flexion. In most works we are advised to pass the index and middle fingers of the left hand at the same time over the child's face, so as to depress the superior maxilla. Dr. Barnes insists that this is quite unnecessary, and that extraction in the manner indicated, by pressure on the occiput, is quite sufficient. Should it not prove so, flexion of the chin may be very effectually assisted by downward pressure on the forehead through the rectum. One or two fingers of the left hand can readily be inserted into the bowel, and the expulsion of the head is thus materially facilitated.

Value of Pressure through the Abdomen.—By far the most powerful aid, however, in hastening delivery of the head, should delay occur, is pressure from above. This has been, strangely enough, almost altogether omitted by writers on the subject. It has been strongly recommended by Professor Penrose, and there can be no question of its utility. Indeed, as the uterus contracts tightly round the head, uterine expression can be applied almost directly to the head itself, and without any fear of deranging its proper relation to the maternal passages. It is very seldom, indeed, that a judicious combination of traction on the part of the accoucheur, with firm pressure through the abdomen applied by an assistant, will fail in effecting delivery of the head before the delay has had time to prove injurious to the child.

Application of the Forceps to the After-coming Head.—Many accoucheurs—among others Meigs and Rigby—advocate the application of the forceps when there is delay in the birth of the after-coming head. If the delay be due to want of expulsive force in a pelvis of normal size, manual extraction, in the manner just described, will be found to be sufficient in almost every case, and preferable, as being more rapid, easier of execution, and safer to the child. The forceps may be quite properly tried if other means have failed, especially if there be some disproportion between the size of the head and the pelvis.

Management of Sacro-posterior Positions.—Difficulties in delivery may also occur in sacro-posterior positions. Up to the time of the birth of the head the labor usually progresses as readily as in sacro-anterior positions. If the forward rotation of the hips do not take place, much subsequent difficulty may be prevented by gently favoring it by traction applied to the breech during the pains, the finger being passed for this purpose into the fold of the groin.

It is after the birth of the shoulders that the absence of rotation is most likely to prove troublesome. It has been recommended that the body should then be grasped, in the interval between the pains, and twisted round so as to bring the occiput forward. It is by no means certain, however, that the head would follow the movement imparted to the body, and there must be a serious danger of giving a fatal twist of the neck by such a manœuvre. The better plan is to direct the face backward toward the cavity of the sacrum, by pressing on the anterior temple during the continuance of a pain. In this way the proper rotation will generally be effected without much difficulty, and the case will terminate in the usual way.

Management of Cases in which Forward Rotation does not Occur.—If rotation of the occiput forward do not occur, it is necessary for the practitioner to bear in mind the natural mechanism of delivery under such circumstances. In the majority of cases the proper plan is to favor flexion of the chin by upward pressure on the occiput, and to exert traction directly backward, remembering that the nape of the neck should be fixed against the anterior margin of the perineum. If this be not remembered, and traction be made in the axis of the pelvic outlet, the delivery of the head will be seriously impeded. In the rare cases in which the head becomes extended and the chin hitches on the upper margin of the pubis, traction directly forward and upward may be required to deliver the head; but before resorting to it care should be taken to ascertain that backward extension of the head has really taken place.

Management of Impacted Breech Presentations.—It remains for us to consider the measures which may be adopted in those very troublesome cases in which the breech refuses to descend and becomes impacted in the pelvic cavity, either from uterine inertia or from disproportion between the breech and the pelvis. Here, unfortunately, the peculiar shape of the presenting part, which is unadapted for the application of the forceps, renders such cases very difficult to manage.

Two measures have been chiefly employed; 1st, bringing down one or both feet, so as to break up the presenting part and convert it into a footling case; 2d, traction on the breech, either by the fingers, a blunt hook, or fillet passed over the groin.

Bringing Down a Foot.—Barnes insists on the superiority of the former plan, and there can be no question that, if a foot can be got down, the accoucheur has a complete control over the progress of the labor which he can gain in no other way. If the breech be arrested at or near the brim, there will generally be no great difficulty in effecting the desired object. It will be necessary to give chloroform to the extent of complete anæsthesia, and to pass the hand over the child's abdomen

in the same manner and with the same precautions as in performing podalic version until a foot is reached, which is seized and pulled down. If the feet be placed in the usual way close to the buttocks, no great difficulty is likely to be experienced. If, however, the legs be extended on the abdomen, it will be necessary to introduce the hand and arm very deeply, even up to the fundus of the uterus—a procedure which is always difficult, and which may be very hazardous. Nor do I think that the attempt to bring down the feet can be safe when the breech is low down and fixed in the pelvic cavity. A certain amount of repression of the breech is possible, but it is evident that this cannot be safely attempted when the breech is at all low down.

Traction on the Groin.—Under such circumstances traction is our only resource, and this is always difficult and often unsatisfactory. Of all contrivances for this purpose, none is better than the hand of the accoucheur. The index finger can generally be slipped over the groin without difficulty, and traction can be applied during the pains. Failing this or when it proves insufficient, an attempt should be made to pass a fillet over the groins. A soft silk handkerchief or a skein of worsted answers best, but it is by no means easy to apply. The simplest plan, and one which is far better than the expensive instruments contrived for the purpose, is to take a stout piece of copper wire and bend it double into the form of a hook. The extremity of this can generally be guided over the hips, and through its looped end the fillet is passed. The wire is now withdrawn, and carries the fillet over the groins. I have found this simple contrivance, which can be manufactured in a few moments, very useful, and by means of such a fillet very considerable tractive force can be employed. The use of a soft fillet is in every way preferable to the blunt hook which is contained in most obstetric bags. A hard instrument of this kind is quite as difficult to apply, and any strong traction employed by it is almost certain to seriously injure the delicate fetal structures over which it is placed. As an auxiliary the employment of uterine expression should not be forgotten, since it may give material aid when the difficulty is only due to uterine inertia. After a difficult breech labor is completed the child should be carefully examined to see that the bones of the thighs and arms have not been injured. Fractures of the thigh are far from uncommon in such cases, and the soft bones of the newly-born child will readily and rapidly unite if placed at once in proper splints.

Embryotomy.—Failing all endeavors to deliver by these expedients, there is no resource left but to break up the presenting part by scissors or by craniotomy instruments; but, fortunately, so extreme a measure is but rarely necessary.

CHAPTER VI.

PRESENTATIONS OF THE FACE.

PRESENTATIONS of the face are by no means rare, and, although in the great majority of cases they terminate satisfactorily by the unassisted powers of nature, yet every now and again they give rise to much difficulty, and then they may be justly said to be amongst the most formidable of obstetric complications. It is therefore essential that the practitioner should thoroughly understand the natural history of this variety of presentation, with the view of enabling him to intervene with the best prospect of success.

Erroneous Views Formerly Held.—The older accoucheurs had very erroneous views as to the mechanism and treatment of these cases, most of them believing that delivery was impossible by the natural efforts, and that it was necessary to intervene by version in order to effect delivery. Smellie recognized the fact that spontaneous delivery is possible, and that the chin turns forward and under the pubes; but it was not until long after his time, and chiefly after the appearance of Mme. La Chapelle's essay on the subject, that the fact that most cases could be naturally delivered was fully admitted and acted upon.

Frequency.—The frequency of face presentation varies curiously in different countries. Thus, Collins found that in the Rotunda Hospital there was only 1 case in 497 labors, although Churchill gives 1 in 249 as the average frequency in British practice; while in Germany this presentation is met with once in 169 labors. The only reasonable explanation of this remarkable difference is, that the dorsal decubitus, generally followed abroad, favors the translocation of vertex presentations into those of the face.

The mode in which this change is effected—sic it can hardly be doubted that in the large majority of cases face presentation is due to a backward displacement of the occiput after labor has actually commenced, but before the head has engaged in the brim—has been made the subject of various explanations.

Mode in which Face Presentations are Produced.—It has generally been supposed that the change is induced by a hitching of the occiput on the hem of the pelvis, so as to produce extension of the head and descent of the face, the occurrence being favored by the oblique position of the uterus so frequently met with in pregnancy. Hecker attaches considerable importance to a peculiarity in the shape of the fetal head generally observed in face presentations, the cranium having the dolicho-cephalous form, prominent posteriorly, with the occiput projecting, which has the effect of increasing the length of the posterior cranial lever arm, and facilitating extension when circumstances favoring it are in action. Dr. Duncan¹ thinks that uterine obliquity has much influence in the production of face presentation, but in a different way to that above referred

¹ *Edin. Med. Journ.*, vol. xx.

to. He points out that when obliquity is very marked a curve in the genital passages is produced, the convexity of which is directed to the side toward which the uterus is deflected. When uterine contraction commences the fetus is propelled downward, and the part corresponding to the concavity of the curve is acted on to the greatest advantage by the propelling force, and tends to descend. Should the occiput happen to lie in the convexity of the curve so formed, the tendency will be for the forehead to descend. In the majority of cases its descent will be prevented by the increased resistance it meets with, in consequence of the greater length of the anterior cranial lever arm; but if the uterine obliquity be extreme this may be counterbalanced, and a face presentation ensues. The influence of this obliquity is corroborated by the observation of Baudelocque, that the occiput in face presentations almost invariably corresponds to the side of the uterine obliquity. A further corroboration is afforded by the fact that in face presentation the occiput is much more frequently directed to the right than to the left, while right lateral obliquity of the uterus is also much more common.

These theories assume that face presentations are produced during labor. In a few cases they certainly exist before labor has commenced. It is possible, however, as we know that uterine contractions exist independently of actual labor, that similar causes may also be in operation, although less distinctly, before the commencement of labor.

Diagnosis.—The diagnosis is often a matter of considerable difficulty at an early period of labor, before the os is fully dilated and the membranes ruptured, and when the face has not entered the pelvic cavity. The finger then impinges on the rounded mass of the forehead, which may very readily be mistaken for the vertex. At this stage the diagnosis may be facilitated by abdominal palpation in the way suggested by Hecker. If the face is presenting at the brim, palpation will enable us to distinguish a hard, firm, and rounded body immediately above the pubes, which is the forehead and sinciput; on the other side will be felt an indistinct, soft substance, corresponding to the thorax and neck. When labor is advanced and the head has somewhat descended, or when the membranes are ruptured, we should be able to make out the nature of the presentation with certainty. The diagnostic marks to be relied on are the edges of the orbits, the prominence of the nose, the nostrils (their orifices showing to which part of the pelvis the chin is turned), and the cavity of the mouth with the alveolar ridges. If these be made out satisfactorily, no mistake should occur. The most difficult cases are those in which the face has been a considerable time in the pelvis. Under such circumstances the cheeks become greatly swollen and pressed together, so as to resemble the nates. The nose might then be mistaken for the genital organs, and the mouth for the anus. The orbits, however, and the alveolar ridges resemble nothing in the breech, and should be sufficient to prevent error. Considerable care should be taken not to examine too frequently and roughly, otherwise serious injury to the delicate structures of the face might be inflicted. When once the presentation has been satisfactorily diagnosed, examinations should be made as seldom as possible, and only to assure ourselves that the case is progressing satisfactorily.

Mechanism.—If we regard face presentations, as we are fully justified in doing, as being generally produced by the extension of the occiput in what were originally vertex presentations, we can readily understand that the position of the face in relation to the pelvis must correspond to that of the vertex. This is, in fact, what is found to be the case, the forehead occupying the position in which the occiput would have been placed had extension not occurred.

The Positions of the Face correspond to those of the Vertex.—The face, then, like the head, may be placed with its long diameter corresponding to almost any of the diameters of the brim, but most generally it lies either in the transverse diameter or between this and the oblique, while, as it descends in the pelvis, it more generally occupies one or other of the oblique diameters. It is common in obstetric works to describe two principal varieties of face presentation—viz. the right and left mento-iliac—according as the chin is turned to one or other side of the pelvis. It is better, however, to classify the positions in accordance with the part of the pelvis to which the chin points. We may therefore describe four positions of the face, each being analogous to one of the ordinary vertex presentations, of which it is the transformation :

First Position.—The chin points to the right sacro-iliac synchondrosis, the forehead to the left foramen ovale, and the long diameter of the face lies in the right oblique diameter of the pelvis. This corresponds to the first position of the vertex, and, as in that, the back of the child lies to the left side of the mother.

Second Position.—The chin points to the left sacro-iliac synchondrosis, the forehead to the right foramen ovale, and the long diameter of the face lies in the left oblique diameter of the pelvis. This is the conversion of the second vertex position.

Third Position.—The forehead (Fig. 112) points to the right sacro-iliac synchondrosis, the chin to the left foramen ovale, and the long diameter of the face lies in the right oblique diameter of the pelvis. This is the conversion of the third vertex position.

Fourth Position.—The forehead points to the left sacro-iliac synchondrosis, the chin to the right foramen ovale, and the long diameter of the face lies in the left oblique diameter of the pelvis. This is the conversion of the fourth vertex position.

The Relative Frequency of these Positions.—The relative frequency of these presentations is not yet positively ascertained. It is certain that there is not the preponderance of first facial that there is of first vertex positions, and this may no doubt be explained by the supposition that an unusual vertex position may of itself facilitate the transformation into a face presentation. Winckel concludes that, *ceteris paribus*, a face presentation is more readily produced when the back of the child lies to the right than when it lies to the left side of the mother, the reason for this being probably the frequency of right lateral obliquity of the uterus. We shall presently see that, with very rare exceptions, it is absolutely essential that the chin should rotate forward under the pubes before delivery can be accomplished; and therefore we may regard the third and fourth face positions, in which the chin from the first points anteriorly, as more favorable than the first and second.

The mechanism of delivery in face is practically the same as in vertex presentations; and we shall have no difficulty in understanding it if we bear in mind that in face cases the forehead takes the place of, and represents the occiput in, vertex presentations. For the purpose of description we will take the first position of the face.

Description of Delivery in the First Position of the Face.—1. *Extension.*—The first step consists in the extension of the head, which is effected by the uterine contractions as soon as the membranes are ruptured. By this the occiput is still more completely pressed back on the nape of the neck, and the fronto-mental, rather than the mento-breg-

FIG. 112.



Third Position in Face Presentations.

matic diameter is placed in relation to the pelvic brim. This corresponds to the stage of flexion in vertex presentations.

The chin descends below the forehead from precisely the same cause as the occiput in vertex presentations. On account of the extended position of the head the presenting face is divided into portions of unequal length in relation to the vertebral column, through which the force is applied, the longer lever arm being toward the forehead. The resistance is, therefore, greatest toward the forehead, which remains behind while the chin descends.


2. *Descent.*—As the pains continue, the head (the chin being still in advance) is propelled through the pelvis. It is generally said that the face cannot descend, like the occiput, down to the floor of the pelvis, its descent being limited by the length of the neck. There is here, however, an obvious misapprehension. The neck from the chin to the sternum, when the head is forcibly extended, measures from $3\frac{1}{2}$ to 4 inches—a length that is more than sufficient to admit of the face descending to the lower pelvic strait. As a matter of fact, the chin is frequently observed in mento-posterior positions to descend so far that it is appar-

ently endeavoring to pass the perineum before rotation occurs. At the brim the two sides of the face are on a level, but as labor advances the right cheek descends somewhat, the caput succedaneum forms on the malar bone, and, if a secondary caput succedaneum form, on the cheek.

3. *Rotation* is by far the most important point in the mechanism of face presentations; for unless it occurs, delivery with a full-sized head and an average pelvis is practically impossible. There are, no doubt, exceptions to this rule, which must be separately considered, but it is certain that the absence of rotation is always a grave and formidable complication of face presentation. Fortunately, it is only very rarely that this is not effected. The mechanical causes are precisely those which produce rotation of the occiput forward in vertex presentations. As it is accomplished the chin passes under the arch of the pubes, and the occiput rotates into the hollow of the sacrum (Fig. 113); and then commences—

4. *Flexion*, a movement which corresponds to extension in vertex cases. The chin passes as far as it can under the pubic arch, and there

FIG. 113.



Rotation Forward of Chin.

becomes fixed. The uterine force is now expended on the occiput, which revolves, as it were, on its transverse axis (Fig. 114), the under surface of the chin resting on the pubes as a fixed point. This movement goes on until, at last, the face and occiput sweep over the distended perineum.

5. *External rotation* is precisely similar to that which takes place in head presentations, and, like it, depends on the movements imparted to the shoulders.

Mento-posterior Positions in which Rotation does Not take Place.—Such is the natural course of delivery in the vast majority of cases; but in order fully to understand the subject it is necessary to study those rare cases in which the chin points backward and forward rotation does not

occur. These may be taken to correspond to the occipito-posterior positions, in which the face is born looking to the pubes; but, unlike

FIG. 114.



Passage of the Head through the External Parts in Face Presentation.

them, it is only very exceptionally that delivery can be naturally completed. The reason of this is obvious, for the occiput gets jammed behind the pubes, and there is no space for the fronto-mental diameter

FIG. 115.



Illustrating the Position of the Head when Forward Rotation of the Chin does Not take Place.

to pass the antero-posterior diameter of the outlet (Fig. 115). Cases are indeed recorded in which delivery has been effected with the chin looking

posteriorly, but there is every reason to believe that this can only happen when the head is either unusually small or the pelvis unusually large. In such cases the forehead is pressed down until a portion appears at the ostium vaginae, when it becomes firmly fixed behind the pubes, and the chin, after many efforts, slips over the perineum. When this is effected flexion occurs, and the occiput is expelled without difficulty. The forehead is probably always on a lower level than the chin.

Dr. Hicks¹ has published a paper in which he attempts to show that this termination of face presentations is not so rare as is generally supposed, and he gives a single instance in which he effected delivery with the forceps; but he practically admits that special conditions are necessary, such as the "antero-posterior diameter of the outlet particularly ample" and a diminished size of the head. When delivery is effected it is probable, as Cazeaux has pointed out, that the face lies in the oblique diameter of the outlet, and that the chin depresses the soft structures at the side of the sacro-ischiatic notch, which yield to the extent of a quarter of an inch or more, and thereby permit the passage of the occipito-mental diameter of the head. It must, however, be borne well in mind that spontaneous delivery in mento-posterior positions is the rare exception, and that, supposing rotation does not occur—and it often does so at the last moment—artificial aid in one form or another will be almost certainly required.

Prognosis of Face Presentations.—As regards the mother, in the great majority of cases the prognosis is favorable, but the labor is apt to be prolonged, and she is therefore more exposed to the risks attending tedious delivery. As regards the child, the prognosis is much more unfavorable than in vertex presentations. Even when the anterior rotation of the chin takes place in the natural way, it is estimated that 1 out of 10 children is stillborn, while if not the death of the child is almost certain. This increased infantile mortality is evidently due to the serious amount of pressure to which the child is subjected, and probably depends in many cases on cerebral congestion, produced by pressure on the jugular veins as the neck lies in the pelvic cavity. Even when the child is born alive, the face is always greatly swollen and disfigured. In some cases the deformity produced in this way is excessive and the features are often scarcely recognizable. This disfiguration passes away in a few days, but the practitioner should be aware of the probability of its occurrence, and should warn the friends, or they might be unnecessarily alarmed, and possibly might lay the blame on him.

Treatment.—After what has been said as to the mechanism of delivery in face presentation, it is obvious that the proper course is to leave the case alone, in the expectation of the natural efforts being sufficient to complete delivery. Fortunately, in the large majority of cases this course is attended by a successful result.

The older accoucheurs, as has been stated, thought active interference absolutely essential, and recommended either podalic version or the attempt to convert the case into a vertex presentation by inserting the hand and bringing down the occiput. The latter plan was recommended by Baudelocque, and is even yet followed by some accoucheurs. Thus, Dr.

¹ *Obst. Trans.*, vol. vii.

Hodge¹ advises it in all cases in which face presentation is detected at the brim; but, although it might not have been attended with evil consequences in his experienced hands, it is certainly altogether unnecessary, and would infallibly lead to most serious results if generally adopted. It may, however, be allowable in certain cases in which the face remains above the brim and refuses to descend into the pelvic cavity. Even then it is questionable whether podalic version should not be preferred, as being easier of performance, giving, when once effected, a much more complete control over delivery, and being less painful to the mother. Version is certainly preferable to the application of the forceps, which are introduced with difficulty in so high a position of the face, and do not take a secure hold.

Rectification by Abdominal Palpation.—Schatz² has more recently suggested the rectification of face presentations at an early stage, before the rupture of the membranes, by manipulation through the abdomen. He raises the fetal body by pressure on the shoulder and breast through the abdominal wall by one hand, while the breech is raised and steadied by the other. By this means the occiput is elevated, and then the breech is pressed downward, when head flexion is produced by the resistance of the pelvic walls. Of this method I have had no practical experience, but it obviously requires an unusual amount of skill and practice in abdominal palpation.

Difficulties from Arrest in the Pelvic Cavity.—When once the face has descended into the pelvis, difficulties may arise from two chief causes—uterine inertia and non-rotation forward of the chin.

The treatment of the former class must be based on precisely the same general principles as in dealing with protracted labor in vertex presentations. The forceps may be applied with advantage, bearing in mind the necessity of getting the chin under the pubes, and, when this has been effected, of directing the traction forward, so as to make the occiput slowly and gradually distend and sweep over the perineum.

Difficulties arising from Non-rotation of Chin Forward.—The second class of difficult face cases are much more important, and may try the resources of the accoucheur to the utmost. Our first endeavor must be, if possible, to secure the anterior rotation of the chin. For this purpose various manœuvres are recommended. By some we are advised to introduce the finger cautiously into the mouth of the child and draw the chin forward during a pain; by others, to pass the finger up behind the occiput and press it backward during the pain. Schroeder points out that the difficulty often depends on the fact of the head not being sufficiently extended, so that the chin is not on a lower level than the forehead, and that rotation is best promoted by pressing the forehead upward with the finger during a pain, so as to cause the chin to descend. Penrose³ believes that non-rotation is generally caused by the want of a *point d'appui* below, on account of the face being unable to descend to the floor of the pelvis, and that if this is supplied rotation will take place. In such cases he applies the hand or the blade of the forceps, so as to press on the posterior cheek. By this means the neces-

¹ *System of Obstetrics*, p. 335.

³ *Amer. Supplement to Obst. Journ.*, April, 1876.

² *Arch. f. Gyn.*, B. v. 313.

sary *point d'appui* is given; and he relates several interesting cases in which this simple manoeuvre was effectual in rapidly terminating a previously lengthy labor. Any or all of these plans may be tried. We must bear in mind, in using them, that rotation is often delayed until the face is quite at the lower pelvic strait, so that we need not too soon despair of its occurring. If, however, in spite of these manoeuvres, it do not take place, what is to be done? If the head be not too low down in the pelvis to admit of version, that would be the simplest and most effectual plan. I have succeeded in delivering in this way when all attempts at producing rotation had failed; but generally the face will be too decidedly engaged to render it possible. An attempt might be made to bring down the occiput by the vectis or by a fillet, but if the face be in the pelvic cavity it is hardly possible for this plan to succeed. An endeavor may be made to produce rotation by the forceps; but it should be remembered that rotation of the face mechanically in this way is very difficult, and much more likely to be attended with fatal consequences to the child than when it is effected by the natural efforts. In using forceps for this purpose, the second or pelvic curve is likely to prove injurious, and a short straight instrument is to be preferred. If rotation be found to be impossible, an endeavor may be made to draw the face downward, so as to get the chin over the perineum and deliver in the mento-posterior position; but, unless the child be small or the pelvis very capacious, the attempt is unlikely to succeed. Finally, if all these means fail there is no resource left but lessening the size of the head by craniotomy—a *dernier ressort* which, fortunately, is very rarely required.

Brow Presentations.—It sometimes happens that the head is partially extended, so as to bring the os frontis into the brim of the pelvis and form what is described as a "*brow presentation*." Should the head descend in this manner, the difficulties, although not insuperable, are apt to be very great, from the fact that the long cervico-frontal diameter of the head is engaged in the pelvic cavity. The diagnosis is not difficult, for the os frontis will be detected by its rounded surface, while the anterior fontanelle is within reach in one direction, the orbit and root of the nose in another.

Spontaneously Converted into Face or Vertex Presentations.—Fortunately, in the large majority of cases brow presentations are spontaneously converted into either vertex or face presentations, according as flexion or extension of the head occurs; and these must be regarded as the desirable terminations and the ones to be favored. For this purpose upward pressure must be made on one or other extremity of the presenting part during a pain, so as to favor flexion or extension; or, if the parts be sufficiently dilated, an attempt may be made to pass the hand over the occiput and draw it down, thus performing cephalic version. The latter is the plan recommended by Hodge, who describes the operation as easy. It is questionable, however, if a well-marked brow presentation be distinctly made out while the head is still at the brim, whether podalic version would not be the easiest and best operation. If the forehead have descended too low for this, and if the endeavor to convert it into either a face or vertex presentation fail, the forceps will probably be required. In such cases the face generally turns toward

the pubes, the superior maxilla becomes fixed behind the pubic arch, and the occiput sweeps over the perineum. Very great difficulty is likely to be experienced, and, if conversion into either a vertex or face presentation cannot be effected, craniotomy is not unlikely to be required.

CHAPTER VII.

DIFFICULT OCCIPITO-POSTERIOR POSITIONS.

A FEW words may be said in this place as to the management of occipito-posterior positions of the head, especially of those in which forward rotation of the occiput does not take place. It has already been pointed out that in the large majority of these cases the occiput rotates forward without any particular difficulty, and the labor terminates in the usual way, with the occiput emerging under the arch of the pubes.

Rotation Forward of the Occiput.—In a certain number of cases such rotation does not occur, and difficulty and delay are apt to follow. The proportion of cases in which face-to-pubes terminations of occipito-posterior positions occur has been variously estimated, and they are certainly more common than most of our text-books lead us to expect. Dr. Uvedale West,¹ who studied the subject with great care, found that labor ended in this way in 79 out of 2585 births, all these deliveries being exceptionally difficult.

Causes of Face-to-Pubes Delivery.—He believed that forward rotation of the head is prevented by the absence of flexion of the chin on the sternum, so that the long occipito-frontal, instead of the short sub-occipito-bregmatic, diameter of the head is brought into contact with the pelvic diameter; hence the occiput is no longer the lowest point, and is not subjected to the action of those causes which produce forward rotation. Dr. Macdonald, who has written a thoughtful paper on the subject,² believes that the non-rotation forward of the occiput is chiefly due to the large size of the head, in consequence of which "the forehead gets so wedged into the pelvis anteriorly that its tendency to slacken and rotate forward does not come into play." Dr. West's interpretation, which has an important bearing on the management of these cases, seems to explain most correctly the non-occurrence of the natural rotation.

The important question for us to decide is, How can we best assist in the management of cases of this kind when difficulties arise and labor is seriously retarded?

Mode of Treatment.—Dr. West, insisting strongly on the necessity of complete flexion of the chin on the sternum, advises that this should be favored by upward pressure on the frontal bone, with the view of causing the chin to approach the sternum and the occiput to descend, and thus to come within the action of the agencies which favor rotation. Supposing

¹ *Cranial Presentations*, p. 33.

² *Edin. Med. Journ.*, Oct., 1874.

the pains to be strong and the fontanelle to be readily within reach, we may in this way very possibly favor the descent of the occiput, and without injuring the mother or increasing the difficulties of the case in the event of the manœuvre failing. The beneficial effects of this simple expedient are sometimes very remarkable. In two cases in which I recently adopted it labor, previously delayed for a length of time without any apparent progress, although the pains were strong and effective, was in each instance rapidly finished almost immediately after the upward pressure was applied. The rotation of the face backward may at the same time be favored by pressure on the pubic side of the forehead during the pains.

Traction on the Occiput.—Others have advised that the descent of the occiput should be promoted by downward traction, applied by the vectis or fillet. The latter is the plan specially advocated by Hodge,¹ and the fillet certainly finds one of its most useful applications in cases of this kind, as being simpler of application and probably more effective than the vectis.

Over-active Endeavors at Assistance should be Avoided.—Although any of these methods may be adopted, a word of caution is necessary against prolonged and over-active endeavors at producing flexion, and rotation when that seems delayed. All who have watched such cases must have observed that rotation often occurs spontaneously at a very advanced period of labor, long after the head has been pressed down for a considerable time to the very outlet of the pelvis, and when it seems to have been making fruitless endeavors to emerge; so that a little patience will often be sufficient to overcome the difficulty.

When Necessary, the Forceps may be Used.—In the event of assistance being absolutely required there is no reason why the forceps should not be used. The instrument is not more difficult to apply than under ordinary circumstances, nor, as a rule, is much more traction necessary. Dr. Macdonald, indeed, in the paper already alluded to, maintains that in persistent occipito-posterior positions there is almost always a want of proportion between the head and the pelvis, and that, therefore, the forceps will be generally required; and he prefers them to any artificial attempts at rectification. Some peculiarities in the mode of delivery are necessary to bear in mind. In most works it is taught that the operator should pay special attention to the rotation of the head, and should endeavor to impart this movement by turning the occiput forward during extraction. Thus, Tyler Smith says: "In delivery with the forceps in occipito-posterior presentations the head should be slowly rotated during the process of extraction, so as to bring the vertex toward the pubic arch, and thus convert them into occipito-anterior presentations." The danger accompanying any forcible attempt at artificial rotation will, however, be evident on slight consideration. It is true that in many cases, when simple traction is applied, the occiput will of itself rotate forward, carrying the instrument with it. But that is a very different thing from forcibly twisting round the head with the blades of the forceps, without any assurance that the body of the child will follow the movement. It is impossible to conceive that such violent interference

¹ *System of Obstetrics*, p. 308.

should not be attended with serious risk of injury to the neck of the child. If rotation do not occur, the fair inference is that the head is so placed as to render delivery with the face to the pubes the best termination, and no endeavor should be made to prevent it. This rule of leaving the rotation entirely to nature, and using traction only, has received the approval of Barnes and most modern authorities, and is the one which recommends itself as the most scientific and reasonable.

Objection to Curved Instruments in such Cases.—There are cases in which the pelvic curve of the forceps is of doubtful utility. When applied in the usual way the convexity of the blades points backward. If rotation accompany extraction, the blades necessarily follow the movement of the head, and their convex edges will turn forward. It certainly seems probable that such a movement would subject the maternal soft parts to considerable risk. I have, however, more than once seen such rotation of the instrument happen without any apparent bad result; but the dangers are obvious. Hence it would be a wise precaution either to use a pair of straight forceps for this particular operation, or to remove the blades and leave the case to be terminated by the natural powers when the head is at the lower strait and rotation seems about to occur. When there is no rotation, more than usual care should be taken with the perineum, which is necessarily much stretched by the rounded occiput. Indeed, the risk to the perineum is very considerable, and even with the greatest care it may be impossible to avoid laceration.

Bearing these precautions in mind, delivery with the forceps in occipito-posterior positions offers no special difficulties or dangers.

[*Version by the Vertex.*—In order to adapt this section to American practice, I addressed letters of inquiry upon the management of occipito-posterior positions to several obstetrical professors and teachers, and have prepared these instructions in accordance with their views.

1. "In *primitive* oblique occipito-posterior positions of the head nature will almost, without exception, cause spontaneous rotation of the occiput to the symphysis pubis; but to favor this movement the bag of waters should be preserved."

2. "Spontaneous rotation, as a rule, does not begin until the head meets with resistance from the floor of the pelvis; hence no effort to force rotation should be made until Nature has proved herself inadequate."

3. Where rotation forward is prevented, it is probably due to the position of the occiput having been originally directly backward, and only becoming oblique after the descent of the head into the pelvis, the position of the child's body preventing the anterior movement of its occiput. That is, the sixth position of Hodge has changed into a fourth or fifth, but will not without assistance become a first or second.

4. If, then, rotation is not spontaneous after the head reaches the floor of the pelvis, version by the vertex will not take place except it be forced by the vectis or forceps.

One professor writes: "I have thus far succeeded so well" (*i. e.* by the vectis and forceps) "that I recall but one instance in which the head was born with the occiput looking to the sacrum." Another says he applies the forceps and lets "the progress of the head determine the mode by which it shall make its exit, not trying to turn by the forceps."

In the *primitive occipito-sacral position changed to oblique*, or in the more rare unchanged sixth position of Hodge, if the head is large or the pelvis in any way obstructed, the case may require to be terminated by craniotomy. It is even possible to rotate the occiput from the sacrum to the pubes and save the child, as this was once done by the late Dr. William Harris of Philadelphia. Of course the body must have partly rotated.

Use of the Hand in Occipito-posterior Positions.—The introduction of the hand for the purpose of effecting version by the vertex, under an anæsthetic, was strongly advocated by the late Dr. John S. Parry¹ of Philadelphia, who certainly used his own, which was small and thin, to very great advantage. Several very small-handed accoucheurs in this city have found their hands of very great value in some cases of obstetrics; and it is said that a celebrated Neapolitan obstetrician owes his great popularity to the advantage thus derived. It will not do to advocate a general use of the hand in obstetric practice, as few have such as it would be safe to use, especially in primiparæ. I have known a primipara labor for hours to deliver herself of a fœtus in an occipito-posterior position, when all that was needed was the assistance of a suitable hand during three labor-pains to bring the occiput fairly under the arch of the pubis.—ED.]

CHAPTER VIII.

PRESENTATIONS OF THE SHOULDER, ARM, OR TRUNK—COMPLEX PRESENTATIONS—PROLAPSE OF THE FUNIS.

In the presentations already considered the long diameter of the fœtus corresponded with that of the uterine cavity, and in all of them the birth of the child by the maternal efforts was the general and normal termination of labor. We have now to discuss those important cases in which the long diameter of the fœtus and uterus do not correspond, but in which the long fetal diameter lies obliquely across the uterine cavity. In the large majority of these it is either the shoulder or some part of the upper extremity that presents: for it is an admitted fact that, although other parts of the body, such as the back or abdomen, may, in exceptional cases, lie over the os at an early period of labor, yet as labor progresses such presentations are almost always converted into those of the upper extremity.

For all practical purposes, we may confine ourselves to a consideration of *shoulder presentations*, the further subdivision of these into *oblique* or *head presentations* being no more necessary than the division of *pelvic presentations* into *breech*, *knee*, and *footling* ones, since the mechanism and management are identical whatever part of the upper extremity presents.

[¹ See *Trans. American Med. Assn.*]

Delivery by the Natural Powers is Quite Exceptional.—There is this great distinction between the presentations we are now considering and those already treated of: that, on account of the relations of the fœtus to the pelvis, delivery by the natural powers is impossible, except under special and very unusual circumstances that can never be relied upon. Intervention on the part of the accoucheur is therefore absolutely essential, and the safety of both the mother and child depends upon the early detection of the abnormal position of the fœtus; for the necessary treatment, which is comparatively easy and safe before labor has been long in progress, becomes most difficult and hazardous if there have been much delay.

Position of the Fœtus.—Presentations of the upper extremity or trunk are often spoken of as “*transverse presentations*” or “*cross-births* ;” but both of these terms are misleading, as they imply that the fœtus is placed transversely in the uterine cavity or that it lies directly across the pelvic brim. As a matter of fact, this is never the case, for the child lies obliquely in the uterus—not indeed in its long axis, but in one intermediate between its long and transverse diameters.

Divided into Dorso-anterior and Dorso-posterior Positions.—Two great divisions of shoulder presentations are recognized: the one in

FIG. 116.



Dorso-anterior Presentation of the Arm.

which the back of the child looks to the abdomen of the mother (Fig. 116), and the other in which the back of the child is turned toward the spine of the mother (Fig. 117). Each of these is subdivided into two subsidiary classes, according as the head of the child is placed in the right or left iliac fossa. Thus, in dorso-anterior positions, if the head lie in the left iliac fossa, the right shoulder of the child presents; if in the right iliac fossa, the left. So in dorso-posterior positions, if the head lie in the left iliac fossa, the left shoulder presents; if in the right, the right. Of the two classes, the dorso-anterior positions are more common, in the proportion, it is said, of two to one.

Causes.—The causes of shoulder presentation are not well known. Amongst those most commonly mentioned are prematurity of the fœtus and excess of liquor amnii; either of these, by increasing the mobility of the fœtus in utero, would probably have considerable influence. The fact that it occurs much more frequently amongst premature births has

FIG. 117.



Dorso-posterior Presentation of the Arm.

long been recognized. Undue obliquity of the uterus has probably some influence, since the early pains might cause the presenting part to hitch against the pelvic brim and the shoulder to descend. An unusually low attachment of the placenta to the inferior segment of the uterine cavity has been mentioned as a predisposing cause. In consequence of this the head does not lie so readily in the lower uterine segment, and is apt to slip up into one of the iliac fossæ. This is supposed to explain the frequency of arm presentation in cases of partial or complete placenta prævia. Danyau and Wigand believe that shoulder presentations are favored by irregularity in the shape of the uterine cavity, especially a relative increase in its transverse diameter. This theory has been generally discredited by writers, and it is certainly not susceptible of proof; but it seems far from unlikely that some peculiarity of shape may exist, not capable of recognition, but sufficient to influence the position of the fœtus. How otherwise are we to explain those remarkable cases, many of which are recorded, in which similar malpositions occurred in many successive labors? Thus, Joulin refers to a patient who had an arm presentation in three successive pregnancies, and to another who had shoulder presentation in three out of four labors. Certainly, such constant recurrences of the same abnormality could only be explained on the hypothesis of some very persistent cause, such as that referred to. Pinard¹ states that shoulder presentations are seven times more common

¹ *Annal. d'Hyg. Pub. et de Mèd.*, Jan., 1879.

in multiparæ than in primiparæ, in consequence, as he believes, of the laxity of the abdominal walls in the former, which allows the uterus to fall forward, and thus prevents the head entering the pelvic brim in the latter weeks of pregnancy. It is probable that merely accidental causes have most influence in the production of shoulder presentation, such as falls or undue pressure exerted on the abdomen by badly-fitting or tight stays. Partially transverse positions during pregnancy are certainly much more common than is generally believed, and may often be detected by abdominal palpation. The tendency is for such malpositions to be righted either before labor sets in or in the early period of labor; but it is quite easy to understand how any persistent pressure, applied in the manner indicated, may perpetuate a position which otherwise would have been only temporary.

Prognosis and Frequency.—According to Churchill's statistics, shoulder presentations occur about once in 260 cases; that is, only slightly less frequently than those of the face. The prognosis to both the mother and child is much more unfavorable; for he estimates that, out of 235 cases, 1 in 9 of the mothers and half the children were lost. The prognosis in each individual case will, of course, vary much with the period of delivery at which the malposition is recognized. If detected early, interference is easy and the prognosis ought to be good; whereas there are few obstetric difficulties more trying than a case of shoulder presentation in which the necessary treatment has been delayed until the presenting part has been tightly jammed into the cavity of the pelvis.

Diagnosis.—Bearing this fact in mind, the paramount necessity of an accurate diagnosis will be apparent; and it is specially important that we should be able not only to detect that a shoulder or arm is presenting, but that we should, if possible, determine which it is, and how the body and head of the child are placed. The existence of a shoulder presentation is not generally suspected until the first vaginal examination is made during labor. The practitioner will then be struck with the absence of the rounded mass of the fetal head, and, if the os be open and the membranes protruding, by their elongated form, which is common to this and to other malpresentations. If the presenting part be too high to reach, as is often the case at an early period of labor, an endeavor should at once be made to ascertain the fetal position by abdominal examination. This is the more important as it is much more easy to recognize presentations of the shoulder in this way than those of the breech or foot; and at so early a period it is often not only possible, but comparatively easy, to alter the position of the fœtus by abdominal manipulation alone, and thus avoid the necessity of the more serious form of version. The method of detecting a shoulder presentation by examination of the abdomen has already been described (p. 124), and need not be repeated. The chief points to look for are the altered shape of the uterus and two solid masses, the head and the breech, one in either iliac fossa. The facility with which these parts may be recognized varies much in different patients. In thin women with lax abdominal parietes they can be easily felt, while in very stout women it may be impossible. Failing this method, we must rely on vaginal examinations, although

before the membranes are ruptured, and when the presenting part is high in the pelvis, it is not always easy to gain accurate information in this way. The difficulty is increased by the paramount importance of retaining the membranes intact as long as possible. It should be remembered, therefore, that when a presentation of the superior extremity is suspected, the necessary examinations should only be made in the intervals between the pains, when the membranes are lax, and never when they are rendered tense by the uterine contractions.

As either the shoulder, the elbow, or the hand may present, it will be best to describe the peculiarities of each separately, and the means of distinguishing to which side of the body the presenting part belongs.

1. *Peculiarities of the Shoulder.*—The shoulder is recognized as a round, smooth prominence, at one point of which may often be felt the sharp edge of the acromion. If the finger can be passed sufficiently high, it may be possible to feel the clavicle and the spine of the scapula. A still more complete examination may enable us to detect the ribs and the intercostal spaces, which would be quite conclusive as to the nature of the presentation, since there is nothing resembling them in any other part of the body. At the side of the shoulder the hollow of the axilla may generally be made out.

Mode of Diagnosing the Position of the Child.—In order to ascertain the position of the child we have to find out in which iliac fossa the head lies. This may be done in two ways: 1st, the head may be felt through the abdominal parietes by palpation; and 2d, since the axilla always points toward the feet, if it point to the left side the head must lie in the right iliac fossa; if to the right, the head must be placed in the left iliac fossa. Again, the spine of the scapula must correspond to the back of the child, the clavicle to its abdomen; and by feeling one or other we know whether we have to do with a dorso-anterior or dorso-posterior position. If we cannot satisfactorily determine the position by these means, it is quite legitimate practice to bring down the arm carefully, provided the membranes are ruptured, so as to examine the hand, which will be easily recognized as right or left. This expedient will decide the point, but it is one which it is better to avoid if possible, for it not only slightly increases the difficulty of turning, although perhaps not very materially, but the arm might possibly be injured in the endeavor to bring it down.

Differential Diagnosis of the Shoulder.—The only part of the body likely to be taken for the shoulder is the breech; but in that its larger size, the groove in which the genital organs lie, the second prominence formed by the other buttock, and the sacral spinous processes are sufficient to prevent a mistake.

2. *The Elbow.*—The elbow is rarely felt at the os, and may be readily recognized by the sharp prominence of the olecranon, situated between two lesser prominences, the condyles. As the elbow always points toward the feet, the position of the fœtus can be easily ascertained.

3. *The Hand.*—The hand is easy to recognize, and can only be confounded with the foot. It can be distinguished by its borders being of the same thickness, by the fingers being wider apart and more readily separated from each other than the toes, and above all by the mobility

of the thumb, which can be carried across the palm and placed in apposition with each of the fingers.

Mode of Detecting which Hand is Presenting.—It is not difficult to tell which hand is presenting. If the hand be in the vagina or beyond the vulva, and within easy reach, we recognize which it is by laying hold of it as if we were about to shake hands. If the palm lie in the palm of the practitioner's hand with the two thumbs in apposition, it is the right hand; if the back of the hand, it is the left. Another simple way is for the practitioner to imagine his own hand placed in precisely the same position as that of the fœtus, and this will readily enable him to verify the previous diagnosis. A simple rule tells us how the body of the child is placed, for, provided we are sure the hand is in a state of supination, the back of the hand points to the back of the child, the palm to its abdomen, the thumb to the head, and the little finger to the feet.

Mechanism.—It is perhaps hardly proper to talk of a mechanism of shoulder presentations, since if left unassisted they almost invariably lead to the gravest consequences. Still, Nature is not entirely at fault even here, and it is well to study the means she adopts to terminate these malpositions.

Terminations.—There are two possible terminations of shoulder presentation. In one, known as "*spontaneous version*," some other part of the fœtus is substituted for that originally presenting; in the other, "*spontaneous evolution*," the fœtus is expelled by being squeezed through the pelvis, without the originally presenting part being withdrawn. It cannot be too strongly impressed on the mind that neither of these can be relied on in practice.

Spontaneous version may occasionally occur before or immediately after the rupture of the membranes, when the fœtus is still readily movable within the cavity of the uterus. A few authenticated cases are recorded in which the same fortunate issue took place after the shoulder had been engaged in the pelvic brim for a considerable time, or even after prolapse of the arm; but its probability is necessarily much lessened under such circumstances. Either the head or the breech may be brought down to the os in place of the original presentation.

The precise mechanism of spontaneous version, or the favoring circumstances, are not sufficiently understood to justify any positive statement with regard to it.

Cazeaux believed that it is produced by partial or irregular contraction of the uterus, one side contracting energetically, while the other remains inert or only contracts to a slight degree. To illustrate how this may effect spontaneous version, let us suppose that the child is lying with the head in the left iliac fossa. Then, if the left side of the uterus should contract more forcibly than the right, it would clearly tend to push the head and shoulder to the right side until the head came to present instead of the shoulder. A very interesting case is related by Geneuil¹ in which he was present during spontaneous version, in the course of which the breech was substituted for the left shoulder more than four hours after the rupture of the membranes. In this case the

¹ *Ann. de Gynécologie*, vol. v., 1876.

uterus was so tightly contracted that version was impossible. He observed the side of the uterus opposite the head contracting energetically, the other remaining flaccid, and eventually the case ended without assistance, the breech presenting. The natural moulding action of the uterus, and the greater tendency of the long axis of the child to lie in that of the uterus, no doubt assist the transformation; and much must depend on the mobility of the foetus in any individual case.

That such changes often take place in the latter weeks of pregnancy, and before labor has actually commenced, is quite certain, and they are probably much more frequent than is generally supposed. When spontaneous version does occur, it is of course a more favorable event; and the termination and prognosis of the labor are then the same as if the head or breech had originally presented.

Spontaneous Evolution.—The mechanism of spontaneous evolution, since it was first clearly worked out by Douglas, has been so often and carefully described that we know precisely how it occurs. Although every now and then a case is recorded in which a living child has been born by this means, such an event is of extreme rarity; and there is no doubt of the accuracy of the general opinion, that spontaneous evolution can only happen when the pelvis is unusually roomy and the child small, and that it almost necessarily involves the death of the foetus, on account of the immense pressure to which it is subjected.

Two varieties are described, in one of which the head is first born, in the other the breech; in both the originally presenting arm remained prolapsed. The former is of extreme rarity, and is believed only to have happened with very premature children, whose bodies were small and flexible, and when traction had been made on the presenting arm. Under such circumstances it can hardly be called a natural process, and we may confine our attention to the latter and more common variety.

What takes place is as follows: The presenting arm and shoulder are tightly jammed down, as far as is possible, by the uterine contractions, and the head becomes strongly flexed on the shoulder. As much of the body of the foetus as the pelvis will contain becomes engaged, and then a movement of rotation occurs which brings the body of the child nearly into the antero-posterior diameter of the pelvis (Fig. 118). The shoulder projects under the arch of the pubes, the head lying above the symphysis and the breech near the sacro-iliac synchondrosis. It is essential that the head should lie forward above the pubes, so that the length of the neck may permit the shoulder to project under the pubic arch, without any part of the head entering the pelvic cavity. The shoulder and neck of the child now become fixed points round which the body of the child rotates, and the whole force of the uterine contractions is expended on the breech. The latter, with the body, therefore becomes more and more depressed, until at last the side of the thorax reaches the vulva, and, followed by the breech and inferior extremities, is slowly pushed out. As soon as the limbs are born the head is easily expelled.

The enormous pressure to which the body is subjected in this process can readily be understood. As regards the practical bearings of this termination of shoulder presentations, all that need be said is, that if we should happen to meet with a case in which the shoulder and thorax

were so strongly depressed that turning was impossible, and in which it seemed that nature was endeavoring to effect evolution, we should be justified in aiding the descent of the breech by traction on the groin before resorting to the difficult and hazardous operation of embryotomy or decapitation.

Treatment.—It is unnecessary to describe specially the treatment of shoulder presentation, since it consists essentially in performing the operation of turning, which is fully described elsewhere. It is only

FIG. 118.



Spontaneous Evolution. (After Chiara of Milan.)

This drawing was made from a patient who died undelivered, the body being frozen and bisected.

needful here to insist on the advisability of performing the operation in the way which involves the least interference with the uterus. Hence if the nature of the case be detected before the membranes are ruptured, an endeavor should be made—and ought generally to succeed—to turn by external manipulation only. If we can succeed in bringing the breech or head over the os in this way, the case will be little more troublesome than an ordinary presentation of these parts. Failing in this, turning by combined external and internal manipulation should be attempted, and the introduction of the entire hand should be reserved for those more troublesome cases in which the waters have long drained away and in which both these methods are inapplicable.

Should all these means fail, we must resort to the mutilation of the child by embryulcia or decapitation, probably the most difficult and dangerous of all obstetric operations. [1]

Complex Presentations.—There are various so-called *complex presentations* in which more than one part of the foetal body presents. Thus we may have a hand or a foot presenting with the head, or a foot and hand presenting simultaneously. The former does not necessarily give rise to any serious difficulty, for there is generally sufficient room for the head to pass. Indeed, it is unlikely that either the hand or foot should enter the pelvic brim with the head unless the head was unusually small or the pelvis more than ordinarily capacious. As regards treatment, it is no doubt advisable to make an attempt to replace the hand or foot by pushing it gently above the head in the intervals between the pains, and maintaining it there until the head be fully engaged in the pelvic cavity. The engagement of the head can be hastened by abdominal pressure, which will prove of great value. Failing this, all we can do is to place the presenting member at the part of the pelvis where it will least impede the labor and be the least subjected to pressure; and that will generally be opposite the temple of the child. As it must obstruct the passage of the head to a certain extent, the application of the forceps may be necessary. When the feet and hands present at the same time, in addition to the confusing nature of the presentation from so many parts being felt together, there is the risk of the hands coming down and converting the case into one of arm presentation. It is the obvious duty of the accoucheur to prevent this by ensuring the descent of the feet, and traction should be made on them, either with the fingers or with a lac, until their descent and the ascent of the hands are assured.

Dorsal Displacement of the Arm.—In connection with this subject may be mentioned the 'curious dorsal displacement of the arm, first

[1] Accoucheurs and surgeons have thought proper to perform the Cesarean operation in the United States in 11 cases where the fetus was dead and impacted in the pelvis in a transverse position, and of these 8 were saved, or 72.7 per cent. In 9 of the cases the arm protruded. In 3 the pelvis was small, but in two of these, small children have since been delivered alive naturally. In no case in the list was there any evidence of pelvic disease. In a twelfth case, where there had been rickets in childhood, and in which the arm protruded, there was likewise a saving of the woman; but this one is excluded from the record, on account of the true cause of dystocia not having been the position of the fetus. Case 3 was in labor 96 hours, 3 days of this time under a midwife, and died of exhaustion in 17 hours. Case 7 was also in the care of a midwife, and died of exhaustion in 12 hours, having been much prostrated at the time of the operation. Case 9 was in a fair way to recover when her husband came home intoxicated, and she arose from her bed to protect her mother from him. This fright, excitement, and exertion caused her death in a few hours three and a half days after the operation.

The celebrated case of Le Bras of Moullyron in 1789, in which the uterus was for the first time sutured, was one of arm protrusion, with impaction, after three days' labor under a midwife: this patient also recovered. Le Bras was much censured for his operation, although Dr. Lacombe had labored a long time to deliver per rits antea, before he was called in, and particularly for the additional risk of inserting a movable suture in the uterus. Certainly the results of these American operations are encouraging in this class of cases. Case 11 was operated upon in June, 1880, and had two silver-wire uterine sutures inserted: she was well in a month, pregnant in two and a half more, and bore a child naturally in twelve and a half months after the operation. Our cautious and conscientious show any more favorable results. —[Ed.]


described by Sir James Simpson,¹ in which the forearm of the child becomes thrown across and behind the neck. The result is the formation of a ridge or bar, which prevents the descent of the head into the pelvis by hitching against the brim (Fig. 119). The difficulty of diagnosis is very great, for the cause of obstruction is too high up to be felt. But if we meet with a case in which the pelvis is roomy and the pains strong, and yet the head does not descend after an adequate time, a full explanation of the cause is essential. For this purpose we would naturally put the patient under chloroform and pass the hand sufficiently high.

FIG. 119.



Dorsal Displacement of the Arm.

FIG. 120.



Dorsal Displacement of the Arm in Footling Presentations. (After Barnes.)

We might then feel the arm in its abnormal position. That was what took place in a case under my own care, in which I failed to get the head through the brim with the forceps, and eventually delivered by turning. The same course was adopted by my friend Mr. Jardine Murray in a similar case.² Simpson advises that the arm should be brought down, so as to convert the case into an ordinary hand-and-head presentation. This, if the arm be above the brim, must always be difficult, and I believe the simpler and more effective plan is podalic version. A similar displacement may cause some difficulty in breech presentations and after turning (Fig. 120). Delay here is easier of diagnosis, since the obstacle to the expulsion will at once lead to careful examination. By carrying the body of the child well backward, so as to enable


¹ *Selected Obstet. Works*, vol. i.

² *Med. Times and Gaz.*, 1861.

the finger to pass behind the symphysis pubis and over the shoulder, it will generally be easy to liberate the arm.

Prolapse of the Umbilical Cord.—It occasionally happens that the umbilical cord falls down past the presenting part (Fig. 121), and is apt to be pressed between it and the walls of the pelvis. The consequence is, that the foetal circulation is seriously interfered with, and the death

FIG. 121.



Prolapse of the Umbilical Cord.

of the child from asphyxia is a common result. Hence prolapse of the funis is a very serious complication of labor in so far as the child is concerned.

Frequency.—Fortunately, it is not a very frequent occurrence. Churchill calculates that out of over 105,000 deliveries it was met with once in 240 cases, and Scanzoni once in 254. Its frequency varies much under different circumstances and in different places. We find from Churchill's figures a remarkable difference in the proportional number of cases observed in France, England, and Germany—viz. 1 in 446½, 1 in 207½, and 1 in 156, respectively. Great as is the proportion referred to Germany in these figures, it has been found to be exceeded in special districts. Thus, Engelmann records 1 case out of 94 labors in the Lying-in Hospital at Berlin, and Michaelis 1 in 90 in that of Kiel. These remarkable differences are at first sight not easy to account for. Dr. Simpson suggests, with considerable show of probability, that the difference in frequency in England, France, and Germany may depend on the varying positions in which lying-in women are placed during labor in each country. In France, where, although the patient is laid on her back, the pelvis is kept elevated, the complication occurs least frequently; in England, where she lies on her side, more often; and in Germany, where she is placed on her back with her shoulders raised,

most often. The special frequency of prolapsed funis in certain districts, as in Kiel, is supposed by Engelmann¹ to depend on the prevalence of rickets, and consequently of deformed pelvis, which we shall presently see is probably one of the most frequent and important causes of the accident.

Prognosis.—With regard to the danger attending prolapsed funis, as far as the mother is concerned it may be said to be altogether unimportant; but the universal experience of obstetricians points to the great risk to which the child is subjected. Scanzoni calculates that 45 per cent. only of the children were saved; Churchill estimated the number at 47 per cent.; thus, under the most favorable circumstances, this complication leads to the death of more than half the children. Engelmann found that out of 202 vertex presentations only 36 per cent. of the children survived. The mortality was not nearly so great in other presentations; 68 per cent. of the cases in which the child presented with the feet were saved, and 50 per cent. in original shoulder presentations. The reason of this remarkable difference is, doubtless, that in vertex presentations the head fits the pelvis much more completely and subjects the chord to much greater pressure, while in other presentations the pelvis is less completely filled and the interference with the circulation in the cord is not so great. Besides, in the latter case the complication is detected early and the necessary treatment sooner adopted.

The fetal mortality is considerably greater in first labors—a result to be expected on account of the greater resistance of the soft parts and the consequent prolongation of the labor.

Causes.—The causes of prolapse of the funis are any circumstances which prevent the presenting part accurately fitting the pelvic brim. Hence it is much more frequent in face, breech, or shoulder than in vertex presentations, and is relatively more common in footling and shoulder presentations than in any other. Amongst occasional accidental predisposing causes may be mentioned early rupture of the membranes, especially if the amount of liquor amnii be excessive, as the sudden escape of the fluid washes down the cord; undue length of the cord itself; or an unusually low placental attachment. Engelmann attaches great importance to slight contraction of the pelvis, and states that in the Berlin Lying-in Hospital, where accurate measurements of the pelvis were taken in all cases, it was almost invariably found to exist. The explanation is evident, since one of the first results of pelvic contraction is to prevent the ready engagement of the presenting part in the pelvic brim.

Diagnosis.—The diagnosis of cord presentation is generally devoid of difficulty, but if the membranes are still unruptured it may not always be quite easy to determine the precise nature of the soft structures felt through them, as they recede from the touch. If the pulsations of the cord can be felt through the membranes, all difficulty is removed. After the membranes are ruptured there is nothing that it can well be mistaken for.

Importance of Determining the Pulsations of the Cord.—The important point to determine in such a case is whether the cord be pulsating or

¹ *Amer. Journ. of Obs.*, vol. vi.

not ; for if pulsations have entirely ceased the inference is that the child is dead, and the case may then be left to nature without further interference. It is of importance, however, to be careful, for if the examination be made during a pain the circulation might be only temporarily arrested. The examination, therefore, should be made during an interval, and a loop of the cord pulled down, if necessary, to make ourselves absolutely certain on this point.

Amount of Cord Prolapsed.—The amount of the prolapse varies much. Sometimes only a knuckle of the cord, so small as to escape observation, is engaged between the pelvis and presenting part. Under such circumstances the child may be sacrificed without any suspicion of danger having arisen. More often the amount prolapsed is considerable—sometimes so as to lie in the vagina in a long loop, or even to protrude altogether beyond the vulva.

Treatment.—In the treatment the great indication is to prevent the cord from being unduly pressed on, and all our endeavors must have this object in view. If the presentation be detected before the full dilatation of the cervix and when the membranes are unruptured, we must try to keep the cord out of the way ; to preserve the membranes intact as long as possible, since the cord is tolerably protected as long as it is surrounded by the liquor amnii ; and to secure the complete dilatation of the os, so that the presenting part may engage rapidly and completely.

Postural Treatment.—Much may be done at this time by the postural treatment, which we chiefly owe to the ingenuity of Dr. T. Gaillard Thomas of New York, whose writings familiarized the profession with it, although it appears that a somewhat similar plan had been occasionally adopted previously. Dr. Thomas's method is based on the principle of causing the cord to slip back into the uterine cavity by its own weight.

FIG. 122.



Postural Treatment of Prolapse of the Cord.

For this purpose the patient is placed on her hands and knees, with the hips elevated and the shoulders resting on a lower level (Fig. 122). The cervix is then no longer the most dependent portion of the uterus, and the anterior wall of the uterus forms an inclined plane down which the

cord slips. The success of this manœuvre is sometimes very great, but by no means always so. It is most likely to succeed when the membranes are unruptured. If, when adopted, the cord slip away and the os be sufficiently dilated, the membranes may be ruptured, and engagement of the head produced by properly-applied uterine pressure. Sometimes the position is so irksome that it is impossible to resort to it. Postural treatment is not even then altogether impossible, for by placing the patient on the side opposite to that of the prolapse, so as to relieve the cord as much as possible from pressure, and at the same time elevating the hips by a pillow, it may slip back. Even after the membranes are ruptured postural treatment in one form or another may succeed; and, as it is simple and harmless, it should certainly be always tried. Attempts at reposition by one or other of the methods described below may also occasionally be facilitated by trying them when the patient is placed in the knee-shoulder position.

Artificial Reposition.—Failing by postural treatment, or in combination with it, it is quite legitimate to make an attempt to place the cord beyond the reach of dangerous pressure by other methods. Unfortunately, reposition is too often disappointing, difficult to effect, and very frequently, even when apparently successful, shortly followed by a fresh descent of the cord. Provided the os be fully dilated and the presenting head engaged in the pelvis (for reposition may be said to be hopeless when any other part presents), perhaps the best way is to attempt it by the hand alone. Probably the simplest and most effectual method is that recommended by McClintock and Hardy, who advise that the patient should lie on the opposite side to the prolapsed cord, which should then be drawn toward the pubes as being the shallowest part of the pelvis. Two or three fingers may then be used to push the cord past the head and as high as they can reach. They must be kept in the pelvis until a pain comes on, and then very gently withdrawn, in the hope that the cord may not again prolapse. During the pain external pressure may very properly be applied to favor descent of the head. This manœuvre may be repeated during several successive pains, and may eventually succeed. The attempt to hook the cord over the foetal limbs or to place it in the hollow of the neck, recommended in many works, involves so deep an introduction of the hand that it is obviously impracticable.

Instruments Used for Reposition.—Various complex instruments have been invented to aid reposition (Fig. 123), but even if we possessed them they are not likely to be at hand when the emergency arises. A simple instrument may be improvised out of an ordinary male elastic catheter by passing the two ends of a piece of string through it, so as to leave a loop emerging from the eye of the catheter. This is passed through the loop of prolapsed cord, and then fixed in the eye of the catheter by means of the stilette. The cord is then pushed up into the uterine cavity by the catheter, and liberated by withdrawing the stilette. Another simple instrument may be made by cutting a hole in a piece of whalebone. A piece of tape is then passed through the loop of the cord and the ends threaded through the eye cut in the whalebone. By tightening the tape the whalebone is held in close apposition to the cord, and the whole is passed as high as possible into the uterine cavity. The tape can

easily be liberated by pulling one end. If preferred, the cord can be tied to the whalebone, which is left in utero until the child is born. Nothing need be said as to the various other methods adopted for keeping up the cord, such as the insertion of pieces of sponge or tying the cord in a bag of soft leather, since they are generally admitted to be quite useless.

FIG. 123.



Braun's Apparatus for
Replacing the Cord.

Treatment when Reposition Fails.—It only too often happens that all endeavors at reposition fail. The subsequent treatment must then be guided by the circumstances of the case. If the pelvis be roomy and the pains strong, especially in a multipara, we may often deem it advisable to leave the case to nature, in the hope that the head may be pushed through before pressure on the cord has had time to prove fatal to the child. Under such circumstances the patient should be urged to bear down, and the descent of the head promoted by uterine pressure, so as to get the second stage completed as soon as possible. If the head be within easy reach, the application of the forceps is quite justifiable, since delay must necessarily involve the death of the child. During this time the cord should be placed, if possible, opposite one or other sacro-iliac synchondrosis, according to the position of the head, as the part of the pelvis where there is the most room, and where the pressure would consequently be least prejudicial. If we have to do with a case in which the head has not descended into the pelvis, and postural treatment and reposition have both failed, provided the os be fully dilated and other circumstances be favorable, turning would undoubtedly offer the best chance to the child. This treatment is strongly advocated by Engelmann, who found that 70 per cent. of the children delivered in this way were saved. There can be no question that, so far as the interests of the child are concerned, it is, under the circumstances indicated, by far the best expedient. Turning, however, is by no means always devoid of a certain risk to the mother, and the performance of the operation in any particular case must be left to the judgment of the practitioner. A fully-dilated os, with membranes unruptured, so that version could be performed by the combined method without the introduction of the hand into the uterus, would be unquestionably the most favorable state. If it be not deemed proper to resort to it, all that can be done is to endeavor to save the cord from pressure as much as possible by one or other of the methods already mentioned.

CHAPTER IX.

PROLONGED AND PRECIPITATE LABORS.

Dystocia arising from Defective or Irregular Action of the Uterus.—Among the difficulties connected with parturition there are none of more frequent occurrence, and none requiring more thorough knowledge of the physiology and pathology of labor, than those arising from deficient or irregular action of the expulsive powers. The importance of studying this class of labors will be seen when we consider the numerous and very diverse causes which produce them.

Evil Effects of Prolonged Labor.—That the mere prolongation of labor is in itself a serious thing is becoming daily more and more an acknowledged axiom of midwifery practice; and that this is so is evident when we contrast the statistical returns of such institutions as the Rotunda Lying-in Hospital of late years with those which were published some twenty or thirty years ago. It may be fairly assumed that the practice of the distinguished heads of that well-known school represents the most advanced and scientific opinion of the day. When we find that less than thirty years ago the forceps were not used more than once in 310 labors, while, according to the report for 1873, the late Master applied them once in 8 labors, it is apparent how great is the change which has taken place.

Causes of Prolonged Labor.—Labor may be prolonged from an immense number of causes, the principal of which will require separate study. Some depend simply on defective or irregular action of the uterus; others act by opposing the expulsion of the child, as, for example, undue rigidity of the parturient passages, tumors, bony deformity, and the like. Whatever the source of delay, a train of formidable symptoms are developed which are fraught with peril both to the mother and the child. As regards the mother, they vary much in degree and in the rapidity with which they become established. In many cases, in which the action of the uterus is slight, it may be long before serious results follow; while in others, in which a strongly-acting organ is exhausting itself in futile endeavors to overcome an obstacle, the worst signs of protraction may come on with comparative rapidity.

The Influence of the Stage of Labor in Protraction.—The stage of labor in which delay occurs has a marked effect in the production of untoward symptoms. It is a well-established fact that prolongation is of comparatively small consequence to either the mother or child in the first stage, when the membranes are still intact, and when the soft parts of the mother, as well as the body of the child, are protected by the liquor amnii from injurious pressure; whereas if the membranes have ruptured, prolongation becomes of the utmost importance to both as soon as the head has entered the pelvis, when the uterus is strongly excited by reflex stimulation, when the maternal soft parts are exposed to continu-

ous pressure, and when the tightly-contracted uterus presses firmly on the fetus and obstructs the placental circulation. It is in reference to the latter class of cases that the change of practice already alluded to has taken place, with the most beneficial results both to the mother and child.

It must not be assumed, however, that prolongation of labor is never of any consequence until the second stage has commenced. The fallacy of such an opinion was long ago shown by Simpson, who proved in the most conclusive way that both the maternal and fetal mortality were greatly increased in proportion to the entire length of the labor; and all practical accoucheurs are familiar with cases in which symptoms of gravity have arisen before the first stage is concluded. Still, relatively speaking, the opinion indicated is undoubtedly correct.

In the present chapter we have to do only with those causes of delay connected with the expulsive powers. Inasmuch, however, as the injurious effects of protraction are similar in kind, whatever be the cause, it will save needless repetition if we consider, once for all, the train of symptoms that arise whenever labor is unduly prolonged.

Delay in the First Stage.—As long as the delay is in the first stage only, with rare exceptions no symptoms of real gravity arise for a length of time, it may be even for days. There is often, however, a partial cessation of the pains, which in consequence of temporary exhaustion of nervous force may even entirely disappear for many consecutive hours. Under such circumstances, after a period of rest, either natural or produced by suitable sedatives, they recur with renewed vigor.

Symptoms of Protraction in the Second Stage.—A similar temporary cessation of the pains may often be observed after the head has passed through the os uteri, to be also followed by renewed vigorous action after rest. But now any such irregularity must be much more anxiously watched. In the majority of cases any marked alteration in the force and frequency of the pains at this period indicates a much more serious form of delay, which in no long time is accompanied by grave general symptoms. The pulse begins to rise, the skin to become hot and dry, the patient to be restless and irritable. The longer the delay and the more violent the efforts of the uterus to overcome the obstacle, the more serious does the state of the patient become. The tongue is loaded with fur, and in the worst cases dry and black; nausea and vomiting often become marked; the vagina feels hot and dry, the ordinary abundant mucous secretion being absent; in severe cases it may be much swollen, and if the presenting part be firmly impacted a slough may even form. Should the patient still remain undelivered, all these symptoms become greatly intensified: the vomiting is incessant, the pulse is rapid and almost imperceptible, low muttering delirium supervenes, and the patient eventually dies with all the worst indications of profound irritation and exhaustion.

So formidable a train of symptoms, or even the slighter degrees of them, should never occur in the practice of the skilled obstetrician; and it is precisely because a more scientific knowledge of the process of parturition has taught the lesson that under such circumstances prevention is better than cure, that earlier interference has become so much more the rule.

Those who taught that nothing should be done until nature had had every possible chance of effecting delivery, and who therefore allowed their patients to drag on in many weary hours of labor, at the expense of great exhaustion to themselves and imminent risk to their offspring, made much capital out of the time-honored maxim that "Meddlesome midwifery is bad." When this proverb is applied to restrain the rash interference of the ignorant, it is of undeniable value; but when it is quoted to prevent the scientific action of the experienced, who know precisely when and why to interfere, and who have acquired the indispensable mechanical skill, it is sadly misapplied.

State of the Uterus in Protracted Labor.—The nature of the pains and the state of the uterus in cases of protracted labor are peculiarly worthy of study, and have been very clearly pointed out by Dr. Braxton Hicks.¹ He shows that when the pains have apparently fallen off and become few and feeble, or have entirely ceased, the uterus is in a state of continuous or tonic contraction, and that the irritation resulting from this is the chief cause of the more marked symptoms of powerless labor. If in a case of the kind the uterus be examined by palpation, it will be found firmly contracted between the pains. The correctness of this observation is beyond question, and it will no doubt often be an important guide in treatment. Under such circumstances instrumental interference is imperatively demanded.

Conditions and Causes affecting the Expulsive Powers.—In considering the causes of protracted labor it will be well first to discuss those which affect the expulsive powers alone, leaving those depending on morbid states of the passages for future consideration; bearing in mind, however, that the results, as regards both the mother and the child, are identical, whatever may be the cause of delay.

Constitution of the Patient.—The general constitutional state of the patient may materially influence the force and efficiency of the pains. Thus, it not unfrequently happens that they are feeble and ineffective in women of very weak constitution or who are much exhausted by debilitating disease. Cazeaux pointed out that the effects of such general conditions are often more than counterbalanced by flaccidity and want of resistance of the tissues, so that there is less obstacle to the passage of the child. Thus, in phthisical patients reduced to the last stage of exhaustion labor is not unfrequently surprisingly easy.

Influence of Tropical Climates.—Long residence in tropical climates causes uterine inertia, in consequence of the enfeebled nervous power it produces. It is a common observation that European residents in India are peculiarly apt to suffer from post-partum hemorrhage from this cause. The general mode of life of patients has an unquestionable effect; and it is certain that deficient and irregular uterine action is more common in women of the higher ranks of society, who lead luxurious, enervating lives, than in women whose habits are of a more healthy character.

Frequent Childbearing.—Tyler Smith lays much stress on frequent childbearing as a cause of inertia, pointing out that a uterus which has been very frequently subjected to the changes connected with pregnancy

¹ *Obst. Trans.*, vol. ix.

is unlikely to be in a typically normal condition. It is doubtful, however, whether the uterus of a perfectly healthy woman is affected in this way; certainly, if childbearing had undermined her general health the labors are likely to be modified also.

Age of Patient.—Age has a decided effect. In the very young the pains are apt to be irregular, on account of imperfect development of the uterine muscle. Labor taking place for the first time in women advanced in life is also apt to be tedious, but not by any means so invariably as is generally believed. The apprehensions of such patients are often agreeably falsified, and where delay does occur it is probably more often referable to rigidity and toughness of the parturient passages than to feebleness of the pains.

Disorders of the Intestines.—Morbid states of the primæ viæ frequently cause irregular, painful, and feeble contractions. A loaded state of the rectum has a remarkable influence, as evidenced by the sudden and distinct change in the character of the labor which often follows the use of suitable remedies. Undue distension of the bladder may act in the same way, more especially in the second stage. When the urine has been allowed to accumulate unduly, the contraction of the accessory muscles of parturition often causes such intense suffering, by compressing the distended viscus, that the patient is absolutely unable to bear down. Hence the labor is carried on by uterine contractions alone, slowly and at the expense of much suffering. A similar interference with the action of the accessory muscles is often produced by other causes. Thus, if labor comes on when the patient is suffering from bronchitis or other chest disease, she may be quite unable to fix the chest by a deep inspiration, and the diaphragm and other accessory muscles cannot act. In the same way they may be prevented from acting when the abdomen is occupied by an ovarian tumor or by ascitic fluid.

Mental conditions have a very marked effect. This is so commonly observed that it is familiar to the merest beginner in midwifery practice. The fact that the pains often diminish temporarily on the entrance of the accoucheur is known to every nurse; and so also undue excitement, the presence of too many people in the room, overmuch talking, have often the same prejudicial effect. Depression of mind, as in unmarried women, and fear and despondency in women who have looked forward with apprehension to their labor, are also common causes of irregular and defective action.

Excessive Amount of Liquor Amnii.—Undue distension of the uterus from an excessive amount of liquor amnii not unfrequently retards the first stage by preventing the uterus from contracting efficiently. When this exists, the pains are feeble and have little effect in dilating the cervix beyond a certain degree. This cause may be suspected when undue protraction of the first stage is associated with an unusually large size and marked fluctuation of the uterine tumor, through which the foetal limbs cannot be made out on palpation. On vaginal examination the lower segment of the uterus will be found to be very rounded and prominent, while the bag of membranes will not bulge through the os during the acme of the pain.

Malpositions of the Uterus.—A somewhat similar cause is undue ob-

liquity of the uterus, which prevents the pains acting to the best mechanical advantage and often retards the entry of the presenting part into the brim. The most common variety is anteversion, resulting from undue laxity of the abdominal parietes, which is especially found in women who have borne many children. Sometimes this is so excessive that the fundus lies over the pubes, and even projects downward toward the patient's knees. The consequence is, that when labor sets in, unless corrective means be taken, the pains force the head against the sacrum, instead of directing it into the axis of the pelvic inlet. Another common deviation is lateral obliquity, a certain degree of which exists in almost all cases, but sometimes it occurs to an excessive degree. Either of these states can readily be detected by palpation and vaginal examination combined. In the former the os may be so high up and tilted so far backward that it may be at first difficult to reach it at all.

Irregular and Spasmodic Pains.—Besides being feeble, the uterine contractions, especially in the first stage, are often irregular and spasmodic, intensely painful, but producing little or no effect on the progress of the labor. This kind of case has been already alluded to in treating of the use of anæsthetics (p. 295), and is very common in highly-nervous and emotional women of the upper classes. Such irregular contractions do not necessarily depend on mental causes alone, and they are often produced by conditions producing irritation, such as loaded bowels, too early rupture of the membranes, and the like. Dr. Trenholme of Montreal¹ believes that such irregular pains most frequently depend on abnormal adhesions between the decidua and the uterine walls, which interfere with the proper dilatation of the os, and he has related some interesting cases in support of this theory.

Treatment.—The mere enumeration of these various causes of protracted labor will indicate the treatment required. Some of them, such as the constitutional state of the patient, age, or mental emotion, it is of course beyond the power of the practitioner to influence or modify; but in every case of feeble or irregular uterine action a careful investigation should be made with a view of seeing if any removable cause exist. For example, the effect of a large enema when we suspect the existence of a loaded rectum is often very remarkable, the pains frequently almost immediately changing in character and a previously lingering labor being rapidly terminated.

Excessive distension of the uterus can only be treated by artificial evacuation of the liquor amnii; and after this is done the character of the pains often rapidly changes. This expedient is indeed often of considerable value in cases in which the cervix has dilated to a certain extent, but in which no further progress is made, especially if the bag of membranes does not protrude through the os during the pains, and the cervix itself is soft and apparently readily dilatable. Under such circumstances rupture of the membranes, even before the os is fully dilated, is often very useful.

Adherent Membranes.—If we have reason to suspect morbid adhesions between the membranes and the uterine walls, an endeavor must be made to separate them by sweeping the finger or a flexible catheter round

¹ *Obst. Trans.*, 1873.

the internal margin of the os or puncturing the sac. The former expedient has been advocated by Dr. Inglis¹ as a means of increasing the pains when the first stage is very tedious, and I have often practised it with marked success. Trenholme's observation affords a rationale of its action. The manœuvre itself is easily accomplished, and, provided the os be not very high in the pelvis, does not give any pain or discomfort to the patient.

Uterine Deviations.—Attention should always be paid to remedying any deviations of the uterus from its proper axis. If this be lateral, the proper course to pursue is to make the patient lie on the opposite side to that toward which the organ is pointing. In the more common anterior deviation she should lie on her back, so that the uterus may gravitate toward the spine, and a firm abdominal bandage should be applied. This prevents the organ from falling forward, while its pressure stimulates the muscular fibres to increased action; hence it is often very serviceable when the pains are feeble, even if there be no anteversion.

Temporary Exhaustion.—In a frequent class of cases, especially in the first stage, the pains diminish in force and frequency from fatigue, and the indication then is to give a temporary rest, after which they recommence with renewed vigor. Hence an opiate, such as 20 minims of Battley's solution, which often acts quickest when given in the form of enema, is frequently of the greatest possible value. If this secure a few hours' sleep, the patient will generally awake much refreshed and invigorated. It is important to distinguish this variety of arrested pain from that dependent on actual exhaustion; and this can be done by attention to the general condition of the patient, and especially by observing that the uterus is soft and flaccid in the intervals between the pains, and that there is none of the tonic contraction indicated by persistent hardness of the uterine parietes. When the pains are irregular, spasmodic, and excessively painful without producing any real effect, opiates are also of great service; and it is under such circumstances that chloral is especially valuable.

Oxytocic Remedies.—Still, a large number of cases will arise in which the absence of all removable causes has been ascertained and in which the pains are feeble and ineffective. We must now proceed to discuss their management. The fault being the want of sufficient contraction, the first indication is to increase the force of the pains. Here the so-called *oxytocic* remedies come into action; and although a large number of these have been used from time to time, such as borax, cinnamon, quinine, and galvanism, practically the only one on which reliance is generally placed is the ergot of rye.

Ergot of Rye.—This has long been the favorite remedy for deficient uterine action, and it is a powerful stimulant of the uterine fibres. It has, however, very serious disadvantages, and it is very questionable whether the risks to both mother and child do not more than counterbalance any advantages attending its use. The ergot is given in doses of 15 or 20 grains of the freshly-powdered drug infused in warm water, or in the more convenient form of the liquid extract in doses of from 20 to 30 minims, or, still better, in the form of ergotin injected hypodermic-

¹ Sydenham Society's Year-Book, 1869.

ally, 3 to 4 minims of the hypodermic solution being used for the purpose. In about fifteen minutes after its administration the pains generally increase greatly in force and frequency, and if the head be low in the pelvis, and if the soft parts offer no resistance, the labor may be rapidly terminated.

Objections to its Use.—Were its use always followed by this effect there would be little or no objection to its administration. The pains, however, are different from those of natural labor, being strong, persistent, and constant. Its effect, indeed, is to produce that very state of tonic and persistent uterine contraction which has been already pointed out as one of the chief dangers of protracted labor. Hence, if, from any cause, the exhibition of the drug be *not* followed by rapid delivery, a condition is produced which is serious to the mother, and which is extremely perilous to the child, on account of the tonic contraction of the muscular fibres obstructing the utero-placental circulation. Dr. Hardy found that soon the foetal pulsations fall to 100, and if delivery be long delayed they commence to intermit. He also observed that when this occurred the child was always born dead, and found that the number of stillborn children after ergot has been exhibited was very large, for out of 30 cases in which he gave it in tedious labor, only 10 of the children were born alive. Nor is its use by any means free from any danger to the mother: a not inconsiderable number of cases of rupture of the uterus have been attributed to its incautious use. Hence, if it is to be given at all, it is obvious that it must be with strict limitations and after careful consideration. It is worthy of note that in the Rotunda Hospital in Dublin the use of ergot as an oxytocic before delivery has been prohibited by the present Master.

Limitations to its Use.—The cardinal point to remember is, that it is absolutely contraindicated unless the absence of all obstacles to rapid delivery has been ascertained. Hence, it is only allowable when the first stage is over and the os fully dilated; when the experience of former labors has proved the pelvis to be of ample size; and when the perineum is soft and dilatable. Perhaps, as has been suggested, the administration of small doses of from 5 to 10 minims of the liquid extract every ten minutes until more energetic action set in might obviate some of these risks.

Quinine as an Oxytocic.—The use of quinine as an oxytocic deserves much more attention than it has generally received. I frequently employ it in lingering labor with marked benefit, and it does not seem to have any of the bad effects of ergot. According to the observations of Dr. Albert H. Smith, in 42 cases of parturition it presented the following peculiar characteristics:

It has no power in itself to excite uterine contractions, but simply acts as a general stimulant and promoter of vital energy and functional activity.

In normal labor at full term its administration in a dose of 15 grains is usually followed in as many minutes by a decided increase in the force and frequency of the uterine contractions, changing in some instances a tedious, exhausting labor into one of rapid energy, advancing to an early completion.

It promotes the permanent tonic contraction of the uterus after the expulsion of the placenta, women that had flooded in former labors escaping entirely, there not having been an instance of post-partum hemorrhage in the whole 42 cases.

It also diminishes the lochial flow where it had been excessive in former labors, the change being remarked upon by the patients, and consequently lessens the severity of the after-pains.

Cinchonism is very rarely observed as an effect of large doses in parturient women.¹

Use of the Faradic Current.—The faradic current applied on either side of the uterine tumor, midway between the anterior-superior spine of the ilium and the umbilicus, has recently been strongly recommended by Dr. Kilner,² not only as a means of increasing uterine action, but of alleviating the sufferings of childbirth. I have tried it in several cases, but am not satisfied as to its possessing the properties attributed to it.

Manual Pressure as a Means of Increasing the Uterine Contractions.—If we had no other means of increasing defective uterine contractions at our disposal, and if the choice lay only between the use of ergot and instrumental delivery, there might not be so much objection to a cautious use of the drug in suitable cases. We have, however, a means of increasing the force of the uterine contractions so much more manageable and so much more resembling the natural process that I believe it to be destined to entirely supersede the administration of ergot. This is the application of manual pressure to the uterus through the abdomen—an expedient that has of late years been much used in Germany, and has begun to be employed in English practice. I believe, therefore, that ergot should be chiefly used for the purpose of exciting uterine contraction after delivery, when its peculiar property of promoting tonic contraction is so valuable, and that it should rarely, if at all, be employed before the birth of the child.

The systematic use of uterine pressure as an oxytocic was first prominently brought under the notice of the profession by Kristeller under the name of "Expressio Fœtus," although it has been used in various forms from time immemorial. Albucasis, for example, was clearly acquainted with its use, and referred to it in the following terms: "Cum ergo vides ista signa, tunc oportet, ut comprimatur uterus ejus ut descendat embryo velociter." There are some curious obstetric customs among various nations which probably arose from a recognition of its value; as, for example, the mode of delivery adopted among the Kalmucks, where the patient sits at the foot of the bed, while a woman seated behind her seizes her round the waist and squeezes the uterus during the pains. Amongst the Japanese, Siamese, North American Indians, and many other nations pressure, applied in various ways, is habitually used.

Kristeller maintains that it is possible to effect the complete expulsion of the child by properly-applied pressure, even when the pains are entirely absent. Strange as this may appear to those who are not familiar with the effects of pressure, I believe that under exceptional circumstances, when the pelvis is very capacious and the soft parts offer but slight resistance, it can be done. I have delivered in this way a patient

¹ *Trans. Coll. Phys. Philada.*, 1875, p. 183.

² *Lancet*, January 1, 1881.

whose friends would not permit me to apply the forceps when I could not recognize the existence of any uterine contraction at all, the fetus being literally squeezed out of the uterus. It is not, however, as replacing absent pains, but as a means of intensifying and prolonging the effects of deficient and feeble ones, that pressure finds its best application.

Its effects are often very remarkable, especially in women of slight build where there is but little adipose tissue in the abdominal walls and not much resistance in the pelvic tissues. If the finger be placed on the head while pressure is applied to the uterus, a very marked descent can readily be felt, and not infrequently two or three applications will force the head on to the perineum. There are, however, certain conditions when it is inapplicable, and the existence of which should contraindicate its use. Thus, if the uterus seem unusually tender on pressure, and, *a fortiori*, if the tonic contraction of exhaustion be present, it is inadmissible. So also if there be any obstruction to rapid delivery, either from narrowing of the pelvis or rigidity of the soft parts, it should not be used. The cases suitable for its application are those in which the head or breech is in the pelvic cavity, and the delay is simply due to a want of sufficiently strong expulsive action.

Mode of Application.—It may be applied in two ways. The better is to place the patient on her back at the edge of the bed, and spread the palms of the hands on either side of the fundus and body of the uterus, and when a pain commences to make firm pressure during its continuance downward and backward in the direction of the pelvic inlet. As the contraction passes off the pressure is relaxed, and again resumed when a fresh pain begins. In this way each pain is greatly intensified and its effect on the progress of the fetus much increased. It is not essential that the patient should lie on her back. A useful although not so great amount of pressure can be applied when she is lying in the ordinary obstetric position on her left side, the left hand being spread out over the fundus, leaving the right free to watch the progress of the presenting part per vaginam.

Special Value of Uterine Pressure.—The special value of this method of treating ineffective pains is, that the amount and frequency of the pressure are completely within the control of the practitioner, and are capable of being regulated to a nicety in accordance with the requirements of each particular case. It has the peculiar advantage of closely imitating the natural means of delivery, and of being absolutely without risk to the child; nor is there any reason to think that it is capable of injuring the mother. At least I may safely say that, out of the large number of cases in which I have used it, I have never seen one in which I had the least reason to think that it had proved hurtful. Of course it is essential not to use undue roughness: firm and even strong pressure may be employed, but that can be done without being rough; and, as its application is always intermittent, there is no time for it to inflict any injury on the uterine tissues.

Pressure is specially valuable when it is desirable to intensify feeble pains. It may be serviceably employed when the pains are altogether absent to imitate and replace them, provided there be nothing but the absence of a *vis a tergo* to prevent speedy delivery. In such cases an

endeavor should be made to imitate the pains as closely as possible by applying the pressure at intervals of four or five minutes, and entirely relaxing it after it has been applied for a few seconds.

Change of Professional Opinion as to Instrumental Delivery.—When all these means fail, we have then left the resource of instrumental aid; and we have now to consider the indications for the use of the forceps under such circumstances. It has been already pointed out that professional opinion on this point has been undergoing a marked change, and that it is now recognized as an axiom by the most experienced teachers that when we are once convinced that the natural efforts are failing and are unlikely to effect delivery, except at the cost of long delay, it is far better to interfere soon rather than late, and thus prevent the occurrence of the serious symptoms accompanying protracted labor. The recent important debate on the use of the forceps at the Obstetrical Society of London remarkably illustrated these statements, for, while there was much difference of opinion as to the advisability of applying the forceps when the head was high in the pelvis, a class of cases not now under consideration, it was very generally admitted that the modern teaching was based on correct scientific grounds. This is, of course, directly opposed to the view so long taught in our standard works, in which instrumental interference was strictly prohibited unless all hope of natural delivery was at an end, and in which the commencement at least, if not the complete establishment, of symptoms of exhaustion was considered to be the only justification for the application of the forceps in lingering labor.

Views of Dr. Johnston on the Use of the Forceps.—The reasons which have led the late distinguished Master of the Rotunda Hospital to a more frequent use of the forceps are so well expressed in his report for 1872 that I venture to quote them as the best justification for a practice that many practitioners of the older school will, no doubt, be inclined to condemn as rash and hazardous. He says:¹ “Our established rule is, that so long as nature is able to effect its purpose without prejudice to the constitution of the patient, danger to the soft parts or the life of the child, we are in duty bound to allow the labor to proceed; but as soon as we find the natural efforts are beginning to fail, and after having tried the milder means for relaxing the parts or stimulating the uterus to increased action, and the desired effects not being produced, we consider we are in duty bound to adopt still prompter measures, and by our timely assistance relieve the sufferer from her distress and her offspring from an imminent death. Why, may I ask, should we permit a fellow-creature to undergo hours of torture when we have the means of relieving her within our reach? Why should she be allowed to waste her strength and incur the risks consequent upon long pressure of the head on the soft parts, the tendency to inflammation and sloughing, or the danger of rupture, not to speak of the poisonous miasm which emanates from an inflammatory state of the passages, the result of tedious labor, and which is one of the fertile causes of puerperal fever and all its direful effects, attributed by some to the influence of being confined in a large maternity, and not to its proper source—i. e. the labor being

¹ *Fourth Clinical Report of the Rotunda Lying-in Hospital for the year ending 1872.*

allowed to continue till inflammatory symptoms appear? The more we consider the benefits of timely interference and the good results which follow it, the more are we induced to pursue the system we have adopted, and to inculcate to those we are instructing the advantages to be gained by such practice, both in saving the life of the child as well as securing the greater safety of the mother." It would be impossible to put the matter in a stronger or clearer light, and I feel confident that these views will be endorsed by all who have adopted the more modern practice.

Effect of Early Interference on Infantile Mortality.—In the first edition of this work I used the statistics of Dr. Hamilton of Falkirk and other modern writers, as proving that a more frequent use of the forceps than had been customary diminished in a remarkable degree the infantile mortality. Dr. Galabin¹ has recently published an admirable paper on this subject, in which, by a careful criticism of these figures, he has, I think, proved that the conclusions drawn from them are open to doubt, and that the saving of infantile life following more frequent forceps delivery is by no means so great as I had supposed. Dr. Roper, in his remarks in the recent debate in the Obstetrical Society, brought forward some strong arguments in support of the same view. This, however, does not in any way touch the main points at issue referred to in the preceding paragraph.

Possible Dangers attending the Use of the Forceps.—It is, of course, right that we should consider the opposite point of view, and reflect on the disadvantages which may attend the interference advocated. Here I should point out that I am now talking only of the use of the forceps in simple inertia, when the head is low in the pelvic cavity, and when all that is wanted is a slight *vis a fronte* to supplement the deficient *vis a tergo*. The use of the instrument when the head is arrested high in the pelvis, or in cases of deformity, or before the os uteri is completely expanded, is an entirely different and much more serious matter, and does not enter into the present discussion. The chief question to decide is if there be sufficient risk to the mother to counterbalance that of delay. It will, of course, be conceded by all that the forceps in the hands of a coarse, bungling, and ignorant practitioner, who has not studied the proper mode of operating, may easily inflict serious damage. The possibility of inflicting injury in this way should act as a warning to every obstetrician to make himself thoroughly acquainted with the proper mode of using the instrument, and to acquire the manual skill which practice and the study of the mechanism of delivery will alone give; but it can hardly be used as an argument against its use. If that were admitted, surgical interference of any kind would be tabooed, since there is none that ignorance and incapacity might not render dangerous.

Assuming, therefore, that the practitioner is able to apply the forceps skilfully, is there any inherent danger in its use? I think all who dispassionately consider the question must admit that in the class of cases alluded to the operation is so simple that its disadvantages cannot for a moment be weighed against those attending protraction and its consequences. Against this conclusion statistics may possibly be quoted, such

¹ *Obstetrical Journal*, December, 1877.

means of unfrequent occurrence. Most obstetric works contain a formidable catalogue of evils that may attend it, such as rupture of the cervix, or even of the uterus itself, from the violence of the uterine action; laceration of the perineum from the presenting part being driven through before dilatation has occurred; fainting from the sudden emptying of the uterus; hemorrhage from the same cause. With regard to the child, it is held that the pressure to which it is subjected, and sudden expulsion while the mother is in the erect position, may prove injurious. Without denying that these results may possibly occur now and again, in the majority of cases over-rapid labor is not attended with any evil effects.

Precipitate labor may generally be traced to one of two conditions, or to a combination of both: excessive force and rapidity of the pains or unusual laxity and want of resistance of the soft parts. The precise causes inducing these it is difficult to estimate. In some cases the former may depend on an undue amount of nervous excitability, and the latter on the constitutional state of the patient, tending to relaxation of the tissues.

Whatever the cause, the extreme rapidity of labor is occasionally remarkable, and one strong pain may be sufficient to effect the expulsion of the child with little or no preliminary warning. I have known a child to be expelled into the pan of a water-closet, the only previous indication of commencing labor being a slight griping pain, which led the mother to fancy that an action of the bowels was about to take place. More often there is what may be described as a storm of uterine contractions, one pain following the other with great intensity until the fetus is expelled. The natural effect of this is to produce a great amount of alarm or nervous excitement, which of itself forms one of the worst results of this class of labor. It is under such circumstances that tem-

per has been lost in the pueral unit Stockman's *Memirs*, vol. i. p. 671. It would be difficult to find a case which more forcibly illustrates the danger of delay in the second stage of labor. "Then what follows?" The uterus, exhausted by the lengthy efforts it should have been spared, fails to contract effectively, nor do we hear of any attempts to produce contraction by pressure. The relaxed organ becomes full of clots, extending up to the umbilicus, and all the most characteristic symptoms of concealed post-partum hemorrhage develop themselves. She complained "of being sick at stomach and of being in her usual degree to be indignant, and her pulse became frequent." Before long other symptoms came on, graphically described by James Stockman, and which seem to point to the formation of a clot in the broad and posterior arteries—a most likely occurrence when such a history. "Baillie sent me word that he wished me to see the patient. I hastened, but as she was with him. She was suffering from spasms of the chest and difficulty of breathing, in great pain, and very restless, and threw herself continually from one side of the bed to the other, speaking now to Baillie, now to Crisp. Baillie said to her, 'Above all, do not breathe a word.' She held out her left hand to me hastily, and pressed mine warmly twice. I felt her pulse; it was going very fast—the heart was beating very rapidly, and incessantly."

There was certainly something different from the formation of hemorrhage, and no one who has witnessed a case of post-partum hemorrhage can fail to recognize in this account an accurate delineation of its essential symptoms.

Never did I witness such one case lead to the conclusion that the misapprehension and unjust criticism, and a variety of the kind as this August, "multitude of misapprehensions" which has so long retarded the progress of science.

[See also.]

W. S. CHAFFIN.

Precipitate Labor less Common than Lingerin.—Undue rapidity of labor is certainly more uncommon than its converse, but still it is by no

soft, but very ineffectual. Toward the evening Sir Richard Croft began to suspect that labor might not terminate without artificial assistance, and a message was despatched for me. I arrived at two on Wednesday morning. The labor was now advancing more favorably, and both Dr. Baillie and myself concurred in the opinion that it would not be advisable to inform her Royal Highness of my arrival. From this time to the end of her labor the progress was uniform, though very slow, the patient in good spirits, the pulse calm, and there never was room to entertain a question about the use of instruments. About six in the afternoon the discharge became of a green color, which led to a suspicion that the child might be dead; still, the giving assistance was quite out of the question, as the pains now became more effectual and the labor proceeded regularly though slowly. The child was born without artificial assistance at nine o'clock in the evening. Attempts were made for a good while to reanimate it by inflating the lungs, friction, hot baths, etc., but without effect; the heart could not be made to beat even once. Soon after delivery Sir Richard Croft discovered that the uterus was contracted in the middle in the hour-glass form, and as some hemorrhage commenced, it was agreed that the placenta should be brought away by introducing the hand. This was done about half an hour after the delivery of the child, with more ease and less blood than usual. Her Royal Highness continued well for about two hours; she then complained of being sick at stomach and of noise in the ears, began to be talkative, and her pulse became frequent; but I understand she was very quiet after this, and her pulse calm. About half-past twelve o'clock she complained of severe pain in the chest, became extremely restless, with rapid, weak, and irregular pulse. At this time I saw her for the first time. It has been said that we had all gone to bed, but that is not a fact: Croft did not leave her room, Baillie retired about eleven, and I went to my bed-chamber and laid down in my clothes at twelve. By dissection, some bloody fluid (two ounces) was found in the pericardium, supposed to be thrown out in *articulo mortis*. The brain and other organs all sound, except the right ovary, which was distended into a cyst the size of a hen's egg. The hour-glass contraction of the uterus still visible, and a considerable quantity of blood in the cavity of the uterus—but those present dispute about the quantity, so much as from twelve ounces to a pound and a half—her uterus extending as high as her navel. The cause of her Royal Highness's death is certainly somewhat obscure; the symptoms were such as attend death from hemorrhage, but the loss of blood did not seem to be sufficient to account for a fatal issue. It is possible that the effusion into the pericardium took place earlier than was supposed, and it does not seem to be quite certain that this might not be the cause. That I did not see her Royal Highness more early was awkward, and it would have been better that I had been introduced before the labor was expected; and it should have been understood that when labor came on I should be sent to without waiting to know whether a consultation was necessary or not. I thought so at the time, but I could not propose such an arrangement to Croft. But this is entirely *entre nous*. I am glad to hear that your son is well, and, with all my family, wish to be remembered to him. We were happy to hear that he was agreeably married.

"I remain, my dear doctor,

"Ever yours, most truly,

"JOHN SIMS, M. D.

"This letter is confidential, as perhaps I might be blamed for writing any particulars without the permission of Prince Leopold."

What are the facts here shown? Here was a delicate young woman prepared for the trial before her, as Baron Stockmar tells us, by "lowering the organic strength of the mother by bleeding, aperients, and low diet," who was allowed to go on in lingering, feeble labor for no less than fifty-two hours after the escape of the liquor amnii! Such was the groundless dread of instrumental interference then prevalent that, although the case dragged on its weary length with feeble, ineffectual pains, every now and then increasing a little in intensity and then falling off again, it is stated "there never was room to entertain a question about the use of instruments;" and even "when the discharge became of a green color, . . . still, the giving assistance was quite out of the question"! Can any reasonable man doubt that if the forceps had been employed hours and hours before—say on Tuesday, when the pains fell off—the result would probably have been very different, and that the life of the child, destroyed by the enormously-prolonged second stage, would have been saved? It must be remembered that early on Tuesday morning delivery was expected, so that the head must

and the state of the patient. In the majority of cases, especially if the membranes be still intact, patience and time are sufficient to overcome the obstacle; but it is often in the power of the accoucheur materially to aid dilatation by appropriate management. Sometimes nature overcomes the obstruction by lacerating the opposing structures, and cases are on record in which even a complete ring of the cervix has been torn off and come away before the head.

Many remedies have been recommended for facilitating dilatation, some of which no doubt act beneficially. Among those most frequently resorted to was venesection, and with it was generally associated the administration of nauseating doses of tartar emetic. Both of these acted by producing temporary depression, under which the resistance of the soft part was lessened. They probably answer best in cases in which there was a rigid and tough cervix; and they might prove serviceable even yet in stout, plethoric women of robust frame. Practically, they are now seldom if ever employed, and other and less debilitating remedies are preferred. The agent, *par excellence*, which is most serviceable is chloral, which is of special value in the more common cases in which rigidity is associated with spasmodic contraction of the muscular fibres of the cervix. Two or three doses of 15 grains, repeated at intervals of twenty minutes, are often of almost magical efficacy, the pains becoming steady and regular and the os gradually relaxing sufficiently to allow the passage of the head. Should the chloral be rejected by the stomach, it may be satisfactorily administered per rectum. Chloroform acts much in the same way, but on the whole less satisfactorily, its effects being often too great; while the peculiar value of chloral is its influence in promoting relaxation of the tissues without interfering with the strength of the pains.

Local Means of Treatment.—Various local means of treatment may be also advantageously used. One is the warm bath, which is much used in France. It is of unquestionable value where there is mere rigidity, and may be used either as an entire bath or as a hip-bath, in which the patient sits from twenty minutes to half an hour. The objection is the fuss and excitement it causes, and for this reason it is an expedient seldom resorted to in this country. A similar effect is produced, and much more easily, by a douche of tepid water upon the cervix. This can be very easily administered, the pipe of a Higginson's syringe being guided up to the cervix by the index finger of the right hand, and a stream of water projected against it for five or ten minutes. Smearing the os with extract of belladonna is advised by continental authorities, but its effects are more than doubtful. Horton¹ advocates the injection into the tissue of the cervix of $\frac{1}{4}$ of a grain of atropine by means of a hypodermic syringe, and speaks very favorably of the practice.

Artificial Dilatation.—Artificial dilatation of the cervix by the finger is recommended, and has been the subject of much discussion, each school, where it was formerly commonly used, being very useful, but it may also do much harm. The class of cases in which

porary mania occurs, produced by the intensity of the suffering, under which the patient may commit acts, her responsibility for which may fairly be open to question.

Treatment.—Little can be done in treating undue rapidity of labor. We can, to some extent, modify the intensity of the pains by urging the patient to refrain from voluntary efforts and to open the glottis by crying out, so that the chest may no longer be a fixed point for muscular action. Opiates have been advised to control uterine action, but it is needless to point out that in most cases there is no time for them to take effect. Chloroform will often be found most valuable, from the rapidity with which it can be exhibited; and its power of diminishing uterine action, which forms one of its chief drawbacks in ordinary practice, will here prove of much service.

u t e a t

CHAPTER X.

LABOR OBSTRUCTED BY FAULTY CONDITION OF THE SOFT PARTS.

Rigidity of the Cervix a Frequent Cause of Protracted Labor.—One of the most frequent causes of delay in the first stage of labor is rigidity of the cervix uteri, which may depend on a variety of conditions. It is often produced by premature escape of the liquor amnii, in consequence of which the fluid wedge which is nature's means of dilating the os is destroyed, and the hard presenting part is consequently brought to bear directly upon the tissues of the cervix, which are thus unduly irritated and thrown into a state of spasmodic contraction. At other times it may be due to constitutional peculiarities, among which there is none so common as a highly nervous and emotional temperament, which renders the patient peculiarly sensitive to her sufferings and interferes with the harmonious action of the uterine fibres. The pains in such cases cause intense agony, are short and cramp-like in character, but have little or no effect in producing dilatation, the os often remaining for many hours without any appreciable alteration, its edges being thin and tightly stretched over the head. Less often—and this is generally met with in stout, plethoric women—the edges of the os are thick and tough.

Effects.—The effects of prolongation of labor from this cause will vary much under different circumstances. If the liquor amnii be prematurely evacuated, the presenting part presses directly upon the cervix, and the case is then practically the same as if the labor were in the second stage. Hence, grave symptoms may soon develop themselves, and early interference may be imperatively demanded. If the membranes be unruptured, delay will be of comparatively little moment, and considerable time may elapse without serious detriment to either the mother or child.

Treatment.—The treatment will naturally vary much with the cause

but the harder forms of cancer may form an insuperable obstacle to delivery.

Occlusion of the Os.—Agglutination of the margins of the os uteri is occasionally met with, and must, of course, have occurred after conception. It is generally the result of some inflammatory affection of the cervix during the early months of gestation; and I have known it to recur in the same woman in two successive pregnancies. Usually it is not associated with any hardness or rigidity, but the entire cervix is stretched over the presenting part, and forms a smooth covering, in which the os may only exist as a small dimple and may be very difficult to detect at all. Occlusion of the os uteri from inflammatory change sometimes so alters the cervix that no sign of the original opening can be discovered; and in two such instances the Cæsarean operation has been performed in the United States, by which the women were saved.¹

Treatment.—Any of these mechanical causes of rigidity may at first be treated in the same way as the more simple cases; and with patience, the use of chloral and chloroform, and of the fluid dilators, sufficient expansion to permit the passage of the head will often take place. But if these methods produce no effect, and symptoms of constitutional irritation are beginning to develop themselves, other and more radical means of overcoming the obstruction may be required.

Incision of the Cervix.—Under such circumstances incision of the cervix may be not only justifiable, but essential, and it frequently answers extremely well. On the Continent it is resorted to much more frequently and earlier than in this country, and with the most beneficial results. The operation offers no difficulties. The simplest way of performing it is to guard the greater portion of the blade of a straight, blunt-pointed bistoury by wrapping lint or adhesive plaster round it, leaving about half an inch cutting edge toward its point. This is guided to the cervix on the under surface of the index finger, and three or four notches are cut in the circumference of the os to about the depth of a quarter of an inch. Very generally, especially when the obstruction is only due to old cicatrices, the pains will now speedily effect complete expansion, which may be very advantageously aided by applying the hydrostatic dilators. When the obstruction is due to carcinomatous infiltration or inflammatory thickening, the case is much more complicated, and will painfully tax the resources of the accoucheur. If it is possible, the disease should be removed as much as can be safely done during pregnancy, which should also be brought to an end before the full period. During labor, incisions should form a preliminary to any subsequent proceedings that may be necessary, as they are, at the worst, not likely to increase in the least the risk the patient has to run, and they may possibly avert more serious operations. In the case of malignant disease the risk of serious hemorrhage, from the great vascularity of the tissues, must not be forgotten, and, if necessary, means must be taken to check this by local styptics, such as perchloride of iron. If incision fail and the state of the patient demands speedy delivery, the forceps may be applied; and Herman thinks they are, as a rule, better

¹ Harris's note to second American edition.

it is most serviceable are those in which the liquor amnii has been long evacuated, and in which the head, covered by the tightly-stretched cervix, has descended low into the pelvic cavity. Under these circumstances, if the finger be passed gently within the os during a pain, and its margin pressed upward and over the head, as it were, while the contraction lasts, the progress of the case may be materially facilitated. This manœuvre is somewhat similar to that which has been already spoken of when the anterior lip of the cervix is caught between the head and the pubic bone, and if properly performed I believe it to be quite safe and often of great value. It is not, however, well adapted for those cases in which the membranes are still intact, or in which the os remains undilated when the head is still high in the pelvis. When there is much delay under these conditions, and interference of some kind seems called for, the dilatation may be much assisted by the use of caoutchouc dilators, described in the chapter on the induction of premature labor, which imitate nature's method of opening up the os, and also act as a direct stimulant to uterine contraction. But it should be remembered that it is precisely in such cases that delay is least prejudicial. If, however, the os be excessively long in opening, its dilatation may be safely and efficiently promoted by passing within it, and distending with water, one of the smallest-sized bags; and after this has been in position from ten to twenty minutes it may be removed and a larger one substituted.

Rigidity depending upon Organic Causes.—Every now and again we meet with cases in which the obstacle depends upon organic changes in the cervix, the most common of which are cicatricial hardening from former lacerations, hypertrophic elongation of the cervix from disease antecedent to pregnancy, or even agglutination and closure of the os uteri. Cicatrices are generally the result of lacerations during former labors. They implicate a portion only of the cervix, which they render hard, rigid, and undilatable, while the remainder has its natural softness. They can readily be made out by the examining finger. A somewhat similar but much more formidable obstruction is occasionally met with in cases of old-standing hypertrophic elongation of the cervix, which is generally associated with prolapse. In most cases of this kind the cervix becomes softened during pregnancy, so that dilatation occurs without any unusual difficulty. But this does not always happen. A good example is related by Mr. Roper in the seventh volume of the *Obstetrical Transactions*, in which such a cervix formed an almost insuperable obstacle to the passage of the child.

Carcinoma of the cervix uteri, which produces extensive thickening and induration of its tissues, and even advanced malignant disease of the uterus, is no bar to conception. The relations of malignant disease to pregnancy and parturition have recently been well studied by Dr. Herman.¹ He concludes that cancer renders the patient inapt to conceive, but that when pregnancy does occur there is a tendency to the intra-uterine death and premature expulsion of the foetus, and the growth of the cancer is accelerated. When delivery is accomplished naturally, there is generally expansion of the cervix by fissuring of its tissue,

¹ *Obst. Trans.*, vol. xx. p. 191.

but the harder forms of cancer may form an insuperable obstacle to delivery.

Occlusion of the Os.—Agglutination of the margins of the os uteri is occasionally met with, and must, of course, have occurred after conception. It is generally the result of some inflammatory affection of the cervix during the early months of gestation; and I have known it to recur in the same woman in two successive pregnancies. Usually it is not associated with any hardness or rigidity, but the entire cervix is stretched over the presenting part, and forms a smooth covering, in which the os may only exist as a small dimple and may be very difficult to detect at all. Occlusion of the os uteri from inflammatory change sometimes so alters the cervix that no sign of the original opening can be discovered; and in two such instances the Cæsarean operation has been performed in the United States, by which the women were saved.¹

Treatment.—Any of these mechanical causes of rigidity may at first be treated in the same way as the more simple cases; and with patience, the use of chloral and chloroform, and of the fluid dilators, sufficient expansion to permit the passage of the head will often take place. But if these methods produce no effect, and symptoms of constitutional irritation are beginning to develop themselves, other and more radical means of overcoming the obstruction may be required.

Incision of the Cervix.—Under such circumstances incision of the cervix may be not only justifiable, but essential, and it frequently answers extremely well. On the Continent it is resorted to much more frequently and earlier than in this country, and with the most beneficial results. The operation offers no difficulties. The simplest way of performing it is to guard the greater portion of the blade of a straight, blunt-pointed bistoury by wrapping lint or adhesive plaster round it, leaving about half an inch cutting edge toward its point. This is guided to the cervix on the under surface of the index finger, and three or four notches are cut in the circumference of the os to about the depth of a quarter of an inch. Very generally, especially when the obstruction is only due to old cicatrices, the pains will now speedily effect complete expansion, which may be very advantageously aided by applying the hydrostatic dilators. When the obstruction is due to carcinomatous infiltration or inflammatory thickening, the case is much more complicated, and will painfully tax the resources of the accoucheur. If it is possible, the disease should be removed as much as can be safely done during pregnancy, which should also be brought to an end before the full period. During labor, incisions should form a preliminary to any subsequent proceedings that may be necessary, as they are, at the worst, not likely to increase in the least the risk the patient has to run, and they may possibly avert more serious operations. In the case of malignant disease the risk of serious hemorrhage, from the great vascularity of the tissues, must not be forgotten, and, if necessary, means must be taken to check this by local styptics, such as perchloride of iron. If incision fail and the state of the patient demands speedy delivery, the forceps may be applied; and Herman thinks they are, as a rule, better

¹ Harris's note to second American edition.

than turning. He also maintains that there is little difference in the risk to the mothers between craniotomy and the Cæsarean section, and that the possibility of saving the child in cases in which incisions have failed should induce us to prefer the latter.

Application of the Forceps within the Cervix.—Before performing craniotomy, when the os is sufficiently open, a cautious application of the forceps is quite justifiable. Steady and careful downward traction, combined with digital expansion, has often enabled a head to pass with safety through an os that has resisted all other means of dilatation, and the destruction of the child has thus been avoided. If, indeed, the os appear to be dilatable, this procedure may advantageously be adopted before incision, and, as a matter of fact, it is commonly practised in the Rotunda Hospital. An operation involving, beyond doubt, of itself some risk, and requiring considerable operative dexterity, should naturally not be lightly and inconsiderately undertaken. But when it is remembered that the alternative is the destruction of the child, the risk of exhaustion, and at least as great mechanical injury to the mother, its difficulty need not stand in the way of its adoption.

Treatment when Occlusion of the Os Exists.—When the os is apparently obliterated, incision is the only resource. Before resorting to it the patient should be placed under chloroform and the entire lower segment of the uterus carefully explored. Possibly, the aperture may be found high up and out of reach of an ordinary examination, or we may detect a depression corresponding to its site. A small crucial incision may then be made at the site of the os, if this can be ascertained; if not, at the most prominent portion of the cervix. Very generally, the pains will then suffice to complete expansion, which may be further aided by the fluid dilators.

Ante-partum Hour-glass Contraction.—Dr. Hosmer¹ has recently drawn attention to a hitherto undescribed species of dystocia, which he terms “*ante-partum hour-glass contraction*,” and which he believes to depend on constriction of the uterine fibres at the site of the internal os uteri. Harris² doubts its limitation to the internal os uteri, and terms it “*tetanoid falciform constriction of the uterus*.” Whatever its site, in the cases recorded difficulties of the most formidable kind arose from this cause. The pelves were normal and the presentations natural, yet out of 7 labors 4 ended fatally, 2 before delivery.^[3] The constriction seems to have grasped the fetus with such force as to have rendered extraction either by the forceps or turning impossible. I have no personal experience of this complication, which must fortunately be very rare. The introduction of the hand, the patient being deeply anesthetized, would probably render diagnosis easy. The treatment must depend on the force and amount of constriction. If the constriction does not relax under chloroform, chloral, or the injection of atropine into the site of constriction, as recommended by Horton in rigidity of the cervix, turning would probably be our best resource. Should this fail, the Cæsarean section may be required to effect delivery, as happened

¹ *Boston Med. and Surg. Journ.*, March and May, 1878.

² Harris's note to second American edition.

^[3] Of 34 labors, 9 ended fatally; 26 children were lost.—ED.]

in a case recorded by Dr. T. A. Foster of Portland, Maine. Gastro-elytrotomy is obviously unsuitable for such cases.

Bands and Cicatrices in the Vagina.—Extreme rigidity of the vagina, or bands and cicatrices in or across its walls, the result of congenital malformation, of injuries in former labors, or of antecedent disease, occasionally obstruct the second stage. There is seldom any really formidable difficulty from this cause, since the obstruction almost always yields to the pressure of the presenting part. If there be any considerable extent of cicatrices in the vagina, artificial assistance may be required. If we should be aware of their existence during pregnancy, and find them to be sufficiently dense and extensive to be likely to interfere with delivery, an endeavor may be made to dilate them gradually by hydrostatic bags or bougies. If they be not detected until labor is in progress, we must be guided in our procedure by the pressure to which they are subjected. It may then be necessary to divide them with a knife and to hasten the passage of the head by the forceps, so as to prevent contusion as much as possible. It is obviously impossible to lay down any positive rules for such rare contingencies, the treatment suitable for which must necessarily vary much with the individual peculiarities of the case.

Extreme rigidity of the perineum is often dependent upon cicatricial hardening from injury in previous labors. This may greatly interfere with its dilatation; and if laceration seems inevitable, we may be quite justified in attempting to avert it by incision of the margins of the perineum, on the principle of a clean cut being always preferable to a jagged tear.

Labor Complicated with Tumor.—Occasionally we meet with very formidable obstacles from tumors connected with the maternal structures. These are most commonly either fibroid or ovarian, although others may be met with, such as malignant growths from the pelvic bones, exostoses, etc.

Fibroid Tumors of the Uterus.—Considering the frequency with which women suffer from fibroid tumors of the uterus, it is perhaps somewhat remarkable that they do not more often complicate delivery. Probably women so affected are not apt to conceive. Occasionally, however, cases of this kind cause much anxiety. Of course, the cases are most grave in which tumors are so situated as to encroach upon the cavity of the pelvis and mechanically obstruct the passage of the child. Even those in which this does not occur are by no means free from danger, for interstitial and subperitoneal fibroids, situated in the upper parts of the uterus and leaving the pelvic cavity quite unimplicated, may interfere with the action of the uterine fibres, prevent subsequent contraction, cause profuse post-partum hemorrhage, or even predispose to rupture of the uterine tissue. Hence, every case in which the existence of uterine fibroids has been ascertained must be anxiously watched. The risk of hemorrhage is perhaps the greatest, for if the tumors be at all large, efficient contraction of the uterus after the birth of the child must be more or less interfered with. Fortunately, it is not so common as might almost be expected. Out of 5 cases recorded in the *Obstetrical Transactions*, 2 of which were in my own practice, no hemorrhage occurred; nor

does it seem to have happened in any of the 26 cases collected by Magdelaine in his thesis on the subject. I recently saw an interesting example of this in a patient whose case was looked forward to with much anxiety in consequence of the existence of several enormous fibroid masses projecting from the fundus and anterior surface of the body of the uterus, and whose labor was, nevertheless, typically normal in every way. Should hemorrhage occur after delivery, the injection of styptic solutions would probably be peculiarly valuable, since the ordinary means of promoting contraction are likely to fail.

It is when the fibroid growths implicate the lower uterine zone and the cervical region that the greatest difficulties are likely to be met with. The practice then to be adopted must be regulated to a great extent by the nature of each individual case. If it be possible to push the tumor above the pelvic brim, out of the way of the presenting part, that, no doubt, is the best course to pursue, as not only clearing the passage in the most effectual way, but removing the tumor from the bruising to which it would otherwise be subjected when pressed between the head and the pelvic walls; which seems to be one of the greatest dangers of this complication. This manœuvre is sometimes possible in what seem to be the most unpromising circumstances. An interesting example is narrated by Spencer Wells,¹ who, called to perform the Cæsarean section, succeeded, although not without much difficulty, in pushing the obstructing mass above the brim, the child subsequently passing with ease. I have myself elsewhere recorded two similar cases² in which I was enabled to deliver the patient by pushing up the obstructing tumor, in both of which the Cæsarean section would have been inevitable had the attempt at reposition failed. Therefore, before resorting to more serious operative procedures a determined effort at pushing the tumor out of the way should be made, the patient being deeply chloroformed, and, if necessary, upward pressure being made by the closed fist passed into the vagina.

Enucleation or Ablation.—Failing this, the possibility of enucleating the tumor, or, if that be not possible, of removing it piecemeal with the *écraseur*, should be considered. On account of the loose attachments of these growths, and the facility with which they can be removed in this way in the non-pregnant state, the expedient seems certainly well worthy of a trial if their site and attachments render it at all feasible. Interesting examples of the successful performance of this operation are recorded by Danyau and Braxton Hicks. Should it be found impracticable, the case must be managed in reference to the amount of obstruction, and the forceps, craniotomy, or even the Cæsarean section, may be necessary.

[The records of Cæsarean delivery show a great mortality in cases where the dystocia is due to an obstruction produced by fibroid tumors. In Dr. Sanger's collection,³ excluding the Porro operations, we find 39 Cæsarean sections, with 31 deaths: 8 of the 39 are credited to the United States, with 3 recoveries. To this I add 1 more recovery and 2 more deaths, giving us 4 saved out of 11, or 42 cases in all, with 9 women

¹ *Obst. Trans.*, vol. ix. p. 73.


² *Ibid.*, vol. xix. p. 101.

³ *Der Kaiserschnitt bei Uterusfibromen*, Leipzig, 1882, pp. 12-23.—Ed.]

saved. An early resort to the knife and suturing the uterine wound promise better results in the future.—ED.]

Tumors of the Ovaries.—The next most common class of obstructing tumors are those of the ovary (Fig. 124); and it is apparently not the largest of these which are most apt to descend into the pelvic cavity. When the tumor is of any considerable size, its bulk is such that it cannot be contained in the true pelvis, and it rises into the abdominal cavity

FIG. 124.



Labor complicated by Ovarian Tumor.

with the uterus. Hence, the existence of the tumor that offers the most formidable obstacle to the delivery is rarely suspected before labor sets in.

In order to estimate the results of the various methods of treatment, I have tabulated 57 cases.¹ In 13 labor was terminated by the natural powers alone, but of these 6 mothers, or nearly one-half, died. In favorable contrast with these we have the cases in which the size of the tumor was diminished by puncture. These are 9 in number, in all of which the mother recovered, 5 out of the 6 children being saved. The reason of the great mortality in the former cases is apparently the bruising to which the tumor, even when small enough to allow the child to be squeezed past it, is necessarily subjected. This is extremely apt to set up a fatal form of diffuse inflammation, the risk of which was long ago pointed out by Ashwell,² who draws a comparison between cases in which such tumors have been subjected to contusion and cases of strangulated hernia; and the cause of death in both is doubtless very similar. This danger is avoided when the tumor is punctured, so as to become flattened between the head and the pelvic walls. On this account I think it should be laid down as a rule that puncture should be performed in all cases of ovarian tumor engaged in front of the presenting part, even when it is of so small a size as not to preclude the possibility of delivery by the natural powers.

¹ *Obst. Trans.*, vol. ix.

² *Guy's Hospital Reports*, vol. ii.

Treatment when Puncture Fails.—In 5 of the 57 cases it was found possible to return the tumor above the pelvic brim, and in these also the termination was very favorable, all the mothers recovering. Should puncture not succeed—and it may fail on account of the gelatinous and semi-solid nature of the contents of the cyst—it may be possible to dispose of the tumor in this way, even when it seems to be firmly wedged down in front of the presenting part and to be hopelessly fixed in its unfavorable position.

Failing either of these resources, it may be necessary to resort to craniotomy, provided the size of the tumor precludes the possibility of delivery by forceps.

The question of the effect on labor of ovarian tumor which does not obstruct the pelvic canal is one of some interest, but there are not a sufficient number of cases recorded to throw much light on it. I am disposed to think that labor generally goes on favorably. What delay there is depends on the inefficient action of the accessory muscles engaged in parturition, on account of the extreme distension of the abdomen.

There are a few other conditions connected with the maternal structures which may impede delivery, but which are of comparatively rare occurrence.

Vaginal Cystocele.—Amongst them is vaginal cystocele, consisting of a prolapse of the distended bladder in front of the presentation, where it forms a tense fluctuating pouch, which has been mistaken for a hydrocephalic head or for the bag of membranes. This complication is only likely to arise when the bladder has been allowed to become unduly distended from want of attention to the voiding of urine during labor. The diagnosis should not offer any difficulty, for the finger will be able to pass behind, but not in front of, the swelling, and reach the presenting part, while the pain and tenesmus will further put the practitioner on his guard. The treatment consists in emptying the bladder; but there may be some difficulty in passing the catheter, in consequence of the urethra being dragged out of its natural direction. A long elastic male catheter will almost always pass if used with care and gentleness. Should it be found impossible to draw off the water—and this is said to have sometimes happened—the tense pouch might be punctured without danger by the fine needle of an aspirator-trocar and its contents withdrawn. When once the viscus is emptied it can easily be pushed above the presenting part in the intervals between the pains.

Vesical Calculus.—In some few cases difficulties have arisen from the existence of a vesical calculus. Should this be pushed down in front of the head, it can readily be understood that the maternal structures would run the risk of being seriously bruised and injured. Should we make out the existence of a calculus—and if the presence of one be suspected the diagnosis could easily be made by means of a sound—an endeavor should be made to push it above the brim of the pelvis. If that be found to be impossible, no resource is left but its removal, either by crushing or by rapid dilatation of the urethra, followed by extraction. Should we be aware of the existence of a calculus during pregnancy, its removal should certainly be undertaken before labor sets in.

to the bleeding point, and the hemorrhage can be controlled without difficulty. This is all the more necessary if spontaneous rupture have taken place, for then the loss of blood is often profuse, and it is of the utmost importance to reach the site of the hemorrhage as nearly as possible.

If the thrombus be not so large as to obstruct delivery, or if it be not detected until after the birth of the child, the question arises whether the case should not be left alone, in the hope that absorption may occur, as in most cases of pelvic hæmatocele. This expectant treatment is advised by Cazeaux, and it seems to be the most rational plan we can adopt. True, it may take a longer time for the patient to convalesce completely than if the coagula were removed at once and the hemorrhage restrained by pressure on the bleeding point; but this disadvantage is more than counterbalanced by the absence of risk from hemorrhage, and of septicæmia from the suppuration that must necessarily follow. Softening and suppuration may in many cases occur in a few days, necessitating operation, but the vessels will then be probably occluded and the risk of hemorrhage much lessened. Dr. Fordyce Barker, however, holds the opposite opinion, and thinks that the proper plan is to open the thrombus early, controlling the hemorrhage in the manner already indicated, unless the thrombus is situated high in the vaginal canal.

Risk of Subsequent Septicæmia.—Whenever the cavity of a thrombus has been opened, either by incision or by spontaneous softening at some time subsequent to its formation, it must not be forgotten that there is considerable risk of septic absorption. To avoid this, care must be taken to use antiseptic dressings freely, such as the glycerin of carbolic acid, applied directly to the part, and frequent vaginal injections of diluted Condy's fluid. Barker lays special stress on the importance of not removing prematurely the coagula formed by the styptic applications for fear of secondary hemorrhage, but of allowing them to come away spontaneously.

[*Polypus.*—Large uterine polypi may act as serious obstacles to delivery. When sufficiently long in pedicle, a polypus may be extruded before the head of the fetus. The tumor may also be detached in its expulsion, or may be removed by an *écraseur* if recognized in time: it may also be pushed up out of the way and secured by bringing down the child. I once replaced a large polypus that was extruded before the head, and the woman carried it two years longer; by which time, being much wasted by the discharge, she made up her mind to have it removed.—ED.]

cervix. It is natural to suppose that a varicose condition of the veins about the vulva would predispose to the accident, but in most of the recorded examples this is not stated to have been the case. Still, if varicose veins exist to any marked degree some anxiety on this point cannot but be felt.

Time of Occurrence.—The thrombus occasionally, though rarely, forms before delivery. Most commonly it first forms toward the end of labor or after the birth of the child. In the latter case it is probable that the laceration in the vessels occurred before the birth of the child, and that the pressure of the presenting part prevented the escape of any quantity of blood at the time of laceration.

Symptoms.—The symptoms are not by any means characteristic. Pain of a tearing character, occasionally very intense, and extending to the back and down the thighs, is very generally associated with the formation of the thrombus. If a careful physical examination be made, the nature of the case can readily be detected. When the blood escapes into the labium, a firm, hard swelling is felt, which has even been mistaken for the fetal head. If the effusion implicate the internal parts only, the diagnosis may not at first be so evident. But even then a little care should prevent any mistake, for the swelling may be felt in the vagina, and may even form an obstacle to the passage of the child. Cazeaux mentions cases in which it was so extensive as to compress the rectum and urethra, and even to prevent the exit of the lochia. In some cases the distension of the tissues is so great that they lacerate, and then hemorrhage, sometimes so profuse as directly to imperil the life of the patient, may occur. The bursting of the skin may take place some time subsequent to the formation of the thrombus. Constitutional symptoms will be in proportion to the amount of blood lost, either by extravasation or externally, after the rupture of the superficial tissues. Occasionally they are considerable, and are the same as those of hemorrhage from any cause.

Terminations.—The terminations of thrombus are either spontaneous absorption, which may occur if the amount of blood extravasated be small; or the tumor may burst, and then there is external hemorrhage; or it may suppurate, the contained coagula being discharged from the cavity of the cyst; or, finally, sloughing of the superficial tissues has occurred.

Treatment.—The treatment must naturally vary with the size of the thrombus and the time at which it forms. If it be met with during labor, unless it be extremely small it will be very apt to form an obstruction to the passage of the child. Under such circumstances it is clearly advisable to terminate the labor as soon as possible, so as to remove the obstacle to the circulation in the vessels. For this purpose the forceps should be applied as soon as the head can be easily reached. If the tumor itself obstruct the passage of the head or if it be of any considerable size, it will be necessary to incise it freely at its most prominent point and turn out the coagula, controlling the hemorrhage at once by filling the cavity with cotton wadding saturated in a solution of perchloride of iron, while at the same time digital compression with the tips of the fingers is kept up. By this means pressure is applied directly

to the bleeding point, and the hemorrhage can be controlled without difficulty. This is all the more necessary if spontaneous rupture have taken place, for then the loss of blood is often profuse, and it is of the utmost importance to reach the site of the hemorrhage as nearly as possible.

If the thrombus be not so large as to obstruct delivery, or if it be not detected until after the birth of the child, the question arises whether the case should not be left alone, in the hope that absorption may occur, as in most cases of pelvic hæmatocele. This expectant treatment is advised by Cazeaux, and it seems to be the most rational plan we can adopt. True, it may take a longer time for the patient to convalesce completely than if the coagula were removed at once and the hemorrhage restrained by pressure on the bleeding point; but this disadvantage is more than counterbalanced by the absence of risk from hemorrhage, and of septicæmia from the suppuration that must necessarily follow. Softening and suppuration may in many cases occur in a few days, necessitating operation, but the vessels will then be probably occluded and the risk of hemorrhage much lessened. Dr. Fordyce Barker, however, holds the opposite opinion, and thinks that the proper plan is to open the thrombus early, controlling the hemorrhage in the manner already indicated, unless the thrombus is situated high in the vaginal canal.

Risk of Subsequent Septicæmia.—Whenever the cavity of a thrombus has been opened, either by incision or by spontaneous softening at some time subsequent to its formation, it must not be forgotten that there is considerable risk of septic absorption. To avoid this, care must be taken to use antiseptic dressings freely, such as the glycerin of carbolic acid, applied directly to the part, and frequent vaginal injections of diluted Condy's fluid. Barker lays special stress on the importance of not removing prematurely the coagula formed by the styptic applications for fear of secondary hemorrhage, but of allowing them to come away spontaneously.

[*Polypus.*—Large uterine polypi may act as serious obstacles to delivery. When sufficiently long in pedicle, a polypus may be extruded before the head of the fetus. The tumor may also be detached in its expulsion, or may be removed by an *écraseur* if recognized in time: it may also be pushed up out of the way and secured by bringing down the child. I once replaced a large polypus that was extruded before the head, and the woman carried it two years longer; by which time, being much wasted by the discharge, she made up her mind to have it removed.—Ed.]

CHAPTER XI.

DIFFICULT LABOR DEPENDING ON SOME UNUSUAL CONDITION OF THE FŒTUS.

Plural Births.—The subject of multiple pregnancy in general having already been fully considered, we have now only to discuss its practical bearing as regards labor. Fortunately, the existence of twins rarely gives rise to any serious difficulty. In the large proportion of cases the presence of a second fœtus is not suspected until the birth of the first, when the nature of the case is at once apparent from the fact of the uterus remaining as large, or nearly as large, as it was before.

There may possibly be some delay in the birth of the first child, inasmuch as the extreme distension of the uterus may interfere with its thoroughly efficient action, while, in addition, the uterine pressure is not directly conveyed to the ovum as in single births, but indirectly through the amniotic sac of the second child (Fig. 125). Such delay is especially apt to arise when the first child presents by the breech, for even if the body be expelled spontaneously, difficulty is likely to occur with the head, since the uterus does not contract upon it, as is ordinarily the case. Hence the intervention of the accoucheur to save the life of the child by the extraction of the head will be almost a matter of necessity.

In the majority of cases, after the birth of the first child there is a temporary lull in the pains, which soon recommence, generally in from ten to twenty minutes, and the second child is rapidly expelled, for on account of the full dilatation of the soft parts there is no obstacle to its delivery. Sometimes there is a considerable interval before the pains recur, and instances are recorded in which even several days have elapsed between the births of the two children.

Treatment.—In most cases the management of twins does not differ from that of ordinary labor. As soon as we are certain of the existence of a second fœtus, we should inform the bystanders, but not necessarily the mother, to whom the news might prove an unpleasant and even dangerous shock. Then, having taken care to tie the cord of the first child for fear of vascular communication between the placentæ, our duty is to wait for a recurrence of the pains. If these come on

FIG. 125.



Twin Pregnancy. Breech and Head Presenting.

rapidly and the presentation of the second fetus be normal, its birth is managed in the usual way.

Management when there is Delay after the Birth of the First Child.—If there be any unusual delay, we have to consider the proper course to pursue, and on this the opinions of authorities differ greatly. Some advise a delay of several hours, and even more, if pains do not recur spontaneously; while others—Murphy, for example—recommend that the second child should be delivered at once. Either extreme of practice is probably wrong, and the safest and best course is doubtless the medium one. The second point to bear in mind is, that in multiple pregnancy, on account of the extreme distension of the uterus, there is a tendency to inertia, and consequently to post-partum hemorrhage, and that, therefore, it is better that the birth of the second child should be delayed, even for a considerable time, rather than the patient should run the risk attending an empty and uncontracted uterus. If, however, uterine action be present, there is an obvious advantage in the delivery of the second child before the dilatation of the passages passes off.


Endeavors should be made to Excite Uterine Action.—The best plan would seem to be if, after waiting a quarter of an hour, labor-pains do not occur, to try and induce them by uterine friction and pressure and by the administration of a dose of ergot, to which, as there can be no obstacle to the rapid birth of the second child, there can be now no objection. The membranes of the second child should always be ruptured at once if easily within reach, as one of the speediest means of inducing contraction. If no progress be made, and speedy delivery be indicated—a necessity which may arise either from the exhausted state of the patient, the presence of hemorrhage, extremely feeble pulsations of the fetal heart (showing that the life of the second child is endangered), or malpresentations of the second fetus—turning is probably the readiest and safest expedient. Under such circumstances the operation is performed with great ease, since the passages are amply dilated. After bringing down the feet, the birth of the body should be slowly effected, with the view of ensuring as complete subsequent contraction as possible. If the head has descended in the pelvis, of course turning is impossible and the forceps must be applied.

Difficulties arising from Locked Twins.—Occasionally, very serious difficulties arise from parts of both fetuses presenting simultaneously, and thus impeding the entrance of either child into the pelvis, or getting locked together, so as to render delivery impossible without artificial aid. Such difficulties are not apt to arise in the more ordinary cases, in which each child has its own bag of membranes, since then the fetuses are kept entirely separate, but in those in which the twins are contained in a common amniotic cavity or in which both sacs have burst simultaneously. They are very puzzling to the obstetrician, and it may be far from easy to discover the cause of the obstruction. Nor is it possible to lay down any positive rules for their management, which must be governed to a considerable extent by the circumstances of each individual case.

Both Heads Presenting Simultaneously.—Sometimes both heads present simultaneously at the brim, and then neither can enter unless they be unusually small or the pelvis very capacious, when both may descend;

or, rather, the first head may descend low into the pelvic cavity, and then the second head enters the brim and gets jammed against the thorax of the first child (Fig. 126). Reimann¹ relates a curious example of this, in which he delivered the head first with the forceps, but found the body would not follow, and on examination a second head was found in the pelvis. He then applied the forceps to the second head; the body of the first child was then born, and afterward that of the second. Such a

FIG. 126.



* Shows Head-Locking, both Children Presenting Head First. (After Barnes.)

mechanism must clearly have been impossible unless the pelvis had been extremely large.

Whenever both heads are felt at the brim, it will generally be found possible to get one out of the way by appropriate manipulation, one hand being passed into the vagina, the other aiding its action from without. Then the forceps may be applied to the other head, so as to engage it at once in the pelvic cavity. If both have actually passed into the pelvis, as in the case just alluded to, the difficulty will be much greater. It will generally be easier to push up the second head, while the lower is drawn out by the forceps, than to deliver the second, leaving the first *in situ*.

Foot or Hand with Head.—In other cases a foot or hand may descend along with the head, and even the four feet may present simultaneously. The rule in the former case is to push the part descending with the head out of the way, and in the latter to disengage one child as soon as possible. Great care is necessary, or we might possibly bring down the limbs of separate children.


Two Heads interlocking in Head and Breech Presentations.—The most common kind of difficulty is when the first child presents by the

¹ *Arch. f. Gynak.*, 1871.

breech and is delivered as far as the head, which is then found to be locked with the head of the second child, which has descended into the pelvic cavity (Fig. 127).

Here it is clear that the obstruction must be very great, and, unless the children are extremely small, insuperable. The first endeavor should be to disentangle the head: this is sometimes feasible if the second be not deeply engaged in the pelvis, and the hand be passed up so as to push it out of the way. This will but rarely succeed: then it may be possible to apply the forceps to the second head and drag it past the body

FIG. 127.



Shows Head-locking, First Child coming Feet First; Impaction of Heads from Wedging in Brim. (After Barnes.)

D. Apex of wedge. E, C. Base of wedge which cannot enter brim. A, B. Line of decapitation to decompose wedge and enable head of second child to pass.

of the first child; and this is the method recommended by Reimann, who has written an excellent paper on the subject.¹ Generally, the sacrifice of one of the children is essential, and as the body of the first child must have been born for some time, it is probable that the pressure to

¹ *American Journal of Obstetrics*, January, 1877.

which it has been subjected will have already imperilled, if it has not destroyed, its life, and therefore the plan usually recommended is to decapitate. This can easily be done with scissors or a wire *écraseur*, after which the second child is expelled without difficulty, leaving the head of the first in utero to be subsequently dealt with.

Another mode of managing these cases is to perforate the upper head and draw it past the lower with the cephalotribe or craniotomy-forceps. This plan has the disadvantage of probably sacrificing both children, since the other child can hardly survive the pressure and delay, whereas the former plan gives the second child a fair chance of being born alive.

Double Monsters.—In connection with the subject of twin labor we may consider those rare cases in which the bodies of the fetuses are partially fused together. The mechanism and management of delivery in cases of double monstrosity have attracted comparatively little attention, no doubt because authors have considered them matters of curiosity merely, rather than of practical importance.

The frequent occurrence of such monstrosities in our museums, and the numerous cases scattered through our periodical literature, are sufficient to show that they are not so very rare as we might be inclined to imagine; and, as they are likely to give rise to formidable difficulties in delivery, it cannot be unimportant to have a clear idea of the usual course taken by nature in effecting such births, with a view of enabling us to assist in the most satisfactory manner should a similar case come under our observation.

Unfortunately, the authors who have placed on record the birth of double monsters have generally occupied themselves more with a description of the structural peculiarities of the fetuses than with the mechanism of their delivery; so that, although the cases to be met with in medical literature are very numerous, comparatively few of them are of real value from an obstetric point of view. Still, I have been able to collect the details of a considerable number¹ in which the history of the labor is more or less accurately described; and doubtless a more extensive research would increase the list.

For obstetric purposes we may confine our attention to four principal varieties of double monstrosity which are met with far more frequently than any others. These are:

A. Two nearly separate bodies united in front, to a varying extent, by thorax or abdomen.

B. Two nearly separate bodies united back to back by the sacrum and lower part of the spinal column.

C. Dicephalous monsters, the bodies being single below, but the heads separate.

D. The bodies separate below, but the heads partially united.

This classification by no means includes all the varieties of monsters that we may meet with. It does, however, include all that are likely to give rise to much difficulty in delivery; and all the cases I have collected may be placed under one of these divisions.

The first point that strikes us in looking over the history of these deliveries is the frequency with which they have been terminated by the

¹ *Obst. Trans.*, vol. viii.

natural powers alone, without any assistance on the part of the accoucheur. Thus, out of the 31 cases, no less than 20 were delivered naturally, and apparently without much trouble. Nothing can better show the wonderful resources of nature in overcoming difficulties of a very formidable kind.

It is pretty generally assumed by authors that the children are necessarily premature, and therefore of small size, and that delivery before the full term is rather the rule than the exception. Dugés states that the children are often dead, and that putrefaction has taken place, which facilitates their expulsion. Both these assumptions seem to me to have been made without sufficient authority, and not to be borne out by the recorded facts. In only 1 of the 31 cases is it mentioned that the children were premature; nor is there any sufficient reason that I can see why labor should commence before the full term of gestation.

Class A.—By far the greatest number are included in the first class—that in which the bodies are nearly separate, but united by some part of the thorax or abdomen. This is the division which includes the celebrated Siamese Twins, an account of whose birth, I may observe, I have not been able to discover.¹ Out of the 31 cases, 19 come under this heading. The details of the labor are briefly as follows: 1 died undelivered; 8 were terminated by the natural powers, in 3 of which the feet, and in 3 the head, presented; in 2 the presentation is doubtful; 6 were delivered by turning or by traction on the lower extremities; 4 were delivered instrumentally.

Footling Presentation is the most Favorable.—The details of the cases in which the feet presented or in which turning was performed clearly show that footling presentation was by far the most favorable, and it is fortunate the feet often present naturally. The inference of course is, that version should be resorted to whenever any other presentation is met with in cases of double monstrosity of this type; but, unfortunately, this rule could rarely be carried into execution, since we possess no means of diagnosing the junction of the fœtuses at a sufficiently early stage of labor to admit of turning being performed. It is only under exceptionally favorable circumstances that this can be done; as, for example, in a case recorded by Molas,² in which both heads presented, but neither would enter the brim of the pelvis.

The Chief Difficulty is in the Delivery of the Heads.—The great difficulty must, of course, be in the delivery of the heads, for in all the recorded cases, with one exception, the bodies have passed through the pelvis parallel to each other with comparative ease until the necks have appeared, and then, as a rule, they could be brought no farther. It is clear that the remainder of the fœtuses could no longer pass simultane-

¹ The mother of these twins was a Chinese half-breed, short, and with a broad pelvis, and had borne several children previously. She stated on several occasions, in conversation with parties in Siam, that the twins were born reversed, the feet of one being followed by the head of the other, and that they were very small and feeble at birth and for several months afterward. The twins confirmed this statement by affirming that they could, when little boys at play on the ground, turn themselves end for end upon the ensiform attachment up to the age of ten or twelve, the attachment being then soft and pliable (Harris's note to second American edition).

² *Mém. de l'Académie*, vol. i.

ously, and were direct traction continued the heads would be inextricably fixed above the brim. In accordance with the direction of the pelvic axes the posterior head must first engage in the inlet; and in order to effect this it will be necessary to carry the bodies of the children well over the abdomen of the mother. This seems to be a point of primary importance. It would also be advisable to see that the bodies are made to pass through the pelvis with their backs in the oblique diameter. By this means more space is gained than if the backs were placed antero-posteriorly, while at the same time there is less chance of the heads hitching against the promontory of the sacrum and symphysis pubis, which otherwise would be very apt to occur.

Mode of Delivery when the Head Presents.—When the head presents, and the labor is terminated by the natural powers, delivery seems to be accomplished in one of two ways.

In the first and more common the head and shoulders of one child are born, its breech and legs being subsequently pushed through the pelvis by a process similar to that of spontaneous evolution; and afterward the second child probably passes footling without much difficulty.

Barkow relates a case in which *both* heads were delivered by the forceps, the bodies subsequently passing simultaneously. Two similar instances are recorded in the third and sixth volumes of the *Obstetrical Transactions*. When delivery takes place in this manner the head of the second child must fit into the cavity formed by the neck of the first, and the pelvis must necessarily be sufficiently roomy to admit of the expulsion of the head of the second child, while its cavity is diminished in size by the presence of the neck and shoulders of the first. Either of these processes must obviously require exceptionally favorable conditions as regards the size of the child and the pelvis, and the difficulty in the way of delivery must be much greater than when the lower extremities present. Therefore, I think the rule should be laid down that when the nature of the case is made out (and for the purpose of accurate diagnosis a complete examination under anæsthesia should be practised) turning should be performed and the feet brought down.

Mutilation of the Fœtuses.—In the event of its being found impossible to effect delivery after a considerable portion of the bodies is born, no resource remains but the mutilation of the body of one child so as to admit of the passage of the other. This was found necessary in one case in which the children presented by the feet and were born as far as the thorax, but could get no farther. The body of the anterior child was removed by a circular incision as far as it had been expelled, which allowed the remaining portion, consisting of the head and shoulders, to re-enter the uterus: after this the posterior child was easily extracted, and the mutilated fœtus followed without difficulty.

Class B.—In Class B, in which the children are united back to back, 3 cases are recorded, all of which were delivered by the natural powers. One of these is the case of Judith and Hélène, the celebrated Hungarian twins, who lived to the age of twenty-one. Hélène was born as far as the umbilicus, and after the lapse of three hours her breech and legs descended. Judith was expelled immediately afterward, her feet descend-

ing first. [¹] Exactly the same process occurred in a case described by M. Norman, the children being also born alive, and dying on the ninth day.

Labor is Easier than in Class A.—It is probable that labor is easier in this class of double monsters than in the former, because the children are so joined that there is no necessity for the bodies to be parallel to each other during birth when the head presents, and after the birth of the head and shoulders of the first child its breech and lower extremities are evidently pushed down and expelled by a process of spontaneous evolution. If the feet originally presented, the mechanism of delivery and the rules to be followed would be the same as in Class A; but the difficulty would probably be greater, since the juncture is not so flexible, and a more complete parallelism of the bodies would be necessary during extraction.

Class C.—In Class C, that of the dicephalous monster, I have found the description of the birth of 8 cases, 3 of which were terminated by the natural powers. In 2 of these the process of evolution was the main agent in delivery, one head being born and becoming fixed under the arch of the pubes, the body being subsequently pushed past it, and the second head following without difficulty. This process failing, the proper course is to decapitate the first-born head, and then bring down the feet of the child, when delivery can be accomplished with ease. This was the course adopted in 2 out of the 8 cases; and it may be done with the less hesitation since, from their structural peculiarities, it is exceedingly improbable that monsters of this kind should survive. In the third case, terminated naturally, the heads were said to have been born simultaneously, but it seems probable that the one head lay in the hollow formed by the neck of the other, and so rapidly followed it. If the feet presented, the case may be managed in the same manner as in Class A.

Class D.—Monstrosities of Class D, in which the heads are united, the bodies being distinct, appear to be the most uncommon of all, and I can find the description of delivery in only 2 cases. One of these gave rise to great difficulty; the labor in the other was easy. We should scarcely anticipate much difficulty in the birth of monsters of this type, for if the head presented and would not pass, we should naturally perform craniotomy; and if the bodies came first, the delivery of the monstrous head could readily be accomplished by perforation.

Result to the Mothers.—The result to the mothers in all these cases seems to have been very favorable. There is only one in which the death of the mother is recorded; and although in many the result is not mentioned, we may fairly assume that recovery took place.

Among difficulties in labor, some of the most important are due to morbid conditions of the fœtus itself.

Intra-uterine Hydrocephalus.—Of these, the most common as well as

[¹ The celebrated colored Carolina twins, born July 11, 1851, and still living, were brought into the world by the same method, but the mother, having a large pelvis, "had a brief and easy" delivery. The larger of the two girls also came first, as in the Tzoni case of 1701. These twins are now (1885) twelve years older than the Hungarian sisters were at death, being 33 years of age.—Ed.]

the most serious is caused by intra-uterine hydrocephalus (giving rise to a collection of watery fluid within the cranium), by which the dimensions of the child's head are enormously increased and the due relations between it and the pelvic cavity entirely destroyed (Fig. 128).

Its Danger both as Regards the Mother and Child.—Fortunately, this disease is of comparatively rare occurrence, for it is one of great gravity both as regards the mother and child. As regards the mother, the serious character of the complication is proved by the statistics of Dr. Keiller of Edinburgh, who found that out of 74 cases no less than 16 were accompanied by rupture of the uterus. The reason of the danger to which the mother is subjected is obvious. In some few cases, indeed, the head is so compressible that, provided the amount of contained fluid

FIG. 128.



Labor Impeded by Hydrocephalus.

be small, it may be sufficiently diminished in size by the moulding to which it is subjected to admit of its being squeezed through the pelvis. In the majority of cases, however, the size of the head is too great for this to occur. The uterus therefore exhausts itself, and may even rupture, in the vain endeavor to overcome the obstacle; while the large and distended head presses firmly on the cervix or on the pelvic tissues if the os be dilated, and all the evil effects of prolonged compression are apt to follow.

Its Diagnosis is not always Easy.—The diagnosis of intra-uterine hydrocephalus is by no means so easy as the description in obstetric works would lead us to believe. It is true that the head is much larger and more rounded in its contour than the healthy foetal cranium, and also that the sutures and fontanelles are more wide, and admit occasionally of fluctuation being perceived through them. Still, it is to be remembered that the head is always arrested above the brim, where it is consequently high up and difficult to reach, and where these peculiarities

are made out with much difficulty. As a matter of fact, the true nature of the case is comparatively rarely discovered before delivery ; thus, Chaussier¹ found that in more than one-half of the cases he collected an erroneous diagnosis had been made.

Method of Diagnosis.—Whenever we meet with a case in which either the history of previous labor or a careful examination convinces us that there is no obstacle due to pelvic deformity, in which the pains are strong and forcing, but in which the head persistently refuses to engage in the brim, we may fairly surmise the existence of hydrocephalus. Nothing, however, short of a careful examination under anæsthesia, the whole hand being passed into the vagina so as to explore the presenting part thoroughly, will enable us to be quite sure of the existence of this complication. Under these circumstances such a complete examination is not only justified, but imperative ; and when it has been made the difficulties of diagnosis are lessened, for then we may readily make out the large round mass, softer and more compressible than the healthy head, the widely-separated sutures, and the fluctuating fontanelles.

Pelvic Prescissions are Frequently met with.—In a considerable proportion of cases—as many, it is said, as 1 out of 5—the fetus presents by the breech. The diagnosis is then still more difficult ; for the labor progresses easily until the shoulders are born, when the head is completely arrested, and refuses to pass with any amount of traction that is brought to bear on it. Even the most careful examination may not enable us to make out the cause of the delay, for the finger will impinge on the comparatively firm base of the skull, and may be unable to reach the distended portion of the cranium. At this time abdominal palpation might throw some light on the case, for, the uterus being tightly contracted round the head, we might be able to make out its unusual dimensions. The wasted and shrivelled appearance of the child's body, which so often accompanies hydrocephalus, would also arouse suspicion as to the cause of delay. On the whole, such cases may be fairly assumed to be less dangerous to the mother than when the head presents ; for in the latter the soft parts are apt to be subjected to prolonged pressure and contusion, while in the former delay does not commence till after the shoulders are born, and then the character of the obstacle would be sooner discovered and appropriate means earlier taken to overcome it.

Treatment.—The treatment is simple, and consists in tapping the head, so as to allow the cranial bones to collapse. There is the less objection to this course since the disease almost necessarily precludes the hope of the child's surviving. The aspirator would draw off the fluid effectually, and would at least give the child a chance of life ; and under certain circumstances the birth of a child who lives for a short time only may be of extreme legal importance. More generally the perforator will be used, and as soon as it has penetrated a gush of fluid will at once verify the diagnosis. Schroeder recommends that after perforation turning should be performed, on account of the difficulty with which the flaccid head is propelled through the pelvis. This seems a very unnecessary complication of an already sufficiently troublesome case. As a rule, when once the fluid has been evacuated, the pains being strong, as

¹ *Gazette médicale*, 1864.

they generally are, no delay need be apprehended. Should the head not come down, the cephalotribe may be applied, which takes a firmer grasp than the forceps, and enables the head to be crushed to a very small size and readily extracted.

Treatment when the Breech Presents.—When the breech presents, the head must be perforated through the occipital bone; and generally this may be accomplished behind the ear without much difficulty. In a case of Tarnier's¹ the vertebral column was divided by a bistoury, and an elastic male catheter introduced into the vertebral canal, through which the intra-cranial fluid escaped, the labor being terminated spontaneously. In any case in which it is found difficult to reach the skull with the perforator this procedure should certainly be tried.

Other forms of dropsical effusion may give rise to some difficulty, but by no means so serious. In a few rare cases the thorax has been so distended with fluid as to obstruct the passage of the child. Ascites is somewhat more common, and occasionally the child's bladder is so distended with urine as to prevent the birth of the body. The existence of any of these conditions is easily ascertained; for the head or breech, whichever happens to present, is delivered without difficulty, and then the rest of the body is arrested. This will naturally cause the practitioner to make a careful exploration, when the cause of the delay will be detected.

The treatment consists in the evacuation of the fluid by puncture. In the case of ascites this should always be done, if possible, by a fine trocar or aspirator, so as not to injure the child. This is all the more important since it is impossible to distinguish a distended bladder from ascites, and an opening of any size into that viscus might prove fatal, whereas aspiration would do little or no harm, and would prove quite as efficacious.

Fœtal Tumors Obstructing Delivery.—Certain fœtal tumors may occasion dystocia, such as malignant growths or tumors of the kidney, liver, or spleen. Cases of this kind are recorded in most obstetric works. Hydro-encephalocele or hydro-rachitis, depending on defective formation of the cranial or spinal bones, with the formation of a large protruding bag of fluid, is not very rare. The diagnosis of all such cases is somewhat obscure, nor is it possible to lay down any definite rules for their management, which must vary according to the particular exigencies. The tumors are rarely of sufficient size to prove formidable obstacles to delivery, and many of them are very compressible. This is specially the case with the spina bifida and similar cystic growths. Puncture, and in the more solid growths of the abdomen or thorax evisceration, may be required.

Other Congenital Deformities.—Other deformities, such as the anencephalous fœtus, or defective development of the thorax or abdominal parietes with protrusion of the viscera, are not likely to cause difficulty, but they may much embarrass the diagnosis by the strange and unusual presentation that is felt. If in any case of doubt a full and careful examination be undertaken, introducing the whole hand if necessary, no serious mistake is likely to be made.

¹ Hergott, *Maladies Fœtales qui peuvent faire obstacle à l'accouchement*, Paris, 1878.

are made out with much difficulty. As a matter of fact, the true nature of the case is comparatively rarely discovered before delivery; thus, Chaussier¹ found that in more than one-half of the cases he collected an erroneous diagnosis had been made.

Method of Diagnosis.—Whenever we meet with a case in which either the history of previous labor or a careful examination convinces us that there is no obstacle due to pelvic deformity, in which the pains are strong and forcing, but in which the head persistently refuses to engage in the brim, we may fairly surmise the existence of hydrocephalus. Nothing, however, short of a careful examination under anaesthesia, the whole hand being passed into the vagina so as to explore the presenting part thoroughly, will enable us to be quite sure of the existence of this complication. Under these circumstances such a complete examination is not only justified, but imperative; and when it has been made the difficulties of diagnosis are lessened, for then we may readily make out the large round mass, softer and more compressible than the healthy head, the widely-separated sutures, and the fluctuating fontanelles.

Pelvic Presentations are Frequently met with.—In a considerable proportion of cases—as many, it is said, as 1 out of 5—the fetus presents by the breech. The diagnosis is then still more difficult; for the labor progresses easily until the shoulders are born, when the head is completely arrested, and refuses to pass with any amount of traction that is brought to bear on it. Even the most careful examination may not enable us to make out the cause of the delay, for the finger will impinge on the comparatively firm base of the skull, and may be unable to reach the distended portion of the cranium. At this time abdominal palpation might throw some light on the case, for, the uterus being tightly contracted round the head, we might be able to make out its unusual dimensions. The wasted and shrivelled appearance of the child's body, which so often accompanies hydrocephalus, would also arouse suspicion as to the cause of delay. On the whole, such cases may be fairly assumed to be less dangerous to the mother than when the head presents; for in the latter the soft parts are apt to be subjected to prolonged pressure and contusion, while in the former delay does not commence till after the shoulders are born, and then the character of the obstacle would be sooner discovered and appropriate means earlier taken to overcome it.

Treatment.—The treatment is simple, and consists in tapping the head, so as to allow the cranial bones to collapse. There is the less objection to this course since the disease almost necessarily precludes the hope of the child's surviving. The aspirator would draw off the fluid effectually, and would at least give the child a chance of life; and under certain circumstances the birth of a child who lives for a short time only may be of extreme legal importance. More generally the perforator will be used, and as soon as it has penetrated a gush of fluid will at once verify the diagnosis. Schroeder recommends that after perforation turning should be performed, on account of the difficulty with which the flaccid head is propelled through the pelvis. This seems a very unnecessary complication of an already sufficiently troublesome case. As a rule, when once the fluid has been evacuated, the pains being strong, as

¹ Gazette médicale, 1864.

CHAPTER XII.

DEFORMITIES OF THE PELVIS.

Importance of the Subject.—Deformities of the pelvis form one of the most important subjects of obstetric study, for from them arise some of the gravest difficulties and dangers connected with parturition. A knowledge, therefore, of their causes and effects, and of the best mode of detecting them, either during or before labor, is of paramount necessity; but the subject is far from easy, and it has been rendered more difficult than it need be from over-anxiety on the part of obstetricians to force all varieties of pelvic deformities within the limits of their favorite classification.

Difficulties of Classification.—Many attempts in this direction have been made, some of which are based on the causes on which the deformities depend, others on the particular kind of deformity produced. The changes of form, however, are so various and irregular, and similar or apparently similar causes so constantly produce different effects, that all such endeavors have been more or less unsuccessful. For example, we find that rickets (of all causes of pelvic deformity the most important) generally produces a narrowing of the conjugate diameter of the brim, while the analogous disease, osteo-malacia, occurring in adult life, generally produces contraction of the transverse diameter, with approximation of the pubic bones and relative or actual elongation of the conjugate diameter. We might, therefore, be tempted to classify the results of these two diseases under separate heads, did we not find that when rickets affects children who are running about and subject to mechanical influences similar to those acting upon patients suffering from osteo-malacia, a form of pelvis is produced hardly distinguishable from that met with in the latter disease, which by some authors is described as the pseudo-osteo-malacic.

Most Simple Classification.—On the whole, therefore, the most simple as well as the most scientific classification is that which takes as its basis the particular seat and nature of the deformity. Let us first glance at the most common causes.

Causes of Pelvic Deformity.—The key to the particular shape assumed by a deformed pelvis will be found in a knowledge of the circumstances which lead to its regular development and normal shape in a state of health. The changes produced may, almost invariably, be traced to the action of the same causes which produce a normal pelvis, but which, under certain diseased conditions of the bones or articulations, induce a more or less serious alteration in form. These have been already described in discussing the normal anatomy of the pelvis; and it will be remembered that they are chiefly the weight of the body transmitted to the iliac bones through the sacro-iliac joints, and counter-pressure on these acting through the acetabula. Sometimes they act in excess on

bones which are healthy, but possibly smaller than usual, and the result may be the formation of certain abnormalities in the size of the various pelvic diameters. At other times they operate on bones which are softened and altered in texture by disease, and which therefore yield to the pressure far more than healthy bones.

The two diseases which chiefly operate in causing deformity are rickets and osteo-malacia. Into the essential nature and symptomatology of these complaints it would be out of place to enter here: it may suffice to remind the reader that they are believed to be pathologically similar diseases, with the important practical distinction that the former occurs in early life before the bones are completely ossified, and that the latter is a disease of adults producing softening in bones that have been hardened and developed. This difference affords a ready explanation of the generally resulting varieties of pelvic deformity.

Effects of Rickets.—Rickets commences very early in life—sometimes, it is believed, even in utero. It rarely produces softening of the entire bones, and only in cases of very great severity of those parts of the bones that have been already ossified. The effects of the disease are principally apparent in the cartilaginous portions of the bones, in which osseous deposit has not yet taken place. The bones, therefore, are not subject to uniform change, and this fact has an important influence in determining their shape. Rickety children also have imperfect muscular development; they do not run about in the same way as other children; they are often continuously in the recumbent or sitting posture, and thus the weight of the trunk is brought to bear, more than in a state of health, on the softened bones. For the same reason, counter-pressure through the acetabula is absent or comparatively slight. When, however, the disease occurs for the first time in children who are able to run about, the latter comes into operation and modifies the amount and nature of the deformity. It is to be observed that in rickety children the bones are not only altered in form from pressure, but are also imperfectly developed; and this materially modifies the deformity. When ossific matter is deposited, the bones become hard and cease to bend under external influences, and retain for ever the altered shape they have assumed.

Effects of Osteo-malacia.—In osteo-malacia, on the contrary, the already hardened bones become softened uniformly through all their textures, and thus the changes which are impressed upon them are much more regular and more easily predicated. It is, however, an infinitely less common cause of pelvic deformity than rickets, as is evidenced by the fact that in the Paris Maternity, in a period of sixteen years, 402 cases of deformity due to rickets occurred to 1 due to osteo-malacia.¹

Their Varying Frequency.—The frequency of both diseases varies greatly in different countries and under different circumstances. Rickets is much more common amongst the poor of large cities, whose children are ill-fed, badly clothed, kept in a vitiated atmosphere, and subjected to unfavorable hygienic conditions. Deformities are therefore more common in them than in the more healthy children of the upper classes or of the rural population. The higher degrees of deformity, necessitating the Cæsarean section or craniotomy, are in this country of extreme rar-

¹ Stanesco, *Recherches cliniques sur les Rétrécissements du Bassin.*

ity, while in certain districts on the Continent they seem to be so frequent that these ultimate resources of the obstetric art have to be constantly employed.

Effects of Ossification of the Pelvic Articulations.—In another class of cases the ordinary shape is modified by weight and counter-pressure operating on a pelvis in which one or more of the articulations is ossified. In this way we have produced the *obliquely-ovate* pelvis of Naegele or the still more uncommon *transversely-contracted* pelvis of Robert.

Other Causes of Pelvic Deformity.—A certain number of deformed pelves cannot be referred to a modification of the ordinary developmental changes of the bones. Amongst these are the deformities resulting from spondylolisthesis, or downward dislocation of the lower lumbar vertebrae; from displacements of the sacrum caused by curvatures of the spinal column, producing the kyphotic and scoliotic pelves; or from diseases of the pelvic bones themselves, such as tumors, malignant growths, and the like.

Equally Enlarged Pelvis.—The first class of deformed pelves to be considered is that in which the diameters are altered from the usual standard without any definite distortion of the bones; and such are often mere congenital variations in size for which no definite explanation can be given. Of this class is the pelvis which is equally enlarged in all its diameters (*pelvis æquabiliter justo major*), which is of no obstetric consequence, except inasmuch as it may lead to precipitate labor, and is not likely to be diagnosed during life.

Equally Contracted Pelvis.—The corresponding diminution of all the pelvic diameters (*pelvis æquabiliter justo minor*) may be met with in women who are apparently well formed in every respect and whose external conformation and previous history gave no indication of the abnormality.[¹] Sometimes the diminution amounts to half an inch or more, and it can readily be understood that such a lessening in the capacity of the pelvis would give rise to serious difficulty in labor. Thus, in 3 cases recorded by Naegele a fatal result followed—in 2 after difficult instrumental delivery, and in the third after rupture of the uterus. The equally lessened pelvis, however, is of great rarity. An unusually small pelvis may be met with in connection with general small size, as in dwarfs. It does not necessarily follow, because a woman is a dwarf, that the pelvis is too small for parturition. On the contrary, many such women have borne children without difficulty.

The Undeveloped Pelvis.—In some cases a pelvis retains its infantile characteristics after puberty (Fig. 129). The normal development of the pelvis has been interfered with, possibly, from premature ossification of

[¹ It is possible for a lady to be tall, erect, weigh 180 pounds, and be conspicuous for her fine appearance, when she has a *justo minor* pelvis of very small interior dimensions. I examined such a patient some years ago, and not only had she the smallest vagina I have ever explored in a well-grown woman, but it has recently been found in labor that her pelvis would not admit of her being delivered of a full-grown living child. In fact, it is doubtful whether she could be delivered of one alive much later than the seventh month. Married twice and unimpregnated for years, I was surprised at her becoming at last pregnant, as an index finger filled her vagina tightly. After a labor of three days, and when the fetus was dead, it was delivered with a crushed head after long and powerful traction, and she made a good recovery. Having large hips, she was under an impression that her pelvis was of corresponding development.—ED.]

bones which are healthy, but possibly smaller than usual, and the result may be the formation of certain abnormalities in the size of the various pelvic diameters. At other times they operate on bones which are softened and altered in texture by disease, and which therefore yield to the pressure far more than healthy bones.

The two diseases which chiefly operate in causing deformity are rickets and osteo-malacia. Into the essential nature and symptomatology of these complaints it would be out of place to enter here: it may suffice to remind the reader that they are believed to be pathologically similar diseases, with the important practical distinction that the former occurs in early life before the bones are completely ossified, and that the latter is a disease of adults producing softening in bones that have been hardened and developed. This difference affords a ready explanation of the generally resulting varieties of pelvic deformity.

Effects of Rickets.—Rickets commences very early in life—sometimes, it is believed, even in utero. It rarely produces softening of the entire bones, and only in cases of very great severity of those parts of the bones that have been already ossified. The effects of the disease are principally apparent in the cartilaginous portions of the bones, in which osseous deposit has not yet taken place. The bones, therefore, are not subject to uniform change, and this fact has an important influence in determining their shape. Rickety children also have imperfect muscular development; they do not run about in the same way as other children; they are often continuously in the recumbent or sitting posture, and thus the weight of the trunk is brought to bear, more than in a state of health, on the softened bones. For the same reason, counter-pressure through the acetabula is absent or comparatively slight. When, however, the disease occurs for the first time in children who are able to run about, the latter comes into operation and modifies the amount and nature of the deformity. It is to be observed that in rickety children the bones are not only altered in form from pressure, but are also imperfectly developed; and this materially modifies the deformity. When ossific matter is deposited, the bones become hard and cease to bend under external influences, and retain for ever the altered shape they have assumed.

Effects of Osteo-malacia.—In osteo-malacia, on the contrary, the already hardened bones become softened uniformly through all their textures, and thus the changes which are impressed upon them are much more regular and more easily predicated. It is, however, an infinitely less common cause of pelvic deformity than rickets, as is evidenced by the fact that in the Paris Maternity, in a period of sixteen years, 402 cases of deformity due to rickets occurred to 1 due to osteo-malacia.¹

Their Varying Frequency.—The frequency of both diseases varies greatly in different countries and under different circumstances. Rickets is much more common amongst the poor of large cities, whose children are ill-fed badly clothed, kept in a vitiated atmosphere, and subjected to unfavorable conditions. Deformities are therefore more common in the healthy children of the upper classes or in the children of higher degrees of deformity, necessitating operation, are in this country of extreme rarity.

sur les Rétrécissements du Bassin.

when more marked it almost invariably is so. When unconnected with rickets, it probably can be traced to some injurious influence before the bones have ossified, such as increased pressure of the trunk from carrying weights in early childhood and the like. By this means the sacrum is unduly depressed and projects forward, so as to slightly narrow the conjugate diameter.

Mode of Production in Rickets.—When caused by rickets, the amount of the contraction varies greatly, sometimes being very slight, sometimes sufficient to prevent the passage of the child altogether, and to necessitate craniotomy or the Cæsarean section. The sacrum, softened by the disease, is pressed vertically downward by the weight of the body, its descent being partially resisted by the already ossified portions of the bone, so that the result is a downward and forward movement of the promontory. The upper portion of the sacral cavity is thus directed more backward; but, as the apex of the bone is drawn forward by the attachment of the perineal muscles to the coccyx and by the sacro-ischiatic ligaments, a sharp curve of its lower part in a forward direction is established.

FIG. 130.



Scolio-rachitic Pelvis. (From a specimen in the Museum of St. Bartholomew's Hospital.)

Occasional Increase of Transverse Diameter.—The depression of the sacral promontory would tend to produce strong traction, through the sacro-iliac ligaments, on the posterior end of the sacro-cotyloid beams, and thus induce expansion of the iliac bones and consequent increase of the transverse diameter of the brim. So an unusual length of the transverse diameter is very often described as accompanying this deformity, but probably it is not so often apparent as might otherwise be expected, on account of the imperfect development of the bones generally accompanying rickets; and Barnes¹ says that in the parts of London where deformities are most rife any enlargement of the transverse diameter is exceedingly rare.

¹ *Lectures on Obst. Operations*, p. 280.

The Sclio-cachitic and Scoliotic Pelvis.—Frequently the sacrum is not only depressed, but displaced more or less to one side, most generally to the left, thus interfering with the regular shape of the deformed brim. This is often the result of a lateral flexion of the spinal column depending on the rachitic diathesis, and when well marked is known as the *sclio-cachitic pelvis* (Fig. 130), in which one side of the pelvis, that corresponding to the direction of the pelvic curve, is asymmetrical and contracted, the ilio-pectineal line being sharply curved inward about the site of the sacro-iliac synchondrosis, the symphysis pubis being displaced toward the opposite side. A somewhat similar but much less marked unilateral asymmetry may exist in cases of scolirosis [?] unconnected with rickets, but rarely to a sufficient degree to interfere materially with labor.

Cavity of the Pelvis is generally not Affected.—In most cases of this kind the cavity of the pelvis is not diminished in size, and is often even more than usually wide. The constant pressure on the ischia which the sitting posture of the child entails tends to force them apart, and also to widen

FIG. 131.



Hickey Pelvis, with Backward Depression of the Symphysis Pubis.

the pubic arch. Considerable advantage results from this in cases in which we have to have perform obstetric operations, as it gives plenty of room for manipulation.

Figure-of-Eight Deformity.—In a few exceptional cases the narrowing of the conjugate diameter is increased by a backward depression of the symphysis pubis, which gives the pelvic brim a sort of figure-of-eight shape (Fig. 131). The most reasonable explanation of this peculiarity

[Although hunchbacks frequently have well-formed pelvis, it is not uncommon to find a deformed spine associated with an asymmetrical pelvis or even a much contracted one. Spinal distortion from caries, especially in the lumbar region, is thus associated, and the pelvic deformity will be increased if there has been catalgia, either double or single, or if from any cause one leg should be shorter than the other. In the records of the Pueri operation we find under "the cause of difficulty," "*pseudio-osteo-malacia*," "*lumbo-dorsal kyphosis*," "*kypho-scoliosis*," etc. *Pseudio-osteo-malacia* is the result of rickets in a walking child, the form of pelvis being changed mechanically, as in *osteo-malacia*. *Lumbo-dorsal kyphosis* may or may not give rise to the kyphotic pelvis, as much will depend upon the extent of vertebral caries. *Scoliosis* is apt to result from rickets, and may be associated with *lordosis*.

Scoliosis, from *scadon*, crooked—a distortion of the spine to one side.

Lordosis, from *luphe*, curved—applied particularly to the forward bending of the spine.

Kyphosis, from *kypos*, gibbous, arched, or vaulted—a hump or backward curvature of the spine.—Ed.]

seems to be that it is the result of the muscular contraction of the recti muscles at their point of attachment, when the centre of gravity of the body is thrown backward on account of the projection of the sacral promontory. Sometimes, also, the antero-posterior diameter of the cavity is unusually lessened by the disappearance of the vertical curvature of the sacrum, which, instead of forming a distinct cavity, is nearly flat (Fig. 132).

Spondylolisthesis.—In a few rare cases, to which attention was first called in 1853 by Kilian of Bonn, a very formidable narrowing of the conjugate diameter of the pelvic brim is produced by a downward displacement of the fourth and fifth lumbar vertebrae, which become dislocated forward, or, if not actually dislocated, at least separated from their several articulations to a sufficient extent to encroach very seriously on the dimensions of the pelvic inlet. This condition is known as *spondylolisthesis* [¹] (Fig. 133).

FIG. 132.



Flatness of Sacrum, with Narrowing of Pelvic Cavity.

FIG. 133.



Pelvis Deformed by Spondylolisthesis. (After Kilian.)

The effect of this is sufficiently obvious, for the projection of the lumbar vertebrae prevents the passage of the child. To such an extent is obstruction thus produced that in the majority of the recorded cases the Cæsarean section was necessary. The true conjugate diameter, that between the promontory of the sacrum and the symphysis pubis, is increased rather than diminished; but, for all practical purposes, the condition is similar to extreme narrowing of the conjugate from rickets, for the bodies of the displaced vertebrae project into and obstruct the pelvic brim.

The cause of this deformity seems to be different in different cases. In some it seems to have been congenital, and in others to have depended on some antecedent disease of the bones, such as tuberculosis or scrofula, producing inflammation and softening of the connection between the last lumbar vertebra and the sacrum, thus permitting downward displacement of the bones. Iambli believed that it generally followed spina

[¹ From σπονδύλος, the vertebra, and ολισθησις, a slipping or sliding.—ED.]

bifida which had become partially cured, but which had produced deformity of the vertebræ and favored their dislocation. Brodhurst,¹ on the other hand, thinks that it most probably depends on rachitic inflammation and softening of the osseous and ligamentous structures, and that it is not a dislocation in the strict sense of the word. This condition has recently been made the subject of special study by Dr. Franz Neugebauer,² who believes that the forward displacement is never the result of antecedent disease of the bones, but depends either on congenital want of development of the vertebral arches or on traumatism, such as fracture of the articular processes, which allows the weight of the trunk to displace the body of the last lumbar vertebra forward, either partially or entirely.

[We are indebted to Kilian of Germany for the first careful investigation of the true character of spondylolisthetic deformity, although the

FIG. 134.



[Spondylolisthesis. (After Neugebauer.)]

credit of initial mention is due to Rokitansky of Austria, who wrote in 1839, antedating the monograph of the former (1853) by fourteen years. No special mention is made of this peculiar lordosis by Rokitansky in his *Manual of Pathological Anatomy* in 1844, but in his *Lehrbuch* (1855) it is given, with due credit to Kilian. During the thirty years that have passed since Kilian prepared his paper from observations made upon three pelvises which had been obtained from subjects in whom the Cæsarean section had proved fatal, one of them after a second operation, there have appeared numerous monographs and descriptions of cases, much the most valuable and extensive of which are those by Dr. Franz Ludwig Neuge-

bauer of Warsaw and Dr. A. Swedelin of St. Petersburg, the latter of whom furnishes the bibliography of the subject. These valuable papers cover 153 and 40 pages, respectively, of the *Archiv für Gynækologie*, Berlin, the former having 87 illustrations and the latter 7, and may be found in vols. xix., xx., xxi., and xxii., for 1882-84.

The cause or causes of spondylolisthesis have not as yet been very satisfactorily determined. The disease in some of its features would appear to be a form of local malacosteon or of *osteitis deformans*, differing from the former in the fact that it may occur without pain, that the general health may not be impaired, and that the subjects may be nulliparous; which last is exceedingly rare in malacosteon. In the lordosis of ordinary malacosteon there is no sliding of the last lumbar vertebra, or this peculiar deformity would be much less infrequent. As it is,

¹ *Obst. Trans.*, vol. vi. p. 97.

² *Contribution à la Pathogénie du Bassin Vicié par le Glissement vertébral*, Paris, 1884.

after an extended search by a number of writers, we are limited to the study of 17 pelves and 25 clinical cases: the photographic process has also made us familiar with the contour of body produced by the disease. The effect of the spinal slipping and anterior curvature of the loins is to shorten the trunk of the woman, cause a pouch-like protrusion of her abdomen, and elevate and widen the hips. As the pelvis is not collapsed, there is no alteration in the pubes or hip-joints, and as many of the subjects are robust, their change of form is quite perceptible even in full dress. The only true test of the condition is the discovery *per vaginam* of the overhanging fifth lumbar vertebra, with which may also be felt the pulsation of the primitive iliacs, and in extreme cases even the bifurcation of the aorta. The shortening of the abdominal cavity forces the pregnant uterus into an extreme anterior obliquity, which, by interfering with the line of its expulsive action, adds to the difficulty produced by the narrowing of the conjugate diameter.

We have a special interest in the study of this deforming malady from the fact that one of the cases on record occurred in San Francisco under the care of Prof. James Blake of Toland Medical College, and was reported by him in the *Pacific Med. and Surg. Journal* of Feb., 1867. The subject of this disease was married at 15 years of age, at which time she weighed 101 pounds, but increased to 199 pounds by the time her first child was born. Her first and second labors were tedious, but the children were born alive; she aborted of another fœtus at four months, and later was delivered at maturity of four others, all dead, the conjugate space in the seventh labor being computed at $3\frac{1}{2}$ inches. This labor was so difficult that it was decided, in the event of another pregnancy, to bring on labor prematurely. She became pregnant for the eighth time at the age of 26, when she weighed 220 pounds. Labor was induced in the seventh month, but the fœtus was lost, as it weighed nearly six pounds and the lumbo-pubic space was reduced to 3 inches. This woman is said to have undergone the change in her vertebræ without pain or sign of ill-health, and to have retained a remarkable activity for her weight. After her eighth delivery she was up in six days, and down stairs in ten. The history of this case would indicate that the deforming process must have been slowly progressing during more than ten years.

In contrast with this painless case in a multipara we have the opposite in a nullipara, reported by Dr. Olshausen of Halle. The disease commenced in his patient when a girl of 18 with severe pains in the sacrum and hips, as in malacosteon. She had not had rickets in childhood, had enjoyed good health up this time, and was quite straight. As her disease progressed she found on awaking one morning that she could not straighten her spine, and was forced to walk with her body bent forward. She was put under medical treatment at the surgical clinic; had no fever, and in time ceased to suffer, and was discharged. Becoming pregnant at the age of 24, Dr. Olshausen delivered her in 1863 by the Cæsarean section: the child lived, but she was lost on the fourth day by peritonitis. The lumbo-pubic diameter was found to measure 3 inches, and the line of the conjugate struck the lower margin of the third lumbar vertebra.

Spondylolisthesis must not be confounded with that form of lumbar *lordosis* in which the fifth vertebra retains its proper articulation with the top of the sacrum, or with that produced by lumbo-sacral caries. In true spondylolisthetic deformity the superior strait is not distorted, but is encroached upon from above by the sagging down of the lumbar vertebræ, and particularly by the slipping forward of the entire body of the fifth. A vertical section of the lumbo-sacral spine will show that the displaced vertebra has in some cases been thinned, and that in others the top of the sacrum has been compressed and new bone thrown out to give additional support to the dislodged vertebra. To accomplish this without paralysis, there must necessarily be some softening of bone and cartilage, with relaxation of ligaments and a gradual giving way, especially at the sacro-vertebral articulation, with an enlargement of the spinal canal in the slipping portion to avoid pinching the spinal nerve. Certainly, the clinical history of some of the cases cannot be explained, except under the belief that there has been some antecedent bone disease affecting at least the articulating processes. Prof. Blake attributed the deformity in his case to a dislocation produced by a rapid increase in weight; but hundreds of women are subjected to the same cause and remain straight; and, besides, his patient must have had a very gradual dislocation. The theory of fracture of the articular processes is not at all satisfactory, as the gravity of the symptoms is not indicative of such an accident having occurred; neither is that of a congenital defect in the said processes, although this may account for some of the cases, especially if we admit that the defective articulations may become insufficient to support the weight of the trunk by reason of a weakening produced by disease.—Ed.]

Spondylolizema.—A somewhat analogous deformity has been described by Hergott¹ under the name of *spondylolizema*. In this the bodies of the lower lumbar vertebræ having been destroyed by caries, the upper lumbar vertebræ sink downward and forward, so as to obstruct the pelvic inlet and prevent the engagement of the fœtus. It thus differs from spondylolisthesis, in which there is dislocation, but not destruction, of the bodies of the lower lumbar vertebræ.

Narrowing of the Oblique Diameter.—The most marked examples of narrowing of both oblique diameters depend on osteo-malacia. In this disease, as has already been remarked, the bones are uniformly softened, and the alterations in form are further influenced by the fact that the disease commences after union of the separate portions of the ossa innominata has been completely effected. The amount of deformity in the worst cases is very great, and frequently renders delivery impossible without the Cæsarean section. Sometimes the softening of the bones proves of service in delivery, by admitting of the dilatation of the contracted pelvic diameter by the pressure of the presenting part or even by the hand. Some curious cases are on record in which the deformity was so great as to apparently require the Cæsarean section, but in which the softened bones eventually yielded sufficiently to render this unnecessary.

Mode of Production in Osteo-malacia.—The weight of the body

¹ Arch. de Tocologie, 1877.

depresses the sacrum in a vertical direction, and at the same time compresses its component parts together, so as to approximate the base and apex of the bone, and narrow the conjugate diameter of the brim by

FIG. 135.

Osteo-malacic Pelvis.[¹]

causing the promontory to encroach upon it. The most characteristic changes are produced by the pushing inward of the walls of the pelvis at the cotyloid cavities in consequence of pressure exerted at these points through the femora. The effect of this is to diminish both oblique diameters, giving the brim somewhat the shape of a trefoil or an ace of clubs. The sides of the pubes are at the same time approximated, and may become almost parallel, and the true conjugate may be even lengthened (Fig. 135). The tuberosities of the ischia are also compressed together with the rest of the lateral pelvic wall, so that the outlet is greatly deformed as well as the brim (Fig. 136).

FIG. 136.



Extreme Degree of Osteo-malacic Deformity.


[*Osteo-malacia not an American Disease.*—In not one of the 134 Cæsarean cases of the United States was the operation performed for this form of deformity. In a few instances the malady has been found

[¹ This form is known as *rostrate* or *beaked*.—ED.]

in foreigners, and the forceps or craniotomy used in their delivery. Many obstetricians of large experience have never seen a case, and I do not know of an instance of extreme rostration of the pelvis having been met with in this country.—ED.]

Obliquely-contracted Pelvis.—That form of deformity in which one oblique diameter only is lessened has received considerable attention,

FIG. 137.



Obliquely-contracted Pelvis.
(After Duncan.)

from having been made the subject of special study by Naegele, and is generally known as the *obliquely-contracted pelvis* (Fig. 137). It is a condition that is very rarely met with, although it is interesting from an obstetric point of view, as throwing considerable light on the mode in which the natural development of the pelvis is effected. It is difficult to diagnose, inasmuch as there is no apparent external deformity, and probably it has never, in fact, been detected before delivery. It has a very serious influence on labor: Litzmann found that out of 28 cases of this

deformity, 22 died in their labors and 5 more in subsequent deliveries. The prognosis, therefore, is very formidable, and renders a knowledge of this distortion, rare though it be, of importance.

Its essential characteristic is flattening and want of development of one side of the pelvis, associated with ankylosis of the corresponding sacro-iliac synchondrosis. The latter is probably always present, and it seems to be most generally a congenital malformation. The lateral half of the sacrum on the same side, and the entire innominate bone, are much atrophied. The promontory of the sacrum is directed toward the diseased side, and the symphysis pubis is pushed over toward the healthy side.

The main agent in the production of this deformity is the absence of the sacro-iliac joint, which prevents the proper lateral expansion of the pelvic brim on that side, and allows the counter-pressure, through the femur, to push in the atrophied os innominatum to a much greater extent than usual. The chief diminution in the length of the pelvic diameter is between the ilio-pectineal eminence of the affected side and the healthy sacro-iliac joint, while the oblique diameter between the ankylosed joint and the healthy os innominatum is of normal length.

[Another form of obliquely-contracted pelvis is the result of coxalgic atrophy of one ilium, produced by hip disease in early childhood, the side of the pelvis being stunted in growth, as well as the whole extremity attached to it. Two Cæsarean operations have been performed in the United States in cases having this form of deformity.—ED.]

Narrowing of the Transverse Diameter.—Transverse contraction of the pelvic brim is very much less common than narrowing of the conjugate diameter. It most frequently depends on backward curvature of the lower parts of the spinal column in consequence of disease of the vertebrae. This form of deformed pelvis is generally known as the *kyphotie* (Fig. 138). The effect of the spinal curvature is to drag the

promontory of the sacrum backward, so that it is high up and out of reach. By this means the antero-posterior diameter of the brim is increased, while the transverse is lessened; the relative proportion between the two is thus reversed. While the upper portion of the

FIG. 138.



Kyphotic Pelvis.

(From a specimen in the Museum of St. Bartholomew's Hospital.)

sacrum is displaced backward, its lower end is projected forward, so that the antero-posterior diameters of the cavity and outlet are considerably diminished. The ischial tuberosities are also nearer to each other, and the pubic arch is narrowed. Obstruction to delivery will be chiefly met with at the lower parts and outlet of the pelvic cavity, for, although the transverse diameter of the brim is narrowed, there is generally sufficient space for the passage of the head.

Robert's Pelvis.—Another form of transversely-contracted pelvis is known as *Robert's pelvis* (Fig. 139), having been first discovered by

Robert of Coblentz. It is in fact a double obliquely-contracted pelvis, depending on ankylosis of both sacro-iliac joints, and consequent defective development of the innominate bones. The shape of the pelvic brim is markedly oblong, and the sides of the pelvis are more or less parallel with each other. The outlet is also much contracted transversely. The amount of obstruction is very great, so that, according to Schroeder, out of 7 well-authenticated cases the Cæsarean section was required in 6.

FIG. 139.



Robert's or Double Obliquely-contracted Pelvis. (After Duncan.)

[The Porro-Cæsarean operation has only been called for in one instance, by reason of this rare condition. Prof. Josef Späth of Vienna operated with success to mother and child on March 7, 1883. The c. v. measured $3\frac{2}{3}$ inches, and the bis-ischiatic only 1 inch.—ED.]

Deformity from Old-standing Hip-joint Disease.—Another cause of transverse deformity occasionally met with is luxation of the head of the femur, depending on old-standing joint-disease. The head of the femur in this case presses on the innominate bone at the site of dislocation, and the result is that the iliac fossa on the affected side, or both if the accident happens on both sides, is pushed inward, the transverse diameter of the brim being lessened. The tuberosity of the ischium is, however, projected outward, so that the outlet of the pelvis is increased rather than diminished.

Deformity from Tumors, Fractures, etc.—Obstruction of the pelvic cavity from exostoses or other forms of tumors growing from the bones

FIG. 140.



Bony Growth from Sacrum obstructing the Pelvic Cavity.

is of great rarity (Fig. 140). It may, however, produce very serious dystocia. Several curious examples are collected in Mr. Wood's article on the pelvis, in some of which the obstruction was so great as to necessitate the Cæsarean section.[¹] Some of these growths were true exostoses, and, according to Stadtfeldt,² these are commonly found in pelves that are otherwise contracted; others, osteo-sarcomatous tumors attached to the pelvic bones, most generally the upper part of the sacrum; and others were malignant. In some cases spiculæ of bone have developed about the linea ilio-pectinea or other parts of the pelvis, which may not be sufficient to produce obstruction, but which may injure the uterus, or even the

fœtal head, when they are pressed upon them. Irregular projections may also arise from the callus of old fractures of the pelvic bones. All such cases defy classification, and differ so greatly in their extent and in their effect on labor that no rules can be laid down for them, and each must be treated on its own merits.

Effects of Contracted Pelvis in Labor.—The effects of pelvic contractions on labor vary, of course, greatly with the amount and nature of the deformity; but they must always give rise to anxiety, and in the graver degrees they produce the most serious difficulties we have to contend with in the whole range of obstetrics.

Nature of Uterine Action in Pelvic Deformity.—In the lesser degrees,

[¹ Ten Cæsarean operations in the United States saved four of the cases and five children. One fatal case had been in labor two weeks; one, four days; two, three days; and one, two days.—ED.]

² *Obstetrical Journal*, July, 1879.

in which the proportion between the presenting part and the pelvis is only slightly altered, we may observe little abnormal beyond a greater intensity of the pains and some protraction of the labor. It is generally observed that the uterine contractions are strong and forcible in cases of this kind, probably because of the increased resistance they have to contend against; and this is obviously a desirable and conservative occurrence, which may of itself suffice to overcome the difficulty. The first stage, however, is not unfrequently prolonged and the pains are ineffective, for the head does not readily engage in the brim, the uterus is more mobile than in ordinary labors, and it probably acts at a disadvantage.

Risk to the Mother.—In the more serious cases the mother is subjected to many risks, directly proportionate to the amount of obstruction and the length of the labor. The long-continued and excessive uterine action produced by the vain endeavors to push the child through the contracted pelvic canal, the more or less prolonged contusion and injury to which the maternal soft parts are necessarily subjected (not unfrequently ending in inflammation and sloughing with all its attendant dangers), and the direct injury which may be inflicted by the measures we are compelled to adopt for aiding delivery (such as the forceps, turning, craniotomy, or Cæsarean section),—all tend to make the prognosis a matter of grave anxiety.

Risk to the Child.—Nor are the dangers less to the child, and a very large proportion of stillbirths will always be met with. The infantile mortality may be traced to a variety of causes, the most important being the protraction of the labor and the continuous pressure to which the presenting part is subjected. For this reason, even in cases in which the contraction is so slight that the labor is terminated by the natural powers, it has been estimated that 1 out of every 5 children is stillborn; and as the deformity increases in amount, so, of course, does the prognosis to the child become more unfavorable.

Frequent Occurrence of Prolapse of the Cord.—Prolapse of the umbilical cord is of very frequent occurrence in cases of pelvic deformity, the tendency to this accident being traceable to the fact of the head not entering and occupying the upper strait of the pelvis as in ordinary labors, and thus leaving a space through which the cord may descend. So frequently is this complication met with in pelvic deformity that Stansesco¹ found it had happened as often as 59 times in 414 labors; and when the dangers of prolapsed funis are added to those of protracted labors, it is hardly a matter of surprise that the occurrence should, under such circumstances, almost always prove fatal to the child.

Injury to Child's Head.—The head of the child is also liable to injury of a more or less grave character from the compression to which it is subjected, especially by the promontory of the sacrum. Independently of the transient effects of undue pressure (temporary alteration of the shape of the bones and bruising of the scalp), there is often met with a more serious depression of the bones of the skull produced by the sacral promontory. This is most marked in cases in which the head has been forcibly dragged past the projecting bone by the forceps or after turning.

¹ *Op. cit.*, p. 94.

The amount of depression varies with the degree of contraction; but sometimes, were it not for the yielding of the bones of the foetal skull in this way, delivery, without lessening the size of the head by perforation, would be impossible. Such depressions are found at the spot immediately opposite the promontory, generally at the side of the skull near the junction of the frontal and parietal bones. Sometimes there is a slight permanent mark, but more often the depression disappears in a few days. The prognosis to the child is, however, grave when the contraction has been sufficient to indent the skull, for it has been found that 50 per cent. of the children thus marked died either immediately or shortly after labor.¹

Course of Labor.—The means which nature takes to overcome these difficulties are well worthy of study, and there are certain peculiarities in the mechanism of delivery, when pelvic deformities exist, which it is of importance to understand, as they guide us in determining the proper treatment to adopt.

Frequency of Malpresentation.—Malpresentations of the foetus are of much more frequent occurrence than in ordinary labors—partly because the head does not engage readily in the brim, but, remaining free above it, is apt to be pushed away by the uterine contractions; and partly because of the frequent alteration of the axis of the uterine tumor. The pendulous condition of the abdomen in cases of pelvic deformity is often very obvious, so that the fundus is sometimes almost in a line with the cervix, and thus transverse or other abnormal positions are very frequently met with. It is to be noted, however, that we cannot regard breech presentations as so unfavorable as in ordinary labors, for the pressure from the contracted pelvis is less likely to be injurious when applied to the body than to the head of the child; and, indeed, as we shall presently see, the artificial production of these presentations is often advisable as a matter of choice.

Mechanism of Delivery in Head Presentations.—The mode in which the head passes naturally through a contracted pelvis is in some respects different from the ordinary mechanism of delivery in head presentations, and has been carefully worked out by Spiegelberg and other German obstetricians.

The means which nature adopts to overcome the difficulty are different in cases in which there is a marked narrowing of the conjugate diameter of the brim, and in those in which there is a generally-contracted pelvis.

In Contracted Brim.—In the former and more common deformity the head lies at the brim with its long occipito-frontal diameter in the transverse diameter of the pelvis, and, as both parietal bones cannot enter the contracted brim, it lies with one parietal bone on a much lower level than the other, in the large majority of cases that nearest the pubes being most depressed, so that the sagittal suture is felt high up near the promontory of the sacrum. As labor advances, if the contraction is not too great to be insuperable, the anterior fontanelle comes much more within reach than in ordinary labor, while at the same time the occipital portion of the head is shoved to the side of the pelvis, so that its narrow

¹ Schroeder, *op. cit.*, p. 256.

bi-temporal diameter engages in the contracted conjugate. At this stage, on examination, it will be found—supposing we have to do with a case in which the occiput points to the left side of the pelvis—that the anterior fontanelle is lower than the posterior, and to the right, the bi-temporal diameter of the head is engaged in the conjugate diameter of the brim (as the smallest diameter of the skull, there is manifest advantage in this), and that the bi-parietal diameter and the largest portion of the head point to the left side. The sagittal suture will be felt running across in the transverse diameter of the brim, but nearer to the sacrum, the head being placed obliquely. As the head is forced down by the uterine contractions, the parietal bone, which is resting on the promontory, is pushed against it, so that the sagittal suture is forced more into the true transverse diameter of the pelvic brim and approaches nearer to the pubes. The next step is the depression of the head, the occiput undergoing a sort of rotation on its transverse axis, so that it reaches a plane below the brim. When this is accomplished, the rest of the head readily passes the obstruction. The forehead now meets with the resistance of the pelvic walls, the posterior fontanelle descends to a lower level, and, as the cavity of the pelvis in cases of antero-posterior contraction of the brim is generally of normal dimensions, the rest of the labor is terminated in the usual way.

In Generally-contracted Pelvis.—In the generally-contracted pelvis the head enters the brim with the posterior fontanelle lowest, and it is after it has engaged in it that the resistance to its progress becomes manifest. The result is, therefore, an exaggeration of what is met with in ordinary cases. The resistance to the anterior or longer arm of the lever is greater than that to the occipital or shorter, and therefore the flexion of the head becomes very marked. The posterior fontanelle is consequently unusually depressed, and the anterior quite out of reach. So the head is forced down as a wedge, and its further progress must depend upon the amount of contraction. If this be not too great, the anterior fontanelle eventually descends, and delivery is completed in the usual way. Should the contraction be too much to permit of this, the head becomes jammed in the pelvis and diminution of its size may be essential.

In cases of deformity of the conjugate diameter, combined with general contraction of the pelvis, the mechanism partakes of the peculiarities of both these classes to a greater or less extent, in proportion to the preponderance of one or other species of deformity.

Diagnosis.—It rarely happens that deformities of the pelvis, except of the gravest kind, are suspected before labor has actually commenced, and therefore we are not often called upon to give an opinion as to the condition of the pelvis before delivery. Should we be, there are various circumstances which may aid us in arriving at a correct conclusion. Prominent among them is the history of the patient in childhood. If she is known to have suffered from rickets in early life, more especially if the disease has left evident traces in deformities of the limbs or in a dwarfed and stunted growth or in curvature of the spine, there will be strong presumptive evidence of pelvic deformity; a markedly pendulous state of the abdomen may also tend to confirm the suspicion. Nothing

short of a careful examination of the pelvis itself will, however, clear up the point with certainty; and even by this means to estimate the precise degree of deformity with accuracy requires considerable skill and practice. The ingenuity of practitioners has been much exercised—it might perhaps be justly said wasted—in the invention of various more or less complicated pelvimeters for aiding us in obtaining the desired object. It is, however, pretty generally admitted by all accoucheurs that the hand forms the best and most reliable instrument for this purpose—at any rate, as regards the interior of the pelvis—although a pair of callipers, such as Baudelocque's well-known instrument, is essential for accurately determining the external measurements. The objections to all internal pelvimeters, even those most simple in their construction, are their cost and complexity and the impossibility of using them without pain or injury to the patient.

External Measurements.—It was formerly thought that by measuring the distance between the spinous processes of the sacrum and the symphysis pubis, and subtracting from it what we judge to be the thickness of the bones and soft parts, we might arrive at an approximate estimate of the measurement of the conjugate diameter of the pelvic brim. It is now admitted that this method can never be depended on, and that, taken by itself, it is practically useless. A change in the relative length of other external measurements of the pelvis is, however, often of great value in showing the existence of deformity internally, although not in judging of its amount. The measurements which are used for this purpose are between the anterior-superior spines of the ilia and between the centres of their crests, averaging respectively 10 and 11 inches. According to Spiegelberg, these measurements may give one of three results:

1. Both may be less than they ought to be, but the relation of one to the other remains unchanged.
2. That between the crests is not, or is at most very little, diminished, but that between the spines is increased.
3. Both are diminished, but at the same time their mutual relation is not normal, the distance between the spines being as long, if not longer, than that between the crests.

No. 1 denotes a uniformly-contracted pelvis; No. 2, a pelvis simply contracted in the conjugate diameter of the brim, and not otherwise deformed; No. 3, a pelvis with narrowed conjugate and also uniformly contracted, as in the severe type of rachitic deformity. If, however, both these measurements are of average length, and the distance between the crests is about 1 inch greater than between the spines, the pelvis is normal.

Besides the above, some information may be obtained by the measurement of the external conjugate diameter, which averages $7\frac{3}{4}$ inches. This may be taken by placing one point of the callipers in the depression below the spine of the last lumbar vertebra, the other at the centre of the upper edge of the symphysis pubis. If the measurement be distinctly below the average—not more, for example, than 6.3 inches—we may conclude that there is a narrowing of the antero-posterior diameter of the brim, the extent of which we must endeavor to ascertain by other means.

For the purpose of making these measurements, Baudelocque's *compas d'épaisseur* can be used, or Dr. Lazarewitch's elegant universal pelvimeter, which can be adopted also for internal pelvimetry; but in the absence of these special contrivances an ordinary pair of callipers, such as are used by carpenters, can be made to answer the desired object.

Internal Measurements.—These external measurements must be corroborated by internal, chiefly of the antero-posterior diameter, by which alone we can estimate the amount of the deformity. We endeavor to find, in the first place, the length of the inclined conjugate between the lower edge of the symphysis pubis and the promontory of the sacrum, which averages about half an inch more than the true conjugate. This is best done by placing the patient on her back with the hips well raised. The index finger of the right hand is then introduced into the vagina, and the perineum is pressed steadily backward, so as to overcome the resistance it offers. If the tip of the finger can reach the promontory of the sacrum, its radial side is raised so as to touch the lower edge of the pubes. A mark is made with the nail of the index of the left hand on that part of the examining finger which rests under the symphysis, and then the distance from this to the tip of the finger, less half an inch, may be taken to indicate the measurement of the true conjugate of the brim. Various pelvimeters have been devised to make the same measurements, such as Lumley Earle's, Lazarewitch's (which is similar in principle), and Van Huevel's; the best and simplest, I think, is that invented by Dr. Greenhalgh (Fig. 141). It consists of a movable rod attached to a flexible band of metal which passes around the palm of the examining hand. At the distal end of the rod is a curved portion which passes over the radial edge of the index finger. The examination is made in the usual way, and when the point of the finger is resting on the promontory of the sacrum the rod is withdrawn until it is arrested by the posterior surface of the symphysis, the exact measurement of the inclined conjugate being then read off the scale.

It is to be remembered that this procedure is useless in the slighter degrees of contraction in which the promontory of the sacrum cannot be easily reached. Dr. Ramsbotham proposed to measure the conjugate by spreading out the index and middle fingers internally, the tip of one resting on the promontory, the other behind the symphysis pubis, and then withdrawing them in the same position and measuring the distance between them. This manoeuvre I believe to be impracticable.

FIG. 141.



Greenhalgh's Pelvimeter.

Whenever, in actual labor, we wish to ascertain the condition of the pelvis accurately, the patient should be anesthetized, and the whole hand introduced into the vagina (which could not otherwise be done without causing great pain), and the proportions of the pelvis and the relations of the head to it thoroughly explored; and if what has been said as to the mechanism of delivery in these cases be borne in mind, this may aid us in determining the kind of deformity existing. In this way contractions about the outlet of the pelvis can also be pretty generally made out.

Mode of Diagnosing the Oblique Pelvis.—The obliquely-contracted pelvis cannot be determined by any of these methods, but certain external measurements, as Naegele has pointed out, will readily enable us to recognize its existence. It will be found that measurements which in the healthy pelvis ought to be equal are unequal in the obliquely-distorted pelvis. The points of measurement are chiefly: (1) From the tuberosity of the ischium on one side to the posterior-superior spine of the ilium on the other; (2) from the anterior-superior iliac spine on the one side to the posterior-superior on the opposite; (3) from the trochanter major of one side to the posterior-superior iliac spine on the other; (4) from the lower edge of the symphysis pubis to the posterior-superior iliac spine on either side; (5) from the spinous process of the last lumbar vertebra to the anterior-superior spine of the ilium on either side.

If these measurements differ from each other by half an inch to an inch, the existence of an obliquely-deformed pelvis may be safely diagnosed. The diagnosis can be corroborated by placing the patient in the erect position and letting fall two plumb-lines—one from the spines of the sacrum, the other from the symphysis pubis. In a healthy pelvis these will fall in the same plane, but in the oblique pelvis the anterior line will deviate considerably toward the unaffected side.

Treatment.—The proper management of labor in contracted pelvis is, even up to this time, one of the most vexed questions in midwifery, notwithstanding the immense amount of discussion to which it has given rise; and the varying opinions of accoucheurs of equal experience afford a strong proof of the difficulties surrounding the subject. This remark applies, of course, only to the lesser degree of deformity, in which the birth of a living child is not hopeless. When the antero-posterior diameter of the brim measures from $2\frac{3}{4}$ to 3 inches, it is universally admitted that the destruction of the child is inevitable, unless the pelvis be so small as to necessitate the performance of the Cæsarean section. But when it is between 3 inches and the normal measurement, the comparative merits of the forceps, turning, and the induction of premature labor form a fruitful theme for discussion. With one class of accoucheurs the forceps is chiefly advocated, and turning admitted as an occasional resource when it has failed; and this indeed, speaking broadly, may be said to have been the general view held in this country. More recently we find German authorities of eminence, such as Schroeder and Spiegelberg, giving turning the chief place, and condemning the forceps altogether in contracted pelvis, or at least restricting its use within very narrow limits. More strangely still, we find of late that the induction of premature labor, on the origination and extension of which British accoucheurs have always prided themselves, is placed without the pale,

and spoken of as injurious and useless in reference to pelvic deformities. To see our way clearly amongst so many conflicting opinions is by no means an easy task, and perhaps we may best aid in its accomplishment by considering separately the three operations in so far as they bear on this subject, and pointing out briefly what can be said for and against each of them.

The Forceps.—In England and in France it is pretty generally admitted that in the slighter degrees of contraction the most reliable means of aiding the patient is by the forceps. It should be remembered that the operation under such circumstances is always much more serious than in ordinary labors simply delayed from uterine inertia, when there is ample room and the head is in the cavity of the pelvis; for the blades have to be passed up very high, often when the head is more or less movable above the brim, and much more traction is likely to be required. For these reasons artificial assistance, when pelvic deformity is suspected, is not to be lightly or hurriedly resorted to. Nor, fortunately, is it always necessary, for if the pains be sufficiently strong, and the contraction not too great to prevent the head engaging at all, after a lapse of time it will become so moulded in the brim as to pass even a considerable obstruction. In all cases, therefore, sufficient time must be given for this; and if no suspicious symptoms exist on the part of the mother—no elevation of temperature, dryness of the vagina, rapid pulse, and the like, and the fetal heart-sounds continue to be normal—labor may be allowed to go on for some hours after the rupture of the membranes, so as to give nature a chance of completing the delivery. When this seems hopeless the intervention of art is called for.

Cases Suitable for the Forceps.—The forceps is generally considered to be applicable in all degrees of contraction, from the standard measurement down to about $3\frac{1}{4}$ inches in the conjugate of the brim. There can be no doubt that in such cases traction with the forceps often enables us to effect delivery when the natural efforts have proved insufficient, and holds out a very fair hope of saving the child. Out of 17 cases in which the high forceps operation was resorted to for pelvic deformity, reported by Stanesco, in 13 living children were born. If the length of the labor and the long-continued compression to which the child has been subjected be borne in mind, this result must be considered very favorable.

Objections that have been Raised to the Forceps.—What are the objections which have been brought against the operation? These have been principally made by Schroeder and other German writers. They are, chiefly, the difficulty of passing the instrument, the risk of injuring the maternal structures, and the supposition that, as the blades must seize the head by the forehead and occiput, their compressive action will diminish its longitudinal and increase its transverse diameter (which is opposed to the contracted part of the brim), and so enlarge the head just where it ought to be smallest. There is little doubt that these writers much exaggerate the compressive power of the forceps. Certainly, with those generally used in this country, any disadvantage likely to accrue from this is more than counterbalanced by the traction on the head; and the fact that minor degrees of obstruction can be thus overcome with safety

both to the mother and child is abundantly proved by the numberless cases in which the forceps has been used.

Not equally Suitable in all Kinds of Deformity.—It is very likely that the forceps does not act equally well in all cases. When the head is loose above the brim; when the contraction is chiefly limited to the antero-posterior diameter, and there is abundance of room at the sides of the pelvis for the occiput to occupy after version; and when, as is usual in these cases, the anterior fontanelle is depressed and the head lies transversely across the brim,—it is probable that turning may be the safer operation for the mother, and the easier performed. When, on the other hand, the head has engaged in the brim and has become more or less impacted, it is obvious that version could not be performed without pushing it back, which may be neither easy nor safe. In the generally-contracted pelvis, in which the head enters in an exaggerated state of flexion and lies obliquely, the posterior fontanelle being much depressed, the forceps is more suitable.

Mechanical Advantage of Turning in Certain Cases.—The special reasons why version sometimes succeeds when the forceps fails, or why it may be elected from the first as a matter of choice, have been by no one better pointed out than by Sir James Simpson. Although the operation was performed by many of the older obstetricians, its revival in modern times and the clear enunciation of its principles can undoubtedly be traced to his writings. He points out that the head of the child is shaped like a cone, its narrowest portion the base of the cranium (Fig. 142, *b b*), measuring, on an average, from $\frac{1}{2}$ to $\frac{2}{3}$ of an inch less than the broadest portion (Fig. 142, *a a*)—viz. the bi-parietal diameter. In




Fig. 142.
Section of Fœtal Cranium, showing its Conical Form.




FIG. 143.
Showing the Greater Breadth of the Bi-parietal Diameter of the Fœtal Cranium. (After Simpson.)

ordinary head presentations the latter part of the head has to pass first; but if the feet are brought down, the narrow apex of the cranial cone is brought first into apposition with the contracted brim, and can be more easily *drawn* through than the broader base can be *pushed* through by the uterine contractions. Nor is this the only advantage, for after turning the narrower bi-temporal diameter (Fig. 143, *b b*)—which measures, on an average, half an inch less than the bi-parietal (Fig. 143, *a a*)—is brought into contact with the contracted conjugate, while the broad parietal lies in the comparatively wide space at the side.

(Fig. 144). These mechanical considerations are sufficiently obvious, and fully explain the success which has often attended the performance of the operation.

Limits of the Operation.—It is generally admitted that it may be possible, for the reasons just mentioned, to deliver a living child by turning

FIG. 144.



Showing the Greater Space for the Bi-parietal Diameter at the Side of the Pelvis in Certain Cases of Deformity. (After Simpson.)

through a pelvis contracted beyond the point which would permit of a living child being extracted by the forceps. Many obstetricians believe that it is possible to deliver a living child by turning in a pelvis contracted even to the extent of $2\frac{3}{4}$ inches in the conjugate diameter. Barnes maintains that, although an unusually compressible head may be drawn through a pelvis contracted to 3 inches, the chance of the child being born alive under such circumstances must necessarily be small, and that from $3\frac{1}{4}$ inches to the normal size must be taken as the proper limits of the operation.

It frequently Succeeds when the Forceps has Failed.—That delivery is often possible by turning after the forceps and the natural powers have failed, and when no other resource is left but the destruction of the child, must, I think, be admitted by all, for the records of obstetrics are full of such cases. To take one example only: Dr. Braxton Hicks¹ records four cases in which the forceps were tried unsuccessfully, in all of which version was used, three of the children being born alive. Here are the lives of three children rescued from destruction within a short period in the practice of one man; and a fact like this would of itself be ample justification of the attempt to deliver by turning when the child was known to be alive and other means had failed. The possibility that craniotomy may still be required is no argument against the operation; for although perforation of the after-coming head is certainly not so easy as perforation of a presenting head, it is not so much more difficult as to justify the neglect of an experiment by which it may possibly be altogether avoided.

Comparative Estimate of the Two Operations.—The original choice of turning is a more difficult question to decide. The most generally received opinion in the present day amongst scientific obstetricians is that in the simply flattened pelvis, with an antero-posterior diameter of not less than $2\frac{3}{4}$ inches, turning is the preferable operation. In every

¹ *Guy's Hospital Reports*, 1870.

case of doubt it is desirable thoroughly to anæsthetize the patient and make a careful examination with the whole hand in the vagina. If we find the sagittal suture lying transversely, one parietal bone on a lower line than the other, and if both fontanelles are easily within reach, and some space exists at the sides of the pelvis beside the forehead and occiput, then turning is the procedure most likely to succeed, and the descent of the head after version can be very materially assisted by strong pressure applied from above by an assistant, as has been well pointed out by Goodell.¹ If, on the other hand, the anterior fontanelle is high up and out of reach, the head being distinctly flexed, we have to do with a generally-contracted pelvis, and the forceps is the preferable operation.

Cases in which Craniotomy or the Cæsarean Section is Required.—When the contraction is below 3 inches in the conjugate, or when the forceps and turning have failed, no resource is left but the destruction of the fœtus or the Cæsarean section.

The induction of premature labor as a means of avoiding the risks of delivery at term, and of possibly saving the life of the child, must now be studied. The established rule in this country is, that in all cases of pelvic deformity, the existence of which has been ascertained either by the experience of former labors or by accurate examination of the pelvis, labor should be induced previous to the full period, so that the smaller and more compressible head of the premature fœtus may pass where that of the fœtus at term could not. The gain is a double one—partly the lessened risk to the mother, and partly the chance of saving the child's life.

Recent Objections to it.—The practice is so thoroughly recognized as a conservative and judicious one that it might be hardly necessary to argue in its favor, were it not that some eminent authorities have of late years tried to show that it is better and safer to the mother to leave the labor to come on at term, and that the risk to the child is so great in artificially-induced labor as to lead to the conclusion that the operation should be altogether abandoned, except, perhaps, in the extreme distortion in which the Cæsarean section might otherwise be necessary. Prominent amongst those who hold these views are Spiegelberg and Litzmann; and they have been supported, in a modified form, by Matthews Duncan. Spiegelberg² tries to show, by a collection of cases from various sources, that the results of induced labor in contracted pelvis are much more unfavorable than when the cases are left to nature—that in the latter the mortality of the mothers is 6.6 per cent., and of the children 28.7 per cent., whereas in the former the maternal deaths are 15 per cent., and the infantile 66.9 per cent. Litzmann³ arrives at not very dissimilar results—namely, 6.9 per cent. of the mothers and 20.3 per cent. of the children in contracted pelvis at term, and 14.7 per cent. of the mothers and 55.8 per cent. of the children in artificially-induced premature labor.

If these statistics were reliable, inasmuch as they show a very decided risk to the mother, there might be great force in the argument that it would be better to leave the cases to run the chance of delivery at term. It is, however, very questionable whether they can be taken, in them-

¹ *Amer. Journ. of Obstet.*, vol. viii.

² *Ib.*, B. ii. S. 169.

³ *Arch. f. Gyn.*, B. i. S. 1.

selves, as being sufficient to settle the question. The fallacy of determining such points by a mass of heterogeneous cases, collected together without a careful sifting of their histories, has over and over again been pointed out; and it would be easy enough to meet them by an equal catalogue of cases in which the maternal mortality is almost nil. The results of the practice of many authorities are given in Churchill's work, where we find, for example, that out of 46 cases of Merriman's not one proved fatal. The same fortunate result happened in 62 cases of Ramsbotham's. His conclusion is that "there is undoubtedly some risk incurred by the mother, but not more than by accidental premature labor;" and this conclusion as regards the mother is that which has long ago been arrived at by the majority of British obstetricians, who undoubtedly have more experience of the operation than those of any other nation. With regard to the child, even if the German statistics be taken as reliable, they would hardly be accepted as contraindicating the operation, inasmuch as it is intended to save the mother from the dangers of the more serious labor at term, and in many cases to give at least a chance to the child, whose life would otherwise be certainly sacrificed. The result, moreover, must depend to a great extent on the method of operation adopted, for many of the plans of inducing labor recommended are certainly, in themselves, not devoid of danger both to the mother and the child. It may, I think, be admitted, as Duncan contends,¹ that the operation has been more often performed than is absolutely necessary, and that the higher degrees of pelvic contraction are much more uncommon than has been supposed to be the case. That is a very valid reason for insisting on a careful and accurate diagnosis, but not for rejecting an operation which has so long been an established and favorite resource.

Determination of Period for Inducing Labor.—When the induction of labor has been determined on, the precise period at which it should be resorted to becomes a question for anxious consideration, since the longer it is delayed the greater, of course, are the dangers for the child. Many tables have been constructed to guide us on this point which are not, on the whole, of so much service as they might appear to be, on account of the difficulty of determining with minute accuracy the amount of contraction. The following, however, which is drawn up by Kiwisch, may serve for a guide in settling this question :

	Inches.	Lines.	
When the sacro-pubic diameter is 2 and 6 or 7,			induce labor at 30th week.
" " 2 " 8 " 9,			" 31st "
" " 2 " 10 " 11,			" 32d "
" " 3 " . . .			" 33d "
" " 3 " 1,			" 33d "
" " 3 " 2 " 3,			" 34th "
" " 3 " 4 " 5,			" 35th "
" " 3 " 5 " 6,			" 36th "

In cases of moderate deformity, when labor-pains have been induced, the further progress of the case may be left to nature; but in more marked cases, as in those below 3 inches, it will often be found necessary to assist delivery by turning or by the forceps, the former being

¹ *Edin. Med. Journ.*, July, 1873, p. 339.

here specially useful, on account of the extreme pliability of the head and the facility with which it may be drawn through the contracted brim. By thus combining the two operations it may be quite possible to secure the birth of a living child even in pelvis very considerably deformed.

Production of Abortion in Extreme Deformity.—When the contraction is so great as to necessitate the induction of the labor before the sixth month—or, in other words, before the child has reached a viable age—it would be preferable to resort to a very early production of abortion. The operation is then indicated, not for the sake of the child, but to save the mother from the deadly risk to which she would otherwise be subjected. As in these cases the mother alone is concerned, the operation should be performed as soon as we have positively determined the existence of pregnancy. No object can be gained by waiting until the development of the child is advanced to any extent, and the less the fetus is developed the less will be the pain and risks the mother has to undergo. There is no amount of deformity, however great, in which we could not succeed in bringing on miscarriage by some of the numerous means at our disposal; and, in spite of Dr. Radford's objections, who maintains that the obstetrician is not justified in sacrificing the life of a human being more than once when the mother knows that she cannot give birth to a viable child, there are few practitioners who would not deem it their duty to spare the mother the terrible dangers of the Cæsarean section.

CHAPTER XIII.

HEMORRHAGE BEFORE DELIVERY; PLACENTA PRÆVIA.

THE hemorrhages which are the result of an abnormal situation of the placenta, partially or entirely, over the internal os uteri have formed a most fruitful theme for discussion. The explanation of the abnormal placental site, the sources of the blood, and the causes of its escape, the means adopted by nature for its arrest, and the proper treatment, have, each and all of them, been the subject of endless controversies, which are not yet by any means settled. It must be admitted, too, that the extreme importance of the subject amply justifies the attention which has been paid to it, for there is no obstetric complication more apt to produce sudden and alarming effects, and none requiring more prompt and scientific treatment.

Definition.—By *placenta prævia* we mean the insertion of the placenta at the lower segment of the uterine cavity, so that a portion of it is situated, wholly or partially, over the internal os uteri. In the former case there is *complete* or *central* placental presentation, in the latter an *incomplete* or *marginal* presentation.

Causes.—The causes of this abnormal placental site are not fully

understood. It was supposed by Tyler Smith to depend on the ovule not having been impregnated until it had reached the lower part of the uterine cavity. Cazeaux suggests that the uterine mucous membrane is less swollen and turgid than when impregnation occurs at the more ordinary place, and that, therefore, it offers less obstruction to the descent of the ovule to the lower part of the uterine cavity. An abnormal size or unusual shape of the uterine cavity may also favor the descent of the impregnated ovule; the former probably explains the fact that placenta prævia more generally occurs in women who have already borne children. Müller believes that it results from uterine contractions occurring shortly after conception, which force the ovum down to the lower part of the uterine cavity. These are merely interesting speculations, having no practical value, the fact being undoubted that in a not inconsiderable number of cases—estimated by Johnson and Sinclair as 1 out of 573—the placenta is grafted partially or entirely over the uterine orifice.

History.—Placenta prævia was not unknown to the older writers, who believed that the placenta had originally been situated at the fundus, from which it had accidentally fallen to the lower part of the uterus. Portal, Levret, Roederer, and especially our own countryman Rigby, were among those whose observations tended to improve the state of obstetrical knowledge as to its real nature. To Rigby we owe the term "*unavoidable hemorrhage*" as a synonym for placenta prævia, and as distinguishing hemorrhage from this source from that resulting from separation of the placenta at its more usual position, termed by him, in contradistinction, "*accidental hemorrhage*." These names, adopted by most writers on the subject, are obviously misleading, as they assume an essential distinction in the etiology of the hemorrhage in the two classes of cases which is not always warranted.

It is of the utmost importance to a right understanding of the nature and treatment of placenta prævia that we should fully understand the source of the hemorrhage and the manner of its production; but we shall be able to discuss this subject better after a description of the symptoms.

Symptoms.—Although the placenta must occupy its unusual site from the earliest period of its formation, it rarely gives rise to appreciable symptoms before the last three months of utero-gestation. It is far from unlikely, however, that such an abnormal situation of the placenta may produce abortion in the earlier months, the site of its attachment passing unobserved.

Sudden Flow of Blood.—The earliest symptom which causes suspicion is the sudden occurrence of hemorrhage without any appreciable cause. The amount of blood lost varies considerably. In some cases the first hemorrhage is comparatively slight, and is soon spontaneously arrested; but if the case be left to itself, the flow after a lapse of time—it may be a few days or it may be weeks—again commences in the same unexpected way, and each successive hemorrhage is more profuse. The losses show themselves at different periods. They rarely begin before the end of the sixth month, more often nearer the full period, and sometimes not until labor has actually commenced. The hemorrhage very

often coincides with what would have been a menstrual period, doubtless on account of the physiological congestion of the uterine organs then present. Should the first loss not show itself until at or near the full time, it may be tremendous, and a few moments may suffice to place the patient's life in jeopardy. Indeed, it may be safely accepted as an axiom that once hemorrhage has occurred the patient is never safe, for excessive losses may occur at any moment without warning and when assistance is not at hand. It often happens that premature labor comes on after one or more hemorrhages.

In any case of placenta prævia when labor has commenced, whether premature or at the full time, the hemorrhage may become excessive, and with each pain fresh portions of placenta may be detached and fresh vessels torn and left open. Under these circumstances the blood often escapes in greater quantity with each successive pain, and diminishes in the interval. This has long been looked upon as a diagnostic mark by which we can distinguish between the so-called "unavoidable" and "accidental" hemorrhage; in the latter the flow being arrested during the pains. The distinction, however, is altogether fallacious. The tendency of uterine contraction in placenta prævia, as in all other forms of uterine hemorrhage, is to constrict the vessels from which the blood escapes, and so to lessen the flow. The apparently increased flow during the pains depends on the pains forcing out blood which has already escaped from the vessels. In one way, up to a certain point, the pains do favor hemorrhage by detaching fresh portions of placenta; but the actual loss takes place chiefly during the intervals, and not during the continuance of contraction.

Results of Vaginal Examination.—On vaginal examination, if the os be sufficiently open to admit the finger—which it generally is, on account of the relaxation produced by the loss of blood—we shall almost always be able to feel some portion of presenting placenta. If it be a central implantation, we shall find the upper aperture of the cervix entirely covered by a thick, boggy mass, which is to be distinguished from a coagulum by its consistence and by its not breaking down under the pressure of the finger. Through the placental mass we may feel the presenting part of the fetus, but not as distinctly as when there is no intervening substance. In partial placental presentations the bag of membranes, and above it the head or other presentation, will be found to occupy a part of the circle of the os, the rest being covered by the edge of the placenta. In marginal presentations we may only be able to make out the thickened edge of the after-birth projecting at the rim of the os. If the cervix be high and the gestation not advanced to term, these points may not be easy to make out on account of the difficulty of reaching the cervix; and, as accurate diagnosis is of the utmost importance, it is proper to introduce two fingers, or even the whole hand, so as thoroughly to explore the condition of the parts. The lower portion of the uterine ovoid may be observed to be more than usually thick and fleshy; and Gendrin has pointed out that ballottement cannot be made out. The accuracy of our diagnosis may be confirmed, in doubtful cases, by finding that the placental bruit is heard over the lower part of the uterine tumor.

Dr. Wallace¹ has suggested that vaginal auscultation may be serviceable in diagnosis, and states that by means of a curved wooden stethoscope the placental bruit may be heard with startling distinctness. This is, however, a manœuvre that can hardly be generally carried out in actual practice.

The Source of Hemorrhage.—It is now generally admitted by authorities that the immediate source of the hemorrhage is the lacerated utero-placental vessels. Only a few years ago Sir James Simpson advocated with his usual energy the theory, sustained by his predecessor, Dr. Hamilton, that the chief if not the only source of hemorrhage was the detached portion of the placenta itself. He argued that the blood flowed from the portion of the placenta which was still adherent into that which was separated, and escaped from the surface of the latter; and on this supposition he based his practice of entirely separating the placenta, having observed that in many cases in which the after-birth had been expelled before the child the hemorrhage had ceased. The fact of the cessation of the hemorrhage, when this occurs, is not doubted, but Simpson's explanation is contested by most modern writers, prominent among whom is Barnes, who has devoted much study to the elucidation of the subject. He points out that the stoppage of the hemorrhage is not due to the separation of the placenta, but to the preceding or accompanying contraction of the uterus, which seals up the bleeding vessels, just as it does in other forms of hemorrhage. The site of the loss was actually demonstrated by the late Dr. Mackenzie in a series of experiments in which he partially detached the placenta in pregnant bitches, and found that the blood flowed from the walls of the uterus, and not from the detached surface of the placenta. The arrangement of the large venous sinuses, opening as they do on the uterine mucous membrane, favors the escape of blood when they are torn across; and it is from them, possibly to some extent also from the uterine arteries, that the blood comes, just as in post-partum hemorrhage, when the whole instead of a part of the placental site is bared.

Causes of Hemorrhage.—Various explanations have been given of the causes of the hemorrhage. It was long supposed to depend on the gradual expansion of the cervix during the latter months of pregnancy, which separated the abnormally placed placenta. It has been seen, however, that this shortening of the cervix is apparent only, and that the cervical canal is not taken up into the uterine cavity during gestation, or, at all events, only during the last week or so. This, therefore, cannot be admitted as an explanation of placental separation. Jacquemier proposed another theory, which has been adopted by Cazeaux. He maintains that during the first six months of utero-gestation the superior portion of the uterus is more especially developed, as shown by the pyriform shape of the fundus during the time, and that, as the placenta is usually attached in that situation, and then attains its maximum of development, its relations to its attachments are undisturbed. During the last three months of pregnancy, on the contrary, the lower segment of the uterus develops more than the upper, while the placenta remains nearly stationary in size, the inevitable result being a loss of

¹ *Edin. Med. Journ.*, Nov., 1872.

proportion between the cervix and the placenta, and the detachment of the latter. There are various objections which can be brought against this theory, the most important being that there is no evidence at all to show that the lower segment of the uterus does expand more in proportion than the upper during the latter months of pregnancy. Barnes's theory is based on the supposition that the loss of relation between the uterus and placenta is caused by excess of growth on the part of the placenta itself over that of the cervix, which is not adapted for its attachment. The placenta, on this hypothesis, grows away from the site of its attachment, and hemorrhage results. It will be observed that neither this theory nor that propounded by Jacquemier is readily reconcilable with the fact that hemorrhage frequently does not begin until labor has commenced at term. Inasmuch as the loss of relation between the placenta and its attachments, which they both presuppose, must exist in every case of placenta prævia, hemorrhage should always occur during some part of the last three months of pregnancy. Matthews Duncan¹ has recently investigated the whole subject at length, and maintains that the hemorrhages are accidental, not unavoidable, being due to causes precisely similar to those which give rise to the occasional hemorrhages when the placenta is normally placed. The abnormal situation of the placenta of course renders these causes more apt to operate; but in their action he believes them to be precisely similar to those of accidental hemorrhage, properly so called. Separation of the placenta from expansion of the cervix he believes to be the cause of hemorrhage after labor has begun, and then it may strictly be called unavoidable; but hemorrhage is comparatively seldom so produced during the continuance of pregnancy. "There are," says Duncan, "four ways in which this kind of hemorrhage may occur:

"1. By the rupture of a utero-placental vessel at or about the internal os uteri.

"2. By the rupture of a marginal utero-placental sinus within the area of spontaneous premature detachment when the placenta is inserted not centrally or covering the internal os, but with a margin at or near the internal os.

"3. By partial separation of the placenta from accidental causes, such as a jerk or fall.

"4. By a partial separation of the placenta the consequence of uterine pains, producing a small amount of dilatation of the internal os. Such cases may be otherwise described as instances of miscarriage commencing, but arrested at a very early stage."

I see no reason to doubt the possibility of hemorrhage being due in many cases to the first three causes, and in its production it would strictly resemble accidental hemorrhage. The fourth heading refers the hemorrhage to partial separation in consequence of commencing dilatation of the cervix, but it explains the dilatation by the supposition of commencing miscarriage. This latter hypothesis seems to be as needless as those which presuppose a want of relation between the placenta and its attachments. We know that, quite independently of commencing miscarriage, uterine contractions are constantly occurring during the

¹ *Edin. Med. Journ.*, Nov

² *Brit. Med. Journ.*, Nov., 1873.

continuance of pregnancy. There is no reason to suppose that these contractions do not affect the cervical as well as the fundal portions of the uterus; and in cases in which the placenta is situated partially or entirely over the os, one or more stronger contractions than usual may at any moment produce laceration of the placental attachments in that neighborhood.

Pathological Changes in the Placenta.—A careful examination of the placenta may show pathological changes at the site of separation, such as have been described by Gendrin, Simpson, and other writers. They probably consist of thromboses in the placental cotyledons and effused blood-clots, variously altered and decolorized according to the lapse of time since separation took place. Changes occur in the portion of the placenta overlying the os uteri whether separation has occurred or not. There may be atrophy of the placental structure in this situation, as well as changes of form, such as complete or partial separation into two lobes, the junction of which overlies the os uteri.¹

Natural Termination when the Placenta Presents.—The history of delivery, if left to nature, is specially worthy of study, as guiding to proper rules of treatment. It sometimes happens, when the pains are very strong and the delivery rapid, that labor is completed without any hemorrhage of consequence. "Although," says Cazeaux, "hemorrhage is usually considered to be inevitable under such circumstances, yet it may not appear even during the labor, and the dilatation of the os uteri may be effected without the loss of a drop of blood." Again, Simpson conclusively showed that when the placenta was expelled before the birth of the child all hemorrhage ceased.

Barnes's theory of placenta prævia, which has been pretty generally adopted, explains satisfactorily both these classes of cases. He describes the uterine cavity as divisible into three zones or regions. When the placenta is situated in the upper or middle of these zones, no separation or hemorrhage need occur during labor. When, however, it is situated partially or entirely in the lower or cervical zone, the expansion of the cervix during labor must produce more or less separation, and consequent loss of blood. As soon as the previous portion of the placenta is sufficiently separated, provided contraction of the uterine tissue be present to seal up the mouths of the vessels, hemorrhage no longer takes place. The placenta may not be entirely detached, but no further hemorrhage occurs, in consequence of the remaining portion being engrafted on the uterus beyond the region of unsafe attachment. In the former, then, of these classes of cases the absence of hemorrhage is explained on this theory by the pains being sufficiently rapid and strong to complete the separation of the placental attachment from the lower cervical zone before flooding had taken place; in the latter, it ceases, not necessarily because the entire placenta is expelled, but because of its detachment from the area of dangerous implantation.

The amount of cervical expansion required for this purpose varies in different cases. Dr. Duncan² estimates the limit of the spontaneous detaching area to be a circle of $4\frac{1}{2}$ inches diameter, and that after the cervix has expanded to that extent no further separation or hemorrhage

¹ Simelius, *Arch. gén. de Méd.*, vol. ii., 1861.

² *Obst. Trans.*, vol. xv.

takes place. To admit of the passage of a full-sized head, Barnes estimates that expansion to about a circle of 6 inches diameter is necessary; on the other hand, he has sometimes observed "that the hemorrhage has completely stopped when the os uteri opened to the size of the rim of a wine-glass, or even less."

It will be seen, then, that in this, as in every other form of puerperal hemorrhage, the tendency of uterine contraction is to check the hemorrhage, and that, provided the pains are sufficiently energetic, Nature may be capable of stopping the flooding without artificial aid. It is but rarely, however, that she can be trusted for the purpose; and we shall presently see that these theoretical views have an important practical bearing on the subject of treatment.

Prognosis.—The prognosis to both the mother and child is certainly grave in all cases of placenta prævia. Read, in his treatise on placenta prævia, estimates the maternal mortality, from the statistics of a large number of cases, as 1 in $4\frac{1}{2}$ cases, and Churchill as 1 in 3. This is unquestionably too high an estimate, and based on statistics the accuracy of which cannot be relied on. The mortality will, of course, greatly depend on the treatment adopted. Doubtless, if cases were left to nature the result would be quite as unfavorable as Read supposes. But if properly managed much more successful results may be safely anticipated. Out of 64 cases recorded by Barnes, the deaths were 6, or 1 in 10 $\frac{2}{3}$. Under any circumstances the risks to the mother are very great. Churchill estimates that more than half the children are lost. The reasons for the great danger to the child are very obvious, subjected as it is to the risk of asphyxia from the loss of the maternal blood, and from its respiration being carried on during labor by a placenta which is only partially attached; many children also perish from prematurity or from malpresentation.

Treatment.—Whenever, in the latter months of pregnancy, a sudden hemorrhage occurs, the possibility of placenta prævia will naturally suggest itself, and by a careful vaginal examination—which under such circumstances should always be insisted on—the existence of this complication will generally be readily ascertained. It is seldom that the os is not sufficiently dilated to enable us to satisfy ourselves whether the placenta is presenting.

Is it Justifiable to Allow the Pregnancy to Continue?—The first question that will arise is, Are we justified in temporizing, using means to check the hemorrhage, and allowing the pregnancy to continue? This is the course which has generally been recommended in works on midwifery. We are told to place the patient on a hard mattress, not to heat or overburden her with clothes, to keep her absolutely at rest, to have the room cooled and well aired, to apply cold cloths to the vulva and lower part of the abdomen, to administer cold and acidulated drinks in abundance, and to prescribe acetate of lead and opium, or gallic acid, on account of their supposed hæmostatic effect. Of late years the judiciousness of these recommendations has been strongly contested. Not long ago an interesting discussion took place at the Obstetrical Society of London¹ on a paper in which Dr. Greenhalgh advised the immediate

¹ *Obst. Trans.*, vol. vi. p. 188.

induction of labor in all cases of placenta prævia. No less than six metropolitan teachers of midwifery took part in it, and, although they differed in details, they all agreed as to the unadvisability of allowing pregnancy to progress when the existence of placenta prævia had been distinctly ascertained. The reasons for this course are obvious and unanswerable. The labor, indeed, very often comes on of its own accord, but should it not do so the patient's life must be considered to be always in jeopardy until the case is terminated, for no one can be sure that most dangerous, or even fatal, flooding may not at any moment come on; and the nearer to term the patient is, the greater the risk to which she is subjected. Nor is the safety of the child likely to be increased by delay. Provided it has arrived at a viable age, the chances of its being born alive may be said to be greater if pregnancy be terminated at once than if repeated floodings occur. I think, therefore, that it may be safely laid down as an axiom that no attempt should be made to prevent the termination of pregnancy, but that our treatment should rather contemplate its conclusion as soon as possible. An exception may, however, be made to this rule when the hemorrhage occurs for the first time before the seventh month of utero-gestation. The chances of the child surviving would then be very small, and if the hemorrhage be not alarming, as at that early period is likely to be the case, the measures indicated above may be employed in the hope of carrying on the pregnancy until there is a prospect of the patient being delivered of a living child. But little benefit is likely to accrue from astringent drugs. Perfect rest in bed is more likely to be beneficial than anything else, and astringent vaginal pessaries of matico or perchloride of iron might be used with advantage as local hæmostatics.

Various Methods of Treatment.—When the period of pregnancy or the urgency of the case determines us to forego any attempt at temporizing, there are various plans of treatment to be considered. These are, chiefly: 1. *Puncture of the membranes*; 2. *Plugging of the vagina*; 3. *Turning*; 4. *Partial or complete separation of the placenta*. It will be well to consider in detail the relative advantages of, and indications for, each of these. It is seldom, however, that we can trust to any one *per se*; in most cases two or more are required to be used in combination.

1. *Puncture of the membranes* is recommended by Barnes as the first measure to be adopted in all cases of placenta prævia sufficient to cause anxiety. "It is," he says, "the most generally efficacious remedy, and it can always be applied." The primary object gained is the increase of uterine contraction by the evacuation of the liquor amnii. Although the first effect of this may be to increase the flow of blood by further separation of the placenta, the flooding can generally be commanded by plugging until the os is sufficiently dilated to permit the passage of the child. As a rule, there is no great difficulty in effecting the puncture, especially if the placental presentation be only partial. A quill or other suitable contrivance, guided by the examining finger, is passed through the cervix and pushed through the membranes. In complete placenta prævia it may not be so easy to effect the evacuation of the liquor amnii; and, although many authorities advise the penetration of the substance of the placenta itself, I am inclined to think that it would be better to

abandon the attempt in such cases and trust to other methods of treatment.

The objections which have been raised to puncture of the membranes are chiefly that it interferes with the gradual dilatation of the os and renders the operation of turning much more difficult. The os is not, however, so regularly dilated by the bag of membranes in cases of placenta prævia as it is in ordinary labors. Moreover, as the cervical tissues are generally relaxed by the hemorrhage, the dilatation is easily effected. Should we desire to dilate the os preparatory to turning, we can readily do so by means of Barnes's bags, which act at the same time as an efficient plug. The objections, therefore, are not so weighty as they might have been before these artificial dilators were used. I am inclined for these reasons to agree with the recommendation that puncture of the membranes should be resorted to in all cases of placenta prævia.

2. *Plugging of the vagina*—or, still better, of the cavity of the cervix itself—is especially serviceable in cases in which the os is not sufficiently dilated to admit of turning or of separation of the placenta, and in which the hemorrhage still continues after the evacuation of the liquor amnii. By means of this contrivance the escape of blood is effectually controlled.

The best way of plugging is to introduce a sponge tent of sufficient size into the cervical canal, and to keep it *in situ* by a vaginal plug; the best material for the latter and the method of introduction are described under the head of Abortion. The sponge tent not only controls the hemorrhage more effectually than any other means, but is at the same time effecting dilatation of the cervix. It cannot be left in many hours, on account of the irritation produced and of the fetor from accumulating vaginal discharges. As long as it is in position we should carefully examine, from time to time, to see that no blood is oozing past it. If preferred, a Barnes's bag may be used for the same purpose.

While the plug is *in situ* other modes of exciting uterine action may be very advantageously employed, such as a firm abdominal bandage, occasional friction over the uterus, and repeated doses of ergot. The last is specially recommended by Dr. Greenhalgh, who used at the same time a plug formed of an oblong india-rubber ball inflated with air and covered with spongio-piline.

On the removal of the plug we may find that considerable dilatation has taken place, perhaps to a sufficient extent to admit of labor being safely concluded by the natural efforts. In that case we shall find that, although the pains continue, no fresh hemorrhage occurs. Should it do so, it will be necessary to adopt further measures.

3. *Turning* has long been considered *the remedy par excellence* in placenta prævia; and it is of unquestionable value in suitable cases. Much harm, however, has been done when it has been practised before the os was sufficiently dilated to admit of the passage of the hand, or when the patient was so exhausted by previous hemorrhage as to be unable to bear the shock of the operation. The records of many fatal cases in the practice of those who taught, as did the large majority of the older writers, that turning at all risks was essential conclusively prove this assertion.

It is most likely to prove serviceable when, either at first or after the use of the tampon, the os is sufficiently dilated to admit the hand, and when the strength of the patient is not much enfeebled. If she have a small, feeble, and thready pulse, it is certainly inapplicable, unless all other methods of arresting the hemorrhage have failed. And even then it would be well to attempt to rally the patient from her exhausted state by stimulants, etc. before the operation is commenced.

Provided the placental presentation be partial, the operation can be performed without difficulty in the usual way. In central implantation the passage of the hand may give rise to some difficulty. Dr. Rigby recommends that it should be pushed through the substance of the placenta until it reaches the uterine cavity. It is hardly possible to conceive how this could be done without completely detaching the placenta, and still less to understand how the fetus could be dragged through the aperture thus made. It will be far better to pass the hand by the border of the placenta, separating it as we do so; and, if we can ascertain to which side of the cervix it is least attached, that should be chosen for the purpose. In all cases in which it is possible turning by the bi-polar method should be preferred. In cases of placenta prævia especially it offers many advantages. The operation can be soon performed, complete dilatation of the os is not so necessary, and it involves less bruising of the cervix, which is likely to be specially dangerous. When once a foot has been brought within the os, the delivery need not be hurried. The foot forms a plug which effectually prevents all further loss, and we may then safely wait until we can excite uterine contraction and terminate the labor with safety. Fortunately, the relaxation of the uterus which is so often present facilitates this manner of performing version, and it can generally be successfully accomplished. Should the case be one which is otherwise suitable for turning, and the requisite amount of dilatation of the cervix not be present, the latter can generally be effected in the space of an hour or more (while at the same time a further loss of blood is effectually prevented) by the use of Barnes's bags.

4. *Entire separation of the placenta* was originally recommended by Simpson in his well-known paper on the subject. The reasons which induced him to recommend it have already been stated. It is a mistake to suppose, however, as is so often done, that he intended to recommend it in all cases alike. This supposition he was always careful to deny. He advised it especially—

1. When the child is dead.
2. When the child is not yet viable.
3. When the hemorrhage is great and the os uteri is not yet sufficiently dilated for safe turning. This was the state in 11 out of 39 cases (Lee).
4. When the pelvic passages are too small for safe and easy turning.
5. When the mother is too exhausted to bear turning.
6. When the evacuation of the liquor amnii fails.
7. When the uterus is too firmly contracted for turning.¹

These are very much the cases in which all modern accoucheurs would exclude the operation of turning; and it was especially when that was unsuitable that Simpson advised extraction of the placenta. As his

¹ *Selected Obst. Works*, p. 68.

theory of the source of hemorrhage is now almost universally disbelieved, so has the practice based on it fallen into disuse, and it need not be discussed at length. It is very doubtful whether the complete separation and extraction of the placenta was a feasible operation; unquestionably, it can be by no means so easy as Simpson's writings would lead us to suppose. The introduction of the hand far enough to remove the placenta in an exhausted patient would probably cause as much shock as the operation of turning itself; and another very formidable objection to the procedure is the almost certain death of the child if any time elapse between the separation of the placenta and the completion of delivery. The modification of this method so strongly advocated by Barnes is certainly much easier of application, and would appear to answer every purpose that Simpson's operation effected. It is impossible to describe it better than in Barnes's own words:¹

"The operation is this: Pass one or two fingers as far as they will go through the os uteri, the hand being passed into the vagina if necessary; feeling the placenta, insinuate the finger between it and the uterine wall; sweep the finger round in a circle so as to separate the placenta as far as the finger can reach: if you feel the edge of the placenta where the membranes begin, tear open the membranes carefully, especially if these have not been previously ruptured; ascertain, if you can, what is the presentation of the child before withdrawing your hand. Commonly, some amount of retraction of the cervix takes place after the operation, and *often the hemorrhage ceases.*"

It will be seen, from what has been said, that no one rule of practice can be definitely laid down for all cases of placenta prævia. Our treatment in each individual case must be guided by the particular conditions that are present; and if only we bear in mind the natural history of the hemorrhage, we may confidently look to a favorable termination.

Summary of Rules for Treatment.—It may be useful, in conclusion, to recapitulate the rules which have been laid down for treatment in the form of a series of propositions:

I. Before the child has reached a viable age temporize, provided the hemorrhage be not excessive, until pregnancy has advanced sufficiently to afford a reasonable hope of saving the child. For this purpose the chief indication is absolute rest in bed, to which other accessory means of preventing hemorrhage, such as cold, astringent pessaries, etc., may be added.

II. In hemorrhage occurring after the seventh month of utero-gestation no attempt should be made to prolong the pregnancy.

III. In all cases in which it can be easily effected the membranes should be ruptured. By this means uterine contractions are favored and the bleeding vessels compressed.

IV. If the hemorrhage be stopped, the case may be left to nature. If flooding continue, and the os be not sufficiently dilated to admit of the labor being readily terminated by turning, the os and the vagina should be carefully plugged, while uterine contractions are promoted by abdominal bandages, uterine compression, and ergot. The plug must not be left in beyond a few hours.

¹ *Obstet. Operations*, 2d ed., p. 417.

V. If, on removal of the plug, the os be sufficiently expanded and the general condition of the patient be good, the labor may be terminated by turning, the bi-polar method being used if possible. If the os be not open enough, it may be advantageously dilated by a Barnes's bag, which also acts as a plug.

VI. Instead of, or before resorting to, turning the placenta may be separated around the site of its attachment to the cervix. This practice is specially to be preferred when the patient is much exhausted and in a condition unfavorable for bearing the shock of turning.

[*Dr. J. Braxton Hicks' bi-manual method of turning*,¹ as tested in Berlin by Drs. Hofmeier, Behm, and Lomer, promises much better results than any other method of treatment in cases of placenta prævia. According to Dr. Lomer's report in the *Am. Journ. of Obstetrics* for December, 1884, Dr. Hofmeier operated upon 37 cases, and saved 36 women and 14 children; Dr. Behm, upon 40 cases, all saved, but lost 31 children; and he himself, with eight other assistants, upon 101 cases, saving 94, with 50 children. This gives 8 deaths of women and 105 of children in 178 cases, or a mortality of 4½ per cent. of the former and 60 per cent. of the latter. Dr. Lomer's directions are as follows: "Turn by the bi-manual method as soon as possible; pull down the leg, and tampon with it and with the breech of the child the ruptured vessels of the placenta. *Do not extract the child then*: let it come by itself, or at least only assist its natural expulsion by gentle and rare tractions. Do away with the plug as much as possible; it is a dangerous thing, for it favors infection and valuable time is lost with its application. Do not wait in order to perform turning until the cervix and the os are sufficiently dilated to allow the hand to pass. Turn as soon as you can pass one or two fingers through the cervix. It is unnecessary to force your fingers through the cervix for this. Introduce the whole hand into the vagina, pass one or two fingers through the cervix, rupture the membranes, and turn by Braxton Hicks' bi-manual method." . . . "If the placenta is in your way, try to rupture the membranes at its margin; but if this is not feasible, do not lose time: perforate the placenta with your finger; get hold of a leg as soon as possible, and bring it down."—ED.]

CHAPTER XIV.

HEMORRHAGE FROM SEPARATION OF A NORMALLY-SITUATED PLACENTA.

Definition.—This is the form of hemorrhage which is generally described in obstetric works as "*accidental*," in contradistinction to the "*unavoidable*" hemorrhage of placenta prævia. In discussing the latter we have seen that the term "*accidental*" is one that is apt to mislead,

[¹ *Lancet*, July, 1860; *Obstetrical Transactions*, vol. v. p. 222.]

and that the causation of the hemorrhage in placenta prævia is, in some cases at least, closely allied to that of the variety of hemorrhage we are now considering.

When, from any cause, separation of a normally-situated placenta occurs before delivery, more or less blood is necessarily effused from the ruptured utero-placental vessels, and the subsequent course of the case may be twofold: 1. The blood, or at least some part of it, may find its way between the membranes and the decidua, and escape from the os uteri. This constitutes the typical "accidental" hemorrhage of authors. 2. The blood may fail to find a passage externally, and may collect internally, giving rise to very serious symptoms, and even proving fatal, before the true nature of the case is recognized. Cases of this kind are by no means so rare as the small amount of attention paid to them by authors might lead us to suppose; and from the obscurity of the symptoms and difficulty of diagnosis they merit special study. Dr. Goodell¹ has collected together no less than 106 instances in which this complication occurred.

Causes and Pathology.—The causes of placental separation may be very various. In a large number of cases it has followed an accident or exertion (such as slipping down stairs, stretching, lifting heavy weights, and the like) which has probably had the effect of lacerating some of the placental attachments. At other times it has occurred without such appreciable cause, and then it has been referred to some change in the uterus, such as a more than usually strong contraction producing separation, or some accidental determination of blood causing a slight extravasation between the placenta and the uterine wall, the irritation of which leads to contraction and further detachment. Causes such as these, which are of frequent occurrence, will not produce detachment except in women otherwise predisposed to it. It generally is met with in women who have borne many children, more especially in those of weakly constitution and impaired health, and rarely in primiparæ. Certain constitutional states probably predispose to it, such as albuminuria or exaggerated anæmia, and, still more so, degenerations and diseases of the placenta itself.

This form of hemorrhage rarely occurs to an alarming extent until the latter months of pregnancy, often not until labor has commenced. The great size of the placental vessels in advanced pregnancy affords a reasonable explanation of this fact.

Symptoms and Diagnosis.—If, after separation of a portion of the placenta, the blood finds its way between the membranes and the decidua, its escape per vaginam, even although in small amount, at once attracts attention and reveals the nature of the accident. It is otherwise when we have to do with a case of concealed hemorrhage, the diagnosis of which is often a matter of difficulty. Then the blood probably at first collects between the uterus and the placenta. Sometimes marginal separation does not occur, and large blood-clots are formed in this situation and retained there. More often the margin of the placenta separates and the blood collects between the membranes and the uterine wall, either toward the cervix, where the presenting part of the child may

¹ *Amer. Journ. of Obst.*, vol. ii.

prevent its escape, or near the fundus. In the latter case especially, the coagula are apt to cause very painful stretching and distension of the uterus. The blood may also find its way into the amniotic cavity, but more frequently it does not do so—probably, as Goodell has pointed out, because, “should the os uteri be closed, the membranes, however delicate, cannot, other things being equal, rupture any sooner from the uterine walls, for the sum of the resistance of the enclosed liquor amnii, being equally distributed, exactly counterbalances the sum of the pressure exerted by the effusion.” This point is of some practical importance, because, after rupture of the membranes, the liquor amnii is frequently found untinged with blood; and this might lead us to suppose ourselves mistaken in our diagnosis if this fact were not borne in mind.

Symptoms of Concealed Accidental Hemorrhage.—The most prominent symptoms in concealed internal hemorrhage are extreme collapse and exhaustion, for which no adequate cause can be assigned. These differ from those of ordinary syncope, with which they might be confounded, chiefly in their persistence and severity, and in the presence of the symptoms attending severe loss of blood, such as coldness and pallor of the surface, great restlessness and anxiety, rapid and sighing respiration, yawning, feeble, quick, and compressible pulse. When there is severe internal, with slight external, hemorrhage, we may be led to a proper diagnosis by observing that the constitutional symptoms are much more severe than the amount of external hemorrhage would account for. Uterine pain is generally present, of a tearing and stretching character, sometimes moderate in amount, more often severe, and occasionally amounting to intolerable anguish. It is often localized, and doubtless depends on the distension of the uterus by the retained coagula. If the distension be marked, there may be an irregularity in the form of the uterus at the site of sanguineous effusion; but this will be difficult to make out, except in women with thin and unusually lax abdominal parietes. A rapid increase in the size of the uterus has been described as a sign by Cazeaux and others. It is not very likely that this will be appreciable toward the end of utero-gestation, as a very large amount of effusion would be necessary to produce it. At an earlier period of pregnancy, at or about the fifth month, I made it out very distinctly in a case in my own practice. It obviously must have occurred to an enormous extent in a case related by Chevalier, in which post-mortem Cæsarean section was performed under the impression that the pregnancy had advanced to term, but only a three months’ fœtus was found, imbedded in coagula which distended the uterus to the size of a nine months’ gestation.¹ Labor-pains may be entirely absent. If present, they are generally feeble, irregular, and inefficient.

Differential Diagnosis.—The only condition, besides ordinary syncope, likely to be confounded with this form of hemorrhage is rupture of the uterus, to which the intense pain and profound collapse induce considerable resemblance. The latter rarely occurs until after labor has been some time in progress and after the escape of the liquor amnii; whereas hemorrhage usually occurs either before labor has commenced or at an early stage. The recession of the presentation and the escape of the

¹ *Journ. de Méd. clin. et pharmac.*, vol. xxi. p. 363.

fœtus into the abdominal cavity in cases of rupture will further aid in establishing the diagnosis.

Prognosis.—The prognosis, when blood escapes externally, is, on the whole, not unfavorable. The nature of the case is apparent, and remedial measures are generally adopted sufficiently early to prevent serious mischief. It is different with the concealed form, in which the mortality is very great. Out of Goodell's 106 cases, no less than 54 mothers died. This excessive death-rate is no doubt partly due to the fact that extreme prostration so often occurs before the existence of hemorrhage is suspected, and partly to the accident generally happening in women of weakly and diseased constitution. The prognosis to the child is still more grave. Out of 107 children, only 6 were born alive. The almost certain death of the child may be explained by the fact that when blood collects between the uterus and the placenta the fœtal portion of the latter is probably lacerated, and the child then also dies from hemorrhage.

Treatment.—In this, as in all other forms of puerperal hemorrhage, the great hæmostatic is uterine contraction, and that we must try to encourage by all possible means. The first thing to be done, whether the hemorrhage be apparent or concealed, is to rupture the membranes. If the loss of blood be only slight, this may suffice to control it, and the case may then be left to nature. A firm abdominal binder should, however, be applied to prevent any risk of blood collecting internally, as there is nothing to prevent its filling the uterine cavity after the membranes are ruptured. Contraction may be further advantageously solicited by uterine compression and by the administration of full doses of ergot. If hemorrhage continue, or if we have any reason to suspect concealed hemorrhage, the sooner the uterus is emptied the better. If the os be sufficiently dilated, the best practice will be to turn without further delay, using the bi-polar method if possible. If the os be not open enough, a Barnes's bag should be introduced, while firm pressure is kept up to prevent uterine accumulation. Should the collapsed condition of the patient be very marked, the mere shock of the operation might turn the scale against her. Under such circumstances it may be better practice to delay further procedure until, by the administration of stimulants, warmth, etc., we have succeeded in producing some amount of reaction, keeping up, in the mean while, firm pressure on the uterus. Should the head be low down in the pelvis, it may be easier to complete labor by means of the forceps.

CHAPTER XV.

HEMORRHAGE AFTER DELIVERY.

Its Importance.—Hemorrhage during or shortly after the third stage of labor is one of the most trying and dangerous accidents connected with parturition. Its sudden and unexpected occurrence just after the labor appears to be happily terminated, and its alarming effect on the patient, who is often placed in the utmost danger in a few moments, tax the presence of mind and the resources of the practitioner to the utmost, and render it an imperative duty on every one who practises midwifery to make himself thoroughly acquainted with its causes and preventive and curative treatment. There is no emergency in obstetrics which leaves less time for reflection and consultation, and the life of the patient will often depend on the prompt and immediate action of the medical attendant.

Frequency of Post-partum Hemorrhage.—Post-partum hemorrhage is one of the most frequent complications of delivery. I do not know of any statistics which enable us to judge with accuracy of its frequency, but I believe it to be an unquestionable fact that, especially in the upper ranks of society, it is very common indeed. This is probably due to the effects of civilization and to the mode of life of patients of that class, whose whole surroundings tend to produce a lax habit of body which favors uterine inertia, the principal cause of post-partum hemorrhage. In the report of the Registrar-General for the five years from 1872 to 1876, 3524 deaths are attributed to flooding. The majority of these must have been caused by post-partum hemorrhage, although some may have been from other forms.

Generally a Preventable Accident.—Fortunately, it is, to a great extent, a preventable accident. I believe this fact cannot be too strongly impressed on the practitioner. If the third stage of labor be properly conducted, if every case be treated—as every case ought to be—as if hemorrhage were impending, it would be much more infrequent than it is. It is a curious fact that post-partum hemorrhage is much more common in the practice of some medical men than in that of others, the reason being that those who meet with it often are careless in their management of their patients immediately after the birth of the child. That is just the time when the assistance of a properly-qualified practitioner is of value—much more so than before the second stage of labor is concluded; hence, when I hear that a medical man is constantly meeting with severe post-partum hemorrhage, I hold myself justified, *ipso facto*, in inferring that he does not know or does not practise the proper mode of managing the third stage of labor.

Causes and Nature's Method of Controlling Hemorrhage after Delivery.—The placenta, as we have seen, is separated by the last pains, and the blood, which in greater or less quantity accompanies the fœtus, prob-

ably comes from the utero-placental vessels which are then lacerated. Almost immediately afterward the uterus contracts firmly, and in a typical labor assumes the hard cricket-ball form which is so comforting to the accoucheur to feel. The result is the compression of all the vascular trunks which ramify in its walls, both arteries and veins, and thus the flow of blood though them is prevented. By referring to what has been said as to the anatomy of the muscular fibres of the gravid uterus, especially at the placental site (p. 65), it will be seen how admirably they are adapted for this purpose. The arrangement of the vessels themselves favors the hæmostatic action of uterine contraction. The large venous sinuses are placed in layers, one above the other, in the thickness of the uterine walls, and they anastomose freely. When the superimposed layers communicate with those immediately below them, the junction is by a falciform or semilunar opening in the floor of the vessel nearest the external surface of the uterus. Within the margins of this aperture there are muscular fibres, the contraction of which probably tends to prevent retrogression of blood from one layer of vessels into the other. The venous sinuses themselves are of a flattened form, and they are intimately attached to the muscular tissues. It is obvious, then, that these anatomical arrangements are eminently adapted to facilitate the closure of the vessels. They are, however, large, and are destitute of valves; and if contraction be absent or if it be partial and irregular, it is equally easy to understand why blood should pour forth in the appalling amount which is sometimes observed.

Importance of Tonic Uterine Contraction.—If uterine action be firm, regular, and continuous, the vessels must be sealed up and hemorrhage effectually prevented. This fact has been doubted by many authorities. Gooch was the first to describe what he called "a peculiar form of hemorrhage" accompanying a contracted womb; similar observations have been made by other writers, such as Velpeau, Rigby, and Gendrin. Simpson says on this point that strong uterine contractions "are not probably so essential a part in the mechanism of the prevention of hemorrhage from the open orifices of the uterine veins as we might, *a priori*, suppose."¹ With regard to Gooch's cases it has been pointed out that his own description proves that, however firmly the uterus may have contracted immediately after the expulsion of the child, it must have subsequently relaxed, for he passed his hand into it to remove retained clots—a manœuvre which he could not have practised had tonic contraction been present. In some of these cases the hemorrhage has been found to come from a laceration of the cervix. Of course, blood may readily escape from mechanical injury of this kind, although the uterus itself be in a satisfactory state of contraction; and the possibility of this occurrence should always be borne in mind. Instances of the successful treatment of this variety of post-partum hemorrhage by sutures applied to the lacerated cervix have been related by Pallen and others.

Although, then, we may admit that post-partum hemorrhage is incompatible with persistent contraction of the uterus, it by no means follows that the converse is true. On the contrary, it is not uncommon to meet with cases in which the uterus is large and apparently quite

¹ *Selected Obst. Works*, p. 234.

flaccid, and in which there is no loss of blood. Alternate relaxation and contraction of the uterus after delivery are also of constant occurrence, and yet hemorrhage during the relaxation does not take place. The explanation, no doubt, is, that immediately after the birth of the child there was sufficient contraction to prevent hemorrhage, and that during its continuance coagula formed in the mouths of the uterine sinuses, by which they were sufficiently occluded to prevent any loss when subsequent relaxation occurred.

In all probability, both uterine contraction and thrombosis are in operation in ordinary cases; and we shall presently see that all the means employed in the treatment of post-partum hemorrhage act by producing one or other of them.

Secondary Causes of Hemorrhage.—Uterine inertia after labor, then, may be regarded as the one great primary cause of post-partum hemorrhage; but there are various secondary causes which tend to produce it, one of the most frequent of which is exhaustion following a protracted labor. The uterus gets worn out by its efforts, and when the fetus is expelled it remains in a relaxed state, and hemorrhage results. Overdistension of the uterus acts in the same way. Hence, hemorrhage is very frequently met with when there has been an excessive amount of liquor amnii or in multiple pregnancies. One of the worst cases I ever met with was after the birth of triplets, the uterus having been of an enormous size. Rapid emptying of the uterus during which there has not been sufficient time for complete separation of the placenta often tends to the same result. This is the reason why hemorrhage so frequently follows forceps delivery, especially if the operation have been unduly hurried, and it is one of the chief dangers to which are referred "prolonged labors." The general condition of the patient may also strongly predispose to it. Thus, it is more often met with in women who have some families, especially if they be weakly in constitution, compared with women who are primiparae, and for the same reason that other parts are more susceptible of the disease—namely, that the uterine relaxation in the former is a relaxing contracta, influenced by the experience of previous labors. The evidence shows that labor pains must be relaxed by the relaxing effects of various causes, and peculiarly prone to it, and it forms one of the chief dangers of childbearing amongst the English middle class.


Influence of Uterine Contraction.—Another important cause of post-partum hemorrhage is partial and temporary relaxation of the uterus. Part of the muscular tissue of the contraction, which sometimes produces rupture of the uterus, may relax, the pressure still being maintained external to the uterus, and the rupture will be the result. This is the case when the placenta is torn from the uterus. The same effect is produced in abortion, when the placenta is torn from the uterus, and the result is a post-partum hemorrhage, which may be very severe, and even fatal. In the ordinary course of the different periods of the contraction, the uterus is relaxed, and the blood is allowed to flow out of the uterus, and the result is a post-partum hemorrhage, which may be very severe, and even fatal. In the ordinary course of the different periods of the contraction, the uterus is relaxed, and the blood is allowed to flow out of the uterus, and the result is a post-partum hemorrhage, which may be very severe, and even fatal.

The treatment of post-partum hemorrhage is, therefore, to be directed to the production of uterine contraction, and to the removal of the cause of relaxation.

so-called "hour-glass contraction." This in reality seems to depend on spasmodic contraction of the internal os uteri, by means of which the placenta becomes encysted in the upper portion of the uterus, which is relaxed. On introducing the hand it first passes through the lax cervical canal until it comes to the closed internal os, with the umbilical cord passing through it, which has generally been supposed to be a circular contraction of a portion of the body of the uterus.

Encystment of the placenta, however, although more rare, unquestionably takes place in a portion only of the body of the uterus (Fig. 145). Then, apparently, the placental site remains more or less para-

FIG. 145.



Irregular Contraction of the Uterus, with Encystment of the Placenta.

lyzed, with the placenta still attached, while the remainder of the body of the uterus contracts firmly, and thus encystment is produced.^[1]

Causes of Irregular Contractions.—These irregular contractions of the uterus are by no means so common as our older authors supposed.

[¹ The central constriction of the right-hand figure is intended to represent the internal os uteri. The condition of the left-hand illustration, with a contraction around one corner, I have very distinctly defined in a case of retained placenta. The adherent placenta was entirely enclosed, and the contracted portion barely sufficed to admit the hand. When the adhesion was broken up, the inert portion of the uterine wall at once contracted: this was no doubt partly the result of the removal of the mechanical obstacle, and partly due to the stimulating effect of the presence and work of the hand. This experience is in correspondence with the opinion of the late Prof. C. D. Meigs, who taught that an *irregularly-contracted uterus* was the effect of an adherent placenta acting as an obstacle to contraction over the seat of union, while the rest of the organ was free to contract. Whether the central muscular fibres of the uterus can ever contract so as to constrict the organ is a question which has long been in dispute. Those who deny its possibility do so upon anatomical reasoning, claiming that the arrangement of the circular muscular fibres is such that a violent linear contraction in the *corpus uteri* is an anatomical impossibility. Those who have had under care cases of "tetanoid constriction of the uterus" are much exercised as to the exact location of the spasmodic band of fibres which has been found both circular and oblique and tightly surrounding the body of the fetus. The discovery of Bandl of Vienna (see p. 433, chap. xvi.) might possibly account for the condition found in one of the cases of Dr. T. A. Foster of Portland, Maine (*Transactions Maine Med. Association*, 1868-69, p. 273), where the child was tightly held by the neck, but for the fact that the uterine walls were thicker than normal. Dr. Thomas C. Smith of Washington, D. C., has, I think, made it clear that tetanoid constriction may occur in the body of the uterus. (See *Am. Journ. of Obstetrics*, April, 1882, pp. 294-322.)—ED.]

When they do occur, I believe them almost invariably to depend on defective management of the third stage of labor. "The most frequent cause," says Rigby,¹ "is from over-anxiety to remove the placenta; the cord is frequently pulled at, and at length the os uteri is excited to contract." While this is being done no attempts are probably being made to excite the fundus to proper action, and therefore the hour-glass contraction is established. Duncan says of this condition, "Hour-glass contraction cannot exist unless the parts above the contraction are in a state of inertia; were the higher parts of the uterus even in moderate action the hour-glass contraction would soon be overcome."² If placental expression were always employed, if it were the rule to effect the expulsion of the placenta by a *vis a tergo* instead of extracting by a *vis a fronte*, I feel confident that these irregular and spasmodic contractions—of the influence of which in producing hemorrhage there can be no question—would rarely, if ever, be met with. It is to be observed that even in these cases it is not because the uterus is in a state of partial contraction, but because it is in a state of partial relaxation, that hemorrhage ensues.

Placental Adhesions.—Adhesions of the placenta to the uterine parietes may cause hemorrhage, especially if they be partial and the remainder of the placenta be detached. The frequency of these has been over-estimated. Many cases believed to be examples of adherent placenta are, in reality, only cases of placenta retained from uterine inertia. The experience of all who see much midwifery will probably corroborate the observation of Braun, that "abnormal adhesion and hour-glass contraction are more frequently encountered in the experience of the young practitioner, and they diminish in frequency in direct ratio to increasing years."³ The cause of adhesion is often obscure, but it most probably results from a morbid state of the decidua, which is produced by antecedent disease of the uterine mucous membrane; then the adhesion is apt to recur in subsequent pregnancies. The decidua is altered and thickened, and patches of calcareous and fibrous degeneration may be often found on the attached surface of the placenta. Most frequently the placenta is only partially adherent; patches of it remain firmly attached to the uterus, while the rest is separated; hence the uterine walls remain relaxed and hemorrhage frequently follows. The diagnosis and management of these very troublesome cases will be found described under the head of Treatment (p. 423).

Constitutional Predisposition to Flooding.—Finally, I think it must be admitted that there are some women who really merit the appellation of "flooders" which has been applied to them, and who, do what we may, have the most extraordinary tendency to hemorrhage after delivery. I do not think that these cases, however, are by any means so common as some have supposed. I have attended several patients who have nearly lost their lives from post-partum hemorrhage in former labors, some who have suffered from it in every preceding confinement, and I have only met with two cases in which the assiduous use of preventive treatment failed to avert it. In these (one of which I have elsewhere

¹ Rigby's *Midwifery*, p. 225.

² *Researches in Obstetrics*, p. 389.

³ Braun's *Lectures*, 1869.

published in detail¹), in spite of all my efforts, I could not succeed in keeping up uterine contraction, and the patients would certainly have lost their lives were it not for the means which modern improvements have fortunately placed at our disposal for producing thrombosis in the mouths of the bleeding vessels. The nature of these rare cases requires further investigation; possibly they may, to some extent, be the subjects of the so-called hemorrhagic diathesis.

Signs and Symptoms.—The loss of blood may commence immediately after the birth of the child, before the expulsion of the placenta, or not until some time afterward, when the contracted uterus has again relaxed. It may commence gradually or suddenly: in the latter case it may begin with a gush, and in the worst form the bedclothes, the bed, and even the floor, are deluged with the blood which, it is no exaggeration to say, is pouring from the patient. If, now, the hand be placed on the abdomen, we shall miss the hard round ball of the contracted uterus, which will be found soft and flabby, or we may even be unable to make out its contour at all. If the hemorrhage be slight or if we succeed in controlling it at once, no serious consequences follow; but if it be excessive or if we fail to check it, the gravest results ensue.

Exhaustion in Extreme Cases.—There are few sights more appalling to witness than one of the worst cases of post-partum hemorrhage. The pulse becomes rapidly affected, and may be reduced to a mere thread or it may become entirely imperceptible. Syncope often comes on—not in itself always an unfavorable occurrence, as it tends to promote thrombosis in the venous sinuses. Or, short of actual syncope, there may be a feeling of intense debility and faintness. Extreme restlessness soon supervenes, the patient throws herself about the bed, tossing her arms wildly above her head; respiration becomes gasping and sighing, the “besoin de respirer” is acutely felt, and the patient cries out for more air; the skin becomes deadly cold and covered with profuse perspiration; if the hemorrhage continue unchecked, we next may have complete loss of vision, jactitation, convulsions, and death.

Formidable as such symptoms are, it is satisfactory to know that recovery often takes place, even when the powers of life seem reduced to the lowest ebb. If we can check the hemorrhage while there is still some power of reaction left, however slight, we may not unreasonably hope for eventual recovery. The constitution, however, may have received a severe shock, and it may be months, or even years, before the patient recovers from the effects of only a few minutes’ hemorrhage. A death-like pallor frequently follows these excessive losses, and the patient often remains blanched and exsanguine for a long time.

Preventive Treatment.—The preventive treatment of post-partum hemorrhage should be carefully practised in every case of labor, however normal. If the practitioner make a habit of never removing his hand from the uterus after the birth of the child until the placenta is expelled, and of keeping up continuous uterine contraction for at least half an hour after delivery is completed—not necessarily by friction on the fundus, but by simply grasping the contracted womb with the palm of the hand and preventing its undue relaxation—cases of post-partum

¹ *Obst. Journ.*, vol. i.

flooding will seldom be met with. As a rule, we should, I think, not apply the binder until at least that time has elapsed. The binder is an effective means of keeping up, but not of producing, contraction, and it should never be trusted to for the latter purpose. If it be put on too soon, the uterus may relax under it, and become filled with clots without the practitioner knowing anything about it; whereas this cannot possibly take place as long as the uterine globe is held in the hollow of the hand. I have seen more than one serious case of concealed hemorrhage result from the too common habit of putting on the binder immediately after the removal of the placenta. I believe also, as I have formerly said, that it is thoroughly good practice to administer a full dose of the liquid extract of ergot in all cases after the placenta has been expelled, to ensure persistent contraction and to lessen the chance of blood-clots being retained in utero.

These are the precautions which should be used in all cases alike; but when we have reason to fear the occurrence of hemorrhage from the history of previous labors or other cause, special care should be taken. The ergot should be given, and preferably in the form of the subcutaneous injection of ergotin, before the birth of the child, when the presentation is so far advanced that we estimate that labor will be concluded in from ten to twenty minutes, as we can hardly expect the drug to produce any effect in less time. Particular attention, moreover, should then be paid to the state of the uterus. Every means should be taken to ensure regular and strong contraction, and it is advisable to rupture the membranes early, as soon as the os is dilated or dilatable, to ensure stronger uterine action. If any tendency to relaxation occur after delivery, a piece of ice should be passed into the vagina or into the uterus. Should coagula collect in the uterus, they may be readily expelled by firm pressure on the fundus, and the finger should be passed occasionally up to the cervix, and any which are felt there should be gently picked away.

We should be specially on our guard in all cases in which the pulse does not fall after delivery. If it beat at 100 or more some ten minutes or a quarter of an hour after the birth of the child, hemorrhage not unfrequently follows; and hence it is a good practical rule, which may save much trouble, that a patient should never be left unless the pulse has fallen to its natural standard.

Curative Treatment.—As there are only two means which nature adopts in the prevention of post-partum hemorrhage, so the remedial measures also may be divided into two classes: 1, those which act by the production of uterine contraction; 2, those which act by producing thrombosis in the vessels. Of these, the first are the most commonly used; and it is only in the worst cases, in which they have been assiduously tried and have failed, that we resort to those coming under the second heading.

Uterine Pressure.—The patient should be placed on her back, in which position we can more readily command the uterus as well as attend to her general state. If the uterus be found relaxed and full of clots, by firmly grasping it in the hand contraction may be evoked, its contents expelled, and further hemorrhage at once arrested. Should this, fortunately, be the case, we must keep up contraction by gently

kneading the uterus until we are satisfied that undue relaxation will not recur.

The powerful influence of friction in promoting contraction cannot be doubted, and nothing will replace it; no doubt it is fatiguing, but as long as it is effectual it must be kept up. No roughness should be used, as we might produce subsequent injury, but it is quite possible to use considerable pressure without any violence.

Another method of applying uterine pressure has been strongly advocated by Dr. Hamilton of Falkirk, and it may be serviceable where there is a constant draining from the uterus and a capacious pelvis. It consists in passing the fingers of the right hand high up into the posterior cul-de-sac of the vagina so as to reach the posterior surface of the uterus, while counter-pressure is exercised by the left hand through the abdomen. The anterior and posterior walls of the uterus are thus closely pressed together.

Administration of Ergot.—During the time that pressure is being applied attention can be paid to general treatment; and in giving his directions to the bystanders the practitioner should be calm and collected, avoiding all hurry and excitement. A full dose of ergot should be administered, and if one have already been given it should be repeated. We cannot, however, look upon ergot as anything but a useful accessory, and it is one which requires considerable time to operate. The hypodermic use of ergotin offers the double advantage, in severe cases, of acting with greater power and much more rapidly than the usual method of administration. It should, therefore, always be used in preference. An aqueous solution of ergotin, $\frac{1}{200}$ th of a grain in 10 minims, has been highly recommended by Chahbazain of Paris as acting more energetically, but of this I have no experience.¹

Stimulants.—The sudden flow will probably have produced exhaustion and a tendency to syncope, and the administration of stimulants will be necessary. The amount must be regulated by the state of the pulse and the degree of exhaustion. There is no more absurd mistake, however, than implicitly relying on the brandy-bottle to check post-partum hemorrhage. In the worst cases absorption is in abeyance, and brandy may be poured down in abundance, the practitioner believing that he is rousing his patient, while he is, in fact, only filling the stomach with a quantity of fluid which is eventually thrown up unaltered. I have more than once seen symptoms produced by the over-free use of brandy in slight floodings which were certainly not those of hemorrhage. I remember on one occasion being summoned by a practitioner, with a view to transfusion, to a patient who was said to be insensible and collapsed from hemorrhage. I found her, indeed, unconscious, but with a flushed face, a bounding pulse, a firmly-contracted uterus, and deep stertorous breathing. On inquiry I ascertained that she had taken an enormous quantity of brandy, which had brought on the coma of profound intoxication, while the hemorrhage had obviously never been excessive.

Hypodermic Injection of Ether.—The phosphoric ether is a remedy of great val

of sulphuric
1 cases

¹ *Obst. Trans.*

in which exhaustion is very great. It has the advantage of acting rapidly and of being capable of administration when the patient is unable to swallow. A fluidrachm may be injected into the nates or thigh, and the injection may be repeated as the state of the patient may require.

Fresh Air, etc.—The window should be thrown widely open, to allow a current of fresh cold air to circulate freely through the room. The pillows should be removed, the head kept low, and the patient should be assiduously fanned.

Emptying of Uterus.—If bleeding continue, or if it commence before the placenta is expelled, the hand should be carefully and gently passed into the uterus and its cavity cleared of its contents. The mere presence of the hand within the uterus is a powerful inciter of uterine action. When the placenta is retained it is the more essential, as the hemorrhage cannot possibly be checked as long as the uterus is distended by it. During the operation the uterus should be supported by the left hand externally, and by using the two hands in concert the chances of injuring the textures are greatly lessened.

Treatment of Hour-glass Contraction.—If the so-called “hour-glass contraction” be present or if the placenta be morbidly adherent, the operation will be more difficult and will require much judgment and care. The spasmodic contraction of the inner os in the former case may generally be overcome by gentle and continuous pressure of the fingers passed within the contraction, while the uterus is supported from without. By this means, too, further hemorrhage can in most cases be controlled until the spasm is sufficiently relaxed to admit of the passage of the hand.

Signs of Adherent Placenta.—There are no very reliable signs to indicate morbid adhesion of the placenta previous to the introduction of the hand. The following are the symptoms as laid down by Barnes, any of which might, however, accompany non-detachment of the placenta unaccompanied by adhesion: “You may suspect morbid adhesion if there have been unusual difficulty in removing the placenta in previous labors; if during the third stage the uterus contracts at intervals firmly, each contraction being accompanied by blood, and yet on following up the cord you feel the placenta *in utero*; if on pulling on the cord, two fingers being pressed into the placenta at the root, you feel the placenta and uterus descend in one mass, a sense of dragging pain being elicited; if during a pain the uterine tumor does not present a globular form, but be more prominent than usual at the place of placental attachment.”¹

Treatment of Adherent Placenta.—The artificial removal of an adherent placenta is always a delicate and anxious operation, which, however carefully performed, must of necessity expose the patient to the risk of injury to the uterine structures, and of leaving behind portions of placental tissue, which may give rise to secondary hemorrhage or septicaemia. The cord will guide the hand to the site of attachment, and the fingers must be very gently insinuated between the lower edge of the placenta and the uterine wall; or, if a portion be already detached, we may commence to peel off the remainder at that spot. Supporting the uterus externally, we carefully pick off as much as possible, proceeding

¹ *Obstetric Operations*, p. 440.

with the greatest caution, as it is by no means easy to distinguish between the placenta and the uterus. At the best, it is far from easy to remove all, and it is wiser to separate only what we readily can than to make too protracted efforts at complete detachment. When it is found to be impossible to detach and remove the whole or a great part of the placenta, we cannot but look upon the further progress of the case with considerable anxiety. The retained portions may be ere long spontaneously detached and expelled, or they may decompose and give rise to a fetid discharge and septic infection. Such cases must be treated by antiseptic intra-uterine injections, so as to lessen the risk of absorption as much as possible; but until the retained masses have been expelled and the discharge has ceased the patient must be considered to be in considerable danger. In a few rare cases there is reason to believe that considerable masses of retained placental tissue have been entirely absorbed. It is difficult to understand so strange a phenomenon, but several well-authenticated cases are recorded in which there seems no reason to doubt that the retained placenta was removed in this way.¹

Excitement of Reflex Action by Cold, etc.—Various means are used for exciting uterine contraction by reflex stimulation. Amongst the most important of these is cold. In patients who are not too exhausted to respond to the stimulus applied it is of extreme value. But to be of use it should be used intermittently and not continuously. Pouring a stream of cold water from a height on the abdomen is a not uncommon, but bad, practice, as it deluges the patient and the bedding in water, which may afterward act injuriously. Flapping the lower part of the abdomen with a wet towel is less objectionable. Ice can generally be obtained, and a piece should be introduced into the uterus. This is a very powerful hæmostatic, and often excites strong action when other means fail. I constantly employ it, and have never seen any bad results follow. A large piece of ice may also be held over the fundus, and removed and reapplied from time to time. Iced water may be injected into the rectum. A very powerful remedy is washing out the uterine cavity with a stream of cold water by means of a vaginal pipe of a Higginson's syringe carried up to the fundus. Another means of applying cold, said to be very effectual, is the application of the ether spray, such as is used for producing local anæsthesia, over the lower part of the abdomen.² All these remedies, however, depend for their good results on the fact of the patient being in a condition to respond to stimulus, and their prolonged use, if they fail to excite contraction rapidly, will certainly prove injurious. Rigby used to look upon the application of the child to the breast as one of the most certain inciters of uterine action. It may be of service, after the hemorrhage has been checked, in keeping up tonic contraction, and should therefore not be omitted; but we certainly cannot waste time in inducing the child to suck in the face of the actual emergency.

Intra-uterine Injections of Hot Water.—Of late, intra-uterine injections of hot water, at a temperature of from 100° to 120°, have been

¹ See an interesting paper by Dr. Thrush on "Retention of the Placenta in Labor at Term," *Amer. Journ. of Obstet.*, July, 1877.

² Griffiths, *Practitioner*, March, 1877.

highly recommended as a powerful means of arresting post-partum hemorrhage, often proving effectual when all other treatment has failed. The number of published cases in which it has proved of great value is now considerable. The present Master of the Rotunda, Dr. Lombe Atthill, has recorded 16 cases¹ in which it checked hemorrhage at once, in many of which ergot, ice, and other means had failed. He speaks of it as especially useful in those troublesome cases in which the uterus alternately relaxes and hardens, and resists all our efforts to produce permanent contraction. My own experience of this treatment is very favorable. I have now used it in several cases, in some of which the tendency to hemorrhage was very great, and in every instance it has at once produced strong uterine action and instantly checked the flow. It is, moreover, much more agreeable to the patient than cold applications. I think it cannot be doubted that we have in these hot irrigations a valuable addition to our methods of treating uterine hemorrhage.

State of the Bladder.—The late Dr. Earle pointed out² that a distended bladder often prevents contraction, and to avoid the possibility of this the catheter should be passed.

Plugging of the Vagina.—Plugging of the vagina has often been used. It is only necessary to mention it for the purpose of insisting on its absolute inapplicability in all cases of post-partum hemorrhage; the only effect it could have would be to prevent the escape of blood externally, which might then collect to any extent in the cavity of the uterus.

Compression of the Abdominal Aorta.—Compression of the abdominal aorta is highly thought of by many continental authorities, but it is little known or practised in this country. It has been objected to by some on the theoretical ground that the hemorrhage is chiefly venous, and not arterial, and that it would only favor the reflux of venous blood into the vena cava. Cazeaux points out that, on account of the close anatomical relations between the aorta and the vena cava, it is hardly possible to compress one vessel within the other. The backward flow of blood, therefore, through the vena cava may also be thus arrested. There is strong evidence in favor of the occasional utility of compression. Its chief recommendation is, that it can be practised immediately, and by an assistant, who can be shown how to apply the pressure. It is most likely to prove useful in sudden and severe hemorrhage, and if it only control the loss for a few moments it gives us time to apply other methods of treatment. As a temporary expedient, therefore, it should be borne in mind and adopted when necessary. It has the great advantage of supplementing, without superseding, other and more radical plans of treatment. The pressure is very easily applied on account of the lax state of the abdominal walls. The artery can readily be felt pulsating above the fundus uteri, and can be compressed against the vertebræ by three or four fingers applied lengthways. Baudelocque, who was a strong advocate of this procedure, states that he has on several occasions controlled an otherwise intractable hemorrhage in this way, and that he on one occasion kept up compression for four consecutive hours. Cazeaux believes that compression of the aorta may have a further advantageous effect in retaining the mass of the blood in the upper part of the body,

¹ *Lancet*, February 9, 1878.

² Earle's *Flooding after Delivery*, p. 163.

and thus lessening the tendency to syncope and collapse. If an aortic tourniquet, such as is used for compressing the vessel in cases of aneurism, could be obtained, it might be used with advantage in such cases.

Faradic Current.—If a battery is at hand, the faradic current may be used, and is, it is said, a very powerful agent in inducing uterine contraction, one pole being introduced into the uterus, the other applied over it through the abdominal parietes.

Bandaging of the Extremities.—When the hemorrhage has been excessive and there is profound exhaustion, firm bandaging of the extremities, by preference with Esmarch's elastic bandages if they can be obtained, may be advantageously adopted, with the view of retaining the blood as much as possible in the trunk, and thus lessening the tendency to syncope. As a temporary expedient in the worst class of cases it may occasionally prove of service.

[Lives of patients in *extremis* have been saved by the expedient of raising the body of the woman and lowering her head, so as to turn the current of blood toward the brain. This may have to be repeated several times in the treatment of a case where attacks of syncope indicate it. A bladder containing ice may be held under the hand of the operator over the abdomen and above the *fundus uteri*, and compression made upon the uterus and aorta at the same time. In one case I was forced, by the long-continued inertia of the uterus and the tendency to a return of hemorrhage, to keep up this form of compression for 6½ hours. The hand of the operator should be protected by a compress of flannel, or he may have an attack of local neuralgia, or possibly rheumatism, in his arm.—ED.]

Injection of Styptics.—Supposing these means fail, and the uterus obstinately refuses to contract in spite of all our efforts—and, do what we may, cases of this kind will occur—the only other agent at our command is the application of a powerful styptic to the bleeding surface to produce thrombosis in the vessels. "The latter," says Dr. Ferguson,¹ alluding to this means of arresting hemorrhage, "appears to be the sole means of safety in those cases of intense flooding in which the uterus flaps about the hand like a wet towel. Incapable of contraction for hours, yet ceasing to ooze out a drop of blood, there is nothing apparently between life and death but a few soft coagula plugging up the sinuses." These form but a frail barrier indeed, but the experience of all who have used the injection of a solution of perchloride of iron in such cases proves that they are thoroughly effectual, and its introduction into practice is one of the greatest improvements in modern midwifery. Although this method of treating these obstinate cases is not new, since it was practised long ago in Germany, its adoption in this country is unquestionably due to the energetic recommendation of Dr. Barnes. Although the dangers of the practice have been strongly insisted on, and with a degree of acrimony that is to be regretted, I know of only one published case in which its use has been followed by any evil effects. Its extraordinary power, however, of instantly checking the most formidable hemorrhage has been demonstrated by the unanimous testimony of all who have tried it. As it is not proposed by any one that this

¹ Preface to Gooch *On Diseases of Women*, p. xlii.

means of treatment should be employed until all ordinary methods of evoking contraction have failed, and as, in cases of this kind, the lives of the patients are of necessity imperilled, we should be fully justified in adopting it even if its possibly injurious effects had been much more certainly proved. It is surely at any time justifiable to avoid a great and pressing peril by running a possible chance of a less one. Whenever, therefore, we have tried the plans above indicated in vain, no time should be lost in resorting to this expedient. No practitioner should attend a case of midwifery without having the necessary styptic with him. The best and most easily obtainable form of using the remedy is the "liquor ferri perchloridi fortior" of the London Pharmacopeia, which should be diluted for use with six times its bulk of water. This is certainly better than a weaker solution. The vaginal pipe of a Higginson's syringe, through which the solution has once or twice been pumped to exclude the air, is guided by the hand to the fundus uteri and the fluid injected gently over the uterine surface. The loose and flabby mucous membrane is instantaneously felt to pucker up, all the blood with which the fluid comes in contact is coagulated, and the hemorrhage is immediately arrested. I think it is of importance to make sure that the uterus and vagina are emptied of clots before injection. In the only cases in which I have seen any bad symptoms follow this precaution had been neglected. The iron hardened all the coagula, which remained in utero, and septicæmia supervened; which, however, disappeared after the clots had been broken up and washed away by intra-uterine antiseptic injections. After we have resorted to this treatment all further pressure on the uterus should be stopped. We must remember that we have now abandoned contraction as a hæmostatic and are trusting to thrombosis, and that pressure might detach and lessen the coagula which are preventing the escape of blood.

Other local astringents may be eventually found to be of use. Tincture of matico possibly might be serviceable, although I am not aware that it has been tried. Dupierris has advocated tincture of iodine, and has recorded 24 cases in which he employed it—in all without accident and with a successful issue. Penrose strongly recommends common vinegar, which has the advantage of being always readily obtainable. But nothing seems likely to act so immediately or so effectually as the perchloride of iron.

Hemorrhage from Laceration of Maternal Structures.—A word may here be said as to the occasional dependence of hemorrhage after delivery on laceration of the cervix or other injury to the maternal soft parts. Duncan has narrated a case in which the bleeding came from a ruptured perineum. If hemorrhage continue after the uterus is permanently contracted, a careful examination should be made to ascertain if any such injury exist. Most generally the source of bleeding is the cervix, and the flow can be readily arrested by swabbing the injured textures with a sponge saturated in a solution of the perchloride.

Secondary Treatment.—The secondary treatment of post-partum hemorrhage is of importance. When reaction commences a train of distressing symptoms often show themselves, such as intense and throbbing headache, great intolerance of light and sound, and general nervous pros-

tration; and when these have passed away we have to deal with the more chronic effects of profuse loss of blood. Nothing is so valuable in relieving these symptoms as opium. It is the best restorative that can be employed, but it must be administered in larger doses than usual. Thirty to forty drops of Battley's solution should be given by the mouth or in an enema. At the same time the patient should be kept perfectly still and quiet in a darkened room and the visits of anxious friends strictly forbidden. Strong beef-essence or gravy soup, milk, or eggs beat up with milk, and similar easily-absorbed articles of diet should be given frequently and in small quantities at a time. Stimulants will be required according to the state of the patient, such as warm brandy and water, port wine, etc. Rest in bed should be insisted on, and continued much beyond the usual time. Eventually, the remedies which act by promoting the formation of blood, such as the various preparations of iron, will be found useful and may be required for a length of time.

Transfusion.—Under the head of Transfusion I have separately treated the application of that last resource in those desperate cases in which the loss of blood has been so excessive as to leave no other hope.

Secondary Post-partum Hemorrhage.—In the majority of cases, if a few hours have elapsed after delivery without hemorrhage, we may consider the patient safe from the accident. It is by no means very rare, however, to meet with even profuse losses of blood coming on in the course of convalescence at a time varying from a few hours or days up to several weeks after delivery. These cases are described as examples of "*secondary hemorrhage*," and they have not received at all an adequate amount of attention from obstetric writers, inasmuch as they often give rise to very serious, and even fatal, results, and are always somewhat obscure in their etiology and difficult to treat. We owe almost all our knowledge of this condition to an excellent paper by Dr. McClintock of Dublin, who has collected characteristic examples from the writings of various authors and accurately described the causes which are most apt to produce it.

Profuse Lochial Discharge.—We must, in the first place, distinguish between true secondary hemorrhage and profuse lochial discharge continued for a longer time than usual. The latter is not a very uncommon occurrence, and is generally met with in cases in which involution of the uterus has been checked, as by too early exertion, general debility, and the like. The amount of the lochial discharge varies in different women. In some patients it habitually continues during the whole puerperal month, and even longer, but not to an extent which justifies us in including it under the head of hemorrhage. In such cases prolonged rest, avoidance of the erect posture, occasional small doses of ergot, and, it may be, after the lapse of some weeks, astringent injections of oak-bark or alum, will be all that is necessary in the way of treatment.

True secondary hemorrhage is often sudden in its appearance and serious in its effects. McClintock mentions 6 fatal cases, and Mr. Bassett of Birmingham¹ has recorded 13 examples which came under his own observation, 2 of which ended fatally.

¹ *Brit. Med. Journ.*, 1872.

Causes.—The causes may be either constitutional or some local condition of the uterus itself.

Constitutional Causes.—Among the former are such as produce a disturbance of the vascular system of the body generally or of the uterine vessels in particular. The state of the uterine sinuses, and the slight barrier which the thrombi formed in them offer to the escape of blood, readily explain the fact of any sudden vascular congestion producing hemorrhage. Thus, mental emotions, the sudden assumption of the erect posture, any undue exertion, the incautious use of stimulants, a loaded condition of the bowels, or sexual intercourse shortly after delivery, may act in this way. McClintock records the case of a lady in whom very profuse hemorrhage occurred on the twelfth day after labor when sitting up for the first time. Feeling faint after sucking, the nurse gave her some brandy, whereupon a gush of blood ensued, “deluging all the bedclothes and penetrating through the mattress so as to form a pool on the floor.” Here the erect position, the exquisite pain caused by nursing, and the stimulating drink, all concurred to excite the hemorrhage. In another instance the flooding was traced to excitement produced by the sudden return of an old lover on the eighth day after labor. Moreau especially dwells on the influence of local congestion produced by a loaded condition of the rectum. Constitutional affections producing general debility and an impoverished state of the blood probably also may have the same effect. Blot specially mentions albuminuria as one of these, and Saboia states that in Brazil secondary hemorrhage is a common symptom of miasmatic poisoning, and can only be cured by change of air and the free use of quinine.¹

Local Causes.—Local conditions seem, however, to be the more frequent factors in the production of secondary hemorrhage. These may be generally classed under the following heads:

1. Irregular and inefficient contraction of the uterus.
2. Clots in the uterine cavity.
3. Portions of retained placenta or membranes.
4. Retroflexion of the uterus.
5. Laceration or inflammatory state of the cervix.
6. Thrombosis or hæmatocele of the cervix or vulva.
7. Inversion of the uterus.
8. Fibroid tumors or polypus of the uterus.

The first four of these need only now be considered, the others being described elsewhere.

Relaxation of, and Clots in, the Uterus.—Relaxation of the uterus and distension of its cavity by coagula may give rise to hemorrhage, although not so readily as immediately after delivery, for coagula of considerable size are often retained in utero for many days after labor. The uterus will be found larger than it ought to be, and tender on pressure. Usually the coagula are expelled with severe after-pains; but this may not take place, and hemorrhage may ensue several days after delivery. Or there may be only a relaxed state of the uterus without retained coagula. Bassett relates four cases traced to these causes, and several illustrations will be found in McClintock's paper. Portions of retained placenta or

¹ Saboia, *Traité des Accouchements*, p. 819.

membranes are more frequent causes. The retention may be due to carelessness on the part of the practitioner, especially if he have removed the placenta by traction and failed to satisfy himself of its integrity. It may, however, often be due to circumstances entirely beyond his control, such as adherent placenta, which it is impossible to remove without leaving portions in utero, or more rarely placenta succenturia. In the latter case there is a small supplementary portion of placental tissue developed entirely separate from the general mass, and it may remain in utero without the practitioner having the least suspicion of its existence. Portions of the membranes are very apt to be left in utero. It is to prevent this that they should be twisted into a rope and extracted very gently after expression of the placenta. Hemorrhage from these causes generally does not occur until at least a week after delivery, and it may not do so until a much longer time has elapsed. In four cases recorded by Mr. Bassett it commenced on the tenth, twelfth, fourteenth, and thirty-second day. It may come on suddenly and continue, or it may stop, and recur frequently at short intervals. In my experience retention of portions of the placenta is very common after abortion, when adhesions are more generally met with than at term. In addition to the hemorrhage there is often a fetid discharge due to decomposition of the retained portion, and possibly more or less marked septicæmic symptoms, which may aid in the diagnosis. The placenta or membranes may simply be lying loose as foreign bodies in the uterine cavity, or they may be organically attached to the uterine walls, when their removal will not be so easily effected.

Retroflexion.—Barnes has especially pointed out the influence of retroflexion of the uterus in producing secondary hemorrhage,¹ which seems to act by impeding the circulation at the point of flexion and thus arresting the process of involution.

In every case in which secondary hemorrhage occurs to any extent careful investigation into the possible causes of the attack, and an accurate vaginal examination, are imperatively required. If it be due to general and constitutional causes only, we must insist on the most absolute rest on a hard bed in a cool room and on the absence of all causes of excitement. The liquid extract of ergot will be very generally useful in ʒj doses repeated every six hours. McClintock strongly recommends the tincture of Indian hemp, which may be advantageously combined with the ergot, in doses of 10 or 15 minims, suspended in mucilage. Astringent vaginal pessaries of matico or perchloride of iron may be used. Special attention should be paid to the state of the bowels, and if the rectum be loaded it should be emptied by enemata. In more chronic cases a mixture of ergot, sulphate of iron, and small doses of sulphate of magnesia will prove very serviceable. This is more likely to be effectual when the bleeding is of an atonic and passive character. McClintock speaks strongly in favor of the application of a blister over the sacrum. When the hemorrhage is excessive more effectual local treatment will be required. Cazeaux advises plugging of the vagina. Although this cannot be considered so dangerous as immediately after delivery, inasmuch as the uterus is not so likely to dilate above the plug, still it is

¹ *Obstetric Operations*, p. 492.

certainly not entirely without risk of favoring concealed internal hemorrhage. If it be used at all, a firm abdominal pad should be applied, so as to compress the uterus; and the abdomen should be examined from time to time to ensure against the possibility of uterine dilatation. With these precautions the plug may prove of real value. In any case of really alarming hemorrhage I should be disposed rather to trust to the application of styptics to the uterine cavity. The injection of fluid in bulk, as after delivery, could not be safely practised, on account of the closure of the os and the contraction of the uterus. But there can be no objection to swabbing out the uterine cavity with a small piece of sponge attached to a handle and saturated in a solution of the perchloride of iron. There are few cases which will resist this treatment.

If we have reason to suspect retained placenta or membranes, or if the hemorrhage continue or recur after treatment, a careful exploration of the interior of the womb will be essential. On vaginal examination we may possibly feel a portion of the placenta protruding through the os, which can then be removed without difficulty. If the os be closed, it must be dilated with sponge or laminaria tents or by a small-sized Barnes's bag, and the uterus can then be thoroughly explored. This ought to be done under chloroform, as it cannot be effectually accomplished without introducing the whole hand into the vagina, which necessarily causes much pain. If the placenta or membranes be loose in the uterine cavity, they may be removed at once, or if they be organically attached, they may be carefully picked off. The uterus should at the same time, and as long as the os remains patulous, be thoroughly washed out with Condy's fluid and water to diminish the risk of septicæmia.

Retroflexion can readily be detected by vaginal examination and the treatment consists in careful reposition with the hand and the application of a large-sized Hodge's pessary.

CHAPTER XVI.

RUPTURE OF THE UTERUS, ETC.

Its Fatality.—Rupture of the uterus is one of the most dangerous accidents of labor, and until of late years it has been considered almost necessarily fatal and beyond the reach of treatment. Fortunately, it is not of very frequent occurrence, although the published statistics vary so much that it is by no means easy to arrive at any conclusion on this point. The explanation is, no doubt, that many of the tables confound partial and comparatively unimportant lacerations of the cervix and vagina with rupture of the body and fundus. It is only in large lying-in institutions, where the results of cases are accurately recorded, that anything like reliable statistics can be gathered, for in private practice

the occurrence of so lamentable an accident is likely to remain unpublished. To show the difference between the figures given by authorities, it may be stated that while Burns calculates the proportion to be 1 in 940 labors, Ingleby fixes it as 1 in 1300 or 1400, Churchill as 1 in 1331, and Lehmann as 1 in 2433. Dr. Jolly of Paris has published an excellent thesis containing much valuable information.¹ He finds that out of 782,741 labors, 230 ruptures, excluding those of the vagina or cervix, occurred—that is, 1 in 3403.

Seat of Rupture.—Lacerations may occur in any part of the uterus—the fundus, the body, or the cervix. Those of the cervix are comparatively of small consequence, and occur to a slight extent in almost all first labors. Only those which involve the supra-vaginal portion are of really serious import. Ruptures of the upper part of the uterus are much less frequent than of the portion near the cervix—partly, no doubt, because the fundus is beyond the reach of the mechanical causes to which the accident can, not unfrequently, be traced, and partly because the lower third of the organ is apt to be compressed between the presenting part and the bony pelvis. The site of placental insertion is said by Madame La Chapelle to be rarely involved in the rupture, but it does not always escape, as numerous recorded cases prove. The most frequent seat of rupture is near the junction of the body and neck, either anteriorly or posteriorly, opposite the sacrum, or behind the symphysis pubis, but it may occur at the sides of the lower segment of the uterus. In some cases the entire cervix has been torn away and separated in the form of a ring.

Rupture may be Partial or Complete.—The laceration may be partial or complete, the latter being the more common. The muscular tissue alone may be torn, the peritoneal coat remaining intact; or the converse may occur, and then the peritoneum is often fissured in various directions, the muscular coat being unimplicated. The extent of the injury is very variable, in some cases being only a slight tear, in others forming a large aperture, sufficiently extensive to allow the fetus to pass into the abdominal cavity. The direction of the laceration is as variable as the size, but it is more frequently vertical than transverse or oblique. The edges of the tear are irregular and jagged, probably on account of the contraction of the muscular fibres, which are frequently softened, infiltrated with blood, and even gangrenous. Large quantities of extravasated blood will be found in the peritoneal cavity, such hemorrhage, indeed, being one of the most important sources of danger.

The causes are divided into *predisposing* and *exciting*; and the progress of modern research tends more and more to the conclusion that the cause which leads to the laceration could only have operated because the tissue of the uterus was in a state predisposed to rupture, and that it would have had no such effect on a perfectly healthy organ. What these predisposing changes are, and how they operate, is yet far from being known, and the subject offers a fruitful field for pathological investigation.

Said to be more Common in Multiparæ.—It is generally believed that lacerations are more common in multiparæ than in primiparæ. Tyler

¹ *Rupture utérine pendant le Travail*, Paris, 1873.

Smith contended that ruptures are relatively as common in first as in subsequent labors, while Bandl¹ found that only 64 cases out of 546 ruptures were in primiparæ. Statistics are not sufficiently accurate or extensive to justify a positive conclusion, but it is reasonable to suppose that the pathological changes presently to be mentioned as predisposing to laceration are more likely to be met with in women whose uteri have frequently undergone the alteration attendant on repeated pregnancies. Age seems to have considerable influence, as a large proportion of cases have occurred in women between thirty and forty years of age.

Alterations in the tissues of the uterus are probably of very great importance in predisposing to the accident, although our information on this point is far from accurate. Among these are morbid states of the muscular fibres, the result of blows and contusions during pregnancy; premature fatty degeneration of the muscular tissues—an anticipation, as it were, of the normal involution after delivery; fibroid tumors or malignant infiltration of the uterine walls, which either produce a morbid state of the tissues or act as an impediment to the expulsion of the fœtus. The importance of such changes has been specially dwelt on by Murphy in this country and by Lehmann in Germany, and it is impossible not to concede their probable influence in favoring laceration. However, as yet these views are founded more on reasonable hypothesis than on accurately-observed pathological facts.

Another and very important class of predisposing causes are those which lead to a want of proper proportion between the pelvis and the fœtus.

Deformity in Pelvis is a Frequent Cause.—Deformity of the pelvis has been very frequently met with in cases in which the uterus has ruptured. Thus, out of 19 cases carefully recorded by Radford,² the pelvis was contracted in 11, or more than one-half. Radford makes the curious observation that ruptures seem more likely to occur when the deformity is only slight; and he explains this by supposing that in slight deformities the lower segment of the uterus engages in the brim, and is therefore much subjected to compression, while in extreme deformity the os and cervix uteri remain above the brim, the body and fundus of the uterus hanging down between the thighs of the mother. This explanation is reasonable, but the rarity with which ruptured uterus is associated with extreme pelvic deformity may rather depend on the infrequency of advanced degrees of contraction.


Views of Bandl.—Bandl, who has made the most important of modern contributions to our knowledge of the subject, points out that rupture nearly always begins in the lower segment of the uterus, which becomes abnormally stretched and distended when from any cause the expulsion of the fœtus is delayed. The upper portion of the uterus becomes at the same time retracted and much thickened (see Fig. 146). As the pains continue the stretching of the lower segment becomes more and more marked, until at last its fibres separate and a laceration is established. The line of demarcation between the thickened body and the distended lower segment, known as the ring of Bandl, can in such cases be occasionally made out by palpation above the pubes.

¹ *Ueber Ruptur der Gebärmutter*, Wien, 1815.

² *Obst. Trans.*, vol. viii.

Malpresentation or Undue Size of the Fœtus.—Amongst causes of disproportion depending on the fœtus is either malpresentation, in which the pains cannot effect expulsion, or undue size of the presenting part. In the latter way may be explained the observation that rupture is more

FIG. 146.



Illustrating the Dangerous Thinning of the Lower Segment of Uterus, owing to Non-descent of Head in a Case of Intra-uterine Hydrocephalus. (After Bandl.)

frequently met with in the delivery of male than of female children, on account, no doubt, of the larger size of the head in the former. The influence of intra-uterine hydrocephalus was first prominently pointed out by Sir James Simpson,¹ who states that out of 74 cases of intra-uterine hydrocephalus, the uterus ruptured in 16. In all such cases of disproportion, whether referable to the pelvis or fœtus, rupture is produced in a twofold manner—either by the excessive and fruitless uterine contractions, which are induced by the efforts of the organ to overcome the obstacle, or by the compression of the uterine tissue between the presenting part and the bony pelvis, leading to inflammation, softening, and even gangrene.

Mechanical Injury.—The proximate cause of rupture may be classed under two heads—mechanical injury and excessive uterine contraction. Under the former are placed those uncommon cases in which the uterus lacerates as the result of some injury in the latter months of pregnancy, such as blows, falls, and the like. Not so rare, unfortunately, are lacerations produced by unskilled attempts at delivery on the part of the

¹ *Selected Obst. Works*, p. 385.

medical attendant, such as by the hand during turning or by the blades of the forceps. Many such cases are on record in which the accoucheur has used force and violence rather than skill in his attempts to overcome an obstacle. That such unhappy results of ignorance are not so uncommon as they ought to be is proved by the figures of Jolly, who has collected 71 cases of rupture during podalic version, 37 caused by the forceps, 10 by the cephalotribe, and 30 during other operations, the precise nature of which is not stated.¹ The *modus operandi* of protracted and ineffectual uterine contractions, as a proximate cause of rupture, is sufficiently evident, and need not be dwelt on. It is necessary to allude, however, to the effect of ergot, incautiously administered, as a producing cause. There is abundant evidence that the injudicious exhibition of this drug has often been followed by laceration of the unduly-stimulated uterine fibres. Thus, Trask, talking of the subject, says that Meigs had seen three cases, and Bedford four, distinctly traceable to this cause. Jolly found that ergot had been administered largely in 33 cases in which rupture occurred.

Premonitory Symptoms.—Some have believed that the impending occurrence of rupture could frequently be ascertained by peculiar premonitory symptoms, such as excessive and acute crampy pains about the lower part of the abdomen, due to the compression of part of the uterine walls. These are far too indefinite to be relied on, and it is certain that the rupture generally takes place without any symptoms that would have afforded reasonable grounds for suspicion.

General Symptoms.—The symptoms are often so distinct and alarming as to leave no doubt as to the nature of the case. Not infrequently, however, especially if the laceration be partial, they are by no means so well marked, and the practitioner may be uncertain as to what has taken place. In the former class of cases a sudden excruciating pain is experienced in the abdomen, generally during the uterine contractions, accompanied by a feeling on the part of the patient of something having given way. In some cases this has been accompanied by an audible sound, which has been noticed by the bystanders. At the same time there is generally a considerable escape of blood from the vagina, and a prominent symptom is the sudden cessation of the previously strong pains. Alarming general symptoms soon develop, partly due to shock, partly to loss of blood, both external and internal. The face exhibits the greatest suffering, the skin becomes deadly cold and covered with a clammy sweat, and fainting, collapse, rapid feeble pulse, hurried breathing, vomiting, and all the usual signs of extreme exhaustion quickly follow.

Results of Abdominal and Vaginal Examinations.—Abdominal palpation and vaginal examination both afford characteristic indications in well-marked cases. If the child, as often happens, has escaped entirely or in great part into the abdominal cavity, it may be readily felt through the abdominal walls; while in the former case the partially-contracted uterus may be found separate from it in the form of a globular tumor resembling the uterus after delivery. Per vaginam it may generally be ascertained that the presenting part has suddenly receded

¹ *Op. cit.*, p. 38.

and can no longer be made out, or some other part of the fœtus may be found in its place. If the rupture be extensive, it may be appreciable on vaginal examination, and sometimes a loop of intestine may be found protruding through the tear. Other occasional signs have been recorded, such as an emphysematous state of the lower part of the abdomen, resulting from the entrance of air into the cellular tissue, or the formation of a sanguineous tumor in the hypogastrium or vagina. These are too uncommon and too vague to be of much diagnostic value.

Symptoms are Sometimes Obscure.—Unfortunately, the symptoms are by no means always so distinct, and cases occur in which most of the reliable indications, such as the sudden cessation of the pains, the external hemorrhage, and the retrocession of the presenting part, may be absent. In some cases, indeed, the symptoms have been so obscure that the real nature of the case has only been detected after death. It is rarely, however, that the occurrence of shock and prostration is not sufficiently distinct to arouse suspicion, even in the absence of the usual marked signs. In not a few cases distinct and regular contractions have gone on after laceration, and the child has even been born in the usual way. Of course in such a case mistake is very possible. So curious a circumstance is difficult of explanation. The most probable way of accounting for it is that the laceration has not implicated the fundus of the uterus, which contracted sufficiently energetically to expel the fœtus. Hence it will be seen that the symptoms are occasionally obscure, and the practitioner must be careful not to overlook the occurrence of so serious an accident because of the absence of the usual and characteristic symptoms.

Prognosis.—The prognosis is necessarily of the gravest possible character, but modern views as to treatment perhaps justify us in saying that it is not so absolutely hopeless as has been generally taught in our obstetric works.

To the Mother.—When we reflect on what has occurred—the profound nervous shock; the profuse hemorrhage, both external, and especially into the peritoneal cavity, where the blood coagulates and forms a foreign body; the passage of the uterine contents into the abdomen, with the inevitable result of inflammation and its consequences if the patient survive the primary shock,—the enormous fatality need cause no surprise. Jolly has found that out of 580 cases 100 recovered—that is, in the proportion of 1 out of 6. This is a far more favorable result than we are generally led to anticipate; and as many of the recoveries happened in apparently the most desperate and unfavorable cases, we should learn the lesson that we need not abandon all hope, and should at least endeavor to rescue the patient from the terrible dangers to which she is exposed.

To the Child.—As regards the child, the prognosis is almost necessarily fatal; and, indeed, the cessation of the fœtal heart-sounds has been pointed out by McClintock as a sign of rupture in doubtful cases. The shock, the profuse hemorrhage, and the time that must necessarily elapse before the delivery of the child are of themselves quite sufficient to explain the fact that the fœtus is almost always dead.

Treatment.—From what has been said of the impossibility of fore-

telling the occurrence of rupture, it must follow that no reliable prophylactic treatment can be adopted beyond that which is a matter of general obstetric principle—viz. timely interference when the uterine contractions seem incapable of overcoming an obstacle to delivery, either on the part of the pelvis or fœtus.

Indications after Rupture has Taken Place.—After rupture the main indications are to effect the removal of the child and the placenta, to rally the patient from the effects of the shock, and, if she survives so long, to combat the subsequent inflammation and its consequences. By far the most important point to decide is the best means to be adopted for the removal of the child; for it is admitted by all that the hopeless expectancy that was recommended by the older accoucheurs—or, in other words, allowing the patient to die without making any effort to save her—is quite inadmissible. If the fœtus be entirely within the uterine cavity, no doubt the proper course to pursue is to deliver at once *per vias naturales*, either by turning, by forceps, or by cephalotripsy. If any part other than the head present, turning will be best, great care being taken to avoid further increase of the laceration. If the head be in the cavity or at the brim of the pelvis, and within easy reach of the forceps, it may be cautiously applied, the child being steadied by abdominal pressure so as to facilitate its application. If there be, as is often the case, some slight amount of pelvic contraction, it may be preferable to perforate and apply the cephalotribe, so as to avoid any forcible attempts at extraction, which might unduly exhaust the already prostrate patient and turn the scale against her. This will be the more allowable since the child is, as we have seen, almost always dead, and we might readily ascertain if it be so by auscultation.

Removal of the Placenta.—After delivery extreme care must be taken in removing the placenta, and for this it will be necessary to introduce the hand. The placenta will generally be in the uterus, for if the rent be not large enough for the child to pass through, it may be inferred that the placenta will not have done so either. If it has escaped from the uterus, very gentle traction on the cord may bring it within reach of the hand, and so the passage of the hand through the tear to search for it will be avoided.

Treatment when the Fœtus has Escaped out of the Uterus.—There can be but little doubt that in the cases indicated such is the proper treatment and that which affords the mother the best chance. Unfortunately, the cases in which the child remains entirely in utero are comparatively uncommon, and generally it will have escaped into the abdomen, along with much extravasated blood. The usual plan of treatment recommended under such circumstances is to pass the hand through the fissure (some have even recommended that it should be enlarged by incision if necessary), to seize the feet of the fœtus, to drag it back through the torn uterus, and then to reintroduce the hand to search for and remove the placenta. Imagine what occurs during the process! The hand gropes blindly among the abdominal viscera, the forcible dragging back of the fœtus necessarily tears the uterus more and more, and, above all, the extravasated blood remains as a foreign body in the peritoneal cavity, and necessarily gives rise to the most serious conse-

quences. It is surely hardly a matter of surprise that there is scarcely a single case on record of recovery after this procedure.

Reasons why Gastrotomy affords a Better Chance of Success.—Of late years a strong feeling has existed that whenever the child has entirely or in great part escaped into the abdominal cavity the operation of gastrotomy affords the mother a far better chance of recovery; and it has now been performed in many cases with the most encouraging results. It is easy to see why the prospects of success are greater. The uterus being already torn and the peritoneum opened, the only additional danger is the incision of the abdominal parietes, which gives us the opportunity of sponging out the peritoneal cavity, as in ovariectomy, and of removing all the extravasated blood, the retention of which so seriously adds to the dangers of the case. Another advantage is that, if the patient be excessively prostrate, the operation may be delayed until she has somewhat rallied from the effects of the shock, whereas delivery by the feet is generally resorted to as soon as the rupture is recognized and when the patient is in the worst possible condition for interference of any kind.

Comparative Results of Various Methods of Treatment.—Jolly has carefully tabulated the results of the various methods of treatment, and, making every allowance for the unavoidable errors of statistics, it seems beyond all question that the results of gastrotomy are so greatly superior to those of other plans that I think its adoption may fairly be laid down as a rule whenever the fœtus is no longer within the uterine cavity:

COMPARATIVE RESULTS OF VARIOUS METHODS OF TREATMENT AFTER RUPTURE OF UTERUS.

Treatment.	No. of Cases.	Deaths.	Recoveries.	Per cent. of Recoveries.
Expectation	144	142	2	1.45
Extraction <i>per vias naturales</i>	382	310	72	19
Gastrotomy	38	12	26	68.4

Of course this table will not justify the conclusion that 68 per cent. of the cases of ruptured uterus in which gastrotomy is performed will recover; but it may fairly be taken as proving that the chances of recovery are at least three or four times as great as when the more usual practice is adopted.

[*American Puerperal Laparotomies.*—After a search of several years I have thus far collected 43 cases in the United States, with 21 women and 2 children saved. One mother and child were saved by an immediate operation with a pocket-knife in 1869. I presume that a general record of American operations, published and unpublished, would show a saving of nearly 50 per cent., which is much lower than that claimed by Trask and Jolly, collected from published reports. Take Trask's foreign cases, 20, and our own 43, and we have, native and foreign, 63, with 37 recoveries and 26 deaths. I look upon our own statistics as much more reliable, because many of the unpublished cases were searched out by correspondence.—ED.]

Lacerations of the Cervix.—Lacerations of the cervix are of very common occurrence. Occasionally, after delivery, they may cause hemorrhage when the uterus itself is firmly contracted, or secondary hemorrhage during the puerperal month. As a rule, they are not recognized, and it is only of late years, chiefly owing to the labors of Emmet, that this important influence in producing various chronic forms of uterine disease has been realized. In the large majority of cases the lacerations are lateral, either on one or both sides of the cervix. If they give rise to hemorrhages, the local application of styptics is probably the best resource. Whether it is advisable to treat severe forms by the immediate application of silver sutures, as recommended by Pallen,¹ is a subject as yet too little understood to justify the expression of an opinion.

Necessity of Care in Performing the Operation.—It is perhaps needless to say that the operation must be performed with the same minute care that has raised ovariectomy to its present pitch of perfection, and that especial attention should be paid to the sponging out of the peritoneum, and the removal of foreign matters.

Recapitulation.—To recapitulate, I think what has been said justifies the following rules of treatment after rupture:

1. If the head or presenting part be above the brim and the fœtus still in utero—forceps, turning, or cephalotripsy, according to circumstances.

2. If the head be in the pelvic cavity—forceps or cephalotripsy.

3. If the fœtus have wholly or in great part escaped into the abdominal cavity—gastrotomy.

Subsequent Treatment.—As to the subsequent treatment little need be said, since in this we must be guided by general principles. The chief indication will be to remove shock and rally the patient by stimulants, etc., and to combat secondary results by opiates and other appropriate remedies.

Lacerations of the vagina occasionally take place, and in the great majority of cases they are produced by instruments, either from a want of care in their introduction or from undue stretching of the vaginal walls during extraction with the forceps. Slight vaginal lacerations are probably much more common after forceps delivery than is generally believed to be the case. As a rule, they are productive of no permanent injury, although it must not be forgotten that every breach of continuity increases the risk of subsequent septic absorption. When the laceration is sufficiently deep to tear through the recto-vaginal septum or the anterior vaginal wall, the passage of the urine or feces is apt to prevent its edges uniting; then that most distressing condition, recto-vaginal or vesico-vaginal fistula, is established.

Fistulæ are Seldom Caused by Mechanical Injury.—It must not be supposed that fistulæ are often the result of injury during operative interference. That is a common but very erroneous opinion both among the profession and the public. In the vast majority of cases the fistulous opening is the consequence of a slough resulting from inflammation, produced by long-continued pressure of the vaginal walls between the child's head and the bony pelvis in cases in which the second stage has

¹ *Transactions of the Intern. Med. Congr.*, v. 4.

been allowed to go on too long. In most of these cases instruments were doubtless eventually used, and they get the blame of the accident; whereas the fault lay, not in their being employed, but rather in their not having been used soon enough to prevent the contusion and inflammation which ended in sloughing.

Proof of this Statement.—When vesico-vaginal fistulæ are the result of lacerations during labor, the urine must escape at once, but this is rarely the case. In the large majority of cases the urine does not pass per vaginam until more than a week after delivery, showing that a lapse of time is necessary for inflammatory action to lead to sloughing. In order to throw some light on these points, on which very erroneous views have been held, I have carefully examined the histories, from various sources, of 63 cases of vesico-vaginal fistula.

1st. In 20 no instruments were employed. Of these, there were in labor—

Under 24 hours	2
From 24 to 48 hours	8 ¹
“ 48 to 70 “	2
“ 70 to 80 “	7
“ 80 hours and upward	1
	<hr/> 20

Therefore, out of these 20 cases, one-half were certainly more than 48 hours in labor, and 6 of the remaining 10 were probably so also. In only 1 of them is the urine stated to have escaped per vaginam immediately after delivery. In 7 it is said to have done so within a week, and in the remainder after the seventh day.

2d. In 34 cases instruments were used, but there is no evidence of their having produced the accident. Of these, there were in labor—

Under 24 hours	2
From 24 to 48 hours	8
“ 48 to 72 “	10
“ 72 hours and upward	14
	<hr/> 34

The urine escaped within 24 hours in 2 cases only, within a week in 16, and after the seventh day in 15. So that here, again, we have the history of unduly-protracted delivery, 24 out of the 34 having been certainly more than 48 hours in labor.

3d. In 9 cases the histories show that the production of the fistula may fairly be ascribed to the unskilled use of instruments. Of these, there were in labor—

Under 24 hours	7
From 24 to 48 hours	1
“ 48 to 72 “	1
	<hr/> 9

The urine escaped at once in 7 cases, and in the remaining 2 after the seventh day.

¹ But of these, in 7 no precise time is stated. 6 of them are marked *very tedious*, therefore they probably exceeded the limit.

These statistics seem to me to prove, in the clearest manner, that in the large majority of cases this unhappy accident may be directly traced to the bad practice of allowing labor to drag on many hours in the second stage without assistance, and not to premature instrumental interference. This question has recently been elaborately studied by Emmet, who gives numerous statistical tables which fully corroborate these views. His conclusion, the result of much practical experience of vesico-vaginal fistulæ, is worthy of being quoted. "I do not hesitate," he says, "to make the statement that I have never met with a case of vesico-vaginal fistula which without doubt could be shown to have resulted from instrumental delivery. On the contrary, the entire weight of evidence is conclusive in showing that the injury is a consequence of delay in delivery."¹

Treatment.—As to the treatment of vaginal laceration, little can be said. In the slighter cases vaginal injections of diluted Condyl's fluid will be useful to lessen the risk of septic absorption; and the graver, when vesico-vaginal or recto-vaginal fistulæ have actually formed, are not within the domain of the obstetrician, but must be treated surgically at some future date.

[*The Rational Treatment of Rupture of the Uterus.*—The three rules given on page 439 are those found in obstetrical works of high authority, but are not based upon the teachings of abdominal surgery as shown by the results of operations recorded within a few years. Reasoning from analogy and the fearful mortality of cases delivered *per vias naturales* after uterine rupture, we are forced to the conclusion that something more is needed than the delivery of the woman and the removal of the placenta if we hope to reduce the proportion of deaths, which is very great except after laparotomy—a method of delivery capable of saving nearly 50 per cent. There is no objection to delivering the fœtus by the natural channel, provided it can be readily done; but we have very little reason to anticipate a favorable result if we rest our efforts here. Children entirely escaped into the abdominal cavity have been drawn back through the rent and delivered by the vagina, and the women have recovered. In one well-authenticated case the woman was thus saved in our own country on four occasions. But we are not to expect such results, as a fatal issue is far more frequent than a recovery under such circumstances. Our object should be to save the life of the mother and, if at all possible, that of the fœtus; and all our efforts should be directed to this end. We may console ourselves with having delivered the woman prior to her death, but to prevent this fatal issue should be our chief aim. The general impression among ovariologists is, that blood is not an innocent fluid in the abdominal cavity; and the remarkable results of the operations of Dr. Keith of Edinburgh are attributed to the care he exercises in preventing the secondary escape of blood into the abdominal cavity. Dr. Ludwig Winckel of Mülheim, Germany, who performed the Cæsarean operation 13 times and laparotomy after rupture of the uterus 4 times, was of the impression that the liquor amnii was innocuous if only a short time in contact with the peritoneum; and the same may be said of blood, ovarian fluid, parova-

¹ *The Principles and Practice of Gynecology*, p. 669.

rian fluid, and, to some degree, also of urine. Rupture of the bladder is now cured by sewing up the rent and carefully cleansing the abdominal cavity of blood and urine. But these fluids are all capable of setting up peritonitis, and blood by its decomposition is particularly apt to give rise to septic poisoning: then why let it remain in the abdominal cavity in cases of ruptured uterus? If it is important to cleanse this cavity from blood and ovarian fluid in ovariectomy, and from blood and amniotic fluid after the Cæsarean section, then why should we be content with delivering the fœtus in cases of rupture of the uterus, when we know that the peritoneal cavity still contains a compound fluid which may destroy the woman if not removed and the parts cleansed? We have also an additional risk in the fact that the uterine rupture may gape and allow the lochia to escape into the peritoneal cavity, thus providing another element for septic poisoning. I am, then, fully persuaded that in all cases of rupture, where it is evident that blood and liquor amnii have escaped into the abdominal cavity, we ought to open the abdomen, cleanse out the cavity, and close the rent by deep-seated and superficial sutures of silver wire or carbolized pure silk. In cervico-vaginal rupture the closure of the rent may not be so important in the sense of safety to the woman, as there is generally a natural drainage into the vagina; neither is laparotomy itself so imperatively demanded as in cases where the fundus or body of the uterus is rent. But it becomes important to close the rent cervix in view of future trouble from ectropium and erosion. As in the Cæsarean operation, promptness of action is all important if we hope to save the patient. I know that these views upon the treatment of ruptured uterus are in advance of those held by British obstetrical writers, but they are certainly logical deductions from the experience of such operators as Dr. Keith, Mr. Lawson Tait, and others, and from the well-known results of promptly-performed laparotomies in rupture accidents in the United States. The removal of the uterus after rupture has as yet only added to the risk, and I do not believe we are justified in resorting to it where there is no pelvic obstruction.—ED.]

CHAPTER XVII.

INVERSION OF THE UTERUS.

Its Formidable Character.—Inversion of the uterus shortly after the birth of the child is one of the most formidable accidents of parturition, leading to symptoms of the greatest urgency, not rarely proving fatal, and requiring prompt and skilful treatment. Hence it has obtained an unusual amount of attention, and there are few obstetric subjects which have been more carefully studied.

An Accident of Great Rarity.—Fortunately, the accident is of great

rarity. It was only observed once in upward of 190,800 deliveries at the Rotunda Hospital since its foundation in 1745, and many practitioners have conducted large midwifery practices for a lifetime without ever having witnessed a case. It is none the less needful, however, that we should be thoroughly acquainted with its natural history and with the best means of dealing with the emergency when it arises.

Division into Acute and Chronic Forms.—Inversion of the uterus may be met with in the acute or chronic form; that is to say, it may come under observation either immediately or shortly after its occurrence, or not until after a considerable lapse of time, when the involution following pregnancy has been completed. The latter falls more properly under the province of the gynaecologist, and involves the consideration of many points that would be out of place in a work on obstetrics. Here, therefore, the acute form alone is considered.

Description of Inversion.—Inversion consists essentially in the enlarged and empty uterus being turned inside out, either partially or entirely; and this may occur in various degrees, three of which are usually described and are practically useful to bear in mind. In the first and slightest degree there is merely a cup-shaped depression of the fundus (Fig. 147); in the second the depression is greater, so that the inverted portion forms an intro-susception, as it were, and projects downward through the os in the form of a round ball, not unlike the body of a polypus, for which, indeed, a careless observer might mistake it; and, thirdly, there is the complete variety, in which the whole organ is turned inside out, and may even project beyond the vulva.

Its Symptoms.—The symptoms are generally very characteristic, although when the amount of inversion is small they may entirely escape observation. They are chiefly those of profound nervous shock—viz. fainting, small, rapid, and feeble pulse, possibly convulsions and vomiting, and a cold clammy skin. Occasionally severe abdominal pain and cramp and bearing-down are felt. Hemorrhage is a frequent accompaniment, sometimes to a very alarming extent, especially if the placenta be partially or entirely detached. The loss of blood depends to a great extent on the condition of the uterine parietes. If there be much contraction on the part that is not inverted, the intro-suscepted part may be sufficiently compressed to prevent any great loss. If the entire organ be in a state of relaxation, the loss may be excessive.

Results of Physical Examination.—The occurrence of such symptoms shortly after delivery would of necessity lead to an accurate examination, when the nature of the case may be at once ascertained. On passing the finger into the vagina we either find the entire uterus forming a glob-

FIG. 147.



Partial Inversion of the Fundus.
(From a preparation in the Museum of
Guy's Hospital.)

ular mass, to which the placenta is often attached, or, if the inversion be incomplete, the vagina is occupied by a firm, round, and tender swelling which can be traced upward through the os uteri. The hand placed on the abdomen will detect the absence of the round ball of the contracted uterus, and bi-manual examination may even enable us to feel the cup-shaped depression at the site of inversion.

Differential Diagnosis.—When such signs are observed immediately after delivery mistake is hardly possible. Numerous instances, however, are recorded in which the existence of inversion was not immediately detected, and the tumor formed by it only observed after the lapse of several days, or even longer, when the general symptoms led to vaginal examination. It is probable that in such cases a partial inversion had taken place shortly after delivery, which as time elapsed became gradually converted into the more complete variety. In a case of this kind, as in a chronic inversion, some care is necessary to distinguish the inversion from a uterine polypus, which it closely resembles. The cautious insertion of the sound will render the diagnosis certain, since its passage is soon arrested in inversion, while if the tumor be polypoid it readily passes in as far as the fundus.

Manner in which Inversion is Produced.—The mechanism by which inversion is produced is well worthy of study, and has given rise to much difference of opinion.

Occasionally Produced by Accidental Mechanical Causes.—A very general theory is that it is caused in many cases by mismanagement of the third stage of labor, either by traction on the cord, the placenta being still adherent, or by improperly-applied pressure on the fundus; the result of both these errors being a cup-shaped depression of the fundus, which is subsequently converted into a more complete variety of inversion. That such causes may suffice to start the inversion cannot be doubted, but it is probable that their frequency has been much exaggerated. Still, there are numerous recorded cases in which the commencement of the inversion can be traced to them. Improperly-applied pressure (as when the whole body of the uterus is not grasped in the hollow of the hand, but when a monthly nurse or other uninstructed person presses on the lower part of the abdomen so as simply to push down the uterus *en masse*) is often mentioned in histories of the accident. Thus, in the *Edinburgh Medical Journal* for June, 1848, a case is related in which the patient would not have a medical man, but was attended by a midwife, who after the birth of the child pulled on the cord, while the patient herself clasped her hands and pushed down her abdomen, at the same time straining forcibly, when the uterus became inverted and the patient died of hemorrhage before assistance could be procured. Here both of the mechanical causes mentioned were in operation. In several cases it is mentioned that the accident occurred while the nurse was compressing the abdomen. That the accident is practically impossible when firm and equable contraction has taken place cannot be questioned. Hence it is of paramount importance that the practitioner should himself carefully attend to the conduct of the third stage of labor.

Often Occurs Spontaneously.—In a large proportion of cases no mechanical causes can be traced, and the occurrence of spontaneous

inversion must be admitted. There are various theories held as to how this occurs. Partial and irregular contraction of the uterus is generally admitted to be an important factor in its production; but it is still a matter of dispute whether the inversion is produced mainly by an active contraction of the fundus and body of the uterus, the lower portion and cervix being in a state of relaxation, or whether the precise reverse of this exists, the fundus being relaxed and in a state of quasi-paralysis, while the cervix and lower portion of the uterus are irregularly contracted. The former is the view maintained by Radford and Tyler Smith, while the latter is upheld by Matthews Duncan.

Evidence in Favor of Duncan's Theory.—There are good clinical reasons for believing that Duncan's view more nearly corresponds with the true facts of the case; for if the fundus and body of the uterus be really in a state of active contraction while the cervix is relaxed, we have, as Duncan points out, the very condition which is normal and desirable after delivery, and that which we do our best to produce. If, however, the opposite condition exist and the fundus be relaxed, while the lower portion is spasmodically contracted, a state exists closely allied to the so-called hour-glass contraction. Supposing now any cause produces a partial depression of the fundus, it is easy to understand how it may be grasped by the contracted portion and carried more and more down, in the manner of an intro-susception, until complete inversion results. That such partial paralysis of the uterine walls often exists, especially about the placental site, was long ago pointed out by Rokitsansky and other pathologists. This theory supposes the original partial depression and relaxation of the fundus. How this is often produced by mismanagement of the third stage has already been pointed out; but, even in the absence of such causes, it may result from strong bearing-down efforts on the part of the patient, or, as Duncan holds, from the absence of the retentive power of the abdomen. Indeed, the incompatibility of an actively-contracted state of the fundus with the partial depression which is essential, according to both views, for the production of inversion, is the strongest argument in favor of Duncan's theory.


Taylor's Theory.—A totally different view has more recently been sustained by Dr. Taylor of New York, who maintains that "spontaneous active inversion of the uterus rests upon prolonged natural and energetic action of the body and fundus: the cervix, the lower part, yielding first, is thus rolled out or everted or doubled up, as there is no obstruction from the contractility of the cervix, which is at rest or functionally paralyzed; the body is gradually, sometimes instantaneously, forced lower and lower, or inverted."¹ That partial inversion may commence at the cervix was pointed out by Duncan in his paper, who depicts it in the accompanying diagram (Fig. 148), and states it to be of not unfrequent occurrence. It is not impossible that occasionally such a state of things should be carried on to complete inversion. But there are serious objections to the acceptance of Dr. Taylor's view that such is the principal cause of inversion, since the process above described would be of necessity a slow and long-continued one, whereas nothing is more certain than that inversion is generally sudden and accompanied by acute symp-

¹ *New York Med. Journ.*, 1872.

toms of shock, and is often attended by severe hemorrhage, which could not occur when such excessive contraction was taking place.

Treatment.—The treatment of inversion consists in restoring the organ to its natural condition as soon as possible. Every moment's delay only

FIG. 148.



Illustrating the Commencement of Inversion at the Cervix. (After Duncan.)

serves to render restoration more difficult, as the inverted portion becomes swollen and strangulated; whereas if the attempt at reposition be made immediately there is generally comparatively little difficulty in effecting it. Therefore it is of the utmost importance that no time should be lost and that we should not overlook a partial or incomplete inversion. Hence the occurrence of any unusual shock, pain, or hemorrhage after delivery without any readily-ascertained cause should always lead to a careful vaginal examination. A want of attention to this rule has too often resulted in the existence of partial inversion being overlooked until its reduction was found to be difficult or impossible.

Mode of Attempting Reduction.—In attempting to reduce a recent inversion the inverted portion of the uterus should be grasped in the hollow of the hand and pushed gently and firmly upward into its natural position, great care being taken to apply the pressure in the proper axis of the pelvis, and to use counter-pressure, by the left hand, on the abdominal walls. Barnes lays stress on the importance of directing the pressure toward one side, so as to avoid the promontory of the sacrum. The common plan of endeavoring to push back the fundus first has been well shown by McClintock¹ to have the disadvantage of increasing the bulk of the mass that has to be reduced; and he advises that while the fundus is lessened in size by compression we should at the same time endeavor to push up first the part that was less inverted—that is to say, the portion nearest the os uteri. Should this be found impossible, some assistance may be derived from the manœuvre, recommended by Merriman and others, of first endeavoring to push up one side or wall of the uterus, and then the other, alternating the upward pressure from one side to the other as we advance. It often happens, as the hand is thus applied, that the uterus somewhat suddenly reinverts itself, sometimes with an audible noise, much as an india-rubber bottle would do under similar circumstances. When reposition has taken place, the hand should be kept for some time in the uterine cavity to excite tonic contraction; or Barnes's suggestion of injecting a weak solution of perchloride of iron may be adopted, so as to constrict the uterine walls and prevent a recurrence of the accident.

It is hardly necessary to point out how much these manœuvres will be facilitated by placing the patient fully under the influence of an anæsthetic.

¹ *Diseases of Women*, p. 79.

Management of the Placenta.—There has been much difference of opinion as to the management of the placenta in cases in which it is still attached when inversion occurs. Should we remove it before attempting reposition, or should we first endeavor to reinvert the organ and subsequently remove the placenta? The removal of the placenta certainly much diminishes the bulk of the inverted portion, and therefore renders reposition easier. On the other hand, if there be much hemorrhage, as is so frequently the case, the removal of the placenta may materially increase the loss of blood. For this reason most authorities recommend that an endeavor should be made at reduction before peeling off the after-birth. But if any delay or difficulty be experienced from the increased bulk, no time should be lost, and it is in every way better to remove the placenta and endeavor to reinvert the organ as soon as possible.

Management of Cases Detected some time after Delivery.—Supposing we met with a case in which the existence of inversion has been overlooked for days, or even for a week or two, the same procedure must be adopted; but the difficulties are much greater, and the longer the delay the greater they are likely to be. Even now, however, a well-conducted attempt at taxis is likely to succeed. Should it fail, we must endeavor to overcome the difficulty by continuous pressure applied by means of caoutchouc bags distended with water and left in the vagina. It is rarely that this will fail in a comparatively recent case, and such only are now under consideration. It is likely that by pressure applied in this way for twenty-four or forty-eight hours, and then followed by taxis, any case detected before the involution of the uterus is completed may be successfully treated.

[Spontaneous Reposition of the Inverted Uterus.]—After all attempts have failed to replace an inverted uterus, already too much contracted to yield to the pressure employed, Nature sometimes accomplishes the work herself, as proved beyond question from quite a number of well-established cases, several of which belong to our own country. A few years ago I saw one of the most remarkable on record. A woman of 29, mother of three children, miscarried at six and a half months from lifting. From the time of her delivery she was subject to weepings of blood, and at times to more or less severe hemorrhages, one of the last of which nearly proved fatal. This condition of disease had lasted three years, when Dr. Walter F. Atlee was called in to relieve her in her worst hemorrhagic attack, and found her uterus inverted, and a nodular growth upon the fundus which gave out an offensive odor. Thinking the disease possibly malignant, and believing, in any event, that to save the woman he would be obliged to remove the uterus, he called a consultation and prepared for the operation; but when the patient was etherized, placed in the knee-elbow position, and Sims's speculum introduced, behold, there was nothing to be seen in the vagina but a soft dilated cervix, the uterus having been replaced by atmospheric pressure, aided perhaps by traction on the uterine attachments within. When explored, the uterus was found to be very soft and thin, and to contain some hard nodular masses, which on removal proved to be portions of an adherent placenta. The hemorrhage ceased upon the reposition and

cleaning out of the uterus, and the patient made a good recovery. She has been again pregnant.

This woman was anæmic to a marked degree, and her abdominal walls so thin that a finger in the uterus could readily be felt above the pubes. There is not the slightest doubt about the inversion, which was proved to exist a short time before the change of posture by Prof. Agnew, who made a finger in the rectum meet another above the pubes, and there was no fundus between them.

Two¹ cases are upon record where reposition was the result of falls, one at eight months and the other after as many years. Drs. Mœhring, C. D. Meigs, H. L. Hodge, and Warrington of this city failed to replace a uterus, and the woman again became pregnant in about six years, aborting with a three months' fœtus under the care of Dr. Warrington. Dr. Meigs saw a second case with Dr. Levis, in which there was violent flooding followed by hemorrhages, which gradually declined. After her return from a journey West she became pregnant and bore a child. Dr. John L. Atlee of Lancaster failed to replace the uterus of a woman, but she recovered spontaneously and bore a child a year afterward.² Dr. Johnson F. Hatch of Kent, Connecticut, reported a case in a letter to Dr. C. D. Meigs in which inversion occurred spontaneously fourteen or fifteen hours after labor. After being under the care of several physicians, she had, at the end of eighteen months, two severe hemorrhagic attacks, after which she improved, and finally, at the end of two years and nine months, bore a child of 9 pounds and 6 ounces.

In all cases spontaneous reposition appears to result from a softening and thinning of the uterine walls as the result of anæmia brought on by hemorrhages. This was particularly noticed by Boivin and Dugès in autopsies of women dying of repeated hemorrhages.—ED.]

[¹ See Dailliez, *Essai sur le Renversement de la Matrice*, Paris, 1805, pp. 105-107.]

[² Meigs's *Obstetrics*, 1852, Philada., p. 608.]

PART IV.

OBSTETRIC OPERATIONS.

CHAPTER I.

INDUCTION OF PREMATURE LABOR.

History of the Operation.—The first of the obstetric operations we have to consider is the *induction of premature labor*—an operation which, like the use of forceps, was first suggested and practised in this country, and the recognition of which, as a legitimate procedure, we also chiefly owe to the labor of our fellow-countrymen, in spite of much opposition both at home and abroad. It is not known with certainty to whom we owe the original suggestion, but we are told by Denman that in the year 1756 there was a consultation of the most eminent physicians at that time in London to consider the advantages which might be expected from the operation. The proposal met with formal approval, and was shortly after carried into practice by Dr. Macaulay, the patient being the wife of a linen-draper in the Strand. From that time it has flourished in Great Britain, the sphere of its application has been largely increased, and it has been the means of saving many mothers and children who would otherwise, in all probability, have perished. On the Continent it was long before the operation was sanctioned or practised. Although recommended by some of the most eminent German practitioners, it was not actually performed until the year 1804. In France the opposition was long continued and bitter. Many of the leading teachers strongly denounced it, and the Academy of Medicine formally discountenanced it so late as the year 1827. The objections were chiefly based on religious grounds, but partly, no doubt, on mistaken notions as to the object proposed to be gained. Although frequently discussed, the operation was never actually carried into practice until the year 1831, when Stoltz performed it with success. Since that time opposition has greatly ceased, and it is now employed and highly recommended by the most distinguished obstetricians of the French schools.

Objects of the Operation.—In inducing premature labor we propose to avoid or lessen the risk to which, in certain cases, the mother is exposed by delivery at term, or to save the life of the child, which might otherwise be endangered. Hence the operation may be indicated either on account of the mother alone or of the child alone, or, as not unfrequently happens, of both together.

Defective Proportion between the Child and Pelvis the most Frequent

Indication.—In by far the largest number of cases the operation is performed on account of defective proportion between the child and the maternal passages, due to some abnormal condition on the part of the mother. This want of proportion may depend on the presence of tumors, either of the uterus or growing from the pelvis. But most frequently it arises from deformity of the pelvis (p. 398), and it is needless to repeat what has been said on that point. I shall therefore only briefly refer to a few more uncommon causes which occasionally necessitate its performance.

Habitually Large Size of the Fœtal Head.—One of these is an habitually large or over-firmly ossified fœtal head. Should we meet with a case in which the labors are always extremely difficult and the head apparently of unusual size, although there is no apparent want of space in the pelvis, the induction of labor would be perfectly justifiable, and in all probability would accomplish the desired object. In such cases the full period of delivery would require to be anticipated by a very short time. A week or a fortnight might make all the difference between a labor of extreme severity and one of comparative ease.

Condition of the Mother's Health calling for the Operation.—There is a large class of cases in which the condition of the mother indicates the operation. Many of these have already been considered when treating of the diseases of pregnancy. Amongst them may be mentioned vomiting which has resisted all treatment, and which has produced a state of exhaustion threatening to prove fatal; chorea, albuminuria, convulsions, or mania; excessive anasarca, ascites, or dyspnœa connected with disease of the heart, lungs, or liver, which may be, in a great measure, caused by the pressure of the enlarged uterus; in fact, any condition or disease affecting the mother, provided only we are convinced that the termination of pregnancy would give the patient relief and that its continuance would involve serious danger. It need hardly be pointed out that the induction of labor for any such causes involves grave responsibility and is decidedly open to abuse: no practitioner would, therefore, be justified in resorting to it, especially if the child have not reached a viable age, without the most anxious consideration. No general rules can be laid down. Each case must be treated on its own merits. It is obvious that the nearer the patient is to the full period, the greater will be the chance of the child surviving, and the less hesitation need then be felt in consulting the interest of the mother.

Conditions Affecting the Safety of the Child alone.—In another class of cases the operation is indicated by circumstances affecting the life of the child alone. Of these the most common are those in which the child dies, in several successive pregnancies, before the termination of uterogestation. This is generally the result of fatty, calcareous, or syphilitic degeneration of the placenta, which is thus rendered incapable of performing its functions. These changes in the placenta seldom commence until a comparatively advanced period of pregnancy, so that if labor be somewhat hastened we may hope to enable the patient to give birth to a living and healthy child. The experience of the mother will indicate the period at which the death of the fœtus has formerly taken place, as she would then have appreciated a difference in her sensations, a dim-

inution in the vigor of the fetal movements, a sense of weight and coldness, and similar signs. For some weeks before the time at which this change has been experienced we should carefully auscultate the fetal heart from day to day, and in most cases the approach of danger will be indicated sufficiently soon, by tumultuous and irregular pulsations or a failure in their strength and frequency, to enable us to interfere with success. On the detection of these, or on the mother feeling that the movements of the child are becoming less strong, the operation should at once be performed. Simpson also induced premature labor with success in a patient who had twice given birth to hydrocephalic children. In the third pregnancy, which he terminated before the natural period, the child was well-formed and healthy.

Induction of Labor when the Mother is Mortally Ill.—Some obstetricians have proposed to induce labor, with the view of saving the child, when the mother was suffering from mortal disease. This indication is, however, so extremely doubtful, from a moral point of view, that it can hardly be considered as ever justifiable.

Various Methods of Inducing Labor: their Mode of Action.—The means adopted for the induction of labor are very numerous. Some of them act through the maternal circulation, as the administration of ergot and other oxytocics; others, by their power of exciting reflex action, or by interfering with the integrity of the ovum, or by a combination of both, as the vaginal douche, separation of the membranes from the uterine walls, puncture of the ovum, dilatation of the os, stimulating enemata, or irritation of the breasts. The former class are never employed in modern obstetric practice. Of the latter, some offer special advantages in particular cases, but none are equally adapted for all emergencies. Often a combination of more methods than one will be found most useful. I shall mention the various methods in use, and discuss briefly the relative advantages and disadvantages of each.

Puncture of Membranes.—The evacuation of the liquor amnii by the puncture of the membranes was the first method practised, and was that recommended by Denman and all the earlier writers. • It is the most certain which can be employed, as it never fails, sooner or later, to induce uterine contractions. There are, however, several disadvantages connected with it which are sufficient to contraindicate its use in the majority of cases. It is uncertain as regards the time taken in producing the desired effect, pains sometimes coming on within a few hours, but occasionally not until several days have elapsed. The contracting walls of the uterus press directly on the body of the child, which, being frail and immature, is less able to bear the pressure than at the full period of pregnancy. Hence it involves great risk to the fetus. Besides, the escape of the water does away with the fluid wedge so useful in dilating the os, and should version be necessary from malpresentation—a complication more likely to occur than in natural labor—the operation would have to be performed under very unfavorable conditions. These objections are sufficient to justify the ordinary opinion that this procedure should not be adopted unless other means have been tried and failed. Every now and then cases are met with in which it is extremely difficult to

...in spite of its ... When the ... that is, before ... it is the sim- ... Indeed, in producing early ... The process should be most simple, ... a suitable instrument to ... of the left hand ... the membranes until ... proposed as a ... should be punctured ... of a gradui ... the risk to the ... he employed a ... which can be projected ... such an inser- ... but better means at our ... When we require to produce ... to puncture the mem- ... The object can be effected ... an ordinary urino- ... and turning it round once or twice.

Administration of Ergot.—The administration of ergot of rye, either alone or combined with beer and cinnamon, has been sometimes resorted to. This practice has been generally advocated by Ramezsius; who was in the habit of administering simple doses of the powdered ergot every fourth hour until delivery took place. Sometimes he found that as many as thirty or forty doses were required to effect the object; occasionally labour commenced after a single dose. Finding that the infantile mortality was very great when this method was followed, he modified it, and administered two or three doses only, and if these proved inefficient he punctured the membranes. There can be no doubt that ergot possesses the power of inducing uterine contractions. The dose in the child is, however, quite as great as when the membranes are punctured, for not only is it subject to injurious pressure from the hypertrophied and irregular contractions which the ergot produces, but the drug itself, when given in large doses, seems to exert a poisonous influence on the fetus. For these reasons ergot may properly be excluded from the available means of inducing labour.

Methods Acting Indirectly on the Uterus.—Various methods have been recommended which act indirectly on the uterus, the source of irritability being at a distance. Thus, M'Quigney used frequently-repeated stimulating frictions and tight bandages. Scanzoni, remembering the intimate connection between the mammae and uterus, and the tendency which irritation of the former has to induce contraction of the latter, recommended the frequent application of cupping-glasses to the breasts. Pothman and others have employed galvanism. Stimulating enemata have been employed. All these methods have occasionally proved successful, and unlike the former plans we have mentioned, they are not attended by any special risk to the child. They are, however, much

too uncertain to be relied on, besides being irksome both to the patient and practitioner.

Artificial Dilatation of the Os Uteri.—The artificial dilatation of the os uteri, in imitation of its natural opening in labor, was first practised by Klüge. He was in the habit of passing within the os a tent made of compressed sponge and allowing it to dilate by imbibition of fluid. If labor were not provoked within twenty-four hours, he removed it, and introduced one of larger dimensions, changing it as often as was necessary until his object was accomplished. Although this operation seldom failed to induce labor, it had the disadvantage of occupying an indefinite time and the irritation produced was often painful and annoying. Dr. Keiller of Edinburgh was the first to suggest the use of caoutchouc bags distended by air as a means of dilating the os. This plan has been perfected by Dr. Barnes in his well-known dilators, which are of great use in many cases in which artificial dilatation of the cervix is necessary. They consist of a series of india-rubber bags of various sizes, with a tube attached (Fig. 149), through which water can be injected by an ordinary Higginson's syringe. They have a small pouch fixed externally, in which a sound can be placed, so as to facilitate their introduction. When distended with water the bags assume somewhat of a fiddle shape, bulging at both extremities, which ensures their being retained within the os. When first introduced into practice as a means of inducing labor, it was thought that this method gave a complete control over the process, so that it could be concluded within a definite time at the will of the operator. The experience of those who have used it much has certainly not justified this anticipation. It is true that occasionally contractions supervene within a few hours after dilatation has been commenced, but, on the other hand, the uterus often responds very imperfectly to this kind of stimulus, and the bags may be inserted for many consecutive hours without the desired result supervening, the puncture of the membranes being eventually necessary in order to hasten the process. Indeed, my own experience would lead me to the conclusion that as a means of evoking uterine contraction cervical dilatation is very unsatisfactory. Dr. Barnes himself has evidently seen reason to modify his original views, for while he at first talked of the bags as enabling us to induce labor with certainty at a given time, he has since recommended that uterine action should be first provoked by other means, the dilators being subsequently used to accelerate the labor thus brought on. The bags, thus employed, find, as I believe, their most useful and a very valuable application; but when used in this way they cannot be considered a means of originating uterine action. A subsidiary objection to the bags is the risk of displacing the presenting part. I have, for example, introduced them when the head was presenting, and on their removal found the shoulder lying over the os. It is not difficult to understand how the continuous pressure of a distended bag in the internal os might easily push away the head, which is so readily

FIG. 149.



Barnes's Bag for Dilating the Cervix.

movable so long as the membranes are unruptured. Still, if labor be in progress and the os insufficiently dilated, the possibility of this occurrence is not a sufficient reason for not availing ourselves of the undoubtedly valuable assistance which the dilators are capable of giving.

Separation of the Membranes.—Some processes for inducing labor act directly on the ovum by separating the membranes, to a greater or less extent, from the uterine walls. The first procedure of the kind was recommended by Dr. Hamilton of Edinburgh, and consisted in the gradual separation of the membranes for one or two inches all around the lower segment of the uterus. To reach them the finger had to be gently insinuated into the interior of the os, which was gradually dilated to a sufficient extent by a series of successive operations repeated at intervals of three or four hours. When this had been accomplished, the fore finger was inserted and swept round between the membranes and the uterus; but it was frequently found necessary to introduce the greater part of the hand to effect the object, and sometimes even this was not sufficient, and a female catheter or other instrument had to be used for the purpose. The method was generally successful in bringing on labor, but it now and then failed, even in Dr. Hamilton's hands. It is certainly based on correct principles, but it is tedious and painful both to the practitioner and the patient, and very uncertain in its time of action. For these reasons it has never been much practised.

Vaginal and Uterine Douches.—In the year 1836, Kiwisch suggested a plan which from its simplicity has met with much approval. It consists in projecting at intervals a stream of warm or cold water against the os uteri. Its action is doubtless complex. Kiwisch himself believed that relaxation of the soft parts through the imbibition of water was the determining cause of labor. Simpson found that the method failed unless the water mechanically separated the membranes from the uterine walls. Besides this effect, it probably directly induces reflex action by distending the vagina and dilating the os. In using it, it has been customary to administer a douche twice daily, and more frequently if rapid effects be desired. The number required varies in different cases. The largest number Kiwisch found it necessary to use was 17, the smallest 5. The average time that elapses before labor sets in is four days. Hence the method is obviously useless when rapid delivery is required.

Dr. Cohen of Hamburg introduced an important modification of the process which has been considerably practised. It consists in passing a silver or gum-elastic catheter some inches within the os, between the membranes and the uterine walls, and injecting the fluid through it directly into the cavity of the uterus. He used creasote or tar-water, and injected without stopping until the patient complained of a feeling of distension. Others have found the plan equally efficacious when they only employed a small quantity of plain water, such as 7 or 8 ounces. Professor Lazarewitch of Charkoff is a strong advocate of this method. He believes that uterine action is evoked much more rapidly and certainly if the water be injected near the fundus, and he has contrived an instrument for the purpose with a long metallic nozzle.

Dangers of these Plans.—So many fatal cases have followed these methods that it cannot be doubted that, in spite of their certainty and

simplicity, there is an element of risk in them that should not be overlooked. Many of these are recorded in Barnes's work, and he comes to the conclusion, which the facts unquestionably justify, that "the douche, whether vaginal or intra-uterine, ought to be absolutely condemned as a means of inducing labor." The precise reason of the danger is not very obvious. Sudden stretching of the uterine walls, producing shock, has been supposed to have caused it; but in many of the fatal cases the symptoms have been rather those attending the passage of air into the veins, and it is easy to understand how air may have been introduced in this way into the large uterine sinuses.

Injection of Carbonic Acid Gas.—Simpson and Scanzoni have both tried with success the injection of carbonic acid gas into the vagina. Fatal results have, however, followed its employment, and Simpson has expressed an opinion that the experiment should not be repeated.

Simpson's Method of Operating.—Simpson originally induced labor by passing the uterine sound within the os and up toward the fundus, and when it has been inserted to a sufficient extent moving it slightly from side to side. He was led to adopt this procedure in the belief that we might thus closely imitate the separation of the decidua which occurs previous to labor at term. Uterine contractions were induced with certainty and ease, but it was found impossible to foretell what time might elapse between the commencement of labor and the operation, which had frequently to be performed more than once. He subsequently modified this procedure by introducing a flexible male catheter without a stilette, which he allowed to remain in the uterus until contractions were excited. This plan is much used in Germany, and is now that which is also most frequently adopted in this country. It is simple and very efficacious, pains coming on almost invariably within twenty-four hours after the catheter or bougie is introduced. A theoretical objection is the possibility of the catheter separating a portion of the placenta and giving rise to hemorrhage; but in practice this has not been found to occur, and the risk might generally be avoided by introducing the catheter at a distance from the placenta, the probable situation of which has been ascertained by auscultation. The more deeply the catheter is introduced, the more certain and rapid is its effect, and not less than seven inches should be pushed up within the os. It is not always easy to insert it so far, especially if a flexible catheter be used, which is apt to be too pliable to pass upward with ease. A solid bougie—male urethral bougie—should therefore be employed; and I have found its introduction greatly facilitated by anesthetizing the patient and passing the greater part of the hand into the vagina. In this way it can be pushed in very gently and without any risk of injury to the uterus. There is some chance of rupturing the membranes while pushing it upward. This accident, indeed, cannot always be avoided, even when the greatest care is taken; but when it occurs the puncture will be at a distance from the os, so that a small portion only of the liquor amnii will escape; and this can scarcely be considered a serious objection. It is always an advantage to allow the pains to come on gradually and in imitation of natural labor. Therefore, if, after the bougie has been inserted for a sufficient time, uterine contractions come on sufficiently strongly, we may leave the case

to be terminated naturally; or if they be comparatively feeble, we may resort to accelerative procedures—viz. dilatation of the cervix by the fluid bags, and subsequently the puncture of the membranes. In this way we have the labor completely under control; and I believe this method will commend itself to those who have experience of it as the simplest and most certain mode of inducing labor yet known, and the one most closely imitating the natural process. Of late I have been in the habit of combining dilatation of the cervix with this method by means of a well-carbolized sponge tent passed into the cervix after the bougie is in position. In ten or twelve hours, when the tent and bougie are removed, the cervix is found well dilated and ready for the passage of the child.

The Child is Immature and Difficult to Rear.—It should not be forgotten that the child is immature, and that unusual care is likely to be required to rear it successfully. We should, therefore, be careful to have at hand all the usual means of resuscitation; and, as the mother may not be able to nurse at once, it would be a good precaution to have a healthy wet-nurse in readiness.^[1]

[Through the kindness of Dr. Cesare Belluzzi of Bologna I have received his two reports, containing the records of 112 cases in which he brought on labor prematurely, with a saving of 104 mothers and the delivery of as many living children. 42 patients were treated in private practice and 70 in the Maternity of Bologna. In 9 patients labor was induced because of disease in the mother; in 1 it was brought on because the foetus had usually died in the ninth month of former pregnancies; and in 102 the pelvis was contracted. Of these 102, 6 died—3 out of 38 in private practice, and 3 out of 64 in the hospital. Of the 9 women operated upon because of serious disease, 7 recovered. 35 out of 42 infants were delivered alive in private practice, and 62 out of 70 in the Maternity.]

Dr. Ludwig Winckel of Mülheim, Germany, has also sent me his record of 25 deliveries in women who were all the subjects of contraction of the pelvis. These patients all recovered: 14 children were still-born and 13 were living; of the latter, only 7 were alive at the end of two weeks. The prolonged vitality of the foetus is largely dependent upon the period in gestation which is chosen for the operation: the later the delivery, the better is the prospect of ultimate safety. But a small proportion of the children reach maturity. Of 32 delivered alive in hospital in a period of less than ten years under Dr. Belluzzi, 27 were dead before the expiration of the first year, and 29 in all within two years of birth.—ED.]

[¹ In some of the European maternities they now have in use a little heated chamber in which the infant is put to sleep; it is kept warm, on the principle of an egg-hatching machine, by a lamp and water-chamber.—ED.]

CHAPTER II.

TURNING.

History of the Operation.—Turning—by which we mean the alteration of the position of the fœtus, and the substitution of some other portion of the body for that originally presenting—is one of the most important of obstetric operations and merits careful study. It is also one of the most ancient, and was evidently known to the Greek and Roman physicians. Up to the fifteenth century, cephalic version—that in which the head of the fœtus is brought over the os uteri—was almost exclusively practised, when Paré and his pupil Guillemeau taught the propriety of bringing the feet down first. It was by the latter physician especially that the steps of the operation were clearly defined; and the French have undoubtedly the merit both of perfecting its performance and of establishing the indications which should lead to its use. Indeed, it was then much more frequently performed than in later times, since no other means of effecting artificial delivery were known which did not involve the death of the child; and practitioners doubtless acquired great skill in its performance, and were inclined to overrate its importance and extend its use to unsuitable cases. An opposite error was fallen into after the invention of the forceps, which for a time led to the abandonment of turning in certain conditions for which it was well adapted and in which it has only of late years been again practised.

Cephalic Version.—Cephalic version has, since Paré wrote, been recommended and practised from time to time, but the difficulty of performing it satisfactorily was so great that it never became an established operation. Dr. Braxton Hicks has perfected a method by which it can be accomplished with greater ease and certainty, and which renders it a legitimate and satisfactory resort in suitable cases. To him we are also indebted for introducing a method of turning without passing the entire hand into the cavity of the uterus, which, under favorable circumstances, is not only easy of performance, but deprives the operation of one of its greatest dangers.

Turning by External and Internal Manipulation.—The possibility of effecting version by external manipulation has long been known, and was distinctly referred to and recommended by Dr. John Pechey¹ so far back as the year 1698. Since that time it has been strongly recommended by Wigand and his followers; and various authors in this country, notably Sir James Simpson, have referred to the advantage to be derived from external manipulation assisting the hand in the interior of the uterus. In 1854, Dr. Wright of Cincinnati advocated the application of the bi-manual method in arm and shoulder presentations, chiefly with the view of effecting cephalic version. To Dr. Hicks, however, incontestably belongs the merit of having been the first dis-

¹ *The Complete Midwife's Practice*, p. 142.

tinely to show the possibility of effecting complete version in all cases in which the operation is indicated by combined external and internal manipulation, of laying down definite rules for its practice, and for thus popularizing one of the greatest improvements in modern midwifery.

Object and Nature of the Operation.—The operation is entirely dependent for success on the fact that the child in utero is freely movable, and that its position may be artificially altered with facility. As long as the membranes are unruptured and the fœtus is floating in the surrounding fluid medium, it is liable to constant changes in position, as may be readily demonstrated in the latter months of pregnancy; and the operation under these circumstances may be performed with the greatest facility. Shortly after the liquor amnii has escaped there is still, as a rule, no great difficulty in effecting version, but, as the body is no longer floating in the surrounding liquid, its rotation must necessarily be attended with some increased risk of injury to the uterus. If the liquor amnii have been long evacuated and the muscular structure of the uterus be strongly contracted, the fœtus may be so firmly fixed that any attempt to move it is surrounded with the greatest difficulties, and may even fail entirely or be attended with such risks to the maternal structures as to be quite unjustifiable.

Cases Suitable for the Operation.—Version may be required either on account of the mother or child alone, or it may be indicated by some condition imperilling both and rendering immediate delivery necessary. The chief cases in which it is resorted to are those of transverse presentation, where it is absolutely essential; accidental or unavoidable hemorrhage; certain cases of contracted pelvis; and some complications, especially prolapse of the funis. The special indications for the operation have been separately discussed under these subjects.

Statistics and Dangers of the Operation.—The ordinary statistical tables cannot be depended on as giving any reliable results as to the risks of the operation. Taking all cases together, Dr. Churchill estimates the maternal mortality as 1 in 16, and the infantile as 1 in 3. Like all similar statistics, they are open to the objection of not distinguishing between the results of the operation itself and of the cause which necessitated interference. Still, they are sufficient to show that the operation is not free from grave hazards, and that it must not be undertaken without due reflection. The principal dangers will be discussed as we proceed. It may suffice to mention here that those to the mother must vary with the period at which the operation is undertaken. If version be performed early, before the rupture of the membranes, or in favorable cases without the introduction of the hand into the interior of the uterus, the risk must of course be infinitely less than in those more formidable cases in which the waters have long escaped and the hand and arm have to be passed into an irritable and contracted uterus. But even in the most unfavorable cases accidents may be avoided if the operator bear constantly in mind that the principal danger consists in laceration of the uterus or vagina from undue force being employed or from the hand and arm not being introduced in the axis of the passages. There is no operation in which gentleness, absence of all hurry, and complete presence of mind are so essential. A certain number of cases

end fatally from shock or exhaustion or from subsequent complications. As regards the child, the mortality is little, if at all, greater than in original breech and footling presentations. Nor is there any good reason why it should be so, seeing that cases of turning, after the feet are brought through the os, are virtually reduced to those of feet presentation, and that the mere version, if effected sufficiently soon, is not likely to add materially to the risk to which the child is exposed.

Version by External Manipulation.—The possibility of effecting *version by external manipulation* has been recognized by various authors, and was made the subject of an excellent thesis by Wigand, who clearly described the manner of performing the operation. In spite of the manifest advantages of the procedure, and the extreme facility with which it can be accomplished in suitable cases, it has by no means become the established custom to trust to it, and probably most practitioners have never attempted it, even under the most favorable conditions. The possibility of the operation is based on the extreme mobility of the fœtus before the membranes are ruptured. After the waters have escaped the uterine walls embrace the fœtus more or less closely, and version can no longer be readily performed in this manner.

Cases Suitable for the Operation.—It may, therefore, be laid down as a rule that it should only be attempted when the abnormal position of the fœtus is detected before labor has commenced, or in the early stage of labor when the membranes are unruptured. It is also unsuitable for any but transverse presentations, for it is not meant to effect complete evolution of the fœtus, but only to substitute the head for the upper extremity. It is useless whenever rapid delivery is indicated, for after the head is brought over the brim the conclusion of the case must be left to the natural powers.

The manner of detecting the presentation by palpation has been already described (p. 121), and the success of the operation depends on our being able to ascertain the positions of the head and breech through the uterine walls. Should labor have commenced and the os be dilated, the transverse presentation may be also made out by vaginal examination. Should the abnormal presentation be detected before labor has actually begun, it is in most cases easy enough to alter it and to bring the fœtus into the longitudinal axis of the uterine cavity. Pinard¹ recommends that after this has been done the fœtus should be maintained in position by a well-fitting elastic abdominal belt. It is seldom, however, discovered until labor has commenced, and, even if it be altered, the child is extremely apt to resume in a short time the faulty position in which it was formerly lying. Still, there can be no harm in making the attempt, since the operation itself is in no way painful, and is absolutely without risk either to the mother or child. When the transverse presentation is detected early in labor, I believe it is good practice to endeavor to remedy it by external manipulation, and if it fail we may at once proceed to other and more certain methods of operating. The procedure itself is abundantly simple. The patient is placed on her back, and the position of the fœtus ascertained by palpation as accurately as possible, in the manner already described. The palms of the hands being then

¹ *De la Version par Manœuvres externes*, Paris, 1878.

placed over the opposite poles of the fœtus, by a series of gentle gliding movements the head is pushed toward the pelvic brim, while the breech is moved in the opposite direction. The facility with which the fœtus may sometimes be moved in this way can hardly be appreciated by those who have never attempted the operation. As soon as the change is effected the long diameters of the fœtus and the uterus will correspond, and vaginal examination will show that the shoulder is no longer presenting and that the head is over the pelvic brim. If the os be sufficiently dilated and labor in progress, the membranes should now be punctured, and the position of the fœtus maintained for a short time by external pressure until we are certain that the cephalic presentation is permanently established. If labor be not in progress, an attempt may at least be made to effect the same object by pads and a binder, one pad being placed on the side of the uterus in the situation of the breech, and another on the opposite side in the situation of the head.

Cephalic Version.—On account of the difficulty of performing *cephalic version* in the manner usually recommended, it has practically scarcely been attempted, and with the exception of some more recent authors it is generally condemned by writers on systematic midwifery. Still, the operation offers unquestionable advantages in those transverse presentations in which rapid delivery is not necessary, and in which the only object of interference is the rectification of malposition; for if successful the child is spared the risk of being drawn footling through the pelvis. The objections to cephalic version are based entirely on the difficulty of performance; and, undoubtedly, to introduce the hand within the uterus, search for, seize, and afterward place the slippery head in the brim of the pelvis, could not be an easy process even under the most favorable circumstances, and must always be attended with considerable risk to the mother. Velpeau, who strongly advocated the operation, was of opinion that it might be more easily accomplished by pushing up the presenting part than by seizing and bringing down the head. Wigand more distinctly pointed out that the head could be brought to a proper position by external manipulation, aided by the fingers of one hand within the vagina. Braxton Hicks has laid down clear rules for its performance which render cephalic version easy to accomplish under favorable conditions, and will doubtless cause it to become a recognized mode of treating malpositions. The number of cases, however, in which it can be performed, must always be limited, since, as in turning by external manipulation alone, it is necessary that the liquor amnii should be still retained, or at least have only recently escaped; that the presentation be freely movable about the pelvic brim; and that there be no necessity for rapid delivery. Dr. Hicks does not believe protrusion of the arm to be a contraindication, and advises that it should be carefully replaced within the uterus. When, however, protrusion of the arm has occurred, the thorax is so constantly pushed down into the pelvis that replacement can neither be safe nor practicable, except under unusually favorable conditions, and podalic version will be necessary.

Method of Performance.—It is impossible to describe the method of performing cephalic version more concisely and clearly than in Dr. Hicks' own words. "Introduce," he says, "the left hand into the vagina,

as in podalic version ; place the right hand on the outside of the abdomen, in order to make out the position of the fœtus and the direction of its head and feet. Should the shoulder, for instance, present, then push it with one or two fingers in the direction of the feet. At the same time pressure with the other hand should be exerted on the cephalic end of the child. This will bring the head down to the os ; then let the head be received on the tips of the inside fingers. The head will play like a ball between the two hands ; it will be under their command, and can be placed in almost any part at will. Let the head then be placed over the os, taking care to rectify any tendency to face presentation. It is as well, if the breech will not rise to the fundus readily after the head is fairly in the os, to withdraw the hand from the vagina, and with it press up the breech from the exterior. The hand which is retaining gently the head from the outside should continue there for some little time till the pains have ensured the retention of the child in its new position and the adaptation of the uterine walls to its new form. Should the membranes be perfect, it is advisable to rupture them as soon as the head is at the os uteri ; during their flow and after, the head will move easily into its proper position."

The procedure thus described is so simple, and would occupy so short a time, that there can be no objection to trying it. Should we fail in our endeavors, we shall not be in a worse position for effecting delivery by podalic version, which can be proceeded with without withdrawing the hand from the vagina or in any way altering the position of the patient.

Podalic Version.—The method of performing podalic version varies with the nature of each particular case. In describing the operation it has been usual to divide the cases into those in which the circumstances are favorable and the necessary manœuvres easily accomplished, and those in which there are likely to be considerable difficulties and increased risk to the mother. This division is eminently practicable, since nothing can be more variable than the circumstances under which version may be required. Before describing the steps of the operation it may be well to consider some general conditions applicable to all cases alike.

Position of the Patient.—In this country the ordinary position on the left side is usually employed. On the Continent and in America the patient is placed on her back, with the legs supported by assistants, as in lithotomy. The former position is preferable, not only as a matter of custom and as involving much less fuss and exposure of the person, but because it admits of both the operator's hands being more easily used in concert. In certain difficult cases, when the liquor amnii has escaped and the back of the child is turned toward the spine of the mother, the dorsal decubitus presents some advantages in enabling the hand to pass more readily over the body of the child ; but such cases are comparatively rare. The patient should be brought to the side of the bed, across which she should be laid with the hips projecting over and parallel to the edge, the knees being flexed toward the abdomen and separated from each other by a pillow or by an assistant. Assistants should also be placed so as to restrain the patient if necessary, and pre-

vent her involuntarily starting from the operator, as this might not only embarrass his movements, but be the cause of serious injury.

Administration of Anæsthetics.—The exhibition of anæsthetics is peculiarly advantageous. There is nothing which tends to facilitate the steps of the process so much as stillness on the part of the patient and the absence of strong uterine contraction. When the vagina is very irritable and the uterus firmly contracted round the body of the child, complete anæsthesia may enable us to effect version, when without it we should certainly fail.

Period when the Operation should be Undertaken.—The most favorable time for operating is when the os is fully dilated, before or immediately after the rupture of the membranes and the discharge of the liquor amnii. The advantage gained by operating before the waters have escaped cannot be overstated, since we can then make the child rotate with great facility in the fluid medium in which it floats. In the ordinary operation, in which the hand is passed into the uterus, it is essential to wait until the os is of sufficient size to admit of its being introduced with safety. This may generally be done when the os is the size of a crown-piece, especially if it be soft and yielding.

Choice of Hand to be Used.—The practice followed with regard to the hand to be used in turning varies considerably. Some accoucheurs always employ the right hand, others the left, and some one or other according to the position of the child. In favor of the right hand it is said that most practitioners have more power with it and are able to use it with greater gentleness and delicacy. In transverse presentations, if the abdomen of the child be placed anteriorly, the right hand is said to be the proper one to use, on account of the greater facility with which it can be passed over the front of the child; and in difficult cases of this kind, when we are operating with the patient on her back, it certainly can be employed with more precision than the left. In all ordinary cases, however, the left hand can be introduced much more easily in the axis of the passages; the back of the hand adapts itself readily to the curve of the sacrum, and, even when the child's abdomen lies anteriorly, it can be passed forward without difficulty so as to seize the feet. These advantages are sufficient to recommend its use, and very little practice is required to enable the practitioner to manipulate with it as freely as with the right. If, in addition, we remember that the right hand is required to operate on the fetus through the abdominal walls—and this is a point which should never be forgotten—we shall have abundant reasons for laying it down as a rule that the left hand should generally be employed. Before passing the hand and arm they should be freely lubricated, with the exception of the palm, which is left untouched to admit of a firm grasp being taken of the foetal limbs. It is also advisable to remove the coat and bare the arm as high as the elbow.


As it should be a cardinal rule to resort to the simplest procedure when practicable, it will be well to consider first the method by combined external and internal manipulation, without passing the hand into the uterus, and subsequently that which involves the introduction of the hand.

Turning by Combined External and Internal Manipulation.—To effect

podalic version by the combined method it is an essential preliminary to ascertain the situation of the fetus as accurately as possible. It will generally be easy in transverse presentation to make out the breech and head by palpation, while in head presentations the fontanelles will show to which side of the pelvis the face is turned. The left hand is then to be passed carefully into the vagina, in the axis of the canal, to a sufficient extent to admit of the fingers passing freely into the cervix. To effect this it is not always necessary to insert the whole hand, three or four fingers being generally sufficient.

If the head lie in the first or fourth position, push it upward and to the left, while the other hand, placed externally on the abdomen, depresses

FIG. 150.




First Stage of Bi-polar Version: Elevation of the Head and Depression of the Breech. (After Barnes.)

the breech toward the right (Fig. 150). By this means we act simultaneously on both extremities of the child's body and easily alter its position. The breech is pushed down gently but firmly by gliding the hand over the abdominal wall. The head will now pass out of reach, and the shoulders will arrive at the os and will lie on the tips of the fingers. This is similarly pushed upward in the same direction as the head (Fig. 151), the breech at the same time being still further depressed until the knee comes within reach of the fingers, when (the membranes being now ruptured, if still unbroken) it is seized and pulled down through the os (Fig. 152). Occasionally the foot comes immediately over the os, when it can be seized instead of the knee. Version may be facilitated by changing the position of the external hand and pushing

the head upward from the iliac fossa, instead of continuing the attempt to depress the breech (Figs. 152 and 153). These manipulations should always be carried on in the intervals, and desisted from when the pains

FIG. 151.



Second Stage of Bi-polar Version : Elevation of the Shoulders and Depression of the Breech. (After Barnes.)

come on ; and when the pains recur with great force and frequency the advantage of chloroform will be particularly apparent. In the second

FIG. 152.



Third Stage of Bi-polar Version : Seizure of the Knee and Partial Elevation of the Head. (After Barnes.)

and third positions the steps of the operation should be reversed : the head is pushed upward and to the right, the breech downward and to the left. When the position cannot be made out with certainty, it is well

to assume that it is the first, since that is the one most frequently met with; and even if it be not, no great inconvenience is likely to occur. If the os be not sufficiently open to admit of delivery being concluded, the lower extremity can be retained in its new position with one finger until dilatation is sufficiently advanced or until the uterus has perma-

FIG. 153.



Fourth Stage of Bi-polar Version: Drawing Down of the Legs and Completion of Version. (After Barnes.)

nently adapted itself to the altered position of the child; either of which results will generally be effected in a short space of time.

In transverse presentations the same means are to be adopted, the shoulder being pushed upward in the direction of the head, while the breech is depressed from without. This is frequently sufficient to bring the knees within reach, especially if the membranes are entire, but version is much facilitated by pressing the head upward from without, alternately with depression of the breech. If the liquor amnii has escaped, and the uterus is firmly contracted round the body of the child, it will be found impossible to effect an alteration in its position without the introduction of the hand, and the ordinary method of turning must be employed. The peculiar advantage of the combined process is that it in no way interferes with the latter, for should it not succeed the hand can be passed on into the uterus without withdrawal from the vagina (provided the os be sufficiently dilated) and the feet or knees seized and brought down.

Podalic Version when the Hand is Introduced into the Uterus.—Turning with the hand introduced into the uterus, provided the waters have not or have only recently escaped and the os be sufficiently dilated, is an operation generally performed with ease.

Introduction of the Hand.—The first step, and one of the most important, is the introduction of the hand and arm. The fingers having been pressed together in the form of a cone, the thumb lying between the rest of the fingers, the hand, thus reduced to the smallest possible dimensions, is slowly and carefully passed into the vagina in the axis of the outlet in an interval between the pains, and passed onward in the same cautious manner and with a semi-rotatory motion until it lies entirely within the vagina, the direction of introduction being gradually changed from the axis of the outlet to that of the brim. If uterine contractions come on, the hand should remain passive until they are over. It should ever be borne in mind as one of the fundamental rules in performing version that we should act only in the absence of pains, and then with the utmost gentleness, all force and violent pushing being avoided. The hand, still in the form of a cone, having arrived at the os, if this be sufficiently dilated may be passed through at once. If the os be not quite open, but dilatable, the points of the fingers may be gently insinuated and occasionally expanded, so as to press it open sufficiently to permit the rest of the hand to pass. While this is being done the uterus should be steadied by the other hand

FIG. 154.



Seizure of the Feet when the Hand is Introduced into the Uterus.

placed externally or by an assistant. If the presentation should not previously have been made out with accuracy, we can now ascertain how to pass the hand onward, so that its palmar surface may correspond with the abdomen of the child.

Rupture of the Membranes.—The membranes should now be ruptured—if possible during the absence of pain—so as to prevent the waters being forced out. The hand and arm form a most efficient plug, and the liquor amnii cannot escape in any quantity. Some practitioners recommend that before rupturing the membranes the hand should be passed onward between them and the uterine walls until we reach the feet. By so doing we run the risk of separating the placenta; besides, we have to introduce the hand much farther than may be necessary, since the knees are often found lying quite close to the os. As soon as the membranes are perforated, the hand can be passed on in search of

FIG. 155.



Drawing Down of the Feet and Completion of Version.

the feet (Fig. 154). At this stage of the operation increased care is necessary to avoid anything like force, and should a pain come on the hand must be kept perfectly flat and still, and rather pressed on the body of the child than on the uterus. If the pains be strong, much inconvenience may be felt from the compression; and were the onward movement continued, or the hand even kept bent in the conical form in which it was introduced, rupture of the uterine walls might easily be caused. This is not likely to occur in the class of cases now under consideration, for it is chiefly when the waters have long escaped that the progress of the hand is a matter of difficulty. Valuable assistance may now be given by pressing the breech downward from without, so as to bring the knees or feet more easily within the reach of the internal hand.

Having arrived at the knees or feet, they may be seized between the fingers and drawn downward in the absence of a pain (Fig. 155). This will cause the fœtus to revolve on its axis, the breech will descend, and at the same time the ascent of the head may be assisted by the right hand from without. It is a question with many accoucheurs which part of the inferior extremities should be seized and brought down. Some recommend us to seize both feet, others prefer one only, while some advise the seizure of one or both knees. In a simple case of turning, before the escape of the waters, it does not matter much which of these plans is followed, since version is accomplished with the greatest ease by any one of them. The seizure of the knee, however, instead of the feet, offers certain advantages which should not be overlooked. It is generally more accessible, affords a better hold (the fingers being inserted in the flexure of the ham), and, being nearer the spine, traction acts more directly on the body of the child. Any danger of mistaking the knee for the elbow may be obviated by remembering the simple rule that the salient angle of the former when the thigh is flexed looks toward the head of the child, of the latter toward its feet. Certain advantages may also be gained by bringing down one foot or knee only, instead of both. When one inferior extremity remains flexed on the body of the child, the part which has to pass through the os is larger than when both legs are drawn down, and consequently the os is more perfectly dilated, and less difficulty is likely to be experienced in the delivery of the rest of the body, so that the risk to the child is materially diminished.

Choice of the Leg to be brought down in Transverse Presentation.—Simpson, whose views have been adopted by Barnes and other writers, recommends the seizing, if possible, in arm presentations, of the knee farthest from and opposite to the presenting arm, as by this means the body is turned round on its longitudinal axis and the presenting arm and shoulder more easily withdrawn from the os. Dr. Galabin has carefully investigated this point in a recent paper,¹ and contends that there is a greater mechanical advantage in seizing the leg which is nearest to, and on the same side as, the presenting arm; and this, moreover, is generally more readily done.


Management of the Case after Version.—As soon as the head has reached the fundus and the lower extremity is brought through the os, the case is converted into a foot or knee presentation, and it comes to be a question whether delivery should now be left to nature or terminated by art. This must depend to a certain extent on the case itself and on the cause which necessitated version, but generally it will be advisable to finish delivery without unnecessary delay. To accomplish this, downward traction is made during the pains, and desisted from in the intervals (Fig. 156). As the umbilical cord appears, a loop should be drawn down; and if the hands be above the head, they must be disengaged and brought over the face in the same manner as in an ordinary footling presentation. The management of the head after it descends into the cavity of the pelvis must also be conducted as in labors of that description.

Turning in Placenta Prævia.—In cases of placenta prævia the os will,

¹ *Obst. Trans.*, vol. xix, 1877.

as a rule, be more easily dilatable than in transverse presentations. Hicks' method offers the great advantage of enabling us to perform version much sooner than was formerly possible, since it only requires the introduction of one or two fingers into the os uteri. Should we not succeed by it, and the state of the patient indicates that delivery is necessary, we have at our command in the fluid dilators a means of artificially dilating the os uteri which can be employed with ease and safety. If we have to do with a case of entire placental presentation, the hand should be passed at that point where the placenta seems to be least attached. This will always be better than attempting to perforate its

FIG. 156.



Showing the Completion of Version. (After Barnes.)

substance—a measure sometimes recommended, but more easily performed in theory than in practice. If the placenta only partially present, the hand should, of course, be inserted at its free border. It will frequently be advisable not to hasten delivery after the feet have been brought through the os, for they form of themselves a very efficient plug and effectually prevent further loss of blood; while, if the patient be much exhausted, she may have her strength recruited by stimulants, etc. before the completion of delivery.

Turning in Abdomino-anterior Positions.—In abdomino-anterior positions in which the waters have escaped, and in which, therefore, some difficulty may be reasonably anticipated, the operation is generally more easily performed with the patient on her back; the right hand is then introduced into the uterus and the left employed externally (Fig. 157).

In this way the internal hand has to be passed a shorter distance and in a less constrained position. The operator then sits in front of the patient, who is supported at the edge of the bed in the lithotomy position

FIG. 157.



Showing the Use of the Right Hand in Abdomino-anterior Position.

with the thighs separated, and the right hand is passed up behind the pubes and over the abdomen of the child.

Difficult Cases of Arm Presentation.—The difficulties of turning culminate in those unfavorable cases of arm presentation in which the membranes have been long ruptured, the shoulder and arm pressed down into the pelvis, and the uterus contracted round the body of the child. The uterus being firmly and spasmodically contracted, the attempt to introduce the hand often only makes matters worse by inducing more frequent and stronger pains. Even if the hand and arm be successfully passed, much difficulty is often experienced in causing the body of the child to rotate; for we have no longer the fluid medium present in which it floated and moved with ease, and the arm of the operator may be so cramped and pained by the pressure of the uterine walls as to be rendered almost powerless. The risk of laceration is also greatly increased, and the care necessary to avoid so serious an accident adds much to the difficulty of the operation.

Value of Anæsthesia in Relaxing the Uterus.—In these perplexing cases various expedients have been tried to cause relaxation of the spasmodically contracted uterine fibres, such as copious venesection in the erect attitude until fainting is induced, warm baths, tartar emetic, and similar depressing agents. None of these, however, are so useful as the free administration of chloroform, which has practically superseded them all, and often answers most effectually when given to its full surgical extent.

Mode of Procedure.—The hand must be introduced with the precautions already described. If the arm be completely protruded into the vagina, we should pass the hand along it as a guide, and its palmar surface will at once indicate the position of the child's abdomen. No advantage is gained by amputation, as is sometimes recommended. When the os is reached the real difficulties of the operation commence, and, if the shoulder be firmly pressed down into the brim of the pelvis, it may not be easy to insinuate the hand past it. It is allowable to repress the presenting part a little, but with extreme caution, for fear of injuring the contracted uterine parietes. It is better to insinuate the hand past the obstruction, which can generally be done by patient and cautious endeavors. Having succeeded in passing the shoulder, the hand is to be pressed forward in the intervals, being kept perfectly flat and still on the body of the fœtus when the pains come on. It is much safer to press on it than on the uterine walls, which might readily be lacerated by the projecting knuckles. When the hand has advanced sufficiently far, it will be better, for the reasons already mentioned, to seize and bring down one knee only.

Management of Cases in which the Foot is brought down, but the Fœtus will not Revolve.—Even when the foot has been seized and brought through the os, it is by no means always easy to make the child revolve on its axis, as the shoulder is often so firmly fixed in the pelvic brim as not to rise toward the fundus. Some assistance may be derived from pushing the head upward from without, which, of course, would raise the shoulder along with it. If this should fail, we may effect our object by passing a noose of tape or wire ribbon round the limb, by which traction is made downward and backward; at the same time the other hand is passed into the vagina to displace the shoulder and push it out of the brim. It is evident that this cannot be done as long as the limb is held by the left hand, as there is no room for both hands to pass into the vagina at the same time. By this manœuvre version may be often completed when the fœtus cannot be turned in the ordinary way. Various instruments have been invented, both for passing a lac round the child's limb and for repressing the shoulder, but none of them can compete, either in facility of use or safety, with the hand of the accoucheur.

If all Attempts at Version Fail, Mutilation of the Fœtus is Necessary.—Should all attempts at version fail, no resource is left but the mutilation of the child, either by evisceration or decapitation. This extreme measure is, fortunately, seldom necessary, as with due care version may generally be effected, even under the most unfavorable circumstances.

CHAPTER III.

THE FORCEPS.

OF all obstetric operations, the most important, because the most truly conservative both to the mother and child, is the application of the forceps. In modern midwifery the use of the instrument is much extended, and it is now applied by some of our most experienced accoucheurs with a frequency which older practitioners would have strongly reprobated. That the injudicious and unskilful use of the forceps is capable of doing much harm no one will for a moment deny. This, however, is not a reason for rejecting the recommendation of those who advise a more frequent resort to the operation, but rather for urging on the practitioner the necessity of carefully studying the manner of performing it, and of making himself familiar with the cases in which it is easy or the reverse. Nothing but practice—at first on the manikin, and afterward in actual cases—can impart the operative dexterity which it should be the aim of every obstetrician to acquire, and without which there can be no assurance of his doing his duty to his patient efficiently.

Description of the Instrument.—The forceps may best be described as a pair of artificial hands by which the foetal head may be grasped and drawn through the maternal passages by a *vis a fronte* when the *vis a tergo* is deficient. This description will impress on the mind the important action of the instrument as a tractor, to which all its other powers are subservient. The forceps consists of two separate blades of a curved form adapted to fit the child's head, a lock by which the blades are united after introduction, and handles which are grasped by the operator and by means of which traction is made. It would be a wearisome and unsatisfactory task to dwell on all the modifications of the instrument which have been made, which are so numerous as to make it almost appear as if no one could practise midwifery with the least pretension to eminence unless he has attached his name to a new variety of forceps.

The Short Forceps.—The original instrument, invented by the Chamberlens, may be looked upon as the type of the short straight forceps, which has been more employed than any other, and which perhaps finds its best representative in the short forceps of Denman (Fig. 158). Indeed, the only essential difference between the two is the lock of the latter, originally invented by Smellie, which is so excellent that it has been adopted in all British forceps, and which, for facility of juncture, is much superior to either the French pivot or the German lock, while for firmness it is, for all practical purposes, as good as either. In this instrument the blades are 7, the handles $4\frac{3}{8}$, inches in length; the extremities of the blades are exactly 1 inch apart, and the space between them at their widest part is $2\frac{7}{8}$ inches. The blades measure $1\frac{3}{4}$ inches at their greatest breadth, and spring with a regular sweep directly from the lock,


there being no shank. The blades are formed of the best and most highly-tempered steel to resist the strain to which they are occasionally subjected, and they are smooth and rounded on their inner surface to obviate the risk of injury to the scalp of the child.

Advantage Claimed for this Form of Instrument.—The special advantage claimed for this form of instrument is that, the two halves being precisely similar, no care or forethought is required on the part of the practitioner as to which blade should be introduced uppermost—an advantage of no great value, since no one should undertake a case of forceps delivery who has not sufficient knowledge of the operation and presence of mind enough to obviate any risk from the introduction of the wrong blade first. On account of its shortness and the want of the second or pelvic curve, it is only adapted for cases in which the head is low down in the pelvis or actually resting on the perineum.

The Pelvic Curve: its Advantages.—The question of the second or pelvic curve is one on which there is much difference of opinion. The forceps we are now considering—and the many modifications formed on the same plan—is constructed solely with reference to its grasp on the child's head, and without regard to the axes of the maternal passages. Consequently, were we to introduce it when the head was at the upper part of the pelvis, we could not fail to expose the soft parts to the risk of contusion and (in consequence of the necessity of drawing more directly backward) unduly stretch and even lacerate the perineum. Hence it is now admitted by obstetricians, with few exceptions, that the second curve is essential before the complete descent of the head, although it is not absolutely so after this has taken place. The only circumstances under which a straight blade can possess any superiority are in certain cases of occipito-posterior position in which it is found necessary to rotate the head round a large extent of the pelvis, when the circular sweep of a strongly-curved instrument might prove injurious. Such cases, however, are of rare occurrence, and need in no way influence the general employment of the pelvic curve.


Ziegler's Forceps.—The short forceps usually employed in Scotland is the invention of the late Dr. Ziegler (Fig. 159), and is useful from the

FIG. 158.



Denman's Short Forceps.

FIG. 159.




Ziegler's Forceps.

facility with which the blades may be introduced in accurate apposition to each other—a point which in practice is of no little value. In general size and appearance it closely resembles Denman's forceps, but the fenestrum of the lower blade is continued down to the handle. In introducing, the lower blade is slipped over the handle of the other blade, already *in situ*, and thus it is guided with great certainty into a proper position, locking itself as it passes on. This instrument has the disadvantage of not having the second curve, but the facility of introduction has rendered it a great favorite with many who have been in the habit of employing it.

The Long Forceps.—For cases in which the head is not on the perineum, or at least not quite low in the pelvis, a longer instrument is essential. To meet this indication Smellie invented the long forceps, which, like the shorter instrument, has been very variously modified. The most perfect instrument of the kind employed in this country is that known as Simpson's forceps (Fig. 160), which combines many excellent points selected from the forceps of various obstetricians, as well as some original additions, and which, as a whole, has never been surpassed until Tarnier's or its modification was invented. The curved portions of the blades are $6\frac{1}{4}$ inches long, the fenestrum measuring $1\frac{1}{4}$ at its widest part. The

FIG. 160.



Simpson's Forceps.

extremities of the blades are 1 inch asunder when the handles are closed, and 3 inches at their widest part. The object of this somewhat unusual width is to lessen the compressing power of the instrument without in any way interfering with its action as a tractor. The pelvic curve is less than in most long forceps, so as to admit of the rotation of the head, when necessary, without the risk of injuring the maternal structures. Between the curve of the blade and the lock is a straight portion or shank measuring $2\frac{3}{8}$ inches, which, before joining the handle, is bent at right angles into a knee. This shank is a useful addition to all forceps, and is essential in the long forceps to ensure the junction of the blades beyond the parts of the mother, which might otherwise be caught in the lock and injured. The knees serve the purpose of preventing the blades from slipping from each other after they have been united. They also admit of one finger being introduced above the lock and used as an aid in traction—a provision which is made in some other varieties of long forceps by a semicircular bend in each shank. The handles, which in most British forceps are too small and smooth to afford a firm grasp, are

serrated at the edge and flattened from before backward, so as to fit the closed fist more accurately. At their extremities, near the lock, there are a pair of projecting rests, over which the fore and middle fingers may be passed in traction, and which greatly increase our power over the instrument. Although this and other varieties of the long forceps are specially constructed for application when the head is high in the pelvis, it answers quite as well as the short forceps—indeed, in most respects better—when the head has descended low down. It is a decided advantage for the practitioner to habituate himself to the use of one instrument, with the application and power of which he becomes thoroughly familiar. It is a mere waste of space and money for him to encumber himself with a number of instruments of various shapes and sizes; and he may be sure that a good pair of long forceps will be suitable for every emergency and in any position of the head.

Disadvantages of a Weak Instrument.—The chief argument against the use of such an instrument in simple cases is its great power. This, however, is entirely based on a misconception. The existence of power does not involve its use, and the stronger instrument can be employed with quite as much delicacy and gentleness as the weaker. The remarks of Dr. Hodge¹ on this point are extremely apposite and are well worthy of quotation. He says: "Certainly, no man ought to apply the forceps who has not sufficient discretion to use no more force than is absolutely requisite for safe delivery. If, therefore, there is more power at command, he is not obliged to use it; while, on the contrary, if much power be demanded, he can, within the bounds of prudence, exercise it by the long forceps, but with the short forceps his efforts might be unavailing. Moreover, in cases of difficulty, the short forceps being used, the practitioner would be forced to make great muscular efforts; while with the long forceps, owing to the great leverage, such effort will be comparatively trifling, and of course the whole force demanded can be much more delicately, and at the same time efficiently, applied, and with more safety to the tissues of the child and its parent."

Continental Forceps.—The forceps usually employed on the Continent and in America differs considerably, both in appearance and construction, from those in use in this country. As a rule, it is a larger and more powerful instrument, joined by a pivot or button joint, and it always possesses the second or pelvic curve. Of late years Simpson's forceps has been much employed in some parts of Germany. The chief objection to the continental instruments is their cumbrousness. This is chiefly in the handles, which in many of them are forged in a piece with the blades, the part introduced within the maternal structures not being materially different from the corresponding part of the English instrument.


Turnier's Forceps.—The forceps invented by Prof. Tarnier (Fig. 161) has recently attracted considerable attention. In this instrument traction is not made on the handles by which the blades are introduced, as in ordinary forceps, but on a supplementary handle (*a*) subsequently attached to the blades near the lower opening of the fenestræ (*b*). The object claimed for this arrangement is that less force is required in trac-

¹ *System of Obstetrics*, p. 242.

tion, which can always be made in the proper axis of the pelvis; that the blades are not likely to slip; and that rotation of the head is not interfered with. The handles of the forceps, moreover, guide the operator to the direction in which he ought to pull, since all that is required is to keep the traction-rods parallel to them. This instrument, however, although theoretically perfect, is somewhat too complicated for general use.

Simpson's Axis-Traction Forceps.—Professor Simpson of Edinburgh has invented a modification of Tarnier's instrument, which he calls the

FIG. 161.



Tarnier's Forceps.

FIG. 162.



Simpson's Axis-Traction Forceps.
a, b. Traction handle. c, f. Line of traction.

“axis-traction forceps” (Fig. 162). The supplementary handles are fixed to the blades, and the whole mechanism is much simpler than in Tarnier's forceps. Dr. Simpson reports very favorably of this forceps, and it is certainly well adapted for the object aimed at. For some years I have used it extensively, and have every reason to be satisfied with it, especially in the high forceps operation, in which it seems to me superior to any other instrument.

Action of the Instrument.—The forceps is generally said to act in three different ways:

- 1st. As a tractor;
- 2d. As a lever;
- 3d. As a compressor.

The Chief Use of the Forceps is as a Tractor.—It is more especially as a tractor that the instrument is of value, and it is used with the greatest advantage when it is employed merely to supplement the action of the uterus, which is insufficient of itself to effect delivery, or when, from some complication, it is necessary to complete labor with greater rapidity than can be accomplished by the unaided powers of nature. In

most cases traction alone is sufficient ; but in order that it may act satisfactorily, and that the instrument may not slip, a proper construction of the forceps and a sufficient curvature of the blades are essential. The want of these is the radical fault of many of the short, straight instruments in common use, which have a tendency to slip during our efforts at extraction.

As a Lever.—The forceps acts also as a lever, but this action has been greatly exaggerated. It is generally described as a lever of the first class, the power being at the handles, the fulcrum at the lock, and the weight at the extremities. There may possibly be some leverage power of this kind when the instrument is first introduced and the handles held so loosely that one blade is able to work on the other. But, as ordinarily used, the handles are held with a sufficiently firm grasp to prevent this movement, and then the two blades practically form a single instrument.

Galabin, who has studied this subject in detail, points out¹ that—"1. The lever is formed by both blades of the forceps and the fetal head united in one immovable mass. As soon as the blades begin to slip over the head the lever is decomposed, and the swaying movement ceases to have any mechanical advantage. 2. The power is applied to the handles in a slanting direction. The resistance or weight does not act at a point either between the former and the fulcrum or beyond the fulcrum, but at a point in a plane nearly at right angles to the line joining these two points, and its direction is a line perpendicular to that plane of the pelvis in which the greatest section of the head is engaged ; that is to say, in the case of straight forceps nearly parallel to the handles. The lever formed does not, therefore, strictly speaking, belong to any one of the three orders into which levers are commonly divided. 3. The fulcrum is fixed partly by friction, partly by the combination of traction with oscillatory movements—in other words, by the power being directed in great measure downward and only slightly to one side."

He further shows that the pendulum motion of the forceps is superfluous in all ordinary forceps operations in which traction alone is amply sufficient for delivery, but that when the head is impacted and great force is required for its extraction, a mechanical advantage may be gained from having recourse to an oscillatory movement, which should, however, be very limited, and only continued if found to effect distinct advance of the head.

As a Compressor.—Regarding the compressive power of the instrument there has been much difference of opinion. There is no doubt that the forceps, especially some of the foreign instruments in which the points nearly approach each other, is capable of exerting considerable compression on the head. It is, however, extremely problematical if this action be of real value. It is to be borne in mind that in cases of protracted labor the head has been already moulded and compressed, and the bones have been made to overlap each other to their utmost extent, by the sides of the pelvis. We can scarcely, therefore, expect to diminish the head much more by the forceps without employing an amount of

¹ Galabin, "Action of Midwifery Forceps as a Lever," *Obstetrical Journal*, November, 1876.

force that will seriously endanger the life of the child. It is in cases of disproportion between the head and the pelvis, depending on slight antero-posterior contraction of the pelvic brim, that diminution of the child's head by compression would be most useful. Then, however, the pressure of the forceps is exerted on that portion of the head which lies in the most roomy diameter of the pelvis, where there is no want of space. If this pressure do not increase the opposite diameter, which is in apposition to the narrower portion of the pelvis, it can at least do nothing toward lessening it, and diminution of any other part of the child's head is not required.

Dynamical Action of the Forceps.—The mere introduction of the forceps sometimes excites increased uterine action through the reflex irritation induced by the presence of a foreign body in the vagina. This has been called the dynamical action of the forceps, but it cannot be looked upon in any other light than that of an occasional accidental result.

The circumstances indicating the use of the forceps have been separately considered elsewhere, and to recapitulate them here would only lead to needless repetition. I shall therefore now merely describe the mode of using the instrument.

Difference between the High and Low Operations.—Before doing so it is well to repeat what has already been said as to the difference between what may be termed the high and low forceps operations. The application of the instrument when the head is low in the pelvis is extremely simple, and when there is no disproportion between the head and the pelvis, and some slight traction is alone required to supplement deficient expulsive power, the operation in the hands of any ordinary well-instructed practitioner ought to be perfectly safe both to the mother and child. It is very different when the head is arrested at the brim or high in the pelvis. Then the application of the forceps is an operation requiring much dexterity for its proper performance, and must never be undertaken without anxious consideration. It is because these two classes of operations have been confused that the use of the instrument is regarded by many with such unreasonable dread.

Preliminary Considerations.—Before attempting to introduce the forceps there are several points to which attention should be directed :

1st. The membranes must, of course, be ruptured.

2dly. For the safe and easy application of the instrument it is also advisable that the os should be fully dilated and the cervix retracted over the head. Still, these two points cannot be regarded, as many have laid down, as being *sine quâ non*. Indeed, we are often compelled to use the instrument when, although the os is fully dilated, the rim of the cervix can be felt at some point of the contour of the head, especially in cases in which the anterior lip is jammed between the head and the pubes. Provided due care be taken to guard the cervical rim with the fingers of one hand as the instrument is slipped past it, there need be no fear of injury from this cause. If the os be not fully dilated, but is sufficiently open to admit of the passage of the forceps, the operation, under urgent circumstances, may be quite justifiable, but it must necessarily be a somewhat anxious one.

3dly. The position of the head should be accurately ascertained by means of the sutures and fontanelles. Unless this be done, the operation will always be haphazard and unsatisfactory, as the practitioner can never be in possession of accurate knowledge of the progress of the case. It may be that the occiput is directed backward, and, although that does not contraindicate the application of the forceps, it involves special precautions being taken.

4thly. The bladder and bowels should be emptied.

Question of Administering Anæsthetics.—Before proceeding to operate the question of anæsthesia will arise. In any case likely to be difficult it is of the greatest assistance to have the patient completely under the influence of an anæsthetic to the surgical degree, so as to have her as still as possible; but, whenever this is deemed necessary, another practitioner should undertake the responsibility of the administration. In simple cases I believe it is better to dispense with anæsthetics altogether—partly because they are apt to stop what pains there are, which is in itself a disadvantage, but chiefly because under partial anæsthesia the patient loses her self-control, is restless, and twists herself into awkward positions, which give rise to the utmost difficulty and inconvenience in the use of the instrument. Moreover, if no anæsthetic be given the patient can assist the operator by placing herself in the most convenient attitude.

Description of the Operation.—In describing the method of applying the forceps I shall assume that we have to do with the simpler variety of the operation, when the head is low in the pelvis. Subsequently I shall point out the peculiarities of the high operation.

Position of the Patient.—As to the position of the patient, I believe there can be no doubt of the superiority of that which is usually adopted in this country. On the Continent and in America the forceps is always employed with the patient lying on her back—a position involving much needless exposure of the person and requiring more assistance from others. In certain cases of unusual difficulty the position on the back is of unquestionable utility, but we may at least commence the operation in the usual way, and subsequently turn the patient on her back if desirable.

Importance of Suitable Position.—Much of the facility with which the blades are introduced depends on the patient's being properly placed. Hence, although it gives rise to a little more trouble at first, I believe that it is always best to pay particular attention to this point, whether the high or low forceps operation be about to be performed. The patient should be brought quite to the side of the bed, with her nates parallel to and projecting somewhat over its edge. The body should lie almost directly across the bed, and nearly at right angles to the hips, with the knees raised toward the abdomen (Fig. 163). In this way there is no risk of the handle of the upper blade, when depressed in introduction, coming in contact with the bed.


The blades should be warmed in tepid water, lubricated with cold cream or carbolic oil, and placed ready to hand.

These preliminaries having been attended to, we proceed to the introduction of the blades, sitting by the side of the bed opposite the nates of the patient.

Direction in which the Blades are to be Introduced.—The important

question now arises, In what direction are the blades to be passed? The almost universal rule in our standard works is, that they must be passed as nearly as possible over the child's ears, without any reference to the pelvic diameters. Hence, if the head have not made its turn, but is lying in one oblique diameter, the blades would require to be passed in

FIG. 163.



Position of Patient for Forceps Delivery, and Mode of Introducing Lower Blade.


the opposite oblique diameter; in short, the position of the forceps, as regards the pelvis, must vary according to the position of the head. Some have even laid down the rule that the forceps is contraindicated unless an ear can be felt—a rule that would very seriously limit its application, as in many cases in which it is urgently required it is a matter of great difficulty, and even impossibility, to feel the ear at all. It is admitted that in the high forceps operation the blades must be introduced in the transverse diameter of the pelvis, without relation to the position of the head. On the Continent it is generally recommended that this rule should be applied to all cases of forceps delivery alike, whether the head be high or low; and I have now for many years adopted this plan, and passed the blades in all cases, whatever be the position of the head, in the transverse diameter of the pelvis, without any attempt to pass them over the bi-parietal diameter of the child's head. Dr. Barnes points out with great force that, do what we will and attempt as we may to pass the blades in relation to the child's head, they find their way to the sides of the pelvis, and that the marks of the fenestra on the head always show that it has been grasped by the brow and side of the occiput. Of the perfect correctness of this observation I have no doubt; hence it is a needless element of complexity to endeavor to vary the position of the blades in each case, and one which only confuses the inexperienced practitioner and renders more difficult an operation which should be simplified as much as possible. While, therefore, it is of importance that the precise position of the head should be ascer-

tained in order that we may have an intelligent notion of its progress, I do not think that it is essential as a guide to the introduction of the forceps.

Method of Introducing the Lower Blade.—As a rule, the lower blade, lightly grasped between the tips of the index and middle fingers and thumb, should be introduced first. Poised in this way, we have perfect command over it, and can appreciate in a moment any obstacle to its passage. Two or more fingers of the left hand are introduced into the vagina and by the side of the head as a guide. The greatest care must be taken, if the cervix be within reach, that they are passed within it, so as to avoid the possibility of injury.

Necessity of Gentleness in Passing the Instrument.—The handle of the instrument has to be elevated, and its point slid gently along the palmar surface of the guiding fingers until it touches the head (Fig. 163). At first the blade should be inserted in the axis of the outlet, but as it progresses the handle must be depressed and carried backward. As it is pushed onward it is made to progress by a slight side-to-side motion, and it is of the utmost importance to bear in mind that the greatest gentleness must always be used. If any obstruction be felt, we are bound to withdraw the instrument partially or entirely, and attempt to manœuvre, not force, the point past it. As the blade is guided on in this way it is made to pass over the convexity of the head, the point being always kept slightly in contact with it, until it finally gains its proper position. When fully inserted the handle is drawn back toward the perineum and

FIG. 164.



Introduction of the Upper Blade.

given in charge to an assistant. The insertion must be carried on only in the intervals between the pains, and desisted from during their occurrence; otherwise there would be a serious risk of injuring the soft parts of the mother.


Introduction of the Upper Blade.—The second blade is passed directly

opposite to the first, and is generally somewhat more difficult to introduce in consequence of the space occupied by the latter. It is passed along two fingers directly opposite the first blade, and with exactly the same precautions as to direction and introduction, except that at first its handle has to be depressed instead of elevated (Fig. 164).

Locking of the Handles.—The handle which was in charge of the assistant is now laid hold of by the operator, and the two handles are drawn together. If the blades have been properly introduced, there should be no difficulty in locking; but should we be unable to join them easily, we must withdraw one or other, either partially or entirely, and reintroduce it with the same precautions as before. We must also assure ourselves that no hairs nor any of the maternal structures are caught in the lock.

Method of Traction.—When once the blades are locked we may commence our efforts at traction. To do this we lay hold of the handles

FIG. 165.



Forceps in Position : Traction in the Axis of the Brim, Downward and Backward.

with the right hand, using only sufficient compression to give a firm grasp of the head and to keep the blades from slipping. The left hand may be advantageously used in assisting and supporting the right during our efforts at extraction, and at a late stage of the operation may be employed in relaxing the perineum when stretched by the head of the child. Traction must always be made in reference to the pelvic axes, being at first backward toward the perineum (Fig. 165) in the direction of the axis of the brim, and as the head descends and the vertex protrudes through the vulva it must be changed to that of the outlet (Fig. 166). If the axis-traction forceps is used, it is to be borne in mind that traction is to be made by the traction-handle only, the handles of the instrument itself being left untouched after they are locked and the traction-

rods are united. By keeping these latter parallel to the handles of the forceps traction can always be made in the proper direction. We must extract only during the pains, and if these should be absent we must imitate them by acting at intervals. This is a point which deserves special attention, for there is no more common error than undue hurry in delivery.

The only valid objection I know of against a more frequent resort to the forceps in lingering labor is that the sudden emptying of the uterus in the absence of pains may predispose to hemorrhage; but it cannot be denied that it is one of some weight. However, if due care be taken to operate slowly, and to allow several minutes to elapse between each tractive effort, while at the same time uterine contractions be stimulated by pressure and support, this need not be considered a contraindication. Besides direct traction we may impart to the instrument a gentle waving motion from handle to handle, which brings into operation its power as a lever, but this must be done only to a very slight extent, and must always be subservient to direct traction.

Descent of the Head.—Proceeding thus in a slow and cautious manner, carefully regulating the force employed according to the exigencies of

FIG. 166.



Last Stage of Extraction. The Handles of the Forceps are being Gradually turned Upward toward the Mother's Abdomen.

the case, we shall perceive that the head begins to descend; and its progress should be determined from time to time by the fingers of the un-employed hand.

The Rotation from the Oblique Diameter.—When the head lies in the oblique diameter, as it descends, in consequence of its perfect adaptation

to the pelvic cavity, it will turn into the antero-posterior diameter without any effort on the part of the operator, provided only that the traction be sufficiently slow and gradual. As the head is about to emerge it is necessary to raise the handles toward the mother's abdomen. More than usual care is required to prevent laceration of the perineum, which is always much stretched (Fig. 166). If, as often happens, the pains have now increased, and the perineum be very thin and tense, it may even be desirable to remove the blades gently, and leave the case to be terminated by the natural powers; but if due precautions are used this need not be necessary.

The peculiarities of forceps delivery in occipito-posterior positions have already been discussed (p. 320), and need not be repeated.

High Forceps Operations.—When the high forceps operation has been decided on, the passage of the blades will be found to be much more difficult from the height of the presenting part, the distance which they must pass, and in some cases from the mobility of the head interfering with their accurate adaptation. The general principles of introduction and of traction are, however, identical. If the operation be attempted before the head has entered the pelvic brim, it must be fixed, as much as possible, by abdominal pressure. In guiding the blades to the head special care must be taken to avoid any injury of the soft parts, especially if the cervix be not completely out of reach. For this purpose it may even be advisable to introduce the entire left hand as a guide, so as to avoid any possibility of injuring the cervix from not passing the instrument under its edge.

Peculiar Method of Introducing the Blades.—Some authors advise that in such cases the blade should be introduced at first opposite the sacrum until the point approaches its promontory. It is then made to sweep round the pelvis, under the protecting fingers, till it reaches its proper position on the head. This plan is advocated by Ramsbotham, Hall Davis, and other eminent practical accoucheurs, and it is certainly of service in some cases of difficulty, especially when, from any reason, it is not possible to draw the nates over the edge of the bed, when the necessary depression of the handle of the upper blade is difficult to effect. It involves, however, a somewhat complicated manœuvre, and it is seldom that the blades cannot be readily introduced in the usual way.

Necessity of Care in Locking.—In locking, the slightest approach to roughness must be carefully avoided, for the extremities of the blades are now within the cavity of the uterus, and serious injury might easily be inflicted. If difficulty be met with, rather than employ any force one of the blades should be withdrawn and reintroduced in a more favorable direction. If the blades have shanks of sufficient length, there should be no risk of including the soft parts of the mother in the lock; which in a badly-constructed instrument is an accident not unlikely to occur.

Method of Traction.—After junction, traction must at first be altogether in the axis of the brim, and to effect this the handles must be pressed well backward toward the perineum. As the head descends it will probably take the usual turn of itself, without effort on the part of the operator, and the direction of the tractive force may be gradually

altered to that of the axis of the outlet. If the pains be strong and regular and there be no indication for immediate delivery, we may remove the forceps after the head has descended upon the perineum, and leave the conclusion of the case to nature. This course may be especially advisable if the perineum and soft parts be unusually rigid, but generally it is better to terminate labor without removing the instrument.

Possible Dangers of Forceps Delivery.—Before concluding this subject reference may be made to the possible dangers of the operation. I would here again insist on the importance of distinguishing between the high and low forceps operations, which have been so unfortunately and unfairly confounded. Reasons have already been given for rejecting the statistics of the risks attending forceps delivery in the latter class of cases (p. 347). A formidable catalogue of dangers, both to mother and child, might easily be gathered from our standard works on obstetrics. Among the former, the principal are lacerations of the uterus, vagina, and perineum; rupture of varicose veins, giving rise to thrombus; pelvic abscess, from contusion of the soft parts; subsequent inflammation of the uterus or peritoneum; tearing asunder of the joints and symphyses; and even fracture of the pelvic bones. A careful analysis of these, such as has been so well made by Drs. Hicks and Philips,¹ proves beyond doubt that the application of the instrument is not so much concerned in their production as the protraction of the labor and the neglect of the practitioner in not interfering sufficiently soon to prevent the occurrence of the evil consequences, afterward attributed to the operation itself. Many of these will be found to arise from the prolonged pressure on the soft parts within the pelvis, and the subsequent inflammation or sloughing. To these causes may be referred with propriety most cases of vesico-vaginal fistula (p. 439), peritonitis, and metritis following instrumental labor.

Some Depend on Ignorance on the Part of the Practitioner.—Lacerations and similar accidents may, however, result from an incautious use of the instrument. Slight lacerations of the mucous membrane of the vagina are probably far from uncommon. But if these cases were closely examined, it would be found that the fault lay not in the instrument, but in the hand that used it. Either the blades were introduced without due regard to the axes of the pelvis, or they were pushed forward with force and violence, or an instrument was employed unsuitable to the case (such as a short straight forceps when the head was high in the pelvis), or undue haste and force in delivery were used. It would be manifestly unfair to lay the blame of such results upon the forceps, which in the hands of a more judicious and experienced practitioner would have effected the desired object with perfect safety. The instrument is doubtless unsafe in the hands of any one who does not understand its use, just as the scalpel or amputating-knife would be in the hands of a rash and inexperienced surgeon. The lesson to be learnt seems to be, clearly, not that the dangers should deter us from the use of the forceps, but that they should induce us to study more carefully the cases in which it is applicable and the method of using it with safety.

Possible Risks to the Child.—The dangers to the child are, principally,

¹ *Obst. Trans.*, vol. xiii.

lacerations of the integuments of the scalp and forehead; contusion of the face; partial but temporary paralysis of the face from pressure of a blade on the facial nerve; depression or fracture of the cranial bones; injury to the brain from undue pressure of the blades. These evils are of rare occurrence, and, when they do happen, generally result from improper management of the operation—such as undue compression, the use of improper instruments, or excessive and ill-directed efforts at traction—and cannot, therefore, be considered as in any way contraindicating the use of the instrument. Many of the more common results, such as slight abrasions of the scalp or paralysis of the face, are transitory in their nature and of no real consequence.

[*The Forceps in America.*—Although the obstetrical forceps was first used in England, other countries in the march of improvement have made great changes, not only in the original forms, but in the manner of use; and various shapes, as well as different positions of the woman in application, have become in a measure national. With the exception of having adopted almost exclusively the French and German dorsal decubitus in making use of the instrument, we have become in a measure eclectic in the selection of the latter; medical schools, accoucheurs, and local obstetrical societies influencing students and the junior members of the profession to adopt the French, German, English, or American style, as the case may be, the forceps themselves bearing the names of the several inventors or compilers; for some are a true compilation—the blade from one contriver; fenestral openings, another; pelvic curve, a third; width, a fourth; shanks, a fifth; method of locking, a sixth; etc. etc. For this reason the late Prof. Hodge named his forceps the *eclectic*, although in some respects entirely original, particularly in the long superimposed shanks—a great improvement for operating at the superior strait and avoiding the painful stretching of the posterior commissure of the vulva. Dr. Hodge expended a great deal of thought and money in perfecting his forceps, and the various steps in the process were marked by a new form, until, from a heavy, clumsy instrument, he gradually evolved what was at one time regarded as a wonderful improvement upon the forceps of France and England.

A contemporary of Prof. Hodge, the late Prof. David D. Davis of London, was equally anxious to perfect the instrument, and turned his attention especially to making the blades light, open, and to fit the sides of the fetal head so as to enable traction to be made without much pressure or leaving any mark on the child's scalp. There is a principle of mechanics involved in his instrument which he studied to perfect by moulding the blades so as to obtain considerable coaptating surface, and thus by increase of friction to avoid undue and dangerous pressure. The Davis blade soon began to effect changes in the form of American forceps, and by the addition of long handles and some alterations of shape, weight, and curve became a leading feature in those bearing the names of William Harris, Prof. Wallace of the Jefferson Medical College, Dr. Bethel, and Albert H. Smith, all of this city. The short Davis instrument was a great favorite with the late Prof. Meigs and Dr. William Harris, both largely engaged in obstetrical practice as well as teaching; and many a delicate woman with wasting forces was aided in

her delivery at their hands, and was surprised to find no mark on the baby's head, and that her own sufferings could be so gently and safely relieved.

Although such was the estimation of the Davis blade, and still is in many parts of our country, it does not appear to have retained its popularity or been adopted, as its mechanical perfection would lead one who appreciates it to suppose it would have been. In Great Britain the favorite forms now in use are but a very slight improvement upon the forceps of a hundred years ago except in finish and material, the open fenestræ and bevelled blades of Davis being declined in favor of the looped fenestræ and flat-edged blades in use when he made his experiments and changes. This appears to have grown out of a practice which has been largely adopted in Germany, Great Britain, and many parts of the United States in applying the forceps to the foetal head, the blades being introduced at the sides of the pelvis, without much reference to the position which the head occupies. As compression is objected to, the blades are made long and widely separated ($3\frac{1}{4}$ to $3\frac{1}{2}$ inches), and the handles short, so as not to allow of much leverage. As the blades do not fit the head, the mechanism of labor as taught by Hodge has been much simplified, as it is not necessary to learn all the oblique fittings of the fenestræ over the parietal protuberances or ears. Dr. Meigs used to tell the students that the forceps was the "*child's instrument*," and should be used as a tractor; and it was as a well-applied mechanical tractor that he advocated the use of the Davis blades against those of Siebold, Levret, Baudelocque, and Haighton, employed generally in our country forty years ago. His language is not very complimentary to what he denominates by distinction "*the mother's instrument*," the form being better adapted for saving the woman than the foetus.¹

At the present day we have two general orders of forceps in use in the United States, under each of which may be placed a vast number of special varieties which are simply changes upon one or the other general type according to the fancy of the inventor. At the head of one type may be placed the long forceps of Prof. Hodge, designed to be adapted to the sides of the child's head in all possible cases; and of the other, those of Prof. Simpson of Edinburgh or their modification by Profs. Elliot and Bedford of New York, intended to be used as tractors, and applied in reference to the sides of the mother's pelvis, rather than to those of the infant's head.

Taking the long forceps of Levret and Baudelocque as improved and modified by Hodge, with the blades of Prof. Davis as a substitute, and handles of less curve than those of Hodge, and we have the long forceps of Prof. Ellerslie Wallace, late of the Jefferson Medical College, the most frequent choice of those who purchase forceps of the manufacturers in Philadelphia. Next in order are the instruments of Hodge, Davis, and Simpson, Elliot, Bedford, and a few others—in all about a dozen forms that vary in popularity. The improvement of the late Prof. Elliot upon the instrument of Simpson consists in narrowing and lengthening the shanks, widening somewhat the fenestræ, elongating the blades, giving greater security against slipping in the handles, and gauging the dis-

[¹ *Obstetrics*, p. 540.]

tance between the blades by a milled-head screw-stop in the end of the handles: the shanks and blades are an exact counterpart of the Miller forceps of England, which appeared about the same time (1858).

The *Hodge forceps* was based in its contrivance upon the following points: 1. The instrument should be shaped to the contour of the foetal head, and have sufficient play to allow of compression where the pelvis is too narrow for the head to pass in its normal condition. 2. The blades should be so arranged in reference to the shanks and handles as to enable them to seize the head of the fœtus in its bi-parietal diameter at the superior strait, and be drawn upon in the direction of the curve of the

FIG. 167.



Hodge Forceps.

FIG. 168.



Wallace Forceps.

FIG. 169.



Davis Forceps.

pelvic canal until the delivery is complete. 3. The long forceps ought to be competent to act either at the superior strait of the pelvis, in its cavity or at its outlet, so as to avoid a multiplicity of instruments and their attendant expense. And, 4. The instrument should not cut the scalp of the child if properly adjusted, or injure the soft parts of the mother.

It would be folly to claim that all this could or has been accomplished, as there must necessarily be exceptional cases in all the points given; hence the contrivance of the forceps of Tarnier and Cleemann for

certain presentations above the superior strait, and the long and short convertible instruments of a few inventors. There are many cases of labor in the higher walks of life where, although there is no obstruction, still the women require manual or instrumental assistance, as they cannot deliver themselves for want of sufficient contractile muscular force. Such women require that the forceps used should be easily introduced—should act simply as tractors, control the movement of the fetal head by being well fitted to its shape, and leave no effect upon the scalp or vulva. Although these requisites may be filled by the Hodge instrument, it is this class of cases that has demanded a lighter and more roomy pair of forceps, such as that devised by Davis.

As the teaching of the Jefferson Medical College under Dr. Meigs favored, as we have stated, the forceps of Davis, so his successor, Prof. Wallace, in carrying out in a measure the same views, combined the blades of the Davis pattern with the long handles of Hodge in contriving what is known as the "*Wallace forceps*," now so much in use by the large number of graduates of this school. As compared with the Hodge instrument, it is 1 inch shorter (15 inches against 16); the blades are of the same length (6 inches); the fenestræ are more open; the shanks are only half the length, giving a much greater compressing power; and the handles are of the same measurement from pivot to hooks. Both have the Siebold lock, over which we believe the broad-topped button and notch to possess some advantages; and the Wallace is somewhat heavier than the Hodge, which should weigh 17 ounces.

The short Davis instrument made for Prof. Meigs under direction of the inventor weighed $10\frac{3}{4}$ ounces and measured 12 inches in length; fenestræ, 5 inches long, 2 inches wide; blades separated $2\frac{3}{4}$ inches; handles, $4\frac{1}{4}$ inches to lock, which was of the Smellie or English pattern. A recently-purchased pair in possession of the editor is $13\frac{1}{4}$ inches long, with 5-inch handles, a button lock, 2-inch close-set shanks, and $6\frac{1}{2}$ -inch blades. I believe the changes are decided improvements, especially the lock and elongated handles. It has answered admirably in adynamic cases, requiring only a few pounds of tractile assistance. The Davis blades have been added to long handles, and the whole made of steel and marvellously light, at the special request of a few accoucheurs, who wished them to aid in some cases of arrest at the perineum.

The late Prof. George T. Elliot of New York, who received much of his practical obstetrical training in the Dublin Lying-in Hospital, imbibed the teachings of the English school, and became impressed with the value of the system as taught by Simpson, upon the principle of whose forceps, modelled somewhat after that of the late Prof. Gunning S. Bedford of New York, he in 1858 presented to the medical profession the instrument that bears his name. The forceps of Prof. Bedford has a traction-ring on each side where the Elliot has a cornu, has a button joint, instead of a Smellie, has no screw-stop, and has diverging instead of superimposed shanks. These points have generally been considered as improvements, and hence the Elliot has taken precedence in large measure over the Bedford instrument in New York, the two being the leading forceps in demand. The instrument of White of Buffalo is

perhaps next, and then Hodge's. But few of Prof. Wallace's forceps, long the leading instrument in Philadelphia sales, are ordered. The *White* is a long forceps, a compound of the Elliot blade, long superimposed shanks of Hodge, Siebold lock, and short corrugated steel handles bowed out like dental forceps and ending in thin blunt hooks.

The Sawyer and Simpson short forceps are said to be about equally in demand in New York. The former is almost unknown in Phila-

FIG. 170.



Elliot Forceps.

delphia, and but comparatively few of the Simpson are asked for, although the system of their application has several advocates in this city.

The Sawyer Forceps.—This is the lightest of all the varieties of the short forceps, weighing but 5 ounces, and measuring $9\frac{3}{4}$ inches in length; the handle being 3 inches, shank $1\frac{1}{2}$, and chord of blade-curve $5\frac{1}{4}$. The blades are $1\frac{1}{2}$ inches wide, with oval fenestræ $\frac{7}{8}$ inch wide, and separated $2\frac{5}{8}$ inches at their widest part and $\frac{3}{4}$ inch at the tips. This instrument was invented eight years ago by Prof. Edw. Warren Sawyer of Rush Medical College, Chicago, and has been highly commended by Prof. Byford and others. The forceps has the blades of Davis, superimposed shanks of Hodge, and lock of Smellie, with hard-rubber plates moulded hot upon the handles. The several parts have been somewhat modified, the object being to secure a tractor for cases of deficient expulsive force where the foetal head is low in the pelvis.

FIG. 171.




Sawyer Forceps.

Professor Sawyer says: "In the labors to which my forceps is applicable it is not necessary for the operator's body to be in line with the pelvic axis. My mode of procedure is the following: The woman is placed upon her back and drawn to the edge of the bed; the outside leg is now flexed; beneath this flexed extremity and the bed-covering I apply the forceps—often using but one hand in the operation. When the instrument is locked, I grasp the handle in such a manner that the palm of the hand looks upward; one hook then rests naturally upon the extensor surface of the first phalanx of the index finger, while the other hook rests upon a corresponding part of the thumb. When thus adjusted, I lift the head from the pelvic outlet, at the same time invoking the pendulum movement if desired. At this moment the advantage of the hooked handle is very apparent to the operator." . . . "All practitioners must have often felt, during

the last moments of labor, when the uterus and the mother seemed fatigued, the need of a little help to the expansive powers. The ordinary instruments are too formidable to be used at the last moment, and it is then that this little forceps is useful."

I have given the names and characters of the various forceps most in use in New York and Philadelphia, and by the large number of graduates of their respective schools, as shown by their preferences in selecting instruments of the leading makers of the two cities. The mechanism of instrumental delivery is much simplified by applying the forceps to whatever parts of the foetal head may be opposite the sides of the pelvis; but it is very questionable whether it is the scientific method or the safer for the child. With one blade over the side of the occiput, and the other over that of the forehead—which is the manner of seizure in oblique positions of the vertex—we certainly have not a very secure

FIG. 172.



Application of the Forceps at the Inferior Strait.

hold and run some risk of injury to the foetus. The advocates of this system claim that they use no compression, only a simple traction; which may be true in one sense, but amounts to the same in effect, else how could Dr. Elliot, by traction with great force, straighten out one

to the pelvic cavity, it will turn into the antero-posterior diameter without any effort on the part of the operator, provided only that the traction be sufficiently slow and gradual. As the head is about to emerge it is necessary to raise the handles toward the mother's abdomen. More than usual care is required to prevent laceration of the perineum, which is always much stretched (Fig. 166). If, as often happens, the pains have now increased, and the perineum be very thin and tense, it may even be desirable to remove the blades gently, and leave the case to be terminated by the natural powers; but if due precautions are used this need not be necessary.

The peculiarities of forceps delivery in occipito-posterior positions have already been discussed (p. 320), and need not be repeated.

High Forceps Operations.—When the high forceps operation has been decided on, the passage of the blades will be found to be much more difficult from the height of the presenting part, the distance which they must pass, and in some cases from the mobility of the head interfering with their accurate adaptation. The general principles of introduction and of traction are, however, identical. If the operation be attempted before the head has entered the pelvic brim, it must be fixed, as much as possible, by abdominal pressure. In guiding the blades to the head special care must be taken to avoid any injury of the soft parts, especially if the cervix be not completely out of reach. For this purpose it may even be advisable to introduce the entire left hand as a guide, so as to avoid any possibility of injuring the cervix from not passing the instrument under its edge.

Peculiar Method of Introducing the Blades.—Some authors advise that in such cases the blade should be introduced at first opposite the sacrum until the point approaches its promontory. It is then made to sweep round the pelvis, under the protecting fingers, till it reaches its proper position on the head. This plan is advocated by Ramsbotham, Hall Davis, and other eminent practical accoucheurs, and it is certainly of service in some cases of difficulty, especially when, from any reason, it is not possible to draw the nates over the edge of the bed, when the necessary depression of the handle of the upper blade is difficult to effect. It involves, however, a somewhat complicated manœuvre, and it is seldom that the blades cannot be readily introduced in the usual way.

Necessity of Care in Locking.—In locking, the slightest approach to roughness must be carefully avoided, for the extremities of the blades are now within the cavity of the uterus, and serious injury might easily be inflicted. If difficulty be met with, rather than employ any force one of the blades should be withdrawn and reintroduced in a more favorable direction. If the blades have shanks of sufficient length, there should be no risk of including the soft parts of the mother in the lock; which in a badly-constructed instrument is an accident not unlikely to occur.

Method of Traction.—After junction, traction must at first be altogether in the axis of the brim, and to effect this the handles must be pressed well backward toward the perineum. As the head descends it will probably take the usual turn of itself, without effort on the part of the operator, and the direction of the tractive force may be gradually

altered to that of the axis of the outlet. If the pains be strong and regular and there be no indication for immediate delivery, we may remove the forceps after the head has descended upon the perineum, and leave the conclusion of the case to nature. This course may be especially advisable if the perineum and soft parts be unusually rigid, but generally it is better to terminate labor without removing the instrument.

Possible Dangers of Forceps Delivery.—Before concluding this subject reference may be made to the possible dangers of the operation. I would here again insist on the importance of distinguishing between the high and low forceps operations, which have been so unfortunately and unfairly confounded. Reasons have already been given for rejecting the statistics of the risks attending forceps delivery in the latter class of cases (p. 347). A formidable catalogue of dangers, both to mother and child, might easily be gathered from our standard works on obstetrics. Among the former, the principal are lacerations of the uterus, vagina, and perineum; rupture of varicose veins, giving rise to thrombus; pelvic abscess, from contusion of the soft parts; subsequent inflammation of the uterus or peritoneum; tearing asunder of the joints and symphyses; and even fracture of the pelvic bones. A careful analysis of these, such as has been so well made by Drs. Hicks and Philips,¹ proves beyond doubt that the application of the instrument is not so much concerned in their production as the protraction of the labor and the neglect of the practitioner in not interfering sufficiently soon to prevent the occurrence of the evil consequences, afterward attributed to the operation itself. Many of these will be found to arise from the prolonged pressure on the soft parts within the pelvis, and the subsequent inflammation or sloughing. To these causes may be referred with propriety most cases of vesico-vaginal fistula (p. 439), peritonitis, and metritis following instrumental labor.


Some Depend on Ignorance on the Part of the Practitioner.—Lacerations and similar accidents may, however, result from an incautious use of the instrument. Slight lacerations of the mucous membrane of the vagina are probably far from uncommon. But if these cases were closely examined, it would be found that the fault lay not in the instrument, but in the hand that used it. Either the blades were introduced without due regard to the axes of the pelvis, or they were pushed forward with force and violence, or an instrument was employed unsuitable to the case (such as a short straight forceps when the head was high in the pelvis), or undue haste and force in delivery were used. It would be manifestly unfair to lay the blame of such results upon the forceps, which in the hands of a more judicious and experienced practitioner would have effected the desired object with perfect safety. The instrument is doubtless unsafe in the hands of any one who does not understand its use, just as the scalpel or amputating-knife would be in the hands of a rash and inexperienced surgeon. The lesson to be learnt seems to be, clearly, not that the dangers should deter us from the use of the forceps, but that they should induce us to study more carefully the cases in which it is applicable and the method of using it with safety.

Possible Risks to the Child.—The dangers to the child are, principally,

¹ *Obst. Trans.*, vol. xiii.

will here be found valuable, as the lock is not introduced or the posterior commissure of the vulva widely stretched. If the head is entirely above the strait, the line of the blades must be changed correspondingly, in order to apply them properly and keep the line of traction within the

FIG. 174.



Direction of the Forceps as the Head is being Delivered.

coccyx; and even then, to draw in the proper direction, the left hand must act at first in a backward direction from the lock, while the right brings the handles downward, forward, and then upward; both hands describing a curve, but that of the right being much the greater. The peculiar forceps of Tarnier or of Cleemann, being designed to meet this form of exigency, may be brought into requisition. These both have the blades of Davis.

In latter years it has become much more common than formerly to introduce the forceps into the uterus before it is fully dilated, in consequence of the success claimed for the plan as carried out in the Dublin Lying-in Hospital. As this should never be done where the os is not readily dilatable, and requires much skill in execution, it is not safe to recommend its general adoption in cases of delay in private practice.

The forceps should not be introduced with any force, but the left blade should be slid in gently, and with a spiral motion, and then the right, care being taken that they should also lock without force, which they will do if properly adjusted. Traction is to be exerted slowly and dur-

ing a pain, the whole movement being made to correspond with the natural as closely as possible.

As the fetal head comes under the arch of the pubes the handles of the forceps must rise more and more from the bed, until at last they are over the abdomen as the head emerges from the perineum. This last movement of instrumental delivery should be a very slow one, for fear of rupture. It has been proposed to remove the blades before delivery is complete; but there is no occasion for this if the forceps is applied to the sides of the head over the parietal protuberances, as, where these protrude and the blades are flat and thin, there is very little additional space required. With such instruments as the old Levret, Baudelocque, and Rohrer forceps, with looped or kite-shaped fenestræ and thick edges, this was a much more imperative direction than with the better instruments of the present day. With a Sawyer forceps the perineum ought to be safer and under better control than without. When the perineum is thought to be in danger, the process of distension should be retarded through two or three pains, or even more if required, instead of drawing the head through at once.

After the head is delivered, if the cord is not around the neck, and therefore in danger from pressure, the body should be allowed to remain until the uterus has well contracted upon it, for fear of hemorrhage after delivery from uterine inertia.—ED.]

CHAPTER IV.

THE VECTIS.—THE FILLET.

The Vectis.—In connection with the subject of instrumental delivery it is essential to say something of the use of the *vectis*, on account of the value which was formerly ascribed to it, which was at one time so great in this country that it became the favorite instrument in the metropolis; Denman saying of it that even those who employed the forceps were "very willing to admit the equal, if not superior, utility and convenience of the *vectis*." Even at the present day there are practitioners of no small experience who believe it to be of occasional great utility, and use it in preference to the forceps in cases in which slight assistance only is required. In spite, however, of occasional attempts to recommend its use, the instrument has fallen into disfavor, and may be said to be practically obsolete.

Nature of the Instrument.—The *vectis*, in its most approved form, consists of a single blade, not unlike that of a short straight forceps, attached to a wooden handle. A variety of modifications exists in its shape and size. The handle has been occasionally manufactured for the convenience of carriage, with a hinge close to the *capitulum*.

blade (Fig. 175), or with a screw at the point where the handle and blade join. The power of the instrument and the facility of introduction depend very much on the amount of curvature of the blade. If

Fig. 175.



Vectis with Hinged Handle.

this be decided, a firmer hold of the head is taken and greater tractive force is obtained, but the difficulty of introduction is increased.

The Vectis is Used either as a Lever or a Tractor.—When employed in the former way the fulcrum is intended to be the hand of the operator; but the risk of using the maternal structures as a *point d'appui*, and the inevitable danger of contusion and laceration which must follow, constitute one of the chief objections to the operation. Its value as a tractor must always be limited and quite inferior to that of the forceps, while it is as difficult to introduce and manipulate.

Cases in which it is Applicable.—The vectis has been recommended in cases in which the low forceps operation is suitable, provided the pains have not entirely ceased. There is no doubt that it may be quite capable of overcoming a slight impediment to the passage of the head. It is applied over various parts of the head, most commonly over the occiput, in the same manner and with the same precautions as one blade of the forceps. Dr. Ramsbotham says, "We shall find it necessary to apply it to different parts of the cranium, and perhaps the face also, successively, in order to relieve the head from its fixed condition and favor its descent." Such an operation obviously requires quite as much dexterity as the application of the forceps; while, if we bear in mind its comparatively slight power and the risk of injury to the maternal structures, we must admit that the disuse of the instrument in modern practice is amply justified.

Is Sometimes of Value in Correcting Malpositions of the Head.—The vectis may, however, find a useful application when employed to rectify malpositions, especially in certain occipito-posterior presentations. This action of the instrument has already been considered (p. 320), and under such circumstances it may prove of service where the forceps is inapplicable. When so employed it is passed carefully over the occiput, and, while the maternal structures are guarded from injury, downward traction is made during the continuance of a pain. So used, its application is perfectly simple and free from danger, and for this purpose it may be retained as part of the obstetric armamentarium.

The Fillet.—The fillet is the oldest of obstetric instruments, having been frequently employed before the invention of the forceps, and even in the time of Smellie it was much used in the metropolis. It has since completely fallen out of favor as a scientific instrument, although its use is every now and again advocated; and it is certainly a favorite instrument with some practitioners. This is to be explained by the apparent simplicity of the operation and the fact that it can generally be performed without the knowledge of the patient. The latter, however, is one strong reason why it should not be used.


Nature of the Instrument.—The fillet consists, in its most improved

form (that which is recommended by Dr. Eardley Wilmot,¹ Fig. 176), of a slip of whalebone fixed into a handle composed of two separate halves, which join into one. The whalebone loop is slipped over either the occiput or face, and traction used at the handle.

Objections to its Use.—When applied over the face after the head has rotated it would probably do no harm, but if it were so placed when the head was high in the pelvis, traction would necessarily produce extension of the chin before the proper time, and would thus interfere with the natural mechanism of delivery. If placed over the occiput, it is impossible to make traction in the direction of the pelvic axes, as the instrument will then infallibly slip. If traction be made in any other direction, there must be a risk of injuring the maternal structures or of changing the position of the head. Hence there is every reason for discarding the fillet as a tractor or as a substitute for the forceps, even in the simplest cases.

Its Use in Certain Malpositions of the Head.—It is quite possible that it may find a useful application in certain cases in which the vectis may also be used—viz. as a rectifier of malposition; and from the comparative facility of its introduction it would probably be the preferable instrument of the two.

FIG. 176.



Wilmot's Fillet.

CHAPTER V.

OPERATIONS INVOLVING DESTRUCTION OF THE FŒTUS.

OPERATIONS involving the destruction and mutilation of the child were among the first practised in midwifery. Craniotomy was evidently known in the time of Hippocrates, as he mentions a mode of extracting the head by means of hooks. Celsus describes a similar operation, and was acquainted with the manner of extracting the fœtus in transverse presentations by decapitation. Similar procedures were also practised and described by Aëtius and others among the ancient writers. The physicians of the Arabian school not only employed perforators for opening the head, but were acquainted with instruments for compressing and extracting it.

Religious Objections to Craniotomy.—Until the end of the seventeenth

¹ *Obst. Trans.*, vol. xv.

century this class of operation was not considered justifiable in the case of living children; it then came to be discussed whether the life of the child might not be sacrificed to save that of the mother. It was authoritatively ruled by the Theological Faculty of Paris that the destruction of the child in any case was mortal sin: "Si l'on ne peut tirer l'enfant sans le tuer, on ne peut sans péché mortel le tirer." This dictum of the Roman Church had great influence on continental midwifery, more especially in France, where, up to a recent date, the leading obstetricians considered craniotomy to be only justifiable when the death of the fœtus had been positively ascertained. Even at the present day there are not wanting practitioners who in their praiseworthy objection to the destruction of a living child counsel delay until the child has died—a practice thoroughly illogical, and only sparing the operator's feelings at the cost of greatly-increased risk to the mother. In England the safety of the child has always been considered subservient to that of the mother; and it has been admitted that in every case in which the extraction of a living fœtus by any of the ordinary means is impossible its mutilation is perfectly justifiable.

Its Unjustifiable Frequency.—It must be admitted that the frequency with which craniotomy has been performed in this country constitutes a great blot on British midwifery. During the mastership of Dr. Labbat at the Rotunda Hospital the forceps was never once applied in 21,867 labors. Even in the time of Clarke and Collins, when its frequency was much diminished, craniotomy was performed three or four times as often as forceps delivery. These figures indicate a destruction of fœtal life which we cannot look back to without a shudder, and which, it is to be feared, justify the reproaches which our continental brethren have cast upon our practice. Fortunately, professional opinion has now completely recognized the sacred duty of saving the infant's life whenever it is practicable to do so; and British obstetricians now teach, as carefully as those of any other nation, the imperative necessity of using every endeavor to avoid the destruction of the fœtus.

Divisions of the Subject.—The operation now under consideration may be necessary—1st, when the head requires either to be simply perforated or afterward more completely broken up and extracted—an operation which has received various names, but is generally known in this country as *craniotomy*, and which may or may not require to be followed by further diminution of the trunk. 2dly, when the arm presents and turning is impossible. This necessitates one of two procedures—*disarticulation* with the separate extraction of the body and head, or *crisection*. In both classes of cases similar instruments are employed, and those generally in use at the present time may be first briefly described.


Description of Instruments Employed.—1. *Perforator.*—The object of the perforator is to pierce the skull of the child, so as to admit of the brain being broken up and the consequent collapse and diminution in size of the cranium. The perforator invented by Denman or some modification of it has been principally employed. It requires the handles to be separated in order to open the blades, and this cannot be done by the operator himself. This difficulty is overcome in the modification of Naegele's perforator used in Edinburgh, in which the handles are so

constructed that they open the points when pressed together, and are separated by a steel rod, with a joint at its centre to prevent their open-

FIG. 177.

FIG. 178.

FIG. 179.




Various Forms of Perforators.

ing too soon. By this arrangement the instrument can be manipulated by one hand only. The sharp-pointed portion has an external cutting edge, with projecting shoulders at its base to prevent its penetrating too far into the cranium. Many modifications of these arrangements have since been contrived (Figs. 177, 178, 179). In some parts of the Continent a perforator is used constructed on the principle of the trephine; but this is vastly more difficult to work, and has the great disadvantage of simply boring a hole in the skull, instead of splitting it up, as is done by the sharp-pointed instrument.

The instruments for extraction are the crotchet and craniotomy forceps.

Crotchets and Craniotomy Forceps.—The crotchet is a sharp-pointed hook of highly-tempered steel, which can be fixed on some portion of the skull, either internal or external, traction being made by the handle. The shank of the instrument is either straight or curved (Figs. 180 and 181), the latter being preferable, and it is either attached to a wooden handle or forged in a single piece of metal. A modification of this instrument is known as *Oldham's vertebral hook*. It consists of a slender hook, measuring, with its handle, 13 inches in length, which is passed through the foramen magnum and fixed in the vertebral canal, so as to secure a firm hold for traction. All forms of crotchets are open to the serious objection of being liable to slip or break through the bone to which they are attached, either

FIGS. 180, 181.



Crotchets.

the soft parts of the mother or the fingers of the operator placed as a guard. Hence they are discountenanced by most recent writers, and may with propriety be regarded as obsolete instruments.

Craniotomy Forceps are Preferable for Extraction.—Their place as tractors is well supplied by the more modern craniotomy forceps (Fig. 182). These are intended to lay hold of the skull, one blade being introduced within the cranium, the other externally, and when a firm grasp has been obtained downward traction is made. A second object it fulfils is to break away and remove portions of the skull when perforation and traction alone are insufficient to effect delivery. Many forms of craniotomy forceps are in use—some armed with formidable teeth; others, of simpler construction, depending on their roughened and serrated internal surfaces for firmness of grasp. For general use there is no better instrument than the *cranioclast* of Sir

James Simpson (Fig. 183), which admirably fulfils both these indications. It consists of two separate blades fastened by a button joint. The extremities of the blades are of a duck-billed shape, and are sufficiently curved to allow of a firm grasp of the skull being taken: the upper blade is deeply grooved to allow the lower to sink into it, and this gives the instrument great power in fracturing the cranial bones when that is found to be necessary. It need not, however, be employed for the latter purpose, and, the blades being serrated on their under surface, form as perfect a pair of craniotomy forceps as any in ordinary use. Provided with it, we are spared the necessity of procuring a number of instruments for extraction.



Craniotomy Forceps.

Simpson's Cranioclast.

Cephalotribe.—Amongst modern improvements in midwifery there are few which have led to more discussion than the use of the *cephalotribe*. This instrument, originally invented by Baudelocque, was long employed on the Continent before it was used in this country, the prejudice against it being no doubt due to its formidable size and appearance. Of late years many of our leading obstetricians have used it in preference either to the crotchet or craniotomy forceps, and have materially modified and improved its construction, so that the most objectionable features of the older instruments are now entirely removed.

Object of the Instrument.—The cephalotribe consists of two powerful solid blades, which are applied to the head after perforation and approximated by means of a screw so as to crush the cranial bones, and after this it may be also used for extraction. The peculiar value of the instrument is that when properly applied it crushes the firm basis of the skull, which is left untouched by craniotomy; or, if it does not,

it at least causes the base to turn edgeways within the blades, so as to be in a more favorable position for extraction. Another and specially valuable property is that it crushes the bones *within* the scalp, which forms a most efficient protective covering to their sharp edges. In this way one of the principal dangers of craniotomy—the wounding of the maternal passages by spiculæ of bone—is entirely avoided.

Some Obstetricians Object to Using it as a Tractor.—The cephalotribe, therefore, acts in two ways—as a crusher and as a tractor. Some obstetricians believe the former to be its more important use, and even maintain that the cephalotribe is unsuited for traction. This view is specially maintained by Pajot, who teaches that after the size of the skull has been diminished by repeated crushings its expulsion should be left to the natural powers. There are some grounds for believing that in the greater degrees of obstruction the tractile power of the instrument should not be called into use; but in the large majority of cases the facility with which the crushed head may be withdrawn by it constitutes one of its chief claims to the attention of the obstetrician. No one who has used it in this way, and experienced the rapid and easy manner in which it accomplishes delivery, can have any doubt on this point.

Its Value.—There is every reason to believe that cephalotripsy will be much extended in this country, and that it will be considered, as I believe it unquestionably deserves to be, the ordinary operation in cases requiring destruction of the fœtus. The comparative merits of cephalotripsy and craniotomy will be subsequently considered.

Description of the Instrument.—The most perfect cephalotribe is probably that known as Braxton Hicks' (Fig. 184), which is a modification of Simpson's. It is not of unwieldy size, but sufficiently powerful for any case, and not extravagant in price. The blades have a slight pelvic curve, which materially facilitates their introduction, yet not sufficiently marked to interfere with their being slightly rotated after application. Dr. Kidd of Dublin prefers a straight blade, while Dr. Matthews Duncan thinks it better to use a somewhat bulkier instrument, modelled on the type of the continental cephalotribes. The principle of action of all these is identical, and their differences are not of very material importance.

Section of the Skull by the Forceps Saw or Écraseur.—Another mode of diminishing the fetal skull is by removing it in sections. The object is aimed at in the *forceps saw* of Van Huevel,

FIG. 184.



Hicks' Cephalotribe.

which consists of two large blades not unlike those of the cephalotribe in appearance. Within these there is a complicated mechanism working a chain-saw from below upward, which cuts through the fetal skull; the separated portions are subsequently withdrawn piecemeal. This instrument is highly spoken of by the Belgian obstetricians, who believe that it affords by far the safest and most effectual way of reducing the bulk of the fetal skull. In this country it is practically unknown, and, although it must be admitted to be theoretically excellent, the complexity and cost of the apparatus have always stood in the way of its being used.

Dr. Barnes has suggested that the same results may be obtained by dividing the head with a strong wire *écraseur*. So far as I know, this suggestion has never yet been carried out in practice, not even by himself, and therefore it is not possible to say much about it. I should imagine, however, that there would be considerable difficulty in satisfactorily passing the loop of wire over the skull in a pelvis in which there is any well-marked deformity.

Cases Requiring Craniotomy.—The most common cause for which craniotomy or cephalotripsy is performed is a want of proper proportion between the head and the maternal passages. This may arise from a variety of causes. The most important, and that most often necessitating the operation, is osseous deformity. This may exist either in the brim, cavity, or outlet, and it is most often met with in the antero-posterior diameter of the brim. Obstetric authorities differ considerably as to the precise amount of contraction which will prevent the passage of a living child at term. Thus, Clarke and Burns believe that a living child cannot pass through a pelvis in which the antero-posterior diameter at the brim is less than $3\frac{1}{4}$ inches; Ramsbotham fixes the limit at 3 inches, and Osborne and Hamilton at $2\frac{3}{4}$ inches. The latter is the extreme limit at which the birth of a living child is possible; but there can be no doubt that under favorable circumstances it may be possible to draw the fœtus, after turning, through a pelvis of that size. The opposite limit of the operation is still more open to discussion. Various authorities have considered it quite possible to draw a mutilated fœtus through a pelvis in which the antero-posterior diameter does not exceed $1\frac{1}{2}$ inches, and, indeed, have succeeded in doing so. But then there must be a fair amount of space in the transverse diameter of the pelvis to admit of the necessary manipulations. If there be a clear space here of 3 inches and upward, it is no doubt possible to deliver *per vias naturales*; but in such extreme deformities the difficulties are so great, and the bruising of the maternal structures so extensive, that it becomes an operation of the greatest possible severity, with results nearly as unfavorable to the mother as the Cæsarean section. Hence some continental authorities have not scrupled to prefer the latter operation in the worst forms of pelvic deformity. The rule in English practice always has been that craniotomy must be performed whenever it is practicable; and there can be no doubt that it is the right one.

Limits of the Operation.—Between from $2\frac{3}{4}$ to 3 inches antero-posterior diameter in one direction, $1\frac{3}{4}$ inches in the other, may be said to be the limits of craniotomy, provided, in the latter case, there be a fair amount

of space in the transverse diameter. The same limits may be laid down with regard to tumors or other sources of obstruction.

Other Causes Justifying Craniotomy.—There are a few other conditions in which craniotomy is justifiable, independently of pelvic contraction, such as certain conditions of the soft parts which are supposed to render the passage of the head peculiarly dangerous to the mother. Among them may be mentioned swelling and inflammation of the vagina from the length of the previous labor, bands and cicatrices of the vagina, and occlusion and rigidity of the os. It is hardly too much to say that with a proper use of the resources of midwifery the destruction of a living fœtus for any of these conditions might be obviated. The most common of them is undoubtedly swelling of the soft parts causing impaction of the head—an occurrence which ought to be invariably prevented by a timely use of the forceps. Should interference unfortunately be delayed until impaction has actually taken place, doubtless no other resource but craniotomy would be left; but such cases, it is to be hoped, are now of rare occurrence in British practice. Undue rigidity of the os can be overcome by dilatation with the caoutchouc bags, or, in more serious cases, by incision, which would certainly be less perilous to the mother than dragging even a mutilated fœtus through the small and rigid aperture. In the case of bands and cicatrices in the vagina, dilatation or incision will generally suffice to remove the obstruction; but even were this not so, here, as in excessive rigidity of the perineum, it would be better that slight lacerations should take place than that the child should be killed.

Certain Complications of Labor are held to Justify Craniotomy.—Certain complications of labor are held to justify craniotomy, such as rupture of the uterus, convulsions, and hemorrhage. The application of the forceps or turning will generally answer our purpose equally well, especially as we have the means of dilating the os sufficiently to admit of one or other of them being performed when the natural dilatation is not sufficient. Craniotomy in rupture of the uterus will also be rarely indicated, as we have seen that gastrotomy appears to afford a better chance to the mother in those cases in which the fœtus has partially or entirely escaped from the uterine cavity.

Excessive Size of the Fœtus may require the Operation.—Want of proportion between the fœtus and the pelvis, depending on undue size of the head, either natural or the result of disease, may render the operation essential. In the former of these cases we shall generally have first attempted delivery with the forceps, and if it has failed there can be no doubt as to the propriety of lessening the bulk of the head by perforation.

Craniotomy when the Child is believed to be Dead.—In most obstetric works we are recommended to perforate rather than apply the forceps when we are convinced that the child has ceased to live. This advice is based on the greater facility with which craniotomy can be performed and its supposed greater safety to the mother. There can be no doubt of the ease with which the child can be extracted after perforation when the pelvis is not contracted; and, if we could always be sure of our diagnosis, the rule might be a good one. Before acting on it, however, we must bear in mind the extreme difficulty of positively ascertaining

the death of the fœtus. Of the signs usually relied on for this purpose there are scarcely any which are not open to fallacy except peeling of the scalp and disintegration of the cranial bones, which do not take place unless the child has been dead for a length of time, and are therefore useless in most instances. Discharge of the meconium constantly takes place when the child is alive; a cold and pulseless prolapsed cord may belong to a twin; and the fœtal heart may become temporarily inaudible although the child is not dead. If, indeed, we have carefully watched the fœtal heart all through the labor, and heard it become more and more feeble, and finally stop altogether, we might be certain that the child has died; but surely such observations would rather indicate an early recourse to the forceps or version, so as to obviate the fatal result we know to be impending.

Perforation of the After-coming Head.—In certain breech presentations or after turning it may be found impossible to extract the head without diminishing its size by perforating behind the ear. In such cases we know to a certainty whether the child be alive or dead before resorting to the operation.

Perforation is an Essential Preliminary both in Craniotomy and Cephalotripsy.—The first step, whether we resort to cephalotripsy or craniotomy, is perforation, which will therefore be first described. In the former the desirability of first perforating the head is not always recognized. To endeavor to crush the head without perforating is needlessly to increase the difficulties of the case, and it should be remembered, as a cardinal rule, that perforation is an essential preliminary to the proper use of the cephalotribe.

Method of Perforation.—Before perforating we must carefully ascertain the exact relation of the os to the presenting part, since in many cases the operation is performed before the os is fully dilated, when there is a risk of wounding the cervix. Two or more fingers of the left hand should be passed up to the head and placed against the most prominent part of the parietal bone. Under these, used as guard (Fig. 185), the perforator should be cautiously introduced until the scalp is reached. It is important to fix on a bony part of the skull, and not on a suture or fontanelle, for puncture, because our object is to break up the vault of the cranium as much as possible, so as to allow the skull to collapse. When the instrument has reached the point we have selected, it should be made to penetrate the scalp and skull with a semi-rotatory boring motion, and advanced until it has sunk up to the rests, which will oppose its further progress. Occasionally considerable force will be necessary to effect penetration, more especially if the scalp be swollen by long-continued pressure; and this stage of the operation will be facilitated by causing an assistant to steady the head by pressure on the fœtus through the abdomen, more especially if it be still free above the pelvic brim. We must then press together the handles of the instrument, which will have the effect of widely separating the cutting portion and making an incision through the bones. After this the point should be turned round, and again opened at right angles to the former incision, so as to make a free crucial opening. During this process care must be taken to bury the perforator in the skull up to the rests, so as to avoid

the possibility of injuring the maternal soft parts. The instrument should now be introduced within the skull and moved freely about, so as to thoroughly and completely break up the brain. Especial care must be taken to reach the medulla oblongata and base of the brain, for if these were not destroyed we might subject ourselves to the distress of extracting a child in whom life was not extinct. If this part of the operation be thoroughly performed, there will be no necessity for washing out the brain by the injection of warm water, as is sometimes recommended, for the broken-up tissue will escape freely through the opening made by the perforator.


Perforation of the After-coming Head.—The perforation of the after-coming head does not generally offer any particular difficulty. It is accomplished in the same manner, the child's body being well drawn out of the way by an assistant. The point of the perforator, carefully guarded by the finger, is guided up to the occiput or behind the ear, where it is inserted.

It is sometimes Useful to Postpone Extraction.—If there be no necessity for very rapid delivery, and the pains be still present, it is often advisable to wait ten minutes or a quarter of an hour before proceeding to extract. This delay will allow the skull to collapse and become moulded to the cavity of the pelvis when forced down by the pains, and possibly the natural effort may suffice to finish the labor in that time; or at least the head will have descended farther, and will be in a better position for extraction. Should perforation be required after having failed to deliver with the forceps—and this is only likely to be the case when the obstruction is comparatively slight—it is certainly a good plan to perforate without removing the forceps, which may then be used as tractors.

We have now to decide on the method of extraction; and our choice generally lies between the cephalotribe and the craniotomy forceps, although in some few cases, in which the pelvic contraction is slight, version may be advantageously employed.

Comparative Merits of Cephalotripsy and Craniotomy.—Those who have used both must, I think, admit that in any ordinary case, in which the obstruction is not great and only a comparatively slight diminution in the size of the head is required, cephalotripsy is infinitely the easier operation. The facility with which the skull can be crushed is some-

FIG. 185.



Perforation of the Skull.

times remarkable, and those who will take the trouble to read the reports of the operation published by Braxton Hicks, Kidd, and others cannot fail to be struck with the rapidity with which the broken-down head may often be extracted. This is far from being the case with the craniotomy forceps, even when the obstruction is moderate only; for it may be necessary to use considerable traction, or the blades may take a proper grasp with difficulty, or it may be essential to break down and remove a considerable portion of the vault of the cranium before the head is lessened sufficiently to pass. During the latter process, however carefully performed, there is a certain risk of injuring the maternal structures, and in the hands of a nervous or inexperienced operator this danger, which is entirely avoided in cephalotripsy, is far from slight. The passage of the blades of the cephalotribe is by no means difficult, and I think it must be admitted that the possible risks attending it are comparatively small. On account, therefore, of its simplicity and safety to the maternal structures, I believe cephalotripsy to be decidedly the preferable operation in all cases of moderate obstruction.

When we approach the lower limit, and have to do with a very marked amount of pelvic deformity, the two operations stand on a more equal footing. Then the deformity may be so great as to render it difficult to pass the blades of even the smallest cephalotribe sufficiently deep to grasp the head firmly, and even when they are passed the space is often so limited as to impede the easy working of the instrument. Besides this, repeated crushings may be required to diminish the skull sufficiently. I attach but little importance to the argument that the diminution of the skull in one diameter increases its bulk in another. The necessity of removing and replacing the blades on another part of the skull, and of repeating this perhaps several times in the manner recommended by Pajot, is a far more serious objection. To do this in a contracted pelvis involves, of necessity, the risk of much contusion. Fortunately, cases of this kind are of extreme rarity—much more so than is generally believed—but when they do occur they tax the resources of the practitioner to the utmost.

On the whole, the conclusion I would be inclined to arrive at with regard to the two operations is, that in all ordinary cases cephalotripsy is safer and easier, whereas in cases with considerable pelvic deformity the advantages of cephalotripsy are not so well marked, and craniotomy may even prove to be preferable.

Description of the Operation.—The first step in using the cephalotribe is the passage of the blades. These are to be inserted in precisely the same manner and with the same precautions as in the high forceps operation. In many cases the os is not fully dilated, and it is absolutely essential to pass the instrument within it. Special care should therefore be taken to avoid any injury to its edges, and for this purpose two or three fingers of the left hand, or even the whole hand, should be passed high up, so as thoroughly to protect the maternal structures. In order that the base of the skull may be reached and effectually crushed, the blades must be deeply inserted; and in doing this great care and gentleness must be used. As the projecting promontory of the sacrum generally tilts the head forward, the handles of the instrument, after

locking, must be well pressed backward toward the perineum. If the blades do not lock easily, or if any obstruction to their passage be experienced, one of them must be withdrawn and reintroduced, just as in forceps operations. Care must be taken, as the instrument is being inserted, to fix and steady the head by abdominal pressure, since it is generally far above the brim, and would readily recede if this precaution were neglected. When the blades are *in situ* we proceed to crush by turning the screw slowly, and as the blades are approximated the bones yield and the cephalotribe sinks into the cranium. The crushed portion then measures, of course, no more than the thickness of the blades—that is, about $1\frac{1}{2}$ inches. This is necessarily accompanied by some bulging of the part of the cranium that is not within the grasp of the instrument (Fig. 186), but in slight deformity this is of no consequence, and we may proceed to extraction, waiting, if possible, for a pain, and drawing at first downward in the axis of the pelvic inlet, as in forceps delivery, then in the axis of the outlet. The site of perforation should be examined to see that no spiculæ of bone are projecting from it, and if so they should be carefully removed. In such cases the head often descends at once and with the greatest ease. Should it not do so, or should the obstruction be considerable, a quarter turn should be given to the handles of the instrument, so as to bring the crushed portion into the narrower diameter and the uncrushed portion into the wider transverse diameter. It may now be advisable to remove the blades carefully, and to reintroduce them with the same precautions, so as to crush the unbroken portion of the skull. This adds materially to the difficulties of the case, since the blades have a tendency to fall into the deep channel already made in the cranium, and so it is by no means always easy to seize the skull in a new direction. Before reapplying them, if the condition of the patient be good and pains be present, it may be well to wait an hour or more, in the hope of the head being moulded and pushed down into the pelvic cavity. This was the plan adopted by Dubois, and, according to Tarnier, was the secret of his great success in the operation. Pajot's method of repeated crushings in the greater degrees of contraction is based on the same idea, and he recommends that the instrument should be introduced at intervals of two, three, or four hours, according to the state of the patient, until the head is thoroughly crushed, no attempts at traction being used, and expulsion being left to the natural powers. This, he says, should always be done when the contraction is below $2\frac{1}{2}$ inches, and he maintains that it is

FIG. 186.



Fœtal Head crushed by the Cephalotribe.

quite possible to effect delivery by this means when there is only $1\frac{1}{2}$ inches in the antero-posterior diameter. The repeated introduction of the blades in this fashion must necessarily be hazardous, except in the hands of a very skilful operator; and I believe that if a second application fail to overcome the difficulty, which will only be very exceptionally the case, it would be better to resort to the measures presently to be described.

Destruction of the Base of the Skull from Within by the Basilyst.—Professor Simpson of Edinburgh¹ has recently suggested the use of an instrument which he calls a “basilyst.” Its object is to

FIG. 187.

Professor Simpson's
Basilyst.

break up the base of the foetal skull from within, after the method originally proposed by Guyon. The screw-like portion of the instrument (Fig. 187), which is inserted through the perforation made in the cranial vault, is driven through the hard base, which is then disintegrated by the separate movable blade. If experience proves that this instrument can be readily worked, it promises to be a valuable addition to our armamentarium, since it will effectually destroy the most resistant portion of the skull without risk of injury to the maternal structures, and thus very materially facilitate extraction.

Extraction by the Craniotomy Forceps.—Should we elect to trust to the craniotomy forceps for extraction, one blade is to be introduced through the perforation, and the other, in apposition to it, on the outside of the scalp. In moderate deformities traction applied during the pains may of itself suffice to bring down the head. Should the obstruction be too great to admit of this, it is necessary to break down and remove the vault of the cranium. For this purpose Simpson's cranioclast answers

better than any other instrument. One of the blades is passed within the cranium, the other, if possible, between the scalp and the skull, and the portion of bone grasped between them is broken off; this can generally be accomplished by a twisting motion of the wrist, without using much force. The separated portion of bone is then extracted, the greatest care being taken to guard the maternal structures during its removal by the fingers of the left hand. The instrument is then applied to a fresh part of the skull, and the same process repeated, until as much of the vault of the cranium as may be necessary is broken up and removed.

Advantages of bringing down the Face in Difficult Cases.—Dr. Braxton Hicks² has conclusively shown that in difficult cases, after the removal of the cranial vault, the proper procedure is to bring down the face, since the smallest measurement of the skull after the removal of the upper part of the cranium is from the orbital ridge to the alveolar edge of the superior maxillary bone. This alteration in the presentation he proposes to effect by a small blunt hook, made for the purpose, which is forced into the orbit, by means of which the face is made to descend. Barnes recommends that this should be done by fixing the craniotomy forceps over the forehead and face and making traction in a backward

¹ *Edin. Med. Journ.*, April, 1880.² *Obst. Trans.*, vol. vii.

direction, so as to get the face past the projecting promontory of the sacrum. The importance of bringing down the face was long ago pointed out by Burns, but it had been lost sight of until Hicks again drew attention to it in the paper referred to. In the class of cases in which this procedure is valuable the risk to the maternal passages from the removal of the fractured portions of bone must always be considerable, and it is of great importance not only to preserve the scalp as entire as possible, so as to protect them, but to use the utmost possible care in removing the broken pieces of bone.

Extraction of the Body.—When the extraction of the head has been effected, either by the cephalotribe or the craniotomy forceps, there is seldom much difficulty with the body. By traction on the head one of the axillæ can easily be brought within reach, and if the body do not readily pass the blunt hook should be introduced and traction made until the shoulder is delivered. The same can then be done with the other arm. If there be still difficulty, the cephalotribe may be used to crush the thorax. The body is, however, so compressible that this is rarely required.

[The craniotomy forceps chiefly in use with us were devised by the late Prof. Charles D. Meigs for his second operation upon Mrs. Reybold of Philadelphia in 1833, and have been used repeatedly since, either as tractors or for reducing the size of the foetal head, in cases of deformity of the pelvis.¹ Some obstetricians prefer the less curved and broader-bladed instrument of Great Britain as a tractor; but for the general purposes of picking away the cranial bones and drawing down the base of the skull in cases of extreme pelvic deformity there is no more simple appliance than that of Dr. Meigs.


To act upon an oval body like the foetal head Dr. M. was obliged to prepare two forms of forceps—straight and curved—to be used as might be required according to the part of the skull to be broken down or drawn upon. These are lightly made, serrated, and $12\frac{1}{2}$ inches in length.—Ed.]

Embryotomy in Transverse Presentations in which Turning is Impossible.—There only remains for us to consider the second class of destructive operations. These may be necessary in long-neglected cases of arm presentation in which turning is found to be impracticable. Here, fortunately, the question of killing the fœtus does not arise, since it will, almost necessarily, have already perished from the continuous pressure. We have two operations to select from—*decapitation* and *evisceration*.

[¹ The illustrations given are taken from the instruments devised by Dr. Meigs as an improvement upon his original pattern, and will be seen to differ from those usually presented in American obstetrical publications.—Ed.]

FIG. 188.

FIG. 189.

Straight
Craniotomy
Forceps.Curved
Craniotomy
Forceps.

Decapitation.—The former of these is an operation of great antiquity, having been fully described by Celsus. It consists in severing the neck, so as to separate the head from the body; the body is then withdrawn by means of the protruded arm, leaving the head in utero to be subsequently dealt with. If the neck can be reached without great difficulty—and in the majority of cases the shoulder is sufficiently pressed down into the pelvis to render this quite possible—there can be no doubt that it is much the simpler and safer operation.

Methods of Dividing the Neck.—The whole question rests on the possibility of dividing the neck. For this purpose many instruments have been invented. The one generally recommended in this country is known as Ramsbotham's hook, and consists of a sharply-curved hook with an internal cutting edge. This is guided over the neck, which is divided by a sawing motion. There is often considerable difficulty in placing the instrument over the neck, although, if this were done, it would doubtless answer well. Others have invented instruments, based on the principle of the apparatus for plugging the nostrils, by means of which a spring is passed round the neck, and to the extremity of the spring a short cord or the chain of an *écraseur* is attached; the spring is then withdrawn and brings the chain or cord into position. The objection to any of these apparatuses is that they are unlikely to be at hand when required, for few practitioners provide themselves with costly instruments which they may never require. It is of importance, therefore, that we should have at our command some means of dividing the neck which is available in the absence of any of these contrivances. Dubois recommends for this purpose a strong pair of blunt scissors. The neck is brought as low as possible by traction on the prolapsed arm, and the blades of the scissors guided carefully up to it. By a series of cautious snipping movements it is then completely divided from below upward. This, if the neck be readily within reach, can generally be effected without any particular difficulty. Dr. Kidd of Dublin,¹ who strongly advocates this operation, recommends that an ordinary male elastic catheter, strongly curved and mounted on a firm stilette, or, still better, on a uterine sound, should be passed round the neck. Previous to introduction a cord should be attached to the extremity of the catheter, which is left round the neck when it is withdrawn. By means of this cord a strong piece of whiplcord or the wire of an *écraseur* can easily be drawn round the neck and used for dividing it. The former, to protect the maternal structures, may be worked through a speculum, and by a series of lateral movements the neck is easily severed. The *écraseur*, however, offers special advantage, since it entirely does away with any risk of injuring the mother.

Withdrawal of the Body and Delivery of the Head.—After the neck is divided the remainder of the operation is easy. The body is withdrawn without difficulty by the arm, and we then proceed to deliver the head. By abdominal pressure this, in most cases, can be pushed down into the pelvis, so as to come easily within reach of the cephalotribe, which is by far the best instrument for extraction. Preliminary perforation is not necessary, since the brain can escape through the sev-

¹ *Dublin Quart. Journ.*, May, 1871.

ered vertebral canal. The secret of doing this easily is to fix and press down the head sufficiently from above, otherwise it would slip away from the grasp of the instrument. The perforator and craniotomy forceps may be used if the cephalotribe be not at hand. Perforation is, however, by no means always easy, on account of the mobility of the head. After it is accomplished one blade of the craniotomy forceps is passed within the skull, the other externally, and the head slowly drawn down.

Evisceration.—The alternative operation of evisceration is a much more troublesome and tedious procedure, and should only be used when the neck is inaccessible. The first step is to perforate the thorax at its most depending part, and to make as wide an opening into it as possible in order to gain access to its contents. Through this the thoracic viscera are removed piecemeal, being first broken up as much as possible by the perforator, and then, the diaphragm being penetrated, those in the abdomen. The object is to allow the body to collapse and the pelvic extremities to descend, as in spontaneous evolution. This can be much facilitated by dividing the spinal column with a strong pair of scissors introduced into the opening made in the thorax, so that the body may be doubled up as on a hinge. Here the crotchet may find a useful application, for it can be passed through the abdominal cavity and fixed on some point in the interior of the child's pelvis, and thus strong traction can be made without any risk of injury to the mother. It can be readily understood that this process is so lengthy and difficult as to render it probably the most trying of obstetric operations; it is certainly inferior in every respect to decapitation, and is only to be resorted to when that is impracticable.

[The Cæsarean operation has been performed in the United States in 11 cases of impaction of the fœtus in a transverse position, because of the great difficulty of accomplishing either decapitation or evisceration, with a saving of 7 women. The 4 deaths were from exhaustion.—ED.]

CHAPTER VI.

THE CÆSAREAN SECTION; PORRO'S OPERATION; SYMPHYSEOTOMY.

History of the Cæsarean Section.—The Cæsarean section has perhaps given rise to more discussion than any other subject connected with midwifery, and there is yet much difference of opinion as to the limits of, and indications for, the operation. The period at which the Cæsarean section was first resorted to is not known with accuracy. It seems to have been practised by the Greeks after the death of the mother, and Pliny mentions that Scipio Africanus and Manlius were born in this way. The name of Cæsar is said to have been given to children so extracted, and afterward to have been assumed as a family patronymic.

These children were dedicated to Apollo; whence arose the practice of things sacred to that god being taken under the special protection of the family of the Cæsars. Many celebrities have been supposed to owe their lives to the operation; among the rest, Æsculapius, Julius Cæsar, and our own Edward VI. Regarding the two latter, there is conclusive proof that the tradition is without foundation. There is no doubt that the operation was constantly practised on women who had died at an advanced period of pregnancy, and indeed it has at various times been enforced by law. Thus, among the Romans it was decreed by Numa that no pregnant woman should be buried until the fœtus had been removed by abdominal section. The Italian laws also made it necessary, and the operation has always received the strong support of the Roman Church. So lately as the middle of the eighteenth century the king of Sicily sentenced to death a physician who had neglected to practise it. The first authentic case in which the operation was performed on a living woman occurred in 1491. It was afterward practised by Nufer in 1500; and in 1581, Rousset published a work on the subject in which a number of successful cases were related. In English works of that time it is not alluded to, although it was undoubtedly performed on the Continent, and to such an extent that its abuse became almost proverbial. We have evidence in Shakespeare, however, that the operation was familiarly known in this country, since he tells us that

"Macduff was from his mother's womb
Untimely ripped." [1]

Paré and Guillemeau, amongst the writers of the period, were noted for their hostility to the operation, while others equally strongly upheld it.

In this country it has scarcely ever been performed in a manner which offers even the faintest hope of success. It has been looked upon as almost necessarily fatal to the mother, and it has therefore been delayed until the patient has arrived at the utmost stage of exhaustion. For example, in looking over the records of British cases it is no uncommon thing to find that the Cæsarean section was resorted to two, three, or even six days after labor had begun, and when the patient was almost moribund. With rare exceptions within the last few years the operation has been performed in what may be called a haphazard way. In many cases long and fruitless attempts at delivery by craniotomy had already been made, so that the passages had been subjected to much contusion and violence. Little or no attempt has been made to obviate the well-known risks of abdominal operations; no care has been taken to prevent blood and other fluids finding their way into the peritoneal cavity, and no means have been adopted subsequently to remove them. It is therefore not so much a matter of surprise that the mortality has been so great, but rather that any cases have recovered.

From what we know of the history of ovariectomy, its early fatality, and the extreme and even apparently exaggerated precautions which are essential to its success, it is fair to conclude that if the Cæsarean section

[1 To my mind, this refers to what had often taken place in ancient wars, where women were ripped open by soldiers with a sword. The expression *untimely* does not indicate that the fœtus had come to maturity or that the woman was in labor.—ED.]

were performed, as it is to be hoped it always will be in future, with the same careful attention to minute details as ovariectomy, the results would not be so disastrous. Making every allowance for these facts, it must be admitted that the Cæsaean section is necessarily almost a forlorn hope; and in making these observations I have no intention of contesting the well-established rule of British practice, that it is not admissible as an operation of election, and must only be resorted to when delivery *per vias naturales* is impossible.

Statistical Returns not Reliable.—The mortality, as given in statistical returns from various sources, differs so greatly as to make them but little reliable. Radford has tabulated the operations performed in this country up to 1868, and the list has been completed by Harris¹ up to 1879.^[2] The cases amount to 118 in all, of which 22 were successful, or rather more than 18 per cent. Michaelis and Kayser found that out of 258 and 338 operations, 54 and 64 per cent. respectively were fatal. These include operations performed under all sorts of conditions, even when the patient was almost moribund; and until we are in possession of a sufficient number of cases performed under conditions showing that the result is obviously due to the operation, in which it was undertaken at an early period of labor and performed with a reasonable amount of care, it is obviously impossible to arrive at any reliable conclusions as to the mortality of the operation. That it is necessarily hopeless is certainly not the case, and we know that on the Continent, where it is resorted to much oftener and earlier in labor than in this country, there are authentic cases in which it has been performed twice, thrice, and even, in one instance, four times, on the same patient. Kayser thinks that a second operation on the same patient affords a better prognosis than a first, probably because peritoneal adhesions resulting from the first operation have shut off the general abdominal cavity from the uterine wound; and he believes that in second operations the mortality is not more than 29 per cent.

The Cæsaean Section in America.—The Cæsaean section has been much more successful in America than in Great Britain. Dr. Harris of Philadelphia, who has paid much attention to the subject, has collected 134 cases occurring in the United States, of which 53 were successful as regards the mother. These favorable results he refers partly to the fact that none of the American cases were the subjects of mollities ossium, rachitic patients forming one-half of the entire number, and partly to the prevalence of habits of beer- and gin-drinking in this country. He also gives some interesting facts showing how remarkably the mortality of the operation is lessened when it is performed soon and the patient is not exhausted by long and fruitless labor. Out of 28 selected cases of this kind, 21, or 75 per cent., were successful. [23 children were delivered alive, and 19 saved.—ED.]

[*Latest Cæsaean Statistics of America.*—For some years we have

¹ "The Cæsaean Operation in the United Kingdom," *Brit. Med. Journ.*, April 3, 1880.

[² The late Dr. Radford in 1880 tabulated the British operations to May, 1879, and presented 132 cases, failing to notice 4, 1 of which recovered. To these I add 2 of a later date, both of which were saved, making in all 138 cases, with 26 recoveries.—ED.]

been very decidedly retrograding in our proportion of cures to deaths and of *timely* to *late* operations, and have been gradually increasing the percentage of deaths in the whole record. Ten years ago, when we had had 101 operations, we counted 45 women saved, or a fraction below 45 per cent. In the ten years we have added 33 cases, with 25 women and 19 children lost, thereby reducing the number of cures below 40 per cent. Worse still than this, the operations of the past ten months in the United States (April 1, 1884, to Feb., 1885), number 6, and all of the women and children were lost. But 2 of these last operations have yet been reported in journals, but all are promised for publication. Of the 33 operations referred to, 24 were performed upon cases regarded as in an unfavorable condition, and only 8 were operated upon within the first twenty-four hours of labor. It is not, then, to be wondered at that 75 per cent. of the women perished. It is folly in an accoucheur to expect a surgeon to save his patient when broken down and exhausted by the length of her labor.

We have had 134 Cæsarean operations in the United States, and the addition made by Mexico and the West Indies increases the number by 9. Of the 134, there were saved 53 women, and 7 of the 9, making, in all, 143 operations, with 60 women saved. Of the 134 cases in the United States, 26 were dwarfs, ranging in height from 3 feet to 4 feet 8 inches, one-half being from 3 to 4 feet; and of these 26 but 7 were saved. The shortest women to recover were respectively 3 feet 9 inches, 3 feet 11½ inches, and 4 feet. Dwarfs should not be allowed to continue in labor more than from two to four hours, as they rapidly become exhausted, and are apt to die of shock, exhaustion, or peritonitis. One dwarf, operated upon and saved after a labor of two hours, was already showing the evidences of exhaustion. By acting very promptly the labor only having lasted 1½ hours, a physician of Brest, France, was successful in saving a woman only 35 inches in height, with her child, a few years ago. He appeared to know the value of time to such a patient, and, when called in, rapidly made his arrangements for using the knife. The term "early" is a relative one, and what might be such as a measure of time in one case would not be in another. "*Timely*" has no measure in hours, but relates more to the strength and condition of the patient. If the strength, pulse, and morale of the woman indicate a favorable condition, and she has not had active labor beyond a few hours, she will in the majority of instances in our country recover.—ED.]

Results to the Child.—The mortality of the children likewise cannot be ascertained from statistical returns, since in the large majority of cases in which dead children were extracted the result had nothing to do with the operation. Indeed, there is nothing in the operation itself which can reasonably be supposed to affect the child. If, therefore, the child be alive when the operation is commenced, there is every probability of its being extracted alive; and Radford's conclusion, that "the risk to infants in Cæsarean births is not much greater than that which is contingent on natural labor, provided correct principles of practice are adopted," probably very nearly represents the truth.

Causes Requiring the Operation.—The Cæsarean section is required

when there is such defective proportion between the child and the maternal passages that even a mutilated fœtus cannot be extracted. This in by far the greatest number of cases is due to deformity of the pelvis arising from rickets or mollities ossium. The latter may occur in a patient who has been previously healthy and who has given birth to living children. It is a more common cause of the extreme varieties of deformity than rickets, and out of 77 British cases, tabulated by Radford, in 43 the deformity was produced by osteo-malacia and in 14 only by rickets.^[1] In certain cases the pelvis itself may be of normal size, but has its cavity obstructed by a solid tumor of the ovary, of the uterus itself, or one growing from the pelvic wall. The obstruction may also depend on morbid conditions of the maternal soft parts, of which the most common is advanced malignant disease of the cervix. Other conditions may, however, render the operation essential. Thus, Dr. Newman² records a case in which he performed it for insurmountable resistance and obstruction of the cervix, which was believed at the time to be caused by malignant disease. The patient recovered, and was subsequently delivered naturally and without anything abnormal being made out. This renders it probable that the disease was not malignant, and it may possibly have been an extensive inflammatory exudation into the tissues of the cervix, subsequently absorbed. I myself was present at a Cæsarean section performed in Calcutta in the year 1857 when the pelvis was so uniformly blocked up with exudation, probably due to extensive pelvic cellulitis or hæmatocele, that the operation was essential.

Limits of Obstruction Justifying the Operation.—Different accoucheurs have fixed on various limits for the operation. Most British authorities are of opinion that it need not be resorted to if the smallest diameter of the pelvis exceed $1\frac{1}{2}$ inches.³ This question has already been considered in discussing craniotomy, and it has been shown that a mutilated fœtus may be drawn through a pelvis of $1\frac{1}{2}$ inches antero-posterior diameter, provided there be a space of 3 inches in the transverse diameter. If sufficient space for using the necessary instruments do not exist, the Cæsarean section may be required even when there is a larger antero-posterior diameter than $1\frac{1}{2}$ inches. This is especially likely to occur when we have to do with deformity arising from mollities ossium, in which the obstruction is in the sides and outlet of the pelvis, the true conjugate being sometimes even elongated. On the Continent the Cæsarean section is constantly practised, as an operation of election, when the smallest diameter measures from 2 to $2\frac{1}{2}$ inches; and when the child is known to be alive some foreign authors recommend it when there is as much as 3 inches in the antero-posterior diameter. In this country, where the life of the child is most properly considered of secondary importance to the safety of the mother, we cannot fix one limit for the operation when the child is living and another when it is dead. Nor, I think, can we admit the desire of the mother to run the risk,

[¹ I must again fill up the record of Dr. Radford. Of his 132 cases, 56 were the subjects of mollities ossium, and in 31 the deformity of the pelvis was attributed to rickets: 14 cases were the subjects of cancer and 10 of tumors.—Ed.]

² *Obst. Trans.*, vol. vii. p. 343.

³ In Dr. Parry's table of 70 craniotomies there are 34 cases of 2 to $2\frac{1}{2}$ inches conjugate.

rather than sacrifice the child, as a justification of the operation, although this is laid down as an indication by Schroeder.¹ Great as are the dangers attending craniotomy in extreme deformity, there can be no doubt that we must perform it whenever it is practicable, and only resort to the Cæsarean section when no other means of delivery are possible.

For this reason I think it unnecessary to discuss the question whether we are justified in destroying the fœtus in several successive pregnancies when the mother knows that it is impossible for her to give birth to a living child. Denman was the first to question the advisability of repeating craniotomy on the same patient. Amongst modern authors, Radford takes the most decided view on this point, and distinctly teaches that even when delivery by craniotomy is possible, it "can be justified on no principle, and is only sanctioned by the dogma of the schools or by usage," and that, therefore, the Cæsarean section should be performed with the view of saving the child. Doubtless, much can be said from this point of view, but, nevertheless, he would be a bold man who would deliberately elect to perform the Cæsarean section on such grounds.² It is to be hoped, however, that in these days the induction of premature labor or abortion would always spare us the necessity of deciding so delicate a point.

Post-mortem Cæsarean Operation.—The Cæsarean section may also be required in cases in which death has occurred during pregnancy or labor. This was the indication for which it was first employed, and it has constantly been performed when a pregnant woman has died at an advanced period of utero-gestation. There is no doubt that a prompt extraction of the child under these circumstances has frequently been the means of saving its life, but by no means so often as is generally supposed. Thus, Schwartz³ showed that out of 107 cases not one living child was extracted. Duer⁴ has written an interesting paper on this subject in which he has tabulated 55 cases of post-mortem Cæsarean sections. In 40 a *living* child was extracted, the time elapsing after the death of the mother being as follows: "Between 1 and 5 minutes, including 'immediately' and 'in a few minutes,' there were 21 cases; between 5 and 10 minutes, none; between 10 and 15 minutes, 13 cases; between 15 and 23 minutes, 2 cases; after 1 hour, 2 cases; and after 2 hours, 2 cases." In those extracted, however, after the lapse of an hour the children did not ultimately survive, and the cases themselves seem open to some doubt.

Want of Success in Post-mortem Operation.—The reason that the want of success has been so great is doubtless the delay that must necessarily occur before the operation is resorted to, for, independently of the fact that the practitioner is seldom at hand at the moment of death, the very time necessary to assure ourselves that life is actually extinct will gen-

¹ *Manual of Midwifery*, p. 202.

² This was done twice successfully by Prof. William Gibson in the case of Mrs. Reybold of Philadelphia in 1835 and 1837, after she had twice been delivered by craniotomy under Prof. Charles D. Meigs, who declined destroying any more children for her. Mrs. R. still lives at the age of seventy-five, and the daughter likewise, with her four children. The son died at 43, leaving two children.—HARRIS.

³ *Monat. f. Geburt*, suppl. vol., 1861, p. 121.

⁴ "Post-mortem Delivery," *Am. Journ. of Obst.*, Jan., 1879.

erally be sufficient to cause the death of the fœtus. Considering the intimate relations between the mother and child, we can scarcely expect vitality to remain in the latter more than a quarter, or at the outside half an hour, after it has ceased in the former. The recorded instances in which a living child was extracted ten, twelve, and even forty hours after death were most probably cases in which the mother fell into a prolonged trance or swoon, during the continuance of which the child must have been removed. A few authentic cases, however, are known in which there can be no reasonable doubt that the operation was performed successfully several hours after the mother was actually dead.

Since, then, there is a chance, however slight, of saving the child's life, we are bound to perform the operation, even when so much time has elapsed as to render the chances of success extremely small. It might be considered almost superfluous to insist on the necessity of assuring ourselves of the mother's death before commencing the necessary incisions; but, unfortunately, numerous instances are known in which mistakes in diagnosis have been made, and in which the first steps of the operation have shown that the mother was still alive. The operation should therefore always be performed with the same care and caution as if the mother were living. If death have occurred during labor, some have advised version as a preferable alternative. This can only be resorted to with any hope of success if the passages be in a condition to admit of delivery with rapidity; otherwise the delay occasioned by dilatation, even when forcibly accomplished, and the drawing of the child through the pelvis, will be almost necessarily fatal. The only argument in favor of version is, that it is less painful to the friends; and, if they manifest a decided objection to the Cæsarean section, there can be no reason why an attempt to save the child in this way should not be made.

Causes of Death after Cæsarean Section.—The causes of death after the Cæsarean section may, speaking generally, be classed under four principal heads: hemorrhage, peritonitis and metritis, shock, septicæmia, and exhaustion from long delay. These are pretty much the same as those following ovariotomy, and the resemblance between the two operations is so great that modern experience as to the best mode of performing ovariotomy, as well as regards the after-treatment, may be taken as a guide in the management of cases of Cæsarean section.

Hemorrhage is Frequent, although Seldom Fatal.—Hemorrhage to an alarming extent is a frequent complication, though seldom the cause of death. Thus, out of 88 operations the particulars of which have been carefully noted, severe hemorrhage occurred in 14, 6 of which terminated successfully, and in 4 only could the fatal result be ascribed to the loss of blood. In 1 of these the source of the hemorrhage is not mentioned, in another it came from the wound in the abdominal wall, and in the other 2 from the uterine incision being made directly over the placenta. In neither of the two latter was the loss of blood immediately fatal, for it was checked by uterine contraction, and only recurred after many hours had elapsed. The divided uterine sinuses and the open mouths of the vessels at the placental site are the most common sources of hemorrhage.

Means of Avoiding the Risk.—Much may be done to diminish the risk of bleeding, but even with every precaution it must be a source of danger. Hemorrhage from the abdominal wall may be best prevented by making the incision as nearly as possible in the line of the linea alba, so as not to wound the epigastric arteries, and by controlling bleeding by pressure-forceps as we proceed, as is done in ovariectomy. The principal loss of blood will be met with in dividing the uterus; and this will be the greatest when the incision is near or over the placental site, where the largest vessels are met with. We are recommended to ascertain the position of the placenta by auscultation, and thus, if possible, to avoid opening the uterus near its insertion. But even if we admit the placental souffle to be a guide to its situation if the placenta be attached to the anterior walls of the uterus, a knowledge of its position would not always enable us to avoid opening the uterus in its immediate vicinity. We must, in the event of its lying under the incision, rather hope to control the hemorrhage by removing it at once from its attachments and rapidly emptying the uterus. When the child has been removed there may be a large escape of blood, but this will generally be stopped by the contraction of the uterus in the same manner as after natural labor. Should contraction not take place, the uterus may be firmly grasped for the purpose of exciting it. This plan is advocated by Winckel, who had a large experience in the operation, and by using free compression in this way, and making a point of not closing the wound until the uterus is firmly contracted, he has never met with any inconvenience from hemorrhage. If bleeding continue, styptic applications may be used, as in a case reported by Hicks, who was obliged to swab out the uterine cavity with a solution of perchloride of iron.

Peritonitis and Metritis are Frequent Causes of Death.—Among the most frequent causes of death are peritonitis and metritis. Kayser attributes the fatal result to them in 77 out of 123 unsuccessful cases. The mere division of the peritoneum will not account for the frequency of this complication, since its occurrence is considerably more frequent than after ovariectomy, in which the injury to the peritoneum is quite as great; and indeed greater, if we take into account the adhesions which have to be divided or torn in that operation.

Degeneration of the Uterine Fibres supposed to be Unfavorable to Repair.—The division of the uterus must be regarded as one source of this danger. Dr. West lays great stress on its unfavorable condition after delivery for reparative action. He believes that the process of involution or fatty degeneration which commences in the muscular fibres previous to delivery renders them peculiarly unfitted to cicatrize; and he points out that on post-mortem examination the edges of the incision have been found dry, of unhealthy color, gaping, and showing no tendency to heal. On this account, Hicks and others have operated ten days or more before the full period of labor, in the hope that the risk from this source might be avoided. It is by no means certain, however, that the change in the uterine fibres is the cause of the wound not healing, and involution will commence at once when the uterus is emptied even if the full period of pregnancy have not arrived. As a point of ethics, moreover, it is questionable if we are justified in anticipating the

date of so dangerous an operation, even by a few weeks, unless the benefit to be derived is very decided indeed.

Escape of Lochia and other Fluids into the Peritoneal Cavity.—One important cause of peritonitis is the escape of the lochia through the uterine incision into the cavity of the peritoneum, which there decompose and act as an unfailing source of irritation. This may be prevented, to a great extent, by seeing that the os uteri is patulous, so as to afford a channel for the escape of discharges, and by closing the uterine wound by sutures. In addition, there is the danger arising from blood and liquor amnii escaping into the peritoneum and subsequently decomposing. There is little evidence that "*la toilette du péritoine*," on which ovariologists now lay so much stress, has ever been particularly attended to in Cæsaean operations.

The Unhealthy Condition of the Patient is the Chief Source of Danger.—The chief predisposing cause of these inflammations, however, must be looked for in the condition of the patient, just as asthenic inflammation in ovariectomy is most frequently met with in those whose general health is broken down by the long continuance of the disease. We are fully justified, therefore, in assuming that peritonitis and metritis will be more likely to occur after the Cæsaean section when that operation has been unnecessarily delayed and when the patient is exhausted by a protracted labor. In proof of this we find that in a large proportion of the cases above mentioned peritonitis occurred when the operation was performed under unfavorable conditions.

Septicæmia.—The sources of septicæmia are abundantly evident—not the least, probably, being absorption by the open vessels in the uterine incision.

Nervous Shock.—The last great danger is general shock to the nervous system. In Kayser's 123 cases, 30 of the deaths are referred to this cause. In the large majority of these the patient was profoundly exhausted before the operation was begun. It is in predisposing to these nervous complications that we should, *a priori*, expect that vacillation and delay would be most hurtful; and in operating when the patient's strength is still unimpaired we afford her the best chance of bearing the inevitable shock of an operation of such magnitude.

Secondary Dangers.—In addition a few cases have been lost from accidental complications, which are liable to occur after any serious operation, and which do not necessarily depend on the nature of the procedure.

Danger to Child from Portions of its Body being Caught by the Contracting Uterus.—There is only one source of danger special to the child which is worthy of attention. As the infant is being removed from the cavity of the uterus the muscular parietes sometimes contract with great rapidity and force, so as to seize and retain some part of its body. This occurred in two of Dr. Radford's cases, and in one of them it is stated that "the child was vigorously alive when first taken hold of, but from the length of time occupied in extracting the head it became so enfeebled as to show only slight signs of life," and subsequently all attempts at resuscitation failed. I have myself seen the head caught in this way, and so forcibly retained that a second incision was required to release it.

In Dr. Radford's cases the placenta happened to be immediately under the incision, and he attributes the inordinate and rapid contraction of the uterus to its premature separation. It is difficult to believe that this was more than a coincidence, because the contraction does not take place until the greater part of the child's body has been withdrawn, and because numerous cases are recorded in which the uterus was opened directly over the placenta, or in which it was lying loose and detached, in none of which this accident occurred. The true explanation may, I think, be found in the varying irritability of the uterus in different cases.

Irrespective of the risk of portions of the child being caught and detained, rapid contraction is a distinct advantage, since the danger of hemorrhage is thereby thus diminished. Serious consequences may be best avoided by removing, when practicable, the head and shoulders of the child first, or by employing both hands in extraction, one being placed near the head, the other seizing the feet. Either of these methods is preferable to the common practice of laying hold of the part that may chance to lie most conveniently near the line of incision. If this point were properly attended to, although the detention of the lower extremities might occasionally occur, the life of the child would not be imperilled.

Whenever it is Possible the Patient should be Prepared for the Operation.—The preparation of the patient for the operation should seriously occupy the attention of the practitioner; and this is the more essential since almost all patients requiring the Cæsarean section are in a wretchedly debilitated condition. If the patient be not seen until she is actually in labor, of course this is out of the question. But this will rarely be the case, since the deformed condition of the patient must generally have attracted attention. Every possible means should be taken, therefore, when practicable, to improve the general health by abundance of simple and nourishing diet, plenty of fresh air, and suitable tonics (amongst which preparations of iron should occupy a prominent place), while the state of the secretions, the bowels, skin, and kidneys should be specially attended to. Whenever it is possible a large, airy apartment should be selected for the operation, which should never be done in a hospital if other arrangements be practicable. These details may seem trivial and unnecessary; but to ensure success in so hazardous an undertaking no care can be considered superfluous, and probably the want of attention to such points has had much to do with increasing the mortality.

Question of Time to be Selected for the Operation.—The question arises whether we should operate before labor has commenced. By selecting our own time, as some have advised, we certainly have the advantage of operating under the most favorable conditions, instead of, possibly, hurriedly. There are, however, numerous advantages in waiting until spontaneous uterine action has commenced which seem to me to more than counterbalance the advantages of choosing our own time. Prominent among these is the partial opening of the os uteri, so as to afford a channel for the escape of the lochia, and the certainty of active contraction of the uterus to arrest hemorrhage. Barnes recommends that premature

labor should be first induced, and then the operation performed. This seems to me to introduce a needless element of complexity ; and, besides, in cases of great deformity it is by no means always easy to reach the cervix with the view of bringing on labor. All needful arrangements should be made, so as to avoid hurry and excitement when the operation is commenced, and we may then wait patiently until labor has fairly set in.

The Administration of Anæsthetics.—The operation itself is simple. The patient should be placed on a table in a good light, and with the temperature of the room raised to about 65°. Chloroform has so frequently been followed by severe vomiting that it is probably better not to administer it. For the same reason Mr. Spencer Wells has long given up using it in ovariectomy, and finds that chloro-methyl answers admirably ; ether also is devoid of the disadvantages of chloroform. In one or two cases local anæsthesia has been used by means of two spray-producers acting simultaneously ; and this plan, if the patient have sufficient fortitude to dispense with general anæsthesia, has the further advantage of stimulating the uterus to powerful contraction.

To ensure as great a measure of success as possible the operation should be performed under carbolic spray and with all the minute precautions used in ovariectomy.

Description of the Operation.—The incision should be made as much as possible in the line of the linea alba, so as to avoid wounding the epigastric arteries. On account of the deformity the configuration of the abdomen is often much altered, and some have advised that the incision should be made oblique or transverse and on the most prominent part of the abdomen. The risk of hemorrhage being thus much increased, the practice is not to be recommended. The incision, commencing a little above the umbilicus, is carried down for about three inches below it. The skin and muscular fibres are carefully divided, layer by layer, until the shining surface of the peritoneum is reached, and any bleeding vessels should be secured as we proceed. A small opening is now made in the peritoneum, which should be laid open along the whole length of the incision upon two fingers of the left hand introduced as a guide. Before incising the uterus an assistant should carefully support it in a proper position, and push it forward by the hands placed on either side of the incision, so as to bring its surface into apposition with the external wound and prevent the escape of the intestines. If we have reason to believe that the placenta is situated anteriorly, we may incise the uterus on one or other side ; otherwise the line of incision should be as nearly as possible central.¹ The substance of the uterus is next divided until the membranes are reached, which are punctured and divided in the same way as the peritoneum. The uterine incision should be of the same length as that in the abdomen, and it should not be made too near the fundus ; for not only is that part more vascular than the body of the uterus, but wounds in that situation are more apt to gape, and do not cicatrize so favorably. After the uterus is opened, Dr. Ludwig Winckel recommends that the fingers of an assistant should be

[¹ See Kehr's plan of incising the uterus transversely for this purpose, on page 524.—Ed.]

placed in the two terminal angles of the wound, so that the ends of the incision may be hooked up and brought into close apposition with the abdominal opening. By this means he prevents not only the escape of blood and liquor amnii into the cavity of the peritoneum, but also the protrusion of the abdominal viscera.

Removal of the Child.—The child should now be carefully removed, the head and shoulders being taken out (if possible) first; the placenta and membranes are afterward extracted. Should the placenta be unfortunately found immediately under the incision, a considerable loss of blood is likely to take place, which can only be checked by removing it from its attachments and concluding the operation as rapidly as possible.

Importance of Securing Uterine Contraction.—As soon as the child and the secundines have been extracted the sooner the uterus contracts the better. It will usually do so of itself, but should it remain lax and flabby, it should be pressed and stimulated by the hand. We are specially warned against handling the uterus by Ramsbotham and others, but there seems no valid reason why we should not restrain hemorrhage in this way, as *after* a natural labor. The intervention of the abdominal incision in their lax condition after delivery can make very little difference between the two cases. Ergotin administered hypodermically will also be useful in promoting efficient contraction.

Closure of the Uterine and Abdominal Wounds.—The advisability of closing the uterine wound by sutures is a mooted point.^[1] The balance of opinion is certainly in favor of this practice, as tending to prevent the escape of the lochia into the peritoneal cavity. Interrupted sutures of silver wire may be used and cut short, or, as successfully practised by Scudder, ² a continuous silk suture may be applied, one end being introduced through the os into the vagina, by which it is subsequently withdrawn. Sutures of ordinary catgut are apt to yield, and are therefore discarded. Some chronic gut or some of the antiseptic guts now prepared may be used with safety. Before closing the uterine wound two fingers should be passed through the cervix to ensure its dilatation. A free escape of the lochia in this direction is of great importance, and Wunkel even advises the placing of a strip of lint around the os, so as to keep up a free exit for the discharge.

Abdominal Wound should not be Closed until Hemorrhage has Stopped.—A point of great importance, and not sufficiently insisted on, is the advisability of not closing the abdominal wound until we are fully assured that hemorrhage is completely stopped, since any blood poured into the peritoneum would very materially lessen the chance of recovery. In a successful case reported by Dr. Newman² the abdominal wound was closed for nearly an hour. Before doing so all blood in the cavity should be carefully removed from the peritoneal cavity

¹ Dr. Simpson has been in use in the United States since June, 1828, but there is no record of its use here until treated from 1828 to 1868, since which time there have been 28 cases reported. Of these, 12 were sutured with silver wire, and 6 recovered; 15 were sutured with catgut, 1 with fine hemp, recovered; 2 with catgut, lost; and 1 was not sutured, recovered, lost. Of the 28 *early operations* in the list of the United States, 12 were sutured, saving 3; and of the balance, 22, there were 18 that recovered.

by clean soft sponges dipped in warm water. The abdominal wound should be closed from above downward by harelip pins, wire or silk sutures, which should be inserted at a distance of an inch from each other and passed entirely through the abdominal walls and the peritoneum at some little distance from the edges of the incision, so as to bring the two surfaces of the peritoneum into contact. By this means we ensure the closure of the peritoneal cavity, the opposed surfaces adhering with great rapidity. If, as should be the case, the operation is performed with full antiseptic precautions, the wound should now be dressed precisely as after ovariectomy.

Subsequent Management.—Into the subsequent treatment it is unnecessary to enter at any length, since it must be regulated by general principles, each symptom being met as it arises. It has been customary to administer opiates freely after the operation, but they seem to have a tendency to produce sickness and vomiting, and ought not to be exhibited unless pain or peritonitis indicates that they are required. In fact, the treatment should in no way differ from that usual after ovariectomy, and the principles that should guide us will be best shown by the following quotation from Mr. Spencer Wells's description of that operation: "The principles of after-treatment are—to obtain extreme quiet, comfortable warmth, and perfectly clean linen to the patient; to relieve pain by warm applications to the abdomen and by opiate enemias; to give stimulants when they are called for by failing pulse or other signs of exhaustion; to relieve sickness by ice or iced drinks; and to allow plain, simple, but nourishing food. The catheter must be used every six or eight hours until the patient can move without pain. The sutures are removed on the third day, unless tympanitic distension of the stomach or intestines endangers reopening of the wound. In such circumstances they may be left for some days longer. The superficial sutures may remain until union seems quite firm."

Improved Methods of Performing Gastro-hysterotomy.—These, with one exception—that of Dr. Garrigues of New York—come from Germany, where there is a disposition to revive the old Cæsaean operation in consequence of that of Porro having saved in that country but a fraction over 32 per cent. of the women. The new antiseptic processes are devised by Cohnstein and Kehrer of Heidelberg, Frank of Cologne, and Säger of Leipzig. Cohnstein's¹ process has not yet been tested. He proposes to turn out the uterus entire; open it vertically through the posterior wall; deliver the fœtus and secundines; replace the organ, and pass a drainage-tube through the Douglas *cul-de-sac* and vagina; and close the abdominal wound. He directs that compression of the aorta shall be made during the opening and contraction of the uterus to check the loss of blood.

*Frank's Process.*²—Wash the abdomen over with ether and with a 5 per cent. solution of carbolic acid; disinfect the vagina by irrigating with the latter fluid. Turn out the uterus, and open it vertically in front, commencing the incision low down in the vesico-uterine excavation; extract the fœtus and secundines; wash the front of the uterus, its

[¹ *Centralblatt für Gynäkol.*, 1881, No. 12, vol. v. p. 290.]

[² *Ibid.*, 1881, No. 25, vol. v. p. 598.]

interior, and the vagina with the 5 per cent. carbolic-acid solution. Pass a large drainage-tube through the abdominal and uterine wounds and out through the vagina. Suture the uterus above the tube; draw the round ligaments together above the uterine wound, and secure them with sutures of Czerny silk, so as to close over and separate the vesico-uterine pouch from the abdominal cavity. Drainage is to be made by three tubes—"one utero-vaginal, one pre-uterine, and a third applied along the uterine wound to the top of the pavilion." This operation has been performed in the interest of the fœtus upon a badly-burned woman, who survived for ten hours; the child was saved.

*Kehrer's Process.*¹—In this form of operation the uterus is to be opened transversely and low down in front. After delivery close the muscular layer of the uterus by from six to ten deep-seated sutures of carbolized silk, and the peritoneal portion by from twelve to twenty-five. Use Listerism in the operation and dressings, abdominal drainage, and vaginal irrigation. This operation has been performed with entire success upon a malacosteon subject of 26 years of age. In a second case it resulted fatally, the wound in the uterus not having been thoroughly closed.

*Sänger's Process.*²—After the abdominal incision is made insert two strong ligatures through the margins of the wound near its upper angle, to be drawn upon after turning out the uterus. Rupture the membranes through the vagina. If practicable, turn out the uterus and hold it vertically. A sheet of caoutchouc, moistened with a 5 per cent. carbolic-acid solution, is to be made to enclose the cervix and cover the abdomen as a protector, and the ligatures are to be drawn while the uterus is being incised vertically in front and evacuated. If the organ is incised *in situ*, manual compression is to be made upon its lower segment as a hæmodynamic. If turned out first, then the same, or the application of clamps to the broad ligaments or of an elastic tube to the cervix. After evacuating the uterus use hæmodynamic pincettes if the edges of the uterine wound bleed. When the uterus is well contracted pass a utero-vaginal drainage-tube, and introduce a carbolized sponge into the uterine cavity; then dissect the peritoneum free from the muscular edges of the uterine wound, and pare from the latter on each side a long slice of tissue of a wedge shape, the thick edge being next to the peritoneal side. Turn in the free edges of the peritoneum over the muscular layer, and unite by deep-seated sutures of silver wire or silk; then bring the serous surfaces of peritoneum in contact, and secure them by numerous superficial sutures.

Three operations under this method, performed by Dr. Leopold of Dresden, saved all the women and children, who continued in good health at the last report. A fourth case, under Dr. Beumer, was lost, as the subject was in bad health from cystitis and pyelo-nephritis, and survived but forty hours; the child was saved. At the late International Medical Congress of Copenhagen, Dr. Leopold expressed the opinion that in future operations he would be able to protect the abdominal cavity against uterine leakage without the resection by folding in the cut edges of the peritoneum in suturing the uterine wound. This was

[¹ *Archiv für Gynäkol.*, 1882, No. 2, vol. xix. p. 180.]

[² *Ibid.*, 1882, No. 3, vol. xix. p. 397.]

proved on autopsy to have been accomplished in the operation of Dr. Garrigues of New York, performed on October 6, 1882.¹ The uterine wound was entirely united, but the woman, who lived 50 hours, never recovered from the effect of a severe ante-partum hemorrhage: her pulse at the time of the operation was 124. Two of the Leopold operations were performed since that of Dr. Garrigues.—Ed.]

[*The Results of the Cæsarean Operation in Great Britain and the United States Compared.*—The fear of, and opposition to, the operation in Great Britain are very natural results of the general fatality which has attended it, even when performed early and by the most skilful hands. It was for a long time claimed that the delay in operating constituted the chief cause of the mortality of the Cæsarean section in the British Isles as compared to that on the Continent; but this can be shown to be an error, as there have been operations enough performed in good season to prove that *gastro-hysterotomy, per se*, is very fatal to British women. I have elsewhere given a record of 28 *early* operations performed in the United States, showing a recovery of 75 per cent. of the women, 23 of the children being delivered alive. By a careful selection from the 138 British cases I have formed a table of 33 operations, 23 of which were performed upon women not over twelve hours in labor, and 10 from twelve to eighteen hours; and the results are as follows:

Labor induced, 1 case; fatal to the mother, child alive.

Labor not commenced, 3 cases; 2 mothers recovered, children all living, 2 premature.

Labor lasting from two to ten hours, inclusive, 11 cases; all fatal, 8 children alive.

Recorded as "early," 1 case; mother saved, child dead.

Labor recorded as having been in progress a few hours; mother and child saved.

Labor from eleven to sixteen hours, inclusive, 8 cases; 1 woman saved, 7 children alive.

Labor from seventeen to eighteen hours, inclusive, 8 cases; 3 women saved, 5 children alive.

Of the 33 cases, 25 averaged in labor eleven and a half hours, and but 4 of them were saved. There were 17 cases not exceeding ten hours in labor, of which 4 were saved; and 16 cases from eleven to eighteen hours in labor, and there were likewise 4 women saved. The mortality of the cases, even when seasonably operated upon, may therefore be set down at 75 per cent., against 25 per cent. of like cases in the United States. The general average of recoveries in our country is now about 40 per cent.; we therefore save more than 2 cases for 1 that recovers in Great Britain; and of seasonable operations the recoveries are as 3 to 1. This difference in our favor can only be accounted for by the difference of physical condition in the subjects. As in the United States and France the women of the farm and village are much more likely to recover than those of the cities, so likewise the better fed and less intemperate poor of our country have a decided advantage over their less-favored sisters in England, where 81 per cent. have perished against 60 per cent. in the United States.—Ed.]

Porro's Operation.—Within the last few years an important modification of the Cæsarean section has been adopted, which is generally known as Porro's operation, from Professor Porro of Pavia, who was

[*American Journ. of Obstetrics*, April, 1883, p. 344.]

the first European surgeon who practised it. In this operation, after the uterus is emptied, the entire organ is drawn out of the abdominal wound and excised, its neck being first constricted so as to suppress hemorrhage, the stump being fixed externally in the manner of the pedicle in ovariectomy. The idea is by no means new. It appears to have been first suggested by an Italian—Dr. Cavallini—in 1768. In 1823 the late Dr. Blundell made the same proposal, and fortified it by numerous experiments on pregnant rabbits, in the course of which he found that he lost all by the Cæsarean section, but saved 3 out of 4 in which he ligatured and amputated the uterus. The suggestion was not, however, carried into actual practice until Dr. Storer of Boston in 1869 removed the uterus in a case of fibroid tumor obstructing the pelvis and impeding delivery.

Since Porro's first case the operation has been frequently performed on the Continent, with results which are, on the whole, encouraging. The cases have been carefully tabulated by Dr. Harris of Philadelphia, and more recently, and very completely, by Dr. Clement Godson,¹ who has collected 138 cases, out of which 61, or 55.8 per cent., were successful as regards the mother. This result is certainly superior to those following the Cæsarean section as ordinarily performed. The obvious advantage of this plan is that instead of leaving the incised uterus with its probably gaping wound and all the attendant risk of septic mischief in the abdominal cavity, it is fixed externally, and in a position where it can be readily dressed.

The objection is that it entirely unsexes the patient, but in the class of women requiring the Cæsarean section from pelvic deformity it is questionable whether this can be fairly considered as a drawback. It is perhaps not justifiable to attempt as yet any positive decision as to the indications for this plan, but it seems beyond doubt that the risks are considerably less than those of the Cæsarean section.

The Operation.—The operation in the successful cases has been performed under the carbolic spray, and the neck of the uterus, after the organ is emptied, carefully secured by ligatures before its body is amputated. Some operators have encircled the neck of the uterus with a chain or wire *écraseur* before removing it, and by this means completely controlled hemorrhage. Richardson² transixed the neck of the uterus with two large pins crossing each other before removing the wire of the *écraseur*, and encircled them with stout carbolized cord. Müller of Berne has recommended that the entire uterus should be turned out of the abdominal cavity through a long incision before it is emptied, so as to avoid the risk of its fluid contents entering the abdomen; but this manœuvre has not always proved feasible. The pedicle has generally been fixed in the lower angle of the abdominal wound and dressed antiseptically. In most cases one or more drainage-tubes have been used, either through Douglas's space or in the abdominal wound.

[*Latest Porro-Cæsarean Statistics*,³—Dr. Clement Godson of London has recently added 15 cases to his former tabular record of 137, and

¹ Porro's Operation, *Brit. Med. Journ.*, Jan. 26, 1884.

² *American Journ. of Med. Science*, July, 1881.

³ *Brit. Med. Journ.*, Jan. 17, 1885, p. 120.]

thereby increased it to 152, and Dr. Ettore Truzzi¹ of Milan has obtained by correspondence 12 more, making the whole 164 cases, saving 69 women and 129 children. The average number of operations per annum for the past five years has been 25; those of 1884, as far as ascertained, amount to 23, with only 9 women saved, although 19 children were living. Of the 164 cases, 109 were operated upon by the method of Porro, many of them very unfavorable, and 46 were saved; 41 by the modification of Müller, with 21 recoveries; and 14 by that of Veit, of dropping in the stump, with 10 deaths. In 6 cases the Müller method failed of completion after making the long incision, and the operations were completed by the plan of Porro, by which 4 were saved. Of 166 children extracted (2 women bore twins), 129 were "living," *but not in a moribund condition*, and 37 were dead or moribund. The Italian operators saved 53 out of 69 children; the Germans saved 18, and lost 10; the Austrians 32, and lost 3; the French 10, and lost 5; English and Scotch 6, and lost 3; the United States 2, and lost 1; Belgians 4, lost 0; and Swiss 2, lost 0.

The operations in private houses number 23, of which 10 recovered and 13 died. The best hospital work was done in Milan, where five operators in Santa Caterina saved 10 women out of 13, and all of the children. In the Krankenhaus of Vienna, 13 women and 23 children were saved by 26 operations. By excluding 3 moribund cases and the 14 Veit experiments from the 164 operations, leaving the Porro and Porro-Müller cases only, we have 147, saving 65 women, or 44 per cent.; which I now regard as the true status of the "Porro operation." These may be subdivided into 90 *favorable* cases, of which 53 recovered, or 58 $\frac{3}{10}$ per cent.; and 57 *unfavorable*, of which only 13 were saved, or less than 23 per cent.

The greatest degree of success has been attained in Austria, where nine operators saved 20 out of 34 women, or nearly 59 per cent. Under forty-three operators in Italy, 28 out of 65 women were saved, or 43 $\frac{1}{2}$ per cent. Germany, under sixteen operators, lost 19 out of 28, and 4 of the 9 saved were by Dr. Fehling of Stuttgart, who lost only 1 out of his 5 cases. France saved 5 out of 15, and Great Britain 1 out of 9. In this enumeration no case is counted where the foetus was *non-viable*. Prof. Breisky of Prague saved all of his 5 cases, and Prof. Porro himself 4 out of 5.—ED.]

Substitute for the Cæsarean Section.—Bearing in mind the great mortality attending the Cæsarean section, it is not surprising that obstetricians should have anxiously considered the possibility of devising a substitute which should afford the mother a better chance of recovery. The first proposal of the kind was one from which great results were at first anticipated. In 1768, Sigault, then a student of medicine in Paris, suggested *symphysectomy*, which consists in the division of the symphysis pubis, with a view of allowing the pubic bones to separate sufficiently to admit of the passage of the child. Although at first strongly opposed, it was subsequently ardently advocated by many obstetricians, and was often performed on the Continent and in a few cases in this country.

The Operation is Admitted to be Useless.—It is generally admitted

[¹ *Annali Universali di Medicina*, Milano, Oct., 1884, p. 387, and Nov., 1884, p. 401.]

that it is quite impossible to make this a substitute for the Cæsarean section, since the utmost gain which even a wide separation of the symphysis pubis would give would be altogether insufficient to admit of the passage of even a mutilated fœtus. Dr. Churchill concludes that even if it were possible to separate it to the extent of four inches, we should only have an increase of from four lines to half an inch in the antero-posterior diameter, in which the obstruction is generally most marked. In the lesser degrees of deformity this might possibly be sufficient to allow the fœtus to pass, but the risk of the operation itself and the subsequent ill-effects altogether contraindicate it in cases of this description.

[*Possibilities of Gastro-hysterotomy.*—Old as it is, the Cæsarean operation is still upon trial, and has by no means attained the minimum of mortality of which it is capable under modern improvements as one of the forms of abdominal surgery. The Porro method has demonstrated the importance of securing the abdominal cavity against the entrance of septic matters escaping from the uterine wound; and its modification by Veit, of dropping in the pedicle, with its far greater fatality, has only tended to confirm our opinion as to the importance of preventing all uterine leakage. But is there no way of rendering the uterine wound as water-tight as that made in the stomach in gastrotomy when secured by the suture of Gely? We believe there is. When the peritoneal coat of the uterus is so stitched, in securing the uterine wound, as to make it form a welt, with its serous surfaces brought in contact, as in the Gely suture, the local adhesive peritonitis which follows will soon effectually secure the wound against the possibility of fluid passing through it. This was demonstrated very satisfactorily in the Garrigues case, already quoted, although the previous condition of the woman gave it a fatal termination; and also in that of Drs. Drysdale in Philadelphia and Jewett in Brooklyn. The results of the three Leopold operations show more decidedly the value of the improvement, having all recovered. As far as ascertained by autopsy, the serous surfaces will unite within thirty hours, provided the uterine muscular tissue is sufficiently firm in texture to enable the deep-seated sutures to hold and prevent gaping. To secure this condition of soundness it is important to operate early, as it is also to save the strength of the patient, and thus secure her against shock and septic peritonitis. The past record of the operation in the United States clearly demonstrates the value of an early use of the knife, and the future ought to show better results when all the additional precautions recently introduced are made use of. In a large proportion of American cases we would not be justified in removing the uterus, as under the Porro method, and it is therefore important to diminish the risks of the old operation. It is also to be considered, in view of past success, whether an improved Cæsarean section will not promise better results than the dangerous expedient of craniotomy and evisceration in cases where the fœtus is impacted in a transverse position. These considerations are purely national, as the records of Great Britain give very little encouragement for performing gastro-hysterotomy. The fact that 25 children were delivered alive from 33 women operated upon within eighteen hours after the commencement of labor

shows that the cases were not lost by delay; yet 25 of these women were lost—a number which we should expect to save in this country.—ED.]

[*The revival of symphyseotomy in Italy*, and its greatly improved results, show that the large mortality of the early cases, and especially of the children, was due to the fœtus having been either turned or forcibly delivered before its head had time to become moulded to the form of the superior strait, the value of which process is set forth by the author on page 397, chap. xii. I do not regard so much the gain in the conjugate diameter as that in the transverse in symphyseotomy: certain it is that the new operation, wherein the fœtus is delivered by the forces of the mother in the large majority of cases, is far less fatal to her and the fœtus than the original one, where the child was turned and forced into the world by traction, to its fatal injury, and the injury of the woman by the strain upon her sacro-iliac symphyses. Under Professors O. Morisani and Novi of Naples 43 women and 42 children were saved by 53 symphyseotomies from 1866 to 1881. "Symphyseotomy can never be made to take the place of the Cæsarean section in cases of extreme deformity, as its advocates are not inclined to recommend it in cases having a conjugate of less measure than 67 millimeters, or 2½ inches."

For a full exposition of the subject see Harris on the "Revival of Symphyseotomy in Italy" in *Am. Journ. of Med. Sci.* for Jan., 1883; also "Una Probabile Risurrezione nel Campo Dell' ostetricia operativa," in the *Annali d'Ostetricia*, anno v., 1883, by Prof. Mangiagalli of the University of Sassari, Italy.—ED.]

CHAPTER VII.

LAPARO-ELYTROTOMY.

IN the former editions of this work laparo-elytrotomy was briefly considered as one of the suggested substitutes for the Cæsarean section which merited careful study and appeared to be of a promising character, but of which too little was known to justify any positive conclusions with regard to it. The subject naturally attracted considerable attention, and several interesting papers have appeared in which its indications, difficulties, and advantages have been carefully considered. Since Thomas's first case was published several operations have been performed, with results so encouraging that I cannot but believe that the operation has a great future before it, and that it will be the duty of the accoucheur to resort to it instead of the more hazardous Cæsarean section, unless some special contraindication exists. Under these circumstances it seems proper no longer to consider it as an addendum to the description of the Cæsarean section, but to study it more in detail in a separate chapter.

History.—The history of the operation is curious and interesting.

The earliest suggestion of a procedure of this character seems to have been made by Joerg in the year 1806, who proposed a modified Cæsarean section, without incision of the uterus, by the division of the linea alba and of the upper part of the vagina, the fœtus being extracted through the cervix. This suggestion was never carried into practice, and it is obvious that it misses the one chief advantage of laparo-elytrotomy, the leaving of the peritoneum intact. In 1820, Ritgen proposed, and actually attempted, an operation much resembling Thomas's, in which section of the peritoneum was avoided. He failed, however, to complete it, and was eventually compelled to deliver his patient by the Cæsarean section. In 1823, Baudelocque the younger independently conceived the same idea, and actually carried it into practice, although without success. Lastly, in 1837, Sir Charles Bell suggested a similar operation, clearly perceiving its advantages. Hence it appears that previous to Thomas's recent work in the matter the operation was independently invented no less than three times. It fell, however, entirely into oblivion, and was only occasionally mentioned in systematic works as a matter of curious obstetric history, no one apparently appreciating the promising character of the procedure.

In the year 1870, Dr. T. Gaillard Thomas of New York read a paper before the Medical Association of the town of Yonkers on the Hudson River, entitled "Gastro-elytrotomy, a Substitute for the Cæsarean Section," in which he described the operation as he had performed it three times on the dead subject and once on a married woman in 1870, with a successful issue as regards the child. It seems beyond doubt that Thomas invented the operation for himself, being ignorant of Ritgen's and Baudelocque's previous attempts, and it is certain, to quote Garrigues,¹ that to him "belongs the glory of having been the first who performed gastro-elytrotomy so as to extract a living child from a living mother in his first operation, and of having brought both mother and child to complete recovery in his second operation."

Since Thomas's first case the operation has been performed three times by Dr. Skene of Brooklyn, and has found its way across the Atlantic, having been twice performed in England, by Hime in Sheffield and by Edis in London.^[2]

Nature of the Operation.—The object of laparo-elytrotomy is to reach the cervix by incision through the lower part of the abdominal wall and upper part of the vagina, and through it to extract the fœtus as may most easily be done.

Advantages over the Cæsarean Section.—If this procedure is found practicable, the enormous advantages it offers over the Cæsarean section are at once apparent, since in dividing the abdomen the abdominal wall

¹ *New York Med. Journ.*, Nov., 1878.

^[2] Thomas operated twice; Skene four times; Charles Jewett of Brooklyn twice; Hime, Edis, Dandridge and Taylor of Cincinnati, and Walter R. Gillette of New York, each once; in all, 12. Women saved, 6; children living, but not moribund, 7; bladder lacerated in 6 cases. In properly calculating the risks of the operation it is fair to exclude the moribund case of Thomas, the intemperate and bedridden one of Hime, and the diseased subject of Edis, who survived, respectively, one hour, two hours, and forty hours. The balance, 9 cases, were favorable in 4 instances and unfavorable in 5: 6 of the 9 women recovered, and 5 children were saved.]

only is incised and the peritoneum is left intact. The vagina is divided, but incision of the uterine parietes, which forms one of the chief risks of the Cæsarean section, is entirely avoided. Now, there is nothing in either of these procedures alarming in itself, and if further experience proves that the practical difficulties of the operation do not stand in the way of its adoption, Dr. Thomas will have introduced, by his able advocacy of the operation, probably the greatest improvement in modern obstetrics.

Cases Suitable for the Operation.—It may be broadly stated that laparo-elytrotomy is applicable in all cases calling for the Cæsarean section when the mother is alive. In post-mortem extractions of the fœtus the Cæsarean section, being the most rapid procedure, would certainly be preferable. Exceptions must be made for certain cases of morbid conditions of the soft parts which render delivery *per vias naturales* impossible, and in which laparo-elytrotomy could not be performed, as in cases of tumor obstructing the pelvic cavity, also in carcinoma or fibroid of the uterus. When the head is firmly impacted in the pelvic brim and cannot be dislodged, the operation would be impossible, as the vagina could not be incised. Unlike the Cæsarean section, the operation cannot be performed twice on the same patient, at least on the same side, since adhesions left by the former incisions would prevent the separation of the peritoneum and division of the vagina. It remains to be seen whether in certain cases of extreme deformity, with pendulous abdomen and distorted thighs, the site of incision might not be so difficult to reach as to render the necessary manœuvres impossible.

It will facilitate the proper comprehension of the operation, and render an avoidance of its possible dangers more easy, if the anatomical relations of the parts concerned are briefly described.

Abdominal Incision.—The abdominal incision extends from a point an inch above the anterior-superior iliac spine, and is carried, with a slight downward curve, parallel to Poupart's ligament until it reaches a point one inch and three-quarters above, and to the outside of, the spine of the pubes. Beyond the latter point it must not extend, so as to avoid the risk of wounding the round ligament and the epigastric artery. In this incision the skin, the aponeurosis of the external oblique, and the fibres of the internal oblique and transversalis muscles are divided. The rectus is not implicated. After the muscles are divided the transversalis fascia is reached. It is fortunately rather dense in this situation, and is separated from the peritoneum by a layer of connective tissue containing fat.

Arteries.—The superficial epigastric artery is necessarily divided, but is too small to give any trouble. The internal epigastric is fortunately not divided, but is so near the inner end of the incision that it may accidentally be so. In one of Dr. Skene's operations it was laid bare. Starting from the external iliac, about a quarter of an inch above Poupart's ligament, it runs downward, forward, and inward to the ligament; thence it turns upward and inward in front of the round ligament and to the inner side of the internal abdominal ring, behind the posterior layer of the sheath of the rectus muscle, which it finally enters. The circumflex iliac artery also rises from the external iliac a little below the

epigastric. It runs between the peritoneum and Poupart's ligament until it reaches the crest of the ilium, to the inner side of which it runs. It thus lies altogether below the line of the incision, and is not likely to be injured.

Peritoneum.—After the transversalis fascia is divided the peritoneum is reached, and is readily lifted up intact, so as to expose the upper part of the vagina, through which the fœtus is extracted. It is fortunate, as facilitating this manœuvre, that the peritoneum is much more lax than in the non-pregnant state, and it has been found very easy to lift it out of the way in all the operations hitherto performed.

Vaginal Incision.—The division of the vagina is the part of the operation likely to give rise to most trouble and risk. It is to be noted that in cases of pelvic contraction calling for this operation the uterus, with its contents, will be abnormally high and altogether above the pelvic brim; the vagina is therefore necessarily elongated and brought more readily within reach. It is enlarged in its upper part during pregnancy, and thrown into folds ready for dilatation during the passage of the child. It is loosely surrounded by the other tissues, and is composed of muscular fibres, easily separable, and an internal mucous layer. Its vascular arrangements are very complex, and the risk of hemorrhage is one of the prominent difficulties of the operation.

In Baudelocque's attempt, in which the vagina was cut instead of torn, the loss of blood was so great as to lead to a discontinuance of the operation. The arteries are numerous, consisting of branches from the hypogastric, inferior vesical, internal pudic, and hemorrhoidal. The veins form a network surrounding the whole canal, but are largest at its extremities, so that it is desirable to open the vagina as low down as possible.

Relations of the Vagina.—Behind the vagina lies the pouch of peritoneum known as Douglas's space, and below that the rectum. In front of it lies the bladder, and the risk of injuring that viscus or the ureter entering it constitutes another of the dangers of the operation. The relations of these parts have been specially studied by Garrigues¹ with the view of facilitating the safe performance of the operation, and I quote his description:

"The anterior-superior surface of the vagina is, in its upper part, bound by loose connective tissue to the bladder on a surface that has the shape of a heart. In the lower or anterior part the boundary-line of this surface runs parallel to and a little outside of the *trigonum vesicale*. In the upper part it follows the outline of the vagina, from which it passes over to the cervix. The distance from the internal opening of the urethra to the neck of the womb is one inch and a quarter (3.2 centimeters). The bladder extends five-eighths of an inch (1.5 centimeters) upon the cervix. It is very liable to be reached by the vaginal rent if the latter is made too high up or too horizontal. The lower part of the antero-superior wall carries in the middle line the *urethra*. In the uppermost part, a little outside of and behind the bladder, lies the *ureter*. In order to avoid the ureter and the bladder, the incision of the vagina should be made nearly an inch and a half (3.8 centimeters) below the

¹ *Loc. cit.*, p. 479.

uterus, and in a direction parallel to the ureter and the boundary-line between the bladder and the vagina."

The Operation.—The operation has hitherto been performed on the right side only. In consequence of the position of the rectum on the left, it seems doubtful if the difficulties of performing it on that side would not render the operation impossible. This point can only be cleared up by experience, and in the mean time the right side should certainly be selected. For the proper performance of the operation four assistants are necessary, besides one who administers the anæsthetic. The patient is placed on her back on the operating-table, with pelvis raised and in the same position as for ovariectomy. In consequence of access of air *per vaginam* strict antiseptic precautions cannot be adopted. Before commencing the operation the cervix is dilated as much as possible by Barnes's bags, assisted, if necessary, by digital dilatation.

The operator stands on the right side of the patient, while an assistant, standing on her left, lays his hand on the uterus and draws it upward and to the left, so as to put the skin on the stretch. The incision is commenced at a point one inch above the anterior-superior spine of the ilium, and is carried inward, in a slightly curved direction, until it reaches a point one and three-quarter inches above and outside the spine of the pubes. The skin and muscular and aponeurotic tissues are carefully divided, layer by layer, any arterial branches being secured as they are severed, until the transversalis fascia is reached. This is raised by a fine tenaculum, and an aperture is made in it, through which a director is introduced, and on this the fascia is divided in the whole length of the superficial incision. The operator now separates the peritoneum from the transversalis and iliac fascia with his fingers, and an assistant, placed on his left, elevates it, as well as the contained intestines, by means of a fine warmed napkin, and keeps it well out of the way during the rest of the operation. A third assistant now introduces a silver catheter into the bladder, and holds it in the position of the boundary-line between it and the vagina and below the uterus.

A blunt wooden instrument like the obturator of a speculum is introduced into the vagina, which is pushed up by it above the ilio-pectineal line. On this an incision is made by Paquelin's thermo-cautery, heated to a red heat only, as far below the uterus as possible, and parallel to the ilio-pectineal line and the catheter felt in the bladder. When the vagina has been burnt through, the index fingers of both hands are pushed through the incision, and the vagina torn through as far forward as is deemed safe by the guide of the catheter in the bladder and as far backward as possible. When this has been done the uterus is depressed to the left and the cervix lifted into the incision by the fingers, and the membranes are ruptured. Through the cervix thus elevated the child is extracted, according to the presentation, either by simple traction, by the forceps, or by turning. Before concluding the operation the bladder should be injected with milk, to make sure that it has not been wounded. Should it be so, the laceration may be at once united by carbolized gut. The principal risk at this stage is hemorrhage from the vaginal vessels, which, however, fortunately did not give rise to much trouble in any of the recent operations. If it occurs, it must be dealt with as best we can,

either by ligature, by the actual cautery, or by thoroughly plugging the vaginal wound with cotton wool both through the incision and *per vaginam*. If the latter be not necessary, the wound should be cleaned by injecting a warm solution of weak carbolized water (2 per cent.), its edges united by interrupted sutures, and dressed as is deemed best. The subsequent treatment must be conducted on general surgical principles, and will much resemble that necessary after other severe abdominal operations, such as ovariectomy. The vagina should be gently syringed two or three times daily with a weak antiseptic lotion. The diet should be light and nutritious, chiefly consisting of milk, beef-tea, and the like. Pain, pyrexia, etc. must be treated as they arise.

[Laparo-elytrotomy has been performed but four times since January, 1880. The skill and number of assistants it requires must necessarily limit its adoption. It may be performed with equal facility on the left side, as was shown in 1878 by the operation of Dr. Hime in England, and in 1883 by that under Drs. Dandridge and Taylor of Cincinnati, in neither of which cases was the bladder injured. Dr. Taylor prefers the left-side operation as more convenient than the right. In 57 of the 134 Cæsarean operations of the United States laparo-elytrotomy would have been impracticable.—Ed.]

CHAPTER VIII.

THE TRANSFUSION OF BLOOD.

THE transfusion of blood in desperate and apparently hopeless cases of hemorrhage offers a possible means of rescuing the patient which merits careful consideration. It has again and again attracted the attention of the profession, but has never become popularized in obstetric practice. The reason of this is not so much the inherent defects of the operation itself—for quite a sufficient number of successful cases are recorded to make it certain that it is occasionally a most valuable remedy—but the fact that the operation has been considered a delicate and difficult one, and that it has been deemed necessary to employ a complicated and expensive apparatus which is never at hand when a sudden emergency arises. Whatever may be the difference of opinion about the value of transfusion, I think it must be admitted that it is of the utmost consequence to simplify the process in every possible way; and it is above all things necessary to show that the steps of the operation are such as can be readily performed by any ordinarily qualified practitioner, and that the apparatus is so simple and portable as to make it easy for any obstetrician to have it at hand. There are comparatively few who would consider it worth while to carry about with them, in ordinary every-day work, cumbrous and expensive instruments which may never be required in a lifelong practice; and hence it is not unlikely that in many cases in which transfusion might have proved useful the

opportunity of using it has been allowed to slip. Of late years the operation has attracted much attention, the method of performing it has been greatly simplified, and I think it will be easy to prove that all the essential apparatus may be purchased for a few shillings, and in so portable a form as to take up little or no room, so that it may be always carried in the obstetric bag ready for any possible emergency.

History of the Operation.—The history of the operation is of considerable interest. In Villari's *Life of Savonarola* it is said to have been employed in the case of Pope Innocent VIII. in the year 1492, but I am not aware on what authority the statement is made. The first serious proposals for its performance do not seem to have been made until the latter half of the seventeenth century. It was first actually performed in France by Denis of Montpellier, although Lower of Oxford had previously made experiments on animals which satisfied him that it might be undertaken with success. In November, 1667, some months after Denis's case, he made a public experiment at Arundel House, in which twelve ounces of sheep's blood were injected into the veins of a healthy man, who is stated to have been very well after the operation; which must therefore have proved successful. These nearly simultaneous cases gave rise to a controversy as to priority of invention which was long carried on with much bitterness.

The idea of resorting to transfusion after severe hemorrhage does not seem to have been then entertained. It was recommended as a means of treatment in various diseased states or with the extravagant hope of imparting new life and vigor to the old and decrepit. The blood of the lower animals only was used; and under these circumstances it is not surprising that the operation, although practised on several occasions, was never established as it might have been had its indications been better understood.

From that time it fell almost entirely into oblivion, although experiments and suggestions as to its applicability were occasionally made, especially by Dr. Harwood, Professor of Anatomy at Cambridge, who published a thesis on the subject in the year 1785. He, however, never carried his suggestions into practice, and, like his predecessors, only proposed to employ blood taken from the lower animals. In the year 1824, Dr. Blundell published his well-known work entitled *Researches, Physiological and Pathological*, which detailed a large number of experiments; and to that distinguished physician belongs the undoubted merit of having brought the subject prominently before the profession and of pointing out the cases in which the operation might be performed with hopes of success. Since the publication of this work transfusion has been regarded as a legitimate operation under special circumstances; but, although it has frequently been performed with success, and in spite of many interesting monographs on the subject, it has never become so established as a general resource in suitable cases as its advantages would seem to warrant. Within the last few years more attention has been paid to the subject, and the writings of Panum, Martin, and De Belina abroad, and of Higginson, McDonnell, Hicks, Aveling, and Schäfer at home, amongst others, have thrown much light on many points connected with the operation.

Nature and Object of the Operation.—Transfusion is practically only employed in cases of profuse hemorrhage connected with labor, although it has been suggested as possibly of value in certain other puerperal conditions, such as eclampsia or puerperal fever. Theoretically, it may be expected to be useful in such diseases, but, inasmuch as little or nothing is known of its practical effects in these diseased states, it is only possible here to discuss its use in cases of excessive hemorrhage. Its action is probably twofold: 1st, the actual restitution of blood which has been lost; 2d, the supply of a sufficient quantity of blood to stimulate the heart to contraction, and thus to enable the circulation to be carried on until fresh blood is formed. The influence of transfusion as a means of restoring lost blood must be trivial, since the quantity required to produce an effect is generally very small indeed, and never sufficient to counterbalance that which has been lost. Its stimulant action is no doubt of far more importance, and if the operation be performed before the vital energies are entirely exhausted the effect is often most marked.

Use of Blood taken from the Lower Animals.—In the earliest operations the blood used was always that of the lower animals, generally of the sheep. It has been thought by Brown-Séquard and others that the blood of some of the lower animals, especially of those in which the corpuscles are of smaller size than in man, as of the sheep, might be used with safety, provided it is not too rich in carbonic acid and too poor in oxygen, and injected in small quantity only. Landois,¹ however, has conclusively proved that the blood of any of the lower animals has a most injurious effect on the human red corpuscles, which rapidly become swollen and decolorized and discharge their coloring matter into the serum. It is certain, therefore, that this plan cannot be adopted in practice.

Difficulties from Coagulation of Fibrin.—The great practical difficulty in transfusion has always been the coagulation of the blood very shortly after it has been removed from the body. When fresh-drawn blood is exposed to the atmosphere, the fibrin commences to solidify rapidly—generally in from three to four minutes, sometimes much sooner. It is obvious that the moment fibrination has commenced the blood is, *ipso facto*, unfitted for transfusion, not only because it can be no longer passed readily through the injecting apparatus, but because of the great danger of propelling small masses of fibrin into the circulation and thus causing embolism. Hence, if no attempt be made to prevent this difficulty, it is essential, no matter what apparatus is used, to hurry on the operation so as to inject before fibrination has begun. This is a fatal objection, for there is no operation in the whole range of surgery in which calmness and deliberation are so essential, the more so as the surroundings of the patient in these unfortunate cases are such as to tax the presence of mind and coolness of the practitioner and his assistants to the utmost.

Methods of Obviating Coagulation.—All the recent improvements have had for their object the avoidance of coagulation; and practically this has been effected in one of three ways: 1st, by immediate transfusion from arm to arm, without allowing the blood to be exposed to the atmosphere,

¹ *Die Transfusion des Blutes*, Leipsic, 1875.

according to the methods proposed by Aveling, Roussel, and Schäfer; 2d, by adding to the blood certain chemical reagents which have the property of preventing coagulation; 3d, removal of the fibrin entirely, by promoting its coagulation and straining the blood, so that the liquor sanguinis and blood-corpuscles alone are injected.

Inasmuch as the success of the operation altogether depends on the method adopted, it will be well, before going farther, to consider briefly the advantages and disadvantages of each of these plans.

Immediate Transfusion: Aveling's Method.—The method of immediate transfusion has been brought prominently before the profession by Dr. Aveling, who has invented an ingenious apparatus for performing it. The apparatus consists essentially of a miniature Higginson's syringe without valves, and with a small silver canula at either end. One canula is inserted into the vein of the person supplying blood, the other into a vein of the patient, and by a curious manipulation of the syringe, subsequently to be described, the blood is carried from one vein into the other. It must be admitted that if there were no practical difficulties this instrument would be admirable; and it is therefore not surprising that it should have met with so much favor from the profession. I cannot but think, however, that the operation is not so simple as at first sight appears, and that therefore it wants one of the essential elements required in any procedure for performing transfusion. One of my objections is that it is by no means easy to work the apparatus without considerable practice. Of this I have satisfied myself by asking members of my class to work it after reading the printed directions, and finding that they are not always able to do so at once. Of course it may be said that it is easy to acquire the necessary manipulative skill; but when the necessity for transfusion arises there is no time left for practising with the instrument, and it is essential that an apparatus to be universally applicable should be capable of being used immediately and without previous experience. Other objections are the necessity of several assistants, the uncertainty of there being a sufficient circulation of blood in the veins of the donor to afford a constant supply, and the possibility of the whole apparatus being disturbed by restlessness or jactitation on the part of the patient. For these reasons it seems to me that this plan of immediate transfusion is not so simple nor so generally applicable as defibrination. Still, it is impossible not to recognize its merits, and it is certainly well worthy of further study and investigation.

Roussel's Method.—Another method of immediate transfusion is that recommended by Roussel,¹ whose apparatus has recently attracted considerable attention. It possesses many undoubted advantages, and is, beyond doubt, a valuable addition to our means of performing the operation. It has, however, the great disadvantage of being costly and complicated, and hence I do not believe that it is likely to come into general use.

Schäfer's Method.—The third method is that recommended by Dr. Schäfer in his recent excellent reports on transfusion submitted to the Obstetrical Society.² Schäfer suggests two methods of performing the operation—one from vein to vein, the other from artery to artery. The

¹ *Obstetrical Transactions*, vol. xviii.

² *Ibid.*, vol. xxi.

latter, he holds, has the advantage of supplying pure oxygenated blood under the best possible conditions for securing the amelioration of a patient suffering from the effects of profuse hemorrhage. The necessary operative proceedings are, however, somewhat complicated, and it seems to me very doubtful if this plan is likely to be at all commonly used. His method of immediate transfusion, however, is very simple, and is well worthy of trial. In his experiments on the lower animals it answered admirably. I am not aware that it has yet been tried on the human subject, but I do not see any practical difficulty in its application. For the description of the operation I have inserted Dr. Schäfer's own directions for the performance of both arterial and venous immediate transfusion.

Addition of Chemical Agents to Prevent Coagulation.—The second plan for obviating the bad effects of clotting is the addition of some substance to the blood which shall prevent coagulation. It is well known that several salts have this property, and the experiments made in the case of cholera patients prove that solutions of some of them may be injected into the venous system without injury. This method has been specially advocated by Dr. Braxton Hicks, who uses a solution of three ounces of fresh phosphate of soda in a pint of water, about six ounces of which are added to the quantity of blood to be injected. He has narrated 4 cases¹ in which this plan was adopted successfully, so far as the prevention of coagulation was concerned. It certainly enables the operation to be performed with deliberation and care, but it is somewhat complicated; and it may often happen that the necessary chemicals are not at hand. A further objection is the bulk of fluid which must be injected; and there is reason to believe that this has in some cases seriously embarrassed the heart's action and interfered with the success of the operation. In many of the successful cases of transfusion the amount of blood injected has been very small, not more than two ounces. Dr. Richardson proposes to prevent coagulation by the addition of liquor ammoniæ to the blood, in the proportion of two minims diluted with twenty minims of water to each ounce of blood.

Defibrination of the Blood.—The last method, and the one which, on the whole, I believe to be the simplest and most effectual, is defibrination. It has been chiefly practised in this country by Dr. McDonnell of Dublin, who has published several very interesting cases in which he employed it, and abroad by Martin of Berlin and De Belina of Paris. The process of removing the fibrin is simple in the extreme, and occupies a few minutes only. Another advantage is that the blood to be transfused may be prepared quietly in an adjoining apartment, so that the operation may be performed with the greatest calmness and deliberation, and the donor is spared the excitement and distress which the sight of the apparently moribund patient is apt to cause, and which, as Dr. Hicks has truly pointed out, may interfere with the free flow of blood. The researches of Panum, Brown-Séquard, and others have proved that the blood-corpuscles are the true vivifying element, and that defibrinated blood acts as well, in every respect, as that containing fibrin. It has been proved that the fibrin is reproduced within a short time,² and the

¹ *Guy's Hospital Reports*, vol. xiv.

² Panum, *Virchow's Arch.*, vol. xxvii.

whole tendency of modern research is to regard it not as an essential element of the blood, but as an excrementitious product, resulting from the degradation of tissue; which may therefore be advantageously removed. Another advantage derived from defibrination is, that the corpuscles are freely exposed to the atmosphere, oxygen is taken up, and carbonic acid given off, and the dangers which Brown-Séquard has shown to arise from the use of blood containing too much carbonic acid are thereby avoided. There can be, therefore, no physiological objection to the removal of the fibrin, which, moreover, takes away all practical difficulty from the operation. The straining to which the defibrinated blood is subjected entirely prevents the possibility of even the most minute particle of fibrin being contained in the injected fluid; the risk from embolism is therefore less than in any of the other processes already referred to. My own experience of this plan is limited to 3 cases, but in 2 it answered so well that I can conceive no reasonable objection to it. I should be inclined to say that transfusion, thus performed, is amongst the simplest of surgical operations—an opinion which the experience of McDonnell and others fully confirms.

Transfusion of Milk.—Recently the intra-venous injection of freshly-drawn warm milk has been recommended as a substitute for blood, chiefly in America. It was first used by Dr. Hodder of Toronto, but has been introduced and strongly advocated by Thomas of New York, who has used it twice after ovariectomy. Brown-Séquard in experimenting on the lower animals found that it answered as well as either fresh or defibrinated blood, and about half an hour after the injection no trace of the milk-corpuscles could be found in the blood. Schäfer, however, found that the action of milk on the blood-corpuscles was highly deleterious, and that it introduces the germs of septic organisms likely to produce very serious results. He therefore pronounces strongly against its use.

Statistical Results.—The number of cases of transfusion are perhaps not sufficient to admit of completely reliable conclusions. It is certain, however, that transfusion has often been the means of rescuing the patient when apparently at the point of death and after all other means of treatment had failed. Professor Martin records 57 cases, in 43 of which transfusion was completely successful, and in 7 temporarily so, while in the remaining 7 no reaction took place. Dr. Higginson of Liverpool has had 15 cases, 10 of which were successful. Figures such as these are encouraging, and they are sufficient to prove that the operation is one which at least offers a fair hope of success, and which no obstetrician would be justified in neglecting when the patient is sinking from the exhaustion of profuse hemorrhage. It is to be hoped also that further experience may prove it to be of value in other cases in which its use has been suggested, but not, as yet, put to the test of experiment.

Possible Dangers of the Operation.—The possible risks of the operation would seem to be the danger of injecting minute particles of fibrin which form emboli, of bubbles of air, or of overwhelming the action of the heart by injecting too rapidly or in too great quantity. These may be, to a great extent, prevented by careful attention to the proper per-

latter, he holds, has the advantage of supplying pure oxygenated blood under the best possible conditions for securing the amelioration of a patient suffering from the effects of profuse hemorrhage. The necessary operative proceedings are, however, somewhat complicated, and it seems to me very doubtful if this plan is likely to be at all commonly used. His method of immediate transfusion, however, is very simple, and is well worthy of trial. In his experiments on the lower animals it answered admirably. I am not aware that it has yet been tried on the human subject, but I do not see any practical difficulty in its application. For the description of the operation I have inserted Dr. Schäfer's own directions for the performance of both arterial and venous immediate transfusion.

Addition of Chemical Agents to Prevent Coagulation.—The second plan for obviating the bad effects of clotting is the addition of some substance to the blood which shall prevent coagulation. It is well known that several salts have this property, and the experiments made in the case of cholera patients prove that solutions of some of them may be injected into the venous system without injury. This method has been specially advocated by Dr. Braxton Hicks, who uses a solution of three ounces of fresh phosphate of soda in a pint of water, about six ounces of which are added to the quantity of blood to be injected. He has narrated 4 cases¹ in which this plan was adopted successfully, so far as the prevention of coagulation was concerned. It certainly enables the operation to be performed with deliberation and care, but it is somewhat complicated; and it may often happen that the necessary chemicals are not at hand. A further objection is the bulk of fluid which must be injected; and there is reason to believe that this has in some cases seriously embarrassed the heart's action and interfered with the success of the operation. In many of the successful cases of transfusion the amount of blood injected has been very small, not more than two ounces. Dr. Richardson proposes to prevent coagulation by the addition of liquor ammoniæ to the blood, in the proportion of two minims diluted with twenty minims of water to each ounce of blood.

Defibrination of the Blood.—The last method, and the one which, on the whole, I believe to be the simplest and most effectual, is defibrination. It has been chiefly practised in this country by Dr. McDonnell of Dublin, who has published several very interesting cases in which he employed it, and abroad by Martin of Berlin and De Belina of Paris. The process of removing the fibrin is simple in the extreme, and occupies a few minutes only. Another advantage is that the blood to be transfused may be prepared quietly in an adjoining apartment, so that the operation may be performed with the greatest calmness and deliberation, and the donor is spared the excitement and distress which the sight of the apparently moribund patient is apt to cause, and which, as Dr. Hicks has truly pointed out, may interfere with the free flow of blood. The researches of Panum, Brown-Séquard, and others have proved that the blood-corpuscles are the true vivifying element, and that defibrinated blood acts as well, in every respect, as that containing fibrin. It has been proved that the fibrin is reproduced within a short time,² and the


¹ *Guy's Hospital Reports*, vol. xiv.

² Panum, *Virchow's Arch.*, vol. xxvii.

of the successful cases an ordinary syringe was used; in one, in the absence of other instruments, a child's toy syringe was employed. I have myself performed it with a simple syringe purchased at the nearest chemist's shop when a special transfusion apparatus failed to act satisfactorily.

Method of Performing Immediate Transfusion.—In immediate transfusion (Fig. 190) the donor is seated close to the patient, and the veins

FIG. 190.



Method of Transfusion by Aveling's Apparatus.

in the arms of each having been opened, the silver canula at either end of the instrument is introduced into them (A B). The tube between the bulb and the donor is now pinched (D), so as to form a vacuum, and the bulb becomes filled with blood from the donor. The finger is now removed so as to compress the distal tube (D'), and, the bulb being compressed (C), its contents are injected into the patient's vein. The bulb is calculated to hold about two drachms, so that the amount injected can be estimated by the number of times it is emptied. The risk of injecting air is prevented by filling the syringe with water, which is injected before the blood.

Schäfer's Directions for Immediate Transfusion.—"Procure two glass canulas of appropriate size and shape (see Fig. 191), and a piece of black india-rubber tubing seven inches long and not less than a quarter of an inch bore, fitted to the canulas. This apparatus could always be improvised.

FIG. 191.



Procedure.—"Place the transfusion-tube in a basin of hot water containing a little carbonate of soda. Put a tape round the arm of the patient just below the place where the vein is to be opened and another just above. Expose the vein by an incision through the skin, which should be made transversely if the position of the vein cannot be made out through the skin. Clear a small piece of the vein with forceps and slip a pointed piece of card underneath it. By a snip with scissors make an oblique opening into the vein, and partly insert a small blunt instrument (such as a wool-needle), so that the aperture is not lost. Remove the upper tape. Next prepare the vein of the

formance of the operation, and it does not clearly appear from the recorded cases that they have ever proved fatal. We must also bear in mind that transfusion is seldom or never likely to be attempted until the patient is in a state which would otherwise almost certainly preclude the hope of recovery, and in which, therefore, much more hazardous proceedings would be fully justified.

Cases Suitable for Transfusion.—The cases suitable for transfusion are those in which the patient is reduced to an extreme state of exhaustion from hemorrhage during or after labor or miscarriage, whether by the repeated losses of placenta prævia or the more sudden and profuse flooding of post-partum hemorrhage. The operation will not be contemplated until other and simpler means have been tried and failed, or until the symptoms indicate that life is on the verge of extinction. If the patient should be deadly pale and cold, with no pulse at the wrist or one that is scarcely perceptible; if she be unable to swallow, or vomits incessantly; if she lie in an unconscious state; if jactitation or convulsions or repeated fainting should occur; if the respiration be laborious or very rapid and sighing; if the pupil do not act under the influence of light,—it is evident that she is in a condition of extreme danger, and it is under such circumstances that transfusion, performed sufficiently soon, offers a fair prospect of success. It does not necessarily follow because one or other of these symptoms is present that there is no chance of recovery under ordinary treatment, and indeed it is within the experience of all that patients have rallied under apparently the most hopeless conditions. But when several of them occur together the prospect of recovery is much diminished, and transfusion would then be fully justified, especially as there is no reason to think that a fatal result has ever been directly traced to its employment. Indeed, like most other obstetric operations, it is more likely to be postponed until too late to be of good service than to be employed too early; and in some of the cases reported as unsuccessful it was not performed until respiration had ceased and death had actually taken place. It has sometimes been said that transfusion should never be employed if the uterus be not firmly contracted, so as to prevent the injected blood again making its escape through the uterine sinuses. The cases in which this is likely to occur are few; and if one were met with the escape of blood could be prevented by the injection into the uterus of the perchloride of iron.

Description of the Operation.—In describing the operation I shall limit myself to an account of Aveling's and Schäfer's method of immediate transfusion and to that of injecting defibrinated blood. I consider myself justified in omitting any account of the numerous instruments which have been invented for the purpose of injecting pure blood, since I believe the practical difficulties are too great ever to render this form of operation serviceable. The great objection to most of them is their cost and complexity, and as long as any special apparatus is considered essential the full benefits to be derived from transfusion are not likely to be realized. The necessity for employing it arises suddenly; it may be in a locality in which it is impossible to procure a special instrument; and it would be well if it were understood that transfusion may be safely and effectually performed by the simplest means. In many

considerable pressure. This would tend to force the tubes apart and might cause copious hemorrhage.

"The transfusion-tube is to be placed as before in carbonate-of-soda solution.

Procedure.—"The artery of the patient must first be exposed. To do this make an incision an inch in length through the skin over the line of the artery, and then divide to an equal extent the subcutaneous tissue and fascia which cover it. About three-quarters of an inch in length of the vessel is to be separated from the ensheathing connective tissue and from its accompanying veins by slipping a blunt instrument, such as an aneurism-needle or the blade of a forceps, underneath and moving it up and down. A small piece of card, cut into a long triangular shape, may then be placed under instead of the needle. A ligature is then tied tightly around the lower end of the piece of artery, another is looped loosely around the middle, and a spring clip is put on close to the upper end. The vessel may now be opened just above the lower ligature by a snip with the scissors.

"If the artery have any branch at the exposed part, this ought to be tied before commencing to isolate the vessel. In the person who is to yield the blood exactly the same process is carried out.

"The transfusion-tube is next filled (by suction) with soda solution, and this is prevented from escaping by one or two spring clips on the tube.

"One of the glass terminals is tied into the artery of the giver and the other into the artery of the patient, the ends of both being directed toward the heart.

"All is now ready for the transfusion. To effect this, remove the clips on the india-rubber tube and open the clip on the artery of the patient; then open—not remove—that on the artery of the giver, and keep it open one minute, or a little longer if it seems advisable. Allow the clips to close again, and if the patient's condition is ameliorated the operation may be ended by tying the arteries—first that of the giver, then that of the patient—just above the clips.

"Finally, cut out and remove the canulas, together with the pieces of artery into which they are tied."

Injection of Defibrinated Blood.—For injecting defibrinated blood various contrivances have been used. McDonnell's instrument is a simple cylinder with a nozzle attached, from which the blood is propelled by gravitation. When the propulsive power is insufficient, increased pressure is applied by breaking forcibly into the open end of the receiver. De Belina's instrument is on the same principle, only atmospheric pressure is supplied by a contrivance similar to Richardson's spray-producer, attached to one end. The idea is simple, but there is some doubt of a gravitation instrument being sufficiently powerful, and it certainly failed in my hands. I have had valves applied to Aveling's instrument, so that it works by compression of the bulb like an ordinary Higginson's syringe. This, with a single silver canula at one end for introduction into the vein, forms a perfect and inexpensive transfusion-apparatus, taking up scarcely any space. If it be not at hand, any small syringe with a tolerably fine nozzle may be used.

Mode of Preparing the Blood.—The first step of the operation is defibrination of the blood, which should, if possible, be prepared in an apartment adjoining the patient's. The blood should be taken from the arm of a strong and healthy man. The quality cannot be unimportant, and in some recorded cases the failure of the operation has been attributed to the fact of the donor having been a weakly female. The supply from a woman might also prove insufficient; and, although it has been shown that blood from two or more persons may be used with safety, yet such a change necessarily causes delay, and should, if possible, be avoided. A vein having been opened, eight or ten ounces of blood are withdrawn, and received into some perfectly clean vessel, such as a dessert finger-glass. As it flows it should be briskly agitated with a clean silver fork or a glass rod, and very shortly strings of fibrin begin to form. It is now strained through a piece of fine muslin, previously dipped in hot water, into a second vessel which is floating in water at a temperature of about 105° . By this straining the fibrin and all air-bubbles resulting from the agitation are removed, and if there be no excessive hurry it might be well to repeat the straining a second time. If the vessel be kept floating in warm water, the blood is prevented from getting cool, and we can now proceed to prepare the arm of the patient for injection.

Mode of Exposing the Veins selected for Transfusion.—This is the most delicate and difficult part of the operation, since the veins are generally collapsed and empty and by no means easy to find. The best way of exposing them is that practised by McDonnell, who pinches up a fold of the skin at the bend of the elbow, and transfixes it with a fine tenotomy-knife or scalpel, so making a gaping wound in the integument, at the bottom of which they are seen lying. A probe should now be passed underneath the vein selected for opening, so as to avoid the chance of its being lost at any subsequent stage of the operation. This is a point of some importance, and from the neglect of this precaution I have been obliged to open another vein than that originally fixed on. A small portion of the vein being raised with the forceps, a nick is made into it for the passage of the canula.

Injection of the Blood.—The prepared blood is now brought to the bedside, and, the apparatus having been previously filled with blood to avoid the risk of injecting any bubbles of air, the canula is inserted into the opening made in the vein and transfusion commenced. It should be constantly borne in mind that this part of the operation should be conducted with the greatest caution, the blood introduced very slowly, and the effect on the patient carefully watched. The injection may be proceeded with until some perceptible effect is produced, which will generally be a return of the pulsation, first at the heart, and subsequently at the wrist, an increase in the temperature of the body, greater depth and frequency of the respirations, and a general appearance of returning animation about the countenance. Sometimes the arms have been thrown about or spasmodic twitchings of the face have taken place. The quantity of blood required to produce these effects varies greatly, but in the majority of cases has been very small. Occasionally 2 ounces have proved sufficient, and the average may be taken as ranging

between 4 and 6, although in a few cases between 10 and 20 have been used. The practical rule is to proceed very slowly with the injection until some perceptible result is observed. Should embarrassed or frequent respiration supervene, we may suspect that we have been injecting either too great a quantity of blood or with too much force and rapidity, and the operation should at once be suspended, and not resumed until the suspicious symptoms have passed away. It may happen that the effects of the transfusion have been highly satisfactory, but that in the course of time there is evidence of returning syncope. This may possibly be prevented by the administration of stimulants, but if these fail there is no reason why a fresh supply of blood should not again be injected, but this should be done before the effects of the first transfusion have entirely passed away.

Secondary Effects of Transfusion.—The subsequent effects in successful cases of transfusion merit careful study. In some few cases death is said to have happened within a few weeks with symptoms resembling pyæmia. Too little is known on this point, however, to justify any positive conclusions with regard to it.

PART V.

THE PUERPERAL STATE.

CHAPTER I.

THE PUERPERAL STATE AND ITS MANAGEMENT.

Importance of Studying the Puerperal State.—The key to the management of women after labor, and to the proper understanding of the many important diseases which may then occur, is to be found in a study of the phenomena following delivery and of the changes going on in the mother's system during the puerperal period. No doubt natural labor is a physiological and healthy function, and during recovery from its effects disease should not occur. It must not be forgotten, however, that none of our patients are under physiologically healthy conditions. The surroundings of the lying-in woman, the effects of civilization, of errors of diet, of defective cleanliness, of exposure to contagion, and of a hundred other conditions which it is impossible to appreciate, have most important influences on the results of childbirth. Hence it follows that labor, even under the most favorable conditions, is attended with considerable risk.

The Mortality of Childbirth.—It is not easy to say with accuracy what is the precise mortality accompanying childbirth in ordinary domestic practice, since the returns derived from the reports of the Registrar-General or from private sources are manifestly open to serious error. The nearest approach to a reliable estimate is that made by Dr. Matthews Duncan,¹ who calculates, from figures derived from various sources, that no fewer than 1 out of every 120 women, delivered at or near the full time, dies within four weeks of childbirth. This indicates a mortality far above that which has been generally believed to accompany childbearing under favorable circumstances. It, however, closely approximates to a similar estimate made by McClintock,² who calculates the mortality in England and Wales as 1 in 126, and in the upper and middle classes alone, where the conditions may naturally be supposed to be more favorable, at 1 in 146; more recently he has come to the conclusion, from his own increased experience and the published results of the practice of others, that 1 in 100 would more correctly represent the rate of puerperal mortality.³ In these calculations there are some obvious sources of error, since they include deaths from all causes within

¹ The "Mortality of Childbed," *Edin. Med. Journ.*, Nov., 1869.

² *Dublin Quarterly Journ.*, Aug., 1869.

³ *Brit. Med. Journ.*, Aug. 10, 1878.

four weeks of delivery, some of which must have been independent of the puerperal state.

But it is not the deaths alone which should be considered. All practitioners know how large a number of their patients suffer from morbid states which may be directly traced to the effects of childbearing. It is impossible to arrive at any statistical conclusion on this point, but it must have a very sensible and important influence on the health of childbearing women.

Alterations in the Blood after Delivery.—The state of the blood during pregnancy, already referred to (p. 140), has an important bearing on the puerperal state. There is hyperinosis, which is largely increased by the changes going on immediately after the birth of the child; for then the large supply of blood which has been going to the uterus is suddenly stopped, and the system must also get rid of a quantity of effete matter thrown into the circulation in consequence of the degenerative changes occurring in the muscular fibres of the uterus. Hence all the depurative channels by which this can be eliminated are called on to act with great energy. If, in addition, the peculiar condition of the generative tract be borne in mind—viz. the large open vessels on its inner surface, the partially bared inner surface of the uterus, and the channels for absorption existing in consequence of slight lacerations in the cervix or vagina—it is not a matter of surprise that septic diseases should be so common.

Condition after Delivery.—It will be well to consider successively the various changes going on after delivery, and then we shall be in a better position for studying the rational management of the puerperal state.

Nervous Shock.—Some degree of nervous shock or exhaustion is observable after most labors. In many cases it is entirely absent, in others it is well marked. Its amount is in proportion to the severity of the labor and the susceptibility of the patient; and it is therefore most likely to be excessive in women who have suffered greatly from pain, who have undergone much muscular exertion, or who have been weakened from undue loss of blood. It is evidenced by a feeling of exhaustion and fatigue, and not uncommonly there is some shivering, which soon passes off, and is generally followed by refreshing sleep. The extreme nervous susceptibility continues for a considerable time after delivery, and indicates the necessity of keeping the lying-in patient as free from all sources of excitement as possible.

Fall of the Pulse.—Immediately after delivery the pulse falls; and the importance of this, as indicating a favorable state of the patient, has already been alluded to. The condition of the pulse has been carefully studied by Blot,¹ who has shown that this diminution, which he believes to be connected with a diminished tension in the arteries due to the sudden arrest of the uterine circulation, continues in a large proportion of cases for a considerable number of days after delivery; and as a matter of clinical import as long as it does the patient may be considered to be in a favorable state. In many instances the slowness of the pulse is remarkable, often sinking to 50, or even 40, beats per minute. Any increase above the normal rate, especially if at all continuous, should

¹ *Arch. gén. de Méd.*, 1864.

always be carefully noted and looked on with suspicion. In connection with this subject, however, it must be remembered that in puerperal women the most trivial circumstances may cause a sudden rise of the pulse. This must be familiar to every practical obstetrician who has constant opportunities of observing this effect after any transient excitement or fatigue. In lying-in hospitals it has generally been observed that the occurrence of any particularly bad case will send up the pulse of all the other patients who may have heard of it.

Temperature in the Puerperal State.—The temperature in the lying-in state affords much valuable information. During and for a short time after labor there is a slight elevation. It soon falls to, or even somewhat below, the normal level. Squire found that the fall occurred within twenty-four hours, sometimes within twelve hours, after the termination of labor.¹ For a few days there is often a slight increase of temperature, especially toward the evening, which is probably caused by the rapid oxidation of tissue in connection with the involution of the uterus. In about forty-eight hours there is a rise connected with the establishment of lactation amounting to one or two degrees over normal level, but this again subsides as soon as the milk is freely secreted. Crédé has also shown² that rapid but transient rises of temperature may occur at any period, connected with trivial causes, such as constipation, errors of diet, or mental disturbances. But if there be any rise of temperature which is at all continuous, especially to over 100° F., and associated with rapidity of the pulse, there is reason to fear the existence of some complication.

The Secretions and Excretions.—The various secretions and excretions are carried on with increased activity after labor. The skin especially acts freely, the patient often sweating profusely. There is also an abundant secretion of urine, but not uncommonly a difficulty of voiding it, either on account of temporary paralysis of the neck of the bladder, resulting from the pressure to which it has been subjected, or from swelling and occlusion of the urethra. For the same reason the rectum is sluggish for a time, and constipation is not infrequent. The appetite is generally indifferent, and the patient is often thirsty.

Secretion of Milk.—Generally in about forty-eight hours the secretion of milk becomes established, and this is occasionally accompanied by a certain amount of constitutional irritation. The breasts often become turgid, hot, and painful. There may or may not be some general disturbance, quickening of pulse, elevation of temperature, possibly slight shivering, and a general sense of oppression, which are quickly relieved as the milk is formed and the breasts emptied by suckling. Squire says that the most constant phenomenon connected with the temperature is a slight elevation as the milk is secreted, rapidly falling when lactation is established. Barker noted elevation either of temperature or pulse in only 4 out of 52 cases which were carefully watched. There can be little doubt that the importance of the so-called "milk fever" has been immensely exaggerated, and its existence as a normal accompaniment of the puerperal state is more than doubtful. It is certain, however, that

¹ "Puerperal Temperatures," *Obstetrical Transactions*, vol. ix.

² *Monat. f. Geburt*, Dec., 1868.

in a small minority of cases there is an appreciable amount of disturbance about the time that the milk is formed. Out of 423 cases, Macan¹ found that in 114, or about 27 per cent., there was no rise in temperature; in 226 the temperature did rise to 100° and over, and of these in 32, or a little over 7 per cent., the only ascertainable cause was a painful or distended condition of the breast. Many modern writers, such as Winckel, Grünewaldt, and D'Espine, entirely deny the connection of this disturbance with lactation, and refer it to a slight and transient septicæmia. Graily Hewitt remarks that it is most commonly met with when the patient is kept low and on deficient diet after delivery, especially when the system is below par from hemorrhage or any other cause. This observation will no doubt account for the comparative rarity of febrile disturbance in connection with lactation in these days, in which the starving of puerperal patients is not considered necessary. It is certain that anything deserving the name of milk fever is now altogether exceptional, and such feverishness as exists is generally quite transient. It is also a fact that it is most apt to occur in delicate and weakly women, especially in those who do not or are unable to nurse. There does not, however, seem to be any sufficient reason for referring it, even when tolerably well marked, to septicæmia. The relief which attends the emptying of the breasts seems sufficient to prove its connection with lactation, and the discomfort which is necessarily associated with the swollen and turgid mammæ is of itself quite sufficient to explain it.

Sugar in the Urine.—In the urine of women during lactation an appreciable amount of sugar may readily be detected. The amount varies according to the condition of the breasts. It increases when they are turgid and congested, and is therefore most abundant in women in whom the breasts are not emptied, as when the child is dead or when lactation is not attempted.

Contraction of the Uterus after Delivery.—Immediately after delivery the uterus contracts firmly, and can be felt at the lower part of the abdomen as a hard, firm mass about the size of a cricket-ball. After a time it again relaxes somewhat, and alternate relaxations and contractions go on at intervals for a considerable time after the expulsion of the placenta. The more complete and permanent the contraction, the greater the safety and comfort of the patient; for when the organ remains in a state of partial relaxation coagula are apt to be retained in its cavity, while for the same reason air enters more readily into it. Hence decomposition is favored and the chances of septic absorption are much increased, while even when this does not occur the muscular fibres are excited to contract and severe after-pains are produced.

Subsequent Diminution in the Size of the Uterus.—After the first few days the diminution in the size of the uterus progresses with great rapidity. By about the sixth day it is so much lessened as to project not more than 1½ or 2 inches above the pelvic brim, while by the eleventh day it is no longer to be made out by abdominal palpation. Its increased size is, however, still apparent per vaginam, and should occasion arise for making internal examination, the mass of the lower segment of the uterus, with its flabby and patulous cervix, can be felt for some

¹ *Dublin Journ. of Med. Science*, May, 1878.

weeks after delivery. This may sometimes be of practical value in cases in which it is necessary to ascertain the fact of recent delivery, and under these circumstances, as pointed out by Simpson, the uterine sound would also enable us to prove that the cavity of the uterus is considerably elongated. Indeed, the normal condition of the uterus and cervix is not regained until six weeks or two months after labor. These observations are corroborated by investigations on the weight of the organ at different periods after labor. Thus, Heschl¹ has shown that the uterus immediately after delivery weighs about 22 to 24 oz., within a week it weighs 19 to 21 oz., and at the end of the second week 10 to 11 oz. only. At the end of the third week it weighs 5 to 7 oz., but it is not until the end of the second month that it reaches its normal weight. Hence it appears that the most rapid diminution occurs during the second week after delivery.

Fatty Transformation of the Muscular Fibres.—The mode in which this diminution in size is effected is by the transformation of the muscular fibres into molecular fat, which is absorbed into the maternal vascular system, which therefore becomes loaded with a large amount of effete material. Heschl has shown that the entire mass of the enlarged uterine muscles are removed, and replaced by newly-formed fibres, which commence to be developed about the fourth week after delivery, the change being complete about the end of the second month. Generally speaking, involution goes on without interruption. It is, however, apt to be interfered with by a variety of causes, such as premature exertion, intercurrent disease, and, very probably, by neglect of lactation. Hence the uterus often remains large and bulky, and the foundation for many subsequent uterine ailments is laid.

Changes in the Uterine Vessels.—Williams has drawn attention to changes occurring in the vessels of the uterus, some of which seem to be permanent, and may, should further observations corroborate his investigations, prove of value in enabling us to ascertain whether a uterus is nulliparous or the reverse—a question which may be of medico-legal importance. After pregnancy he found all the vessels enlarged in calibre. The coats of the arteries are thickened and hypertrophied, and this he has observed even in the uteri of aged women who have not borne children for many years. The venous sinuses, especially at the placental site, have their walls greatly thickened and convoluted, and contain in their centre a small clot of blood (Fig. 192). This thickening attains its greatest dimensions in the third month after gestation, but traces of it may be detected as late as ten or twelve weeks after labor.

Changes in the Uterine Mucous Membrane.—The changes going on in the lining membrane of the uterus immediately after delivery are of great importance in leading to a knowledge of the puerperal state, and have already been discussed when describing the decidua (p. 104). Its cavity is covered with a reddish-gray film formed of blood and fibrin. The open mouths of the uterine sinuses are still visible, more especially over the site of the placenta, and thrombi may be seen projecting from them. The placental site can be distinctly made out in the form of an irregularly oval patch, where the lining membrane is thicker than elsewhere.

¹ *Researches on the Conduct of the Human Uterus after Delivery.*

Contraction of the Vagina, etc.—The vagina soon contracts, and by the time the puerperal month is over it has returned to its normal dimensions, but after childbearing it always remains more lax and less rugose than in nulliparæ. The vulva, at first very lax and much distended, soon regains its former state. The abdominal parietes remain

FIG. 192.



Section of a Uterine Sinus from a Placental Site Nine Weeks after Delivery. (After Williams.)

loose and flabby for a considerable time, and the white streaks produced by the distension of the cutis very generally become permanent. In some women, especially when proper support by bandaging has not been given, the abdomen remains permanently loose and pendulous.

The Lochial Discharge.—From the time of delivery up to about three weeks afterward a discharge escapes from the interior of the uterus known as the *lochia*. At first this consists almost entirely of pure blood, mixed with a variable amount of coagula. If efficient uterine contraction has not been secured after the expulsion of the placenta, coagula of considerable size are frequently expelled with the lochia for one or two days after delivery. In three or four days the distinctly bloody character of the lochia is altered. They have a reddish watery appearance, and are known as the *lochia rubra* or *cruenta*. According to the researches of Wertheimer,¹ they are at this time composed chiefly of blood-corpuscles, mixed with epithelium scales, mucous corpuscles, and the débris of the decidua. The change in the appearance of the discharge progresses gradually, and about the seventh or eighth day it has no longer a red color, but is a pale-greenish fluid with a peculiar sickening and disagreeable odor, and is familiarly described as the “green

¹ *Virchow's Arch.*, 1861.

waters." It now contains a small quantity of blood-corpuscles, which lessen in amount from day to day, but a considerable number of pus-corpuscles, which remain the principal constituent of the discharge until it ceases. Besides these, epithelial scales, fatty granules, and crystals of cholesterin are observed. Occasionally a small infusorium, which has been named the "*Trichomena vaginalis*," has been detected, but it is not of constant occurrence.

Variation in its Amount and Duration.—The amount of the lochia varies much, and in some women it is habitually more abundant than in others. Under ordinary circumstances it is very scanty after the first fortnight, but occasionally it continues somewhat abundant for a month or more without any bad results. It is apt again to become of a red color and to increase in quantity in consequence of any slight excitement or disturbance. If this red discharge continues for any undue length of time, there is reason to suspect some abnormality, and it may not unfrequently be traced to slight lacerations about the cervix which have not healed properly. This result may also follow premature exertion, interfering with the proper involution of the uterus; and the patient should certainly not be allowed to move about as long as much colored discharge is going on.

Occasional Fetor of the Discharge.—Occasionally the lochia have an intensely fetid odor. This must always give rise to some anxiety, since it often indicates the retention and putrefaction of coagula and involves the risk of septic absorption. It is not very rare, however, to observe a most disagreeable odor persist in the lochia without any bad results. The fetor always deserves careful attention, and an endeavor should be made to obviate it by directing the nurse to syringe out the vagina freely night and morning with Condy's fluid and water, while, if it be associated with quickened pulse and elevated temperature, other measures, to be subsequently described, will be necessary.

The After-Pains.—The *after-pains*, which many childbearing women dread even more than the labor-pains, are irregular contractions occurring for a varying time after delivery, and resulting from the efforts of the uterus to expel coagula which have formed in its interior. If, therefore, special care be taken to secure complete and permanent contraction after labor, they rarely occur, or to a very slight extent. Their dependence on uterine inertia is evidenced by the common observation that they are seldom met with in primiparæ, in whom uterine contraction may be supposed to be more efficient, and are more frequent in women who have borne many children. They are a preventable complication, and one which need not give rise to any anxiety: they are, indeed, rather salutary than the reverse, for if coagula be retained in utero, the sooner they are expelled the better. The after-pains generally begin a few hours after delivery, and continue in bad cases for three or four days, but seldom longer. They are generally increased when the mammæ are irritated by suction. When at their height they are often relieved by the expulsion of the coagula. In some severe cases they are apparently neuralgic in character, and do not seem to depend on the retention of coagula. They may be readily distinguished from pains due to more serious causes by feeling the enlarged uterus harden under their influ-

ence, by the uterus not being tender on pressure, and by the absence of any constitutional symptoms.

Management of Women after Delivery.—The management of women after childbirth has varied much at different times according to fashion or theory. The dread of inflammation long influenced the professional mind, and caused the adoption of a strictly antiphlogistic diet, which led to a tardy convalescence. The recognition of the essentially physiological character of labor has resulted in more sound views, with manifest advantage to our patients. The main facts to bear in mind with regard to the puerperal woman are—her nervous susceptibility, which necessitates quiet and absence of all excitement; the importance of favoring involution by prolonged rest; and the risk of septicæmia, which calls for perfect cleanliness and attention to hygienic precautions.

The Administration of Opiates is generally Unadvisable.—As soon as we are satisfied that the uterus is perfectly contracted and that all risk of hemorrhage is over, the patient should be left to sleep. Many practitioners administer an opiate, but, as a matter of routine, this is certainly not good practice, since it checks the contractions of the uterus and often produces unpleasant effects. Still, if the labor have been long and tedious and the patient be much exhausted, 15 or 20 drops of Batley's solution may be administered with advantage.

Attention to the State of the Pulse, Bladder, and Uterus.—Within a few hours the patient should be seen, and at the first visit particular attention should be paid to the state of the pulse, the uterus, and the bladder. The pulse during the whole period of convalescence should be carefully watched, and if it be at all elevated the temperature should at once be taken. If the pulse and temperature remain normal, we may be satisfied that things are going on well; but if the one be quickened and the other elevated, some disturbance or complication may be apprehended. The abdomen should be felt to see that the uterus is not unduly distended and that there is no tenderness. After the first day or two this is no longer necessary.

Treatment of Retention of Urine.—Sometimes the patient cannot at first void the urine, and the application of a hot sponge over the pubes may enable her to do so. If the retention of urine be due to temporary paralysis of the bladder, three or four 20-minim doses of the liquid extract of ergot, at intervals of half an hour, may prove successful. Many hours should not be allowed to elapse without relieving the patient by the catheter, since prolonged retention is only likely to make matters worse. Subsequently, it may be necessary to empty the bladder night and morning until the patient regain her power over it or until the swelling of the urethra subsides, and this will generally be the case in a few days. Occasionally the bladder becomes largely distended, and is relieved to some degree by dribbling of urine from the urethra. Such a state of things may deceive the patient and nurse, and may produce serious consequences by causing cystitis. Attention to the condition of the abdomen will prevent the practitioner from being deceived, for in addition to some constitutional disturbance a large, tender, and fluctuating swelling will be found in the hypogastric region, distinct from the uterus, which it displaces to one or other side. The

catheter will at once prove that this is produced by distension of the bladder.

Treatment of Severe After-Pains.—If the after-pains be very severe an opiate may be administered, or if the lochia be not over-abundant a linseed-meal poultice sprinkled with laudanum or with the chloroform and belladonna liniment may be applied. If proper care have been taken to induce uterine contraction, they will seldom be sufficiently severe to require treatment. In America quinine, in doses of 10 grains twice daily, has been strongly recommended, especially when opiates fail and when the pains are neuralgic in character; and I have found this remedy answer extremely well. The quinine is best given in solution with 10 or 15 minims of hydrobromic acid, which materially lessens the unpleasant head symptoms often accompanying the administration of such large doses.

Diet and Regimen.—The diet of the puerperal patient claims careful attention, the more so as old prejudices in this respect are as yet far from exploded, and as it is by no means rare to find mothers and nurses who still cling tenaciously to the idea that it is essential to prescribe a low regimen for many days after labor. The erroneousness of this plan is now so thoroughly recognized that it is hardly necessary to argue the point. There is, however, a tendency in some to err in the opposite direction, which leads them to insist on the patient's consuming solid food too soon after delivery, before she has regained her appetite, thereby producing nausea and intestinal derangement. Our best guide in this matter is the feelings of the patient herself. If, as is often the case, she be disinclined to eat, there is no reason why she should be urged to do so. A good cup of beef-tea, some bread and milk, or an egg beat up with milk may generally be given with advantage shortly after delivery, and many patients are not inclined to take more for the first day or so. If the patient be hungry, there is no reason why she should not have some more solid but easily-digested food, such as white fish, chicken, or sweetbread; and after a day or two she may resume her ordinary diet, bearing in mind that, being confined to bed, she cannot with advantage consume the same amount of solid food as when she is up and about. Dr. Oldham, in his presidential address to the Obstetrical Society,¹ has some apposite remarks on this point which are worthy of quotation: "A puerperal month under the guidance of a monthly nurse is easily drawn out, and it is well if a love of the comforts of illness and the persuasion of being delicate, which are the infirmities of many women, do not induce a feeble life which long survives after the occasion of it is forgotten. I know no reason why, if a woman is confined early in the morning, she should not have her breakfast of tea and toast at nine, her luncheon from some digestible meat at one, her cup of tea at five, her dinner with chicken at seven, and her tea again at nine, or the equivalent, according to the variation of her habits of living. Of course, there is the common-sense selection of articles of food, guarding against excess and avoiding stimulants. But gruel and slops and all intermediate feeding are to be avoided." No one who has seen both methods adopted can fail to have been struck with the more rapid and satisfactory conva-

¹ *Obstet. Trans.*, vol. vi.

lescence which takes place when the patient's strength is not weakened by an unnecessarily low diet. Stimulants, as a rule, are not required, but if the patient be weakly and exhausted, or if she be accustomed to their use, there can be no reasonable objection to their judicious administration. [As a rule, in the United States puerperal women have not been accustomed to the use of stimulants, and such are not advisable during their convalescence. The old system of starvation for three days has been done away with, but the opposite is also to be avoided. I believe in a spare diet for the robust and in feeding up the delicate. *Where much blood has been lost* I have found great benefit from the use of essence of beef, given in severe cases quite largely. As a rule, that made from three pounds of beef is to be given daily for two weeks, but I have used in a very extreme case as high as eleven pounds. The removal of the anæmic pallor is sometimes very decided within fifteen days under this diet. Of course the patient is to have a regular diet in addition to the essence of beef.—ED.]

Attention to Cleanliness, etc.—Immediately after delivery a warm napkin is applied to the vulva, and after the patient has rested a little the nurse removes the soiled linen from the bed and washes the external genitals. It is impossible to pay too much attention during the subsequent progress of the case to the maintenance of perfect cleanliness. Perfectly antiseptic midwifery is no doubt an impossibility, but a near approach to it may be made, and the greater the care taken the more certainly will the safety of the patient be ensured.¹ It will be a wise precaution to advise the nurse never to touch the genitals for the first few days unless her hands have been moistened in a 1-in-20 solution of carbolic acid or a 1-in-1000 solution of perchloride of mercury, or lubricated with carbolized oil. The linen should be frequently changed, and

¹ The following rules I have for the past year or two distributed to the monthly nurses attending my own patients, with the result, I believe, of a marked improvement in their comfort and a more generally satisfactory convalescence:

ANTISEPTIC RULES FOR MONTHLY NURSES.

1. Two bottles are supplied to each patient. One contains a mixture of phenol or pure carbolic acid, of the strength of one part to twenty of water (called the 1-in-20 solution), the other carbolized oil (1-in-8).

2. A small basin containing the 1-in-20 solution must always stand by the bedside of the patient, and the nurse must *thoroughly* rinse her hands in it every time she touches the patient in the neighborhood of the genital organs, for washing or any other purpose whatsoever, before or during labor and for a week after delivery.

3. All sponges, vaginal and rectal pipes, catheters, etc. must be dipped in the 1-in-20 solution before being used. The surfaces of slippers, bedpans, etc. should also be sponged with it.

4. Vaginal pipes, enema-tubes, catheters, etc. should be smeared with the carbolized oil before use.

5. Unless express directions are given to the contrary, the vagina should be syringed twice daily after delivery with the 1-in-20 solution, with an equal quantity of hot water added to it. Occasionally the patient finds that this smarts a little, in which case the quantity of warm water may be slightly increased.

6. All water used for washing should have sufficient Condly's fluid dropped into it to give it a pale pink color.

7. All soiled linen, diapers, etc. should be immediately removed from the bedroom.

N. B.—These rules are for the purpose of protecting the patient from the risk arising from accidental contamination of the hands, sponges, etc. It is therefore hoped that they will be faithfully and minutely adhered to.

all dirty linen and discharges immediately removed from the apartment. The vulva should be washed daily with Condy's fluid and water, and the patient will derive great comfort from having the vagina syringed gently out once a day with the same solution. The remarkable diminution of mortality which has followed such antiseptic precautions in certain lying-in hospitals in Germany well shows the importance of these measures. The room should be kept tolerably cool and fresh air freely admitted.

Action of the Bowels.—It is customary on the morning of the second or third day to secure an action of the bowels; and there is no better way of doing this than by a large enema of soap and water. If the patient object to this and the bowels have not acted, some mild aperient may be administered, such as a small dose of castor oil, a few grains of colocynth and henbane pill, or the popular French aperient the "Tamar Indien."

Lactation.—The management of suckling and of the breasts forms an important part of the duties of the monthly nurse which the practitioner should himself superintend. This will be more conveniently discussed under the head of Lactation.

Importance of Prolonged Rest.—The most important part of the management of the puerperal state is the securing to the patient prolonged rest in the horizontal position, in order to favor proper involution of the uterus. For the first few days she should be kept as quiet and still as possible, not receiving the visits of any but her nearest relatives, thus avoiding all chance of undue excitement. It is customary among the better classes for the patient to remain in bed for eight or ten days; but, provided she be doing well, there can be no objection to her lying on the outside of the bed or slipping on to a sofa somewhat sooner. After ten days or a fortnight she may be permitted to sit on a chair for a little; but I am convinced that the longer she can be persuaded to retain the recumbent position, the more complete and satisfactory will be the progress of involution; and she should not be allowed to walk about until the third week, about which time she may also be permitted to take a drive. If it be borne in mind that it takes from six weeks to two months for the uterus to regain its natural size, the reason for prolonged rest will be obvious. The judicious practitioner, however, while insisting on this point, will take measures at the same time not to allow the patient to lapse into the habits of an invalid or to give the necessary rest the semblance of disease.

Subsequent Treatment.—Toward the termination of the puerperal month some slight tonic, such as small doses of quinine with phosphoric acid, may be often given with advantage, especially if convalescence be tardy. Nothing is so beneficial in restoring the patient to her usual health as change of air, and in the upper classes a short visit to the seaside may generally be recommended, with the certainty of much benefit.

CHAPTER II.

MANAGEMENT OF THE INFANT, LACTATION, ETC.

Commencement of Respiration.—Almost immediately after its expulsion a healthy child cries aloud, thereby showing that respiration is established; and this may be taken as a signal of its safety. The first respiratory movements are excited partially by reflex action resulting from the contact of the cold external air with the cutaneous nerves, and partly by the direct irritation of the medulla oblongata in consequence of the circulation through it of blood no longer oxygenated in the placenta.

Apparent Death of the New-born Child.—Not infrequently the child is born in an apparently lifeless state. This is especially likely to be the case when the second stage of labor has been unduly prolonged, so that the head has been subjected to long-continued pressure. The utero-placental circulation is also apt to be injuriously interfered with before the birth of the child when a tardy labor has produced tonic contraction of the uterus and consequent closure of the uterine sinuses, or, more rarely, from such causes as the injudicious administration of ergot, premature separation of the placenta, or compression of the umbilical cord. In any of these cases it is probable that the arrest of the utero-placental circulation induces attempts at inspiration which are necessarily fruitless, since air cannot reach the lungs, and the foetus may die asphyxiated; the existence of the respiratory movement being proved on post-mortem examination by the presence in the lungs of liquor amnii, mucus, and meconium, and by the extravasation of blood from the rupture of their engorged vessels.

Appearance of the Child in such Cases.—In most cases, when the child is born in a state of apparent asphyxia, its face is swollen and of a dark livid color. It not infrequently makes one or two feeble and gasping efforts at respiration, without any definite cry; on auscultation the heart may be heard to beat weakly and slowly. Under such circumstances there is a fair hope of its recovery. In other cases the child, instead of being turgid and livid in the face, is pale, with flaccid limbs, and no appreciable cardiac action; then the prognosis is much more unfavorable.

Treatment of Apparent Death.—No time should be lost in endeavoring to excite respiration; and at first this must be done by applying suitable stimulants to the cutaneous nerves in the hope of exciting reflex action. The cord should be at once tied and the child removed from the mother, for the final uterine contractions have so completely arrested the utero-placental circulation as to render it no longer of any value. If the face be very livid, a few drops of blood may with advantage be allowed to flow from the cord before it is tied, with the view of relieving the embarrassed circulation. Very often some slight stimulus, such as one or two sharp slaps on the thorax or rapidly rubbing the body with

brandy poured into the palms of the hands, will suffice to induce respiration. Failing this, nothing acts so well as the sudden and instantaneous application of heat and cold. For this purpose extremely hot water is placed in one basin, and quite cold water in another. Taking the child by the shoulders and legs, it should be dipped for a single moment into the hot water and then into the cold; and these alternate applications may be repeated once or twice as occasion requires. The effect of this measure is often very marked, and I have frequently seen it succeed when prolonged efforts at artificial respiration had been made in vain.

Artificial Respiration.—If these means fail an endeavor must be at once made to carry on respiration artificially. The Sylvester method is, on the whole, that which is most easily applied, and, on account of the compressibility of the thorax, it is peculiarly suitable for infants. The child being laid on its back with the shoulders slightly elevated, the elbows are grasped by the operator and alternately raised above the head and slowly depressed against the sides of the thorax, so as to produce the effect of inspiration and expiration. If this do not succeed, the Marshall Hall method may be substituted, and one or more of the plans of exciting reflex action through the cutaneous nerves may be alternated with it.

Insufflation of the Lungs.—Other means of exciting respiration have been recommended. One of them, much used abroad, is the artificial insufflation of the lungs by means of a flexible catheter guided into the glottis. It is not difficult to pass the end of a catheter into the glottis, using the little finger as a guide; and, once in position, it may be used to blow air gently into the lungs, which is expelled by compression on the thorax, the insufflation being repeated at short intervals of about ten seconds. One advantage of this plan is that it allows the liquor amnii and other fluids, which may have been drawn into the lungs in the premature efforts at respiration before birth, to be sucked up into the catheter, and so removed from the lungs. The same effect may be produced, but less perfectly, by placing the hand over the nostrils of the child, blowing into its mouth, and immediately afterward compressing the thorax.[¹] One of these two methods should certainly be tried if all other means have failed. Faradization along the course of the phrenic nerves is a promising means of inducing respiration which should be used if the proper apparatus can be procured. Encouragement to persevere in our endeavors to resuscitate the child may be derived from the numerous authenticated instances of success after the lapse of a considerable time, even of an hour or more. As long as the cardiac pulsations continue, however feebly, there is no reason to despair.

Washing and Dressing of the Child.—When the child cries lustily from the first, it is customary for the nurse to wash and dress it as soon as her immediate attendance on the mother is no longer required. For this purpose it is placed in a bath of warm water and carefully soaped and sponged from head to foot. With the view of facilitating the removal of the unctuous material with which it is covered, it is usual to anoint it with cold cream or olive oil, which is washed off in the

[¹ The œsophagus must be closed by placing the thumb and fingers on either side of the larynx and pressing it back, or you will inflate the stomach instead of the lungs.—Ed.]

bath. Nurses are apt to use undue roughness in endeavoring to remove every particle of the vernix caseosa, small portions of which are often firmly adherent. This mistake should be avoided, as these particles will soon dry up and become spontaneously detached. The cord is generally wrapped in a small piece of charred linen, which is supposed to have some slight antiseptic property, and this is renewed from day to day until the cord has withered and separated. This generally occurs within a week, and a small pad of soft linen is then placed over the umbilicus, and supported by a flannel belly-band placed round the abdomen, which should not be too tight, for fear of embarrassing the respiration. By this means the tendency to umbilical hernia is prevented. [As the vernix caseosa is readily miscible with pure lard, and can be easily removed by its means, it has become the practice with many obstetricians in the United States to order the infant to be well anointed, and then wiped from head to foot with soft rags, until all the vernix disappears, and the skin retains a slight oily trace, not enough to soil the clothing. By this means water is avoided, and with it much of the risk of taking cold, and the skin is left less sensitive after the sudden change which it is made to endure at birth than when subjected to hot water and soap. In the hot months water is preferable at the first dressing.—ED.]

Clothing, etc.—The clothing of the infant varies according to fashion and the circumstances of the parents. The important points to bear in mind are that it should be warm (since newly-born children are extremely susceptible to cold), and at the same time light and sufficiently loose to allow free play to the limbs and thorax. All tight bandaging and swaddling, such as is so common in some parts of the Continent, should be avoided, and the clothes should be fastened by strings or by sewing, and no pins used. At the present day it is customary not to use caps, so that the head may be kept cool. The utmost possible attention should be paid to cleanliness, and the child should be regularly bathed in tepid water—at first once daily, and after the first few weeks both night and morning. After drying, the flexures of the thighs and arms and the nates should be dusted with violet powder or fuller's earth, to prevent chafing of the skin. The excrements should be received in napkins wrapped round the hips, and great care is required to change the napkins as often as they are wet or soiled, otherwise troublesome irritation will arise. A neglect of this precaution and the washing of the napkins with coarse soap or soda are among the principal causes of the eruptions and excoriations so common in badly cared-for children. When washed and dressed the child may be placed in its cradle and covered with soft blankets or an eider-down quilt.

Application of the Child to the Breast.—As soon as the mother has rested a little it is advisable to place the child to the breast. This is useful to the mother by favoring uterine contraction. Even now there is in the breasts a variable quantity of the peculiar fluid known as *colostrum*. This is a viscid yellowish secretion, different in appearance from the thin bluish milk which is subsequently formed. Examined under the microscope, it is found to contain some milk-globules and a number of large granular and small fat-corpuscles. It has a purgative property, and soon produces, with less irritation than any of the laxa-

tives so generally used, a discharge of the meconium with which the bowels are loaded. Hence the accoucheur should prohibit the common practice of administering castor oil or other aperient within the first few days after birth, although there can be no objection to it in special cases if the bowels appear to act inefficiently and with difficulty.

Over-frequent Suckling should be Avoided.—For the first few days, and until the secretion of milk is thoroughly established, the child should be put to the breast at long intervals only. Constant attempts at sucking an empty breast lead to nothing but disappointment, both to the mother and child, and, by unduly irritating the mammae, sometimes to positive harm. Therefore, for the first day or two it is sufficient if the child be applied to the breast twice, or at most three times, in the twenty-four hours. Nor is it necessary to be apprehensive, as many mothers naturally are, that the child will suffer from want of food. A few spoonfuls of milk and water being given from time to time, the child may generally wait without injury until the milk is secreted. This is generally about the third day, when the secretion is found to be a whitish fluid, more watery in appearance than cow's milk, and showing under the microscope an abundance of minute spherical globules, refracting light strongly, which are abundant in proportion to the quality of the milk. A certain number of granular corpuscles may also be observed shortly after the birth of the child, but after the first month these should have almost altogether disappeared. The reaction of human milk is decidedly alkaline, and the taste much sweeter than that of cow's milk.

Importance of Nursing when Practicable.—The importance to the mother of nursing her own child whenever her health permits, on account of the favorable influence of lactation in promoting a proper involution of the uterus, has already been insisted on. Unless there be some positive contraindication, such as a marked strumous cachexia, an hereditary phthisical tendency, or great general debility, it is the duty of the accoucheur to urge the mother to attempt lactation, even if it be not carried on more than a month or two. It is, however, the fact that in the upper classes of society a large number of patients are unable to nurse, even though willing and anxious to do so. In some there is hardly any lacteal secretion at all; in others there is at first an overabundance of watery and innutritious milk, which floods the breasts and soon dies away altogether.

When the Mother cannot Nurse, a Wet-nurse should be Procured.—Whenever the mother cannot or will not nurse the question will arise as to the method of bringing up the child. From many causes there is an increasing tendency to resort to bottle-feeding, instead of procuring the services of a wet-nurse, even when the question of expense does not come into consideration. No long experience is required to prove that hand-feeding is a bad and imperfect substitute for nature's mode, and one which the practitioner should discourage whenever it lies in his power to do so. It is true that in many cases bottle-fed children do well, but there is good reason to believe that, even when apparently most successful, the children are not so strong in after-life as they would have been had they been brought up at the breast. When, in addition, it is borne in mind how much of the success of hand-feeding depends

on intelligent care on the part of the nurse, what evils are apt to accrue from the injurious selection of the food and from ignorance of the commonest laws of dietetics, there is abundant reason for urging the substitution of a wet-nurse whenever the mother is unable to undertake the suckling of her child. It must be admitted that good hand-feeding is better than bad wet-nursing, and the success of the latter hinges on the proper selection of a wet-nurse. As this falls within the duties of the practitioner, it will be well to point out the qualities which should be sought for in a wet-nurse, before proceeding to discuss the mode of rearing the child at the breast.

Selection of a Wet-nurse.—In selecting a wet-nurse we should endeavor to choose a strong, healthy woman, who should not be over 30, or 35 years of age at the outside, since the quality of the milk deteriorates in women who are more advanced in life. For a similar reason a very young woman of 16 or 17 should be rejected. It is needless to say that care must be taken to ascertain the absence of all traces of constitutional disease, especially marks of scrofula or enlarged cervical or inguinal glands, which may possibly be due to antecedent syphilitic taint. If the nurse be of good muscular development, healthy-looking, with a clear complexion, and sound teeth (indicating a generally good state of health), the color of the hair and eyes is of secondary importance. It is commonly stated that brunettes make better nurses than blondes, but this is by no means necessarily the case, and, provided all the other points be favorable, fairness of skin and hair need be no bar to the selection of a nurse. The breasts should be pear-shaped, rather firm, as indicating an abundance of gland-tissue, and with the superficial veins well marked. Large, flabby breasts owe much of their size to an undue deposit of fat, and are generally unfavorable. The nipple should be prominent, not too large, and free from cracks and erosions, which, if existing, might lead to subsequent difficulties in nursing. On pressing the breast the milk should flow from it easily in a number of small jets, and some of it should be preserved for examination. It should be of a bluish-white color, and when placed under the microscope the field should be covered with an abundance of milk-corpuscles and the large granular corpuscles of the colostrum should have entirely disappeared. If the latter be observed in any quantity in a woman who has been confined five or six weeks, the inference is that the milk is inferior in quality. It is not often that the practitioner has an opportunity of inquiring into the moral qualities of the nurse, although much valuable information might be derived from a knowledge of her previous character. An irascible, excitable, or highly-nervous woman will certainly make a bad nurse, and the most trivial causes might afterward interfere with the quality of her milk. Particular attention should be paid to the nurse's own child, since its condition affords the best criterion of the quality of her milk. It should be plump, well-nourished, and free from all blemishes. If it be at all thin and wizened, especially if there be any snuffling at the nose, or should any eruption exist affording the slightest suspicion of a syphilitic taint, the nurse should be unhesitatingly rejected.

Management of Suckling.—The management of suckling is much the same whether the child is nursed by the mother or by a wet-nurse. As

soon as the supply of milk is sufficiently established, the child must be put to the breast at short intervals—at first of about two hours, and in about a month or six weeks of three hours. From the first few days it is a matter of the greatest importance, both to the mother and child, to acquire regular habits in this respect. If the mother get into the way of allowing the infant to take the breast whenever it cries as a means of keeping it quiet, her own health must soon suffer, to say nothing of the discomfort of being incessantly tied to the child's side; while the child itself has not sufficient rest to digest its food, and very shortly diarrhoea or other symptoms of dyspepsia are pretty sure to follow. After a month or two the infant should be trained to require the breast less often at night, so as to enable the mother to have an undisturbed sleep of six or seven hours. For this purpose she should arrange the times of nursing so as to give the breast just before she goes to bed, and not again until the early morning. If the child should require food in the interval, a little milk and water from the bottle may be advantageously given.

Diet of Nursing Women.—The diet of the nursing woman should be arranged on ordinary principles of hygiene. It should be abundant, simple, and nutritious, but all rich and stimulating articles of food should be avoided. A common error in the diet of wet-nurses is over-feeding, which constantly leads to deterioration of the milk. Many of these women before entering on their functions have been living on the simplest and even sparest diet, and not uncommonly, in the better class of houses, they are suddenly given heavy meat meals three and even four times a day, and often three or four glasses of stout. It is hardly a matter of astonishment that under such circumstances their milk should be found to disagree. For a nursing woman in good health two good meat meals a day, with two glasses of beer or porter, and as much milk and bread and butter as she likes to take in the intervals, should be amply sufficient. Plenty of moderate exercise should be taken, and the more nurse and child are out in the open air, provided the weather be reasonably fine, the better it is for both. [As a rule, American wet-nurses have been much better fed than those here described, and have not been in the habit of using malt drinks. A healthy woman will usually nurse well on her ordinary diet, which should be largely farinaceous. If she can drink milk, there is nothing equal to it in furnishing a lacteal supply.—ED.]

Signs of Successful Lactation.—Carried on methodically in this manner, wet-nursing should give but little trouble. In the intervals between its meals the child sleeps most of its time, and wakes with regularity to feed; but if the child be wakeful and restless, cry after feeding, have disordered bowels, and, above all, if it do not gain week by week in weight (a point which should be from time to time ascertained by the scales), we may conclude that there is either some grave defect in the management of suckling or that the milk is not agreeing. Should this unsatisfactory progress continue in spite of our endeavors to remedy it, there is no resource left but the alteration of the diet, either by changing the nurse or by bringing up the child by hand. The former should be preferred whenever it is practicable, and in the upper ranks of life it

is by no means rare to have to change the wet-nurse two or three times before one is met with whose milk agrees perfectly. If the child have reached six or seven months of age, it may be preferable to wean it altogether, especially if the mother have nursed it, as hand-feeding is much less objectionable if the infant have had the breast for even a few months.

Period of Weaning.—As a rule, weaning should not be attempted until dentition is fairly established, that being the sign that nature has prepared the child for an alteration of food; and it is better that the main portion of the diet should be breast milk until at least six or seven teeth have appeared. This is a safer guide than any arbitrary rule taken from the age of the child, since the commencement of dentition varies much in different cases. About the sixth or seventh month it is a good plan to commence the use of some suitable artificial food once a day, so as to relieve the strain on the mother or nurse and prepare the child for weaning, which should always be a very gradual process. In this way a meal of rusks of the entire wheat flour, or of beef- or chicken-tea with bread-crumbs in it, may be given with advantage; and as the period for weaning arrives a second meal may be added, and so eventually the child may be weaned without distress to itself or trouble to the nurse.

The Disorders of Lactation.—The disorders of lactation are numerous, and, as they frequently come under the notice of the practitioner, it is necessary to allude to some of the most common and important.

Means of Arresting the Secretion of Milk.—The advice of the accoucheur is often required in cases in which it has been determined that the patient is not to nurse, when we desire to get rid of the milk as soon as possible, or when, at the time of weaning, the same object is sought. The extreme heat and the distension of the breasts in the former class of cases often give rise to much distress. A smart saline aperient will aid in removing the milk, and for this purpose a double Seidlitz powder or frequent small doses of sulphate of magnesia act well, while, at the same time, the patient should be advised to take as small a quantity of fluid as possible. Iodide of potassium in large doses of 20 or 25 grains, repeated twice or thrice, has a remarkable effect in arresting the secretion of milk. This observation was first empirically made by observing that the secretion of milk was arrested when this drug was administered for some other cause; and I have frequently found it answer remarkably well. The distension of the breasts is best relieved by covering them with a layer of lint or cotton wool soaked in a spirit lotion or eau de Cologne and water, over which oiled silk is placed, and by directing the nurse to rub them gently with warm oil whenever they get hard and lumpy. Breast-pumps and similar contrivances only irritate the breasts, and do more harm than good. The local application of belladonna has been strongly recommended as a means for preventing lacteal secretion. As usually applied, in the form of belladonna plaster, it is likely to prove hurtful, since the breast often enlarges after the plasters are applied, and the pressure of the unyielding leather on which they are spread produces intense suffering. A better way of using it is by rubbing down a drachm of the extract of belladonna with an ounce of glycerin and applying this on lint. In some cases it answers extremely

well, but it is very uncertain in its action, and frequently is quite useless.

Defective Secretion of Milk.—A deficiency of milk in nursing mothers is a very common source of difficulty. In a wet-nurse this drawback is, of course, an indication for changing the nurse; but to the mother the importance of nursing is so great that an endeavor must be made either to increase the flow of milk or to supplement it by other food. Unfortunately, little reliance can be placed on any of the so-called galactagogues. The only one which in recent times has attracted attention is the leaves of the castor-oil plant, which, made into poultices and applied to the breast, are said to have a beneficial effect in increasing the flow of milk. More reliance must be placed in the sufficiency of nutritious food, especially such as contains phosphatic elements; stewed eels, oysters, and other kinds of shellfish, and the Revalenta Arabica, are recommended by Dr. Routh, who has paid some attention to this point,¹ as peculiarly appropriate. If the amount of milk be decidedly deficient, the child should be less often applied to the breast, so as to allow milk to collect, and properly prepared cow's milk from a bottle should be given alternately with the breast. This mixed diet generally answers well, and is far preferable to pure hand-feeding. [There is no diet equivalent to milk for a nursing mother, where it agrees with her. This I have tested repeatedly in women who had failed entirely in former attempts to nurse their infants. One lady who had lost her milk three times at the end of a month, and had nursed two babies into starvation, was enabled to nurse her fourth while on a milk diet for eighteen months, and gained while doing so 19 pounds. Another gained 65 pounds while nursing, and her son was very large for his age. A third lost a child by hand-feeding, and nursed the next infant on a milk diet, at the same time becoming fatter than she had ever been. A decided advantage in the use of milk is, that it prevents the exhausted feeling so common with delicate nursing mothers. I have had a patient of 86 pounds weight use two quarts of milk a day, and at the same time eat her usual measure of food, which had always been of small amount.—Ed.]

Depressed Nipples.—A not uncommon source of difficulty is a depressed condition of the nipples, which is generally produced by the constant pressure of the stays. The result is, that the child, unable to grasp the nipple and wearied with ineffectual efforts, may at last refuse the breast altogether. An endeavor should be made to elongate the nipple before putting it into the child's mouth, either by the fingers or by some form of breast-pump, which here finds a useful application. In the worst class of cases, when the nipple is permanently depressed, it may be necessary to let the child suck through a glass nipple shield to which is attached an india-rubber tube, similar to that of a sucking-bottle; this it is generally well able to do. [In some instances this anatomical defect appears to be beyond remedy, unless a proposed surgical operation can be made effective. I have tried to prepare primiparæ for several months before labor, and then failed as soon as the breasts filled with milk. In some cases there is absolutely no nipple, and as a shield is of no value in protection, the escaping milk produces an eczema over the waist and

¹ Routh on *Infant-feeding*.

upper part of the abdomen. This condition I have seen associated with a most obstinate galactorrhœa lasting several months.—ED.]

Fissures and Excoriations of the Nipples.—Fissures and excoriations of the nipples are common causes of suffering, in some cases leading to mammary abscess. Whenever the practitioner has the opportunity, he should advise his patient to prepare the nipple for nursing in the latter months of pregnancy; and this may best be done by daily bathing it with a spirituous or astringent lotion, such as eau de Cologne and water or a weak solution of tannin. After nursing has begun great care should be taken to wash and dry the nipple after the child has been applied to it, and as long as the mother is in the recumbent position she may, if the nipples be at all tender, use zinc nipple-shields with advantage when she is not nursing. In this way these troublesome complications may generally be prevented. The most common forms are either an abrasion on the surface of the nipple, which, if neglected, may form a small ulcer, or a crack at some part of the nipple, most generally at its base. In either case the suffering when the child is put to the breast is intense, sometimes indeed amounting to intolerable anguish, causing the mother to look forward with dread to the application of the child. Whenever such pain is complained of, the nipple should be carefully examined, since the fissure or sore is often so minute as to escape superficial examination. The remedies recommended are very numerous and not always successful. Amongst those most commonly used are astringent applications, such as tannin or weak solutions of nitrate of silver, or cauterizing the edges of the fissure with solid nitrate of silver, or applying the flexible collodion of the Pharmacopœia. Dr. Wilson of Glasgow speaks highly of a lotion composed of ten grains of nitrate of lead in an ounce of glycerin, which is to be applied after suckling, the nipple being carefully washed before the child is again put to the breast. I have myself found nothing answer so well as a lotion composed of half an ounce of sulphurous acid, half an ounce of the glycerin of tannin, and an ounce of water, the beneficial effects of which are sometimes quite remarkable. Relief may occasionally be obtained by inducing the child to suck through a nipple-shield, especially when there is only an excoriation; but this will not always answer, on account of the extreme pain which it produces.

Excessive Flow of Milk.—An excessive flow of milk, known as *galactorrhœa*, often interferes with successful lactation. It is by no means rare in the first weeks after delivery for women of delicate constitution, who are really unfit to nurse, to be flooded with a superabundance of watery and innutritious milk, which soon produces disordered digestion in the child. Under such circumstances the only thing to be done is to give up an attempt which is injurious both to the mother and child. At a later stage the milk, secreted in large quantities, is sufficiently nourishing to the child, but the drain on the mother's constitution soon begins to tell on her. Palpitation, giddiness, emaciation, headache, loss of sleep, spots before the eyes, indicate the serious effects which are being produced and the absolute necessity of at once stopping lactation. Whenever, therefore, a nursing woman suffers from such symptoms, it is far better at once to remove the cause, otherwise a very serious and

permanent deterioration of health might result. When, under such circumstances, nursing is unwisely persevered in, most serious results may follow. Should any diathetic tendency exist, especially when there is a predisposition to phthisis, nothing is so likely to develop it as the debility produced by excessive lactation. Certain diseases of the eye are then specially apt to occur, such as severe inflammation of the cornea, leading to opacity and even sloughing, and certain forms of choroiditis; also impairment of accommodation due to defective power of the ciliary muscle.¹

Mammary Abscess.—There is no more troublesome complication of lactation than the formation of abscess in the breast—an occurrence by no means rare, and which, if improperly treated, may, by long-continued suppuration and the formation of numerous sinuses in and about the breast, produce very serious effects on the general health. The causes of breast abscesses are numerous, and very trivial circumstances may occasionally set up inflammation, ending in suppuration. Thus it may follow exposure to cold, a blow or other injury to the breast, some temporary engorgement of the lacteal tubes, or even sudden or depressing mental emotions. The most frequent cause is irritation from fissures or erosions of the nipples, which must therefore always be regarded with suspicion, and cured as soon as possible.

Signs and Symptoms.—The abscess may form in any part of the breast or in the areolar tissue below it; in the latter case the inflammation very generally extends to the gland-structure. Abscess is usually ushered in by constitutional symptoms, varying in severity with the amount of the inflammation. Pyrexia is always present; elevated temperature, rapid pulse, and much malaise and sense of feverishness, followed in many cases by distinct rigor when deep-seated suppuration is taking place. On examining the breast it will be found to be generally enlarged and very tender, while at the site of the abscess an indurated and painful swelling may be felt. If the inflammation be chiefly limited to the sub-glandular areolar tissue, there may be no localized swelling felt, but the whole breast will be acutely sensitive and the slightest movement will cause much pain. As the case progresses the abscess becomes more and more superficial, the skin covering it is red and glazed, and if left to itself it bursts. In the more serious cases it is by no means rare for multiple abscesses to form. These, opening one after the other, lead to the formation of numerous fistulous tracts, by which the breast may become completely riddled. Sloughing of portions of the gland-tissue may take place, and even considerable hemorrhage from the destruction of blood-vessels. The general health soon suffers to a marked degree, and, as the sinuses continue to suppurate for many successive months, it is by no means uncommon for the patient to be reduced to a state of profound and even dangerous debility.

Treatment.—Much may be done by proper care to prevent the formation of abscess, especially by removing engorgement of the lacteal ducts, when threatened, by gentle hand-friction in the manner already indi-

¹ See Foerster of Breslau in Graefe and Saemisch's *Handbuch des Gesammten Augenheilkunde*, and Power on "The Diseases of the Eye in Connection with Pregnancy," *Lancet*, May 8, 1880, *et seq.*

cated. When the general symptoms and the local tenderness indicate that inflammation has commenced, we should at once endeavor to moderate it, in the hope that resolution may occur without the formation of pus. Here general principles must be attended to, especially giving the affected part as much rest as possible. Feverishness may be combated by gentle salines, minute doses of aconite, and large doses of quinine, while pain should be relieved by opiates. The patient should be strictly confined in bed, and the affected breast supported by a suspensory bandage. Warmth and moisture are the best means of relieving the local pain, either in the form of hot fomentations or of light poultices of linseed meal or bread and milk, and the breast may be smeared with extract of belladonna rubbed down with glycerin, or the belladonna liniment sprinkled over the surface of the poultices. The local application of ice in india-rubber bags has been highly extolled as a means of relieving the pain and tension, and is said to be much more effectual than heat and moisture.¹ Generally, the pain and irritation produced by putting the child to the breast are so great as to contraindicate nursing from the affected side altogether, and we must trust to relieving the tension by poultices, suckling being in the mean time carried on by the other breast alone. In favorable cases this is quite possible for a time, and it may be that, if the inflammation do not end in suppuration or if the abscess be small and localized, the affected breast is again able to resume its functions. Often this is not possible, and it may be advisable in severe cases to give up nursing altogether.

Pus should be Removed as Soon as Possible.—The subsequent management of the case consists in the opening of the abscess as soon as the existence of pus is ascertained, either by fluctuation or, if the site of the abscess be deep seated, by the exploring-needle. It may be laid down as a principle that the sooner the pus is evacuated the better, and nothing is to be gained by waiting until it is superficial. On the contrary, such delay only leads to more extensive disorganization of tissue and the further spread of inflammation.

Antiseptic Treatment of Mammary Abscess.—The method of opening the abscess is of primary importance. It has always been customary simply to open the abscess at its most dependent part, without using any precaution against the admission of air, and afterward to treat secondary abscesses in the same way. The results are well known to all practical accoucheurs, and the records of surgery fully show how many weeks or months generally elapse in bad cases before recovery is complete. The antiseptic treatment of mammary abscess, in the way first pointed out by Lister, affords results which are of the most remarkable and satisfactory kind. Instead of being weeks and months in healing, I believe that the practitioner who fairly and minutely carries out Sir Joseph Lister's directions may confidently look for complete closure of the abscess in a few days; and I know of nothing in the whole range of my professional experience that has given me more satisfaction than the application of this method to abscesses of the breast. The plan I first used is that recommended by Lister in the *Lancet* for 1867, but which is now superseded by his improved methods, which of course will

¹ Carson, *Amer. Journ. of Obstet.*, Jan., 1881.

be used in preference by all who have made themselves familiar with the details of antiseptic surgery. The former, however, is easily within the reach of every one, and is so simple that no special skill or practice is required in its application, whereas the more perfected antiseptic appliances will probably not be so readily obtained and are much more difficult to use. I therefore insert Sir Joseph Lister's original directions, which he assures me are perfectly antiseptic, for the guidance of those who may not be able to obtain the more elaborate dressings: "A solution of one part of crystallized carbolic acid in four parts of boiled linseed oil having been prepared, a piece of rag from four to six inches square is dipped into the oily mixture and laid upon the skin where the incision is to be made. The lower edge of the rag being then raised while the upper edge is kept from slipping by an assistant, a common scalpel or bistoury dipped in the oil is plunged into the cavity of the abscess, and an opening about three-quarters of an inch in length is made; and the instant the knife is withdrawn the rag is dropped upon the skin as an antiseptic curtain, beneath which the pus flows out into a vessel placed to receive it. The cavity of the abscess is firmly pressed, so as to force out all existing pus as nearly as may be (the old fear of doing mischief by rough treatment of the pyogenic membrane being quite ill-founded); and if there be much oozing of blood, or if there be considerable thickness of parts between the abscess and the surface, a piece of lint dipped in the antiseptic oil is introduced into the incision to check bleeding and prevent primary adhesion, which is otherwise very apt to occur. The introduction of the lint is effected as rapidly as may be, and under the protection of the antiseptic rag. Thus the evacuation of the original contents is accomplished with perfect security against the introduction of living germs. This, however, would be of no avail unless an antiseptic dressing could be applied that would effectually prevent the decomposition of the stream of pus constantly flowing out beneath it. After numerous disappointments I have succeeded with the following, which may be relied upon as absolutely trustworthy: About six teaspoonfuls of the above-mentioned solution of carbolic acid in linseed oil are mixed up with common whiting (carbonate of lime) to the consistence of a firm paste, which is, in fact, glazier's putty with the addition of a little carbolic acid. This is spread upon a piece of common tin-foil about six inches square, so as to form a layer about a quarter of an inch thick. The tin-foil, thus spread with putty, is placed upon the skin so that the middle of it corresponds to the position of the incision, the antiseptic rag used in opening the abscess being removed the instant before. The tin is then fixed securely by adhesive plaster, the lowest edge being left free for the escape of the discharge into a folded towel placed over it and secured by a bandage. The dressing is changed, as a general rule, once in twenty-four hours, but if the abscess be a very large one it is prudent to see the patient twelve hours after it has been opened, when, if the towel should be much stained with discharge, the dressing should be changed, to avoid subjecting its antiseptic virtues to too severe a test. But after the first twenty-four hours a single daily dressing is sufficient. The changing of the dressing must be methodically done, as follows: A second similar piece of tin-foil having been

spread with the putty, a piece of rag is dipped in the oily solution and placed on the incision the moment the first tin is removed. This guards against the possibility of mischief occurring during the cleansing of the skin with a dry cloth and pressing out any discharge which may exist in the cavity. If a plug of lint was introduced when the abscess was opened, it is removed under cover of the antiseptic rag, which is taken off at the moment when the new tin is to be applied. The same process is continued daily until the sinus closes."

Treatment of Long-continued Suppuration and Fever.—If the case come under our care when the abscess has been long discharging or when sinuses have formed, the treatment is directed mainly to procuring a cessation of suppuration and closure of the sinuses. For this purpose methodical strapping of the breast with adhesive plaster, so as to afford steady support and compress the composing pyogenic surfaces, will give the best results. It may be necessary to lay open some of the sinuses, or to inject tinct. iodi or other stimulating lotions so as to moderate the discharge, the subsequent surgical treatment varying according to the requirements of each case. In such neglected cases Billroth recommends that after the patient has been anaesthetized the openings should be dilated so as to admit the finger, by which the septa between the various sinuses should be broken down and a large single abscess-cavity made. This should then be thoroughly irrigated with a 3 per cent. solution of carbolic acid, a drainage-tube introduced, and the ordinary antiseptic dressings applied. As the drain on the system is great and the constitutional debility generally pronounced, much attention must be paid to general treatment, and abundance of nourishing food, appropriate stimulants, and such medicines as iron and quinine will be indicated.

Hand-Feeding.—In a considerable number of cases the inability of the mother to nurse the child, her invincible repugnance to a wet-nurse, or inability to bear the expense renders hand-feeding essential. It is therefore of importance that the accoucheur should be thoroughly familiar with the best method of bringing up the child by hand, so as to be able to direct the process in the way that is most likely to be successful.

Causes of Mortality in Hand-fed Children.—Much of the mortality following hand-feeding may be traced to unsuitable food. Among the poorer classes especially there is a prevalent notion that milk alone is insufficient, and hence the almost universal custom of administering various farinaceous foods, such as corn flour or arrowroot, even from the earliest period. Many of these consist of starch alone, and are therefore absolutely unsuited for forming the staple of diet on account of the total absence of nitrogenous elements. Independently of this, it has been shown that the saliva of infants has not the same digestive property on starch that it subsequently acquires, and this affords a further explanation of its so constantly producing intestinal derangement. Reason, as well as experience, abundantly proves that the object to be aimed at in hand-feeding is to imitate as nearly as possible the food which nature supplies for the new-born child, and therefore the obvious course is to use milk from some animal, so treated as to make it resemble human milk as nearly as may be.

Ass's Milk.—Of the various milks used, that of the ass, on the whole,

most closely resembles human milk, containing less casein and butter and more saline ingredients. It is not always easy to obtain, and in towns is excessively expensive. Moreover, it does not always agree with the child, being apt to produce diarrhœa. We can, however, be more certain of its being unadulterated, which in large cities is in itself no small advantage, and it may be given without the addition of water or sugar.

Goat's Milk.—Goat's milk in this country is still more difficult to obtain, but it often succeeds admirably. In many places the infant sucks the teat directly, and certainly thrives well on the plan.

[We reverse the order in this country, where the ass is seldom seen and the goat is quite common, particularly in the suburbs of our large cities, where its milk is most required. I have seen marvellous results from feeding sick infants with its milk freshly drawn and diluted with hot water. I do not believe it is as suitable as that of the cow, but it has the advantage that it can be obtained freshly drawn in a city by keeping the animal in the yard or on a vacant lot. The goat should be fed upon grass and other suitable diet, and not permitted to run at large, as it eats with impunity stramonium and other noxious weeds.—ED.]

Cow's Milk, and its Preparation.—In a large majority of cases we have to rely on cow's milk alone. It differs from human milk in containing less water, a larger amount of casein and solid matters, and less sugar. Therefore, before being given it requires to be diluted and sweetened. A common mistake is over-dilution, and it is far from rare for nurses to administer one-third cow's milk to two-thirds water. The result of this excessive dilution is that the child becomes pale and puny and has none of the firm and plump appearance of a well-fed infant. The practitioner should therefore ascertain that this mistake is not being made; and the necessary dilution will be best obtained by adding to pure fresh cow's milk one-third hot water, so as to warm the mixture to about 96°, the whole being slightly sweetened with sugar of milk or ordinary crystallized sugar. After the first two or three months the amount of water may be lessened, and pure milk, warmed and sweetened, given instead. Whenever it is possible the milk should be obtained from the same cow, and in towns some care is requisite to see that the animal is properly fed and stabled. Of late years it has been customary to obviate the difficulties of obtaining good fresh milk by using some of the tinned milks now so easily to be had. These are already sweetened, and sometimes answer well if not given in too weak a dilution. One great drawback in bottle-feeding is the tendency of the milk to become acid, and hence to produce diarrhœa. This may be obviated to a great extent by adding a tablespoonful of lime-water to each bottle, instead of an equal quantity of water.

[The milk of the Alderney cow contains too much butter to make a good substitute for human milk in feeding young infants, and will often disagree with them when that of the common cow will be digested. The milk of one cow, which should neither be young or old, is to be preferred, but it must be borne in mind that the *special* can is often filled from the general supply.—ED.]

Artificial Human Milk.—An admirable plan of treating cow's milk,

so as to reduce it to almost absolute chemical identity with human milk, has been devised by Professor Frankland, to whom I am indebted for permission to insert the recipe. I have followed this method in many cases, and find it far superior to the usual one, as it produces an exact and uniform compound. With a little practice nurses can employ it with no more trouble than the ordinary mixing of cow's milk with water and sugar. The following extract from Dr. Frankland's work¹ will explain the principles on which the preparation of the artificial human milk is founded: "The rearing of infants who cannot be supplied with their natural food is notoriously difficult and uncertain, owing chiefly to the great difference in the chemical composition of human milk and cow's milk. The latter is much richer in casein and poorer in milk-sugar than the former, whilst ass's milk, which is sometimes used for feeding infants, is too poor in casein and butter, although the proportion of sugar is nearly the same as in human milk. The relations of the three kinds of milk to each other are clearly seen from the following analytical numbers, which express the percentage amounts of the different constituents:

	Woman.	Ass.	Cow.
Casein	27	1.7	4.2
Butter	3.5	1.3	3.8
Milk-sugar	5.0	4.5	3.8
Salts2	.5	.7

These numbers show that by the removal of one-third of the casein from cow's milk, and the addition of about one-third more milk-sugar, a liquid is obtained which closely approaches human milk in composition, the percentage amounts of the four chief constituents being as follows:

Casein	2.8
Butter	3.8
Milk-sugar	5.0
Salts7

The following is the mode of preparing the milk: Allow one-third of a pint of new milk to stand for about twelve hours; remove the cream, and add to it two-thirds of a pint of new milk, as fresh from the cow as possible. Into the one-third of a pint of blue milk left after the abstraction of the cream put a piece of rennet about one inch square. Set the vessel in warm water until the milk is fully curdled—an operation requiring from five to fifteen minutes, according to the activity of the rennet, which should be removed as soon as the curdling commences and put into an egg-cup for use on subsequent occasions, as it may be employed daily for a month or two. Break up the curd repeatedly, and carefully separate the whole of the whey, which should then be rapidly heated to boiling in a small tin pan placed over a spirit or gas lamp. During the heating a further quantity of casein, technically called 'flectings,' separates, and must be removed by straining through muslin. Now dissolve 110 grains of powdered sugar of milk in the hot whey, and mix it with the two-thirds of a pint of new milk to which the cream from the other third of a pint was added as already described. The

¹ Frankland's *Experimental Researches in Chemistry*, p. 843.

artificial milk should be used within twelve hours of its preparation, and it is almost needless to add that all the vessels employed in its manufacture and administration should be kept scrupulously clean."¹

Method of Hand-feeding.—Much of the success of bottle-feeding must depend on minute care and scrupulous cleanliness—points which cannot be too strongly insisted on. Particular attention should be paid to preparing the food fresh for every meal, and to keeping the feeding-bottle and tubes constantly in water when not in use, so that minute particles of milk may not remain about them and become sour. A neglect of this is one of the most fertile sources of the thrush from which bottle-fed infants often suffer. The particular form of bottle used is not of much consequence. Those now commonly employed, with a long india-rubber tube attached, are preferable to the older forms of flat bottle, as they necessitate strong suction on the part of the infant, thus forcing it to swallow the food more slowly. Care must be taken to give the meals at stated periods, as in breast-feeding, and these should be at first about two hours apart, the intervals being gradually extended. The nurse should be strictly cautioned against the common practice of placing the bottle beside the infant in its cradle and allowing it to suck to repletion—a practice which leads to over-distension of the stomach and consequent dyspepsia. The child should be raised in the arms at the proper time, have its food administered, and then be replaced in the cradle to sleep. In the first few weeks of bottle-feeding constipation is very common, and may be effectually remedied by placing as much phosphate of soda as will lie on a threepenny-piece in the bottle two or three times in the twenty-four hours.

Other Kinds of Food.—If this system succeed, no other food should be given until the child is six or seven months old, and then some of the various infant's foods may be cautiously commenced. Of these there are an immense number in common use, some of which are good articles of diet, others are unfitted for infants. In selecting them we have to see that they contain the essential elements of nutrition in proper combination. All those, therefore, that are purely starchy in character, such as arrowroot, corn flour, and the like, should be avoided, while those that contain nitrogenous as well as starch elements may be safely given. Of the latter the entire wheat flour, which contains the husks ground down with the wheat, generally answers admirably; and of the same character are rusks, tops and bottoms, Nestle's or Liebig's infant's

¹ The following recipe yields the same results, but the method is easier, and I find that nurses prepare the milk with less difficulty when it is followed: "Take half a pint of skimmed milk, heat it to about 96°, and put into the warmed milk a piece of rennet about an inch square. Set the milk to stand in the fender or over a lamp until it is quite warm. When it is set, take the rennet out and break up the curd quite small with a knife, and let it stand ten or fifteen minutes, when the curd will sink. Then pour the whey into a saucepan, and let it boil quickly. Measure one-third of a pint of this whey, and dissolve in it, when hot, a powder containing 110 grains of sugar of milk. When this third of a pint of whey is quite cold add to it two-thirds of a pint of new milk and two teaspoonfuls of cream, stirring the whole together. The food should be made fresh every twelve hours, and warmed as required. The piece of rennet when taken out can be kept in an egg-cup and used for ten days or a fortnight."—N. B. It is often advisable during the first month to use rather more than a third of a pint of whey, as the milk is apt to be rather too rich for a newly-born child.

food, and many others. If the child be pale and flabby, some more purely animal food may often be given twice a day, and great benefit may be derived from a single meal of beef, chicken, or veal tea, with a little bread-crumbs in it, especially after the sixth or seventh month. Milk, however, should still form the main article of diet, and should continue to do so for many months.


Management when Milk Disagrees.—If the child be pale, flabby, and do not gain flesh, more especially if diarrhœa or other intestinal disturbance be present, we may be certain that hand-feeding is not answering satisfactorily and that some change is required. If the child be not too old and will still take the breast, that is certainly the best remedy, but if that be not possible it is necessary to alter the diet. When milk disagrees, cream, in the proportion of one tablespoonful to three of water, sometimes answers as well. Occasionally also Liebig's or Mellin's infant's food, when carefully prepared, renders good service. Too often, however, when once diarrhœa or other intestinal disturbance has set in, all our efforts may prove unavailing, and the health, if not the life, of the infant becomes seriously imperilled. It is not, however, within the scope of this work to treat of the disorders of infants at the breast, the proper consideration of which requires a large amount of space, and I therefore refrain from making any further remarks on the subject.

CHAPTER III.

PUERPERAL ECLAMPSIA.

By the term *puerperal eclampsia* is meant a peculiar kind of epileptiform convulsions which may occur in the latter months of pregnancy or during or after parturition, and it constitutes one of the most formidable diseases with which the obstetrician has to cope. The attack is often so sudden and unexpected, so terrible in its nature, and attended with such serious danger both to the mother and child, that the disease has attracted much attention.

Its Doubtful Etiology.—The researches of Lever, Braun, Frerichs, and many other writers who have shown the frequent association of eclampsia with albuminuria have, of late years, been supposed to clear up to a great extent the etiology of the disease, and to prove its dependence on the retention of urinary elements in the blood. While the urinary origin of eclampsia has been pretty generally accepted, more recent observations have tended to throw doubt on its essential dependence on this cause, so that it can hardly be said that we are yet in a position to explain its true pathology with certainty. These points will require separate discussion, but it is first necessary to describe the character and history of the attack.



Considerable confusion exists in the description of puerperal convulsions from the confounding of several essentially distinct diseases under the same name. Thus, in most obstetric works it has been customary to describe three distinct classes of convulsion—the *epileptic*, the *hysterical*, and the *apoplectic*. The two latter, however, come under a totally different category. A pregnant woman may suffer from hysterical paroxysms, or she may be attacked with apoplexy, accompanied with coma and followed by paralysis. But these conditions in the pregnant or parturient woman are identical with the same diseases in the non-pregnant, and are in no way special in their nature. True eclampsia, however, is different in its clinical history from epilepsy, although the paroxysms, while they last, are essentially the same as those of an ordinary epileptic fit.

Premonitory Symptoms.—An attack of eclampsia seldom occurs without having been preceded by certain more or less well-marked precursory symptoms. It is true that in a considerable number of cases these are so slight as not to attract attention, and suspicion is not aroused until the patient is seized with convulsions. Still, subsequent investigations will very generally show that some symptoms did exist, which, if observed and properly interpreted, might have put the practitioner on his guard and possibly have enabled him to ward off the attack. Hence a knowledge of them is of real practical value. The most common are associated with the cerebrum, such as severe headache, which is the one most generally observed and is sometimes limited to one side of the head. Transient attacks of dizziness, spots before the eyes, loss of sight, or impairment of the intellectual faculties are also not uncommon. These signs in a pregnant woman are of the gravest import, and should at once call for investigation into the nature of the case. Less marked indications sometimes exist in the form of irritability, slight headache or stupor, and a general feeling of indisposition. Another important premonitory sign is œdema of the subcutaneous cellular tissue, especially of the face or upper extremities, which should at once lead to an examination of the urine.

Symptoms of the Attack.—Whether such indications have preceded an attack or not, as soon as the convulsion comes on there can no longer be any doubt as to the nature of the case. The attack is generally sudden in its onset, and in its character is precisely that of a severe epileptic fit or of the convulsions in children. Close observation shows that there is at first a short period of tonic spasm, affecting the entire muscular system. This is almost immediately succeeded by violent clonic contractions, generally commencing in the muscles of the face, which twitch violently; the expression is horribly altered; the globes of the eyes are turned up under the eyelids, so as to leave only the white sclerotics visible; and the angles of the mouth are retracted and fixed in a convulsive grin. The tongue is at the same time protruded forcibly, and, if care be not taken, is apt to be lacerated by the violent grinding of the teeth. The face, at first pale, soon becomes livid and cyanosed, while the veins of the neck are distended and the carotids beat vigorously. Frothy saliva collects about the mouth, and the whole appearance is so changed as to render the patient quite unrecognizable. The convulsive move-

ments soon attack the muscles of the body. The hands and arms, at first rigidly fixed, with the thumbs clenched into the palms, begin to jerk, and the whole muscular system is thrown into rapidly-recurring convulsive spasms. It is evident that the involuntary muscles are implicated in the convulsive action, as well as the voluntary. This is shown by a temporary arrest of respiration at the commencement of the attack, followed by irregular and hurried respiratory movements, producing a peculiar hissing sound. The occasional involuntary expulsion of urine and feces indicates the same fact. During the attack the patient is absolutely unconscious, sensibility is totally suspended, and she has afterward no recollection of what has taken place. Fortunately, the convulsion is not of long duration, and at the outside does not last more than three or four minutes, generally not so long. In most cases, after an interval there is a recurrence of the convulsion, characterized by the same phenomena, and the paroxysms are repeated with more or less force and frequency according to the severity of the attack. Sometimes several hours may elapse before a second convulsion comes on; at others the attacks may recur very often, with only a few minutes between them. In the slighter forms of eclampsia there may not be more than two or three paroxysms in all; in the more serious as many as fifty or sixty have been recorded.

Condition between the Attacks.—After the first attack the patient generally soon recovers her consciousness, being somewhat dazed and somnolent, with no clear conception of what has occurred. If the paroxysms be frequently repeated, more or less profound coma continues in the intervals between them; which no doubt depends upon intense cerebral congestion, resulting from the interference with the circulation in the great veins of the neck, produced by spasmodic contraction of the muscles. The coma is rarely complete, the patient showing signs of sensibility when irritated, and groaning during the uterine contractions. In the worst class of cases the torpor may become intense and continuous, and in this state the patient may die. When the convulsions have entirely stopped and the patient has completely regained her consciousness, and is apparently convalescent, recollection of what has taken place during and some time before the attack may be entirely lost; and this condition may last for a considerable time. A curious instance of this once came under my notice in a lady who had lost her brother, to whom she was greatly attached, in the week immediately preceding her confinement, and in whom the mental distress seemed to have had a good deal to do in determining the attack. It was many weeks before she recovered her memory, and during that time she recollected nothing about the circumstances connected with her brother's death, the whole of that week being as it were blotted out of her recollection.

Relation of the Attacks to Labor.—If the convulsions come on during pregnancy, we may look upon the advent of labor as almost a certainty; and if we consider the severe nervous shock and general disturbance, this is the result we might reasonably anticipate. If they occur, as is not uncommon, for the first time during labor, the pains generally continue with increased force and frequency, since the uterus partakes of the convulsive action. It has not rarely happened that the pains have

gone on with such intensity that the child has been born quite unexpectedly, the attention of the practitioner being taken up with the patient. In many cases the advent of fresh paroxysms is associated with the commencement of a pain, the irritation of which seems sufficient to bring on the convulsion.

Results to the Mother and Child.—The results of eclampsia vary according to the severity of the paroxysms. It is generally said that about 1 in 3 or 4 cases dies. The mortality has certainly lessened of late years, probably in consequence of improved knowledge of the nature of the disease and more rational modes of treatment. This is well shown by Barker,¹ who found in 1855 a mortality of 32 per cent. in cases occurring before and during labor, and 22 per cent. in those after labor, while since that date the mortality has fallen to 14 per cent. The same conclusion is arrived at by Dr. Phillips,² who has shown that the mortality has greatly lessened since the practice of repeated and indiscriminate bleeding, long considered the sheet-anchor in the disease, has been discontinued and the administration of chloroform substituted.

Cause of Death.—Death may occur during the paroxysm, and then it may be due to the long continuance of the tonic spasm producing asphyxia. It is certain that as long as the tonic spasm lasts the respiration is suspended, just as in the convulsive disease of children known as laryngismus stridulus; and it is possible also that the heart may share in the convulsive contraction which is known to affect other involuntary muscles. More frequently death happens at a later period, from the combined effects of exhaustion and asphyxia. The records of post-mortem examinations are not numerous; in those we possess the principal changes have been an anæmic condition of the brain, with some oedematous infiltration. In a few rare cases the convulsions have resulted in effusion of blood into the ventricles or at the base of the brain. The prognosis as regards the child is also serious. Out of 36 children, Hall Davis found 26 born alive, 10 being stillborn. There is good reason to believe that the convulsion may attack the child *in utero*—of this several examples are mentioned by Cazeaux—or it may be subsequently attacked with convulsions, even when apparently healthy at birth.

Pathology of the Disease.—The precise pathology of eclampsia cannot be considered by any means satisfactorily settled. When, in the year 1843, Lever first showed that the urine in patients suffering from puerperal convulsions was generally highly charged with albumen—a fact which subsequent experience has amply confirmed—it was thought that a key to the etiology of the disease had been found. It was known that chronic forms of Bright's disease were frequently associated with retention of urinary elements in the blood and not rarely accompanied by convulsions. The natural inference was drawn that the convulsions of eclampsia were also due to toxæmia resulting from the retention of urea in the blood, just as in the uræmia of chronic Bright's disease; and this view was adopted and supported by the authority of Braun, Frerichs, and many other writers of eminence, and was generally received as a satisfactory explanation of the facts. Frerichs modified it so far that he held that the true toxic element was not urea as such, but carbonate

¹ *The Puerperal Diseases*, p. 125.

² *Guy's Hospital Reports*, 1870.

of ammonia, resulting from its decomposition, and experiments were made to prove that the injection of this substance into the veins of the lower animals produced convulsions of precisely the same character as eclampsia. Dr. Hammond¹ of Maryland subsequently made a series of counter-experiments, which were held as proving that there was no reason to believe that urea ever did become decomposed in the blood in the way that Frerichs supposed, or that the symptoms of uræmia were ever produced in this way. Others have believed that the poisonous elements retained in the blood are not urea or the products of its decomposition, but other extractive matters which have escaped detection. As time elapsed, evidence accumulated to show that the relation between albuminuria and eclampsia was not so universal as was supposed, or at least that some other factors were necessary to explain many of the cases. Numerous cases were observed in which albumen was detected in large quantities, without any convulsion following, and that not only in women who had been the subject of Bright's disease before conception, but also when the albuminuria was known to have developed during pregnancy. Thus, Imbert Goubeyre found that, out of 164 cases of the latter kind, 95 had no eclampsia, and Blot, out of 41 cases, found that 34 were delivered without untoward symptoms. It may be taken as proved, therefore, that albuminuria is by no means necessarily accompanied by eclampsia. Cases were also observed in which the albumen only appeared after the convulsion, and in these it was evident that the retention of urinary elements could not have been the cause of the attack; and it is highly probable that in them the albuminuria was produced by the same cause which induced the convulsion. Special attention has been called to this class of cases by Braxton Hicks,² who has recorded a considerable number of them. He says that the nearly simultaneous appearance of albuminuria and convulsion—and it is admitted that the two are almost invariably combined—must then be explained in one of three ways:

1st. That the convulsions are the cause of the nephritis.

2dly. That the convulsions and the nephritis are produced by the same cause—*e. g.* some detrimental ingredient circulating in the blood, irritating both the cerebro-spinal system and other organs at the same time.

3dly. That the highly-congested state of the venous system, induced by the spasm of the glottis in eclampsia, is able to produce the kidney complication.

Theory of Traube and Rosenstein.—More recently, Traube and Rosenstein have advanced a theory of eclampsia purporting to explain these anomalies. They refer the occurrence of eclampsia to acute cerebral anæmia resulting from changes in the blood incident to pregnancy. The primary factor is the hydræmic condition of the blood which is an ordinary concomitant of the pregnant state, and of course when there is also albuminuria the watery condition of the blood is greatly intensified; hence the frequent association of the two states. Accompanying this condition of the blood there is increased tension of the arterial system, which is favored by the hypertrophy of the heart which is known to be

¹ *Amer. Journ.*, 1861.

² *Obstet. Trans.*, vol. viii.

a normal occurrence in pregnancy. The result of these combined states is a temporary hyperæmia of the brain, which is rapidly succeeded by serous effusion into the cerebral tissues, resulting in pressure on its minute vessels and consequent anæmia. There is much in this theory that accords with the most recent views as to the etiology of convulsive disease; as, for example, the researches of Kussmaul and Tenner, who had experimentally proved the dependence of convulsion on cerebral anæmia, and of Brown-Séquard, who showed that an anæmic condition of the nerve-centres preceded an epileptic attack. It explains also very satisfactorily how the occurrence of labor should intensify the convulsions, since during the acme of the pains the tension of the cerebral arterial system is necessarily greatly increased. There are, however, obvious difficulties against its general acceptance. For example, it does not satisfactorily account for those cases which are preceded by well-marked precursory symptoms, and in which an abundance of albumen is present in the urine. Here the premonitory signs are precisely those which precede the development of uræmia in chronic Bright's disease, the dependence of which on the retention in the blood of urinary elements can hardly be doubted. Moreover, it has been shown by Löhlein and others that on post-mortem examination the brain does not, as a rule, exhibit the œdema, anæmia, and flattened convolutions which this theory assumes.

Views of MacDonald.—MacDonald¹ has published an interesting paper on this subject, in which he describes two very careful post-mortem examinations. In these he found extreme anæmia of the cerebro-spinal centres, with congestion of the meninges, but no evidence of œdema. He inclines to the belief that eclampsia is caused by irritation of the vaso-motor centre in consequence of an anæmic condition of the blood produced by the retention in it of excrementitious matters which the kidneys ought to have removed, this over-stimulation resulting in anæmia of the deeper-seated nerve-centres and consequent convulsion.

Excitability of Nervous System.—This key to the liability of the puerperal woman to convulsive attacks is no doubt to be found in the peculiar excitable condition of the nervous system in pregnancy—a fact which was clearly pointed out by the late Dr. Tyler Smith and by many other writers. Her nervous system is in this respect not unlike that of children, in whom the predominant influence and great excitability of the nervous system are well-established facts, and in whom precisely similar convulsive seizures are of common occurrence on the application of a sufficiently exciting cause.

Exciting Causes.—Admitting this, we require some cause to set the predisposed nervous system into morbid action; and this we may have either in a toxæmic or in an extremely watery condition of the blood, associated with albuminuria; or along with these, or sometimes independently of them, in some excitement, such as strong emotional disturbance. It is highly probable, however, that extreme anæmia is one of the actual conditions of the nerve-centres—a fact of much practical importance in reference to treatment.

¹ See his volume of collected essays entitled *Heart Disease during Pregnancy*, London, 1878.

Treatment.—The management of cases in which the occurrence of suspicious symptoms has led to the detection of albuminuria, has already been fully discussed (p. 209). We shall therefore here only consider the treatment of cases in which convulsions have actually occurred.

Venesection.—Until quite recently venesection was regarded as the sheet-anchor in the treatment, and blood was always removed copiously, and, there is sufficient reason to believe, with occasional remarkable benefit. Many cases are recorded in which a patient in apparently profound coma rapidly regained her consciousness when blood was extracted in sufficient quantity. The improvement, however, was often transient, the convulsions subsequently recurring with increased vigor. There are good theoretical grounds for believing that bloodletting can only be of merely temporary use, and may even increase the tendency to convulsion. These are so well put by Schroeder that I cannot do better than quote his observations on this point. "If," he says, "the theory of Traube and Rosenstein be correct, a sudden depletion of the vascular system, by which the pressure is diminished, must stop the attacks. From experience it is known that after venesection the quantity of blood soon becomes the same through the serum taken from all the tissues, while the quality is greatly deteriorated by the abstraction of blood. A short time after venesection we shall expect to find the former blood-pressure in the arterial system, but the blood far more watery than previously. From this theoretical consideration it follows that abstraction of blood, if the above-mentioned conditions really cause convulsions, must be attended by an immediate favorable result, and under certain circumstances the whole disease may surely be cut short by it. But if all other conditions remain the same the blood-pressure will after some time again reach its former height. The quality of blood has in the mean time been greatly deteriorated, and consequently the danger of the disease will be increased."

These views sufficiently well explain the varying opinions held with regard to this remedy, and enable us to understand why, while the effects of venesection have been so lauded by certain authors, the mortality has admittedly been much lessened since its indiscriminate use has been abandoned. It does not follow because a remedy, when carried to excess, is apt to be hurtful, that it should be discarded altogether; and I have no doubt that in properly-selected cases and judiciously employed, venesection is a valuable aid in the treatment of eclampsia, and that it is specially likely to be useful in mitigating the first violence of the attack and in giving time for other remedies to come into action. Care should, however, be taken to select the cases properly, and it will be specially indicated when there is marked evidence of great cerebral congestion and vascular tension, such as a livid face, a full bounding pulse, and strong pulsation in the carotids. The general constitution of the patient may also serve as a guide in determining its use, and we shall be the more disposed to resort to it if the patient be a strong and healthy woman; while, on the other hand, if she be feeble and weak, we may wisely discard it and trust entirely to other means. In any case, it must be looked upon as a temporary expedient only, useful in warding off immediate danger to the cerebral tissues, but never as the main agent in treatment.

Nor can it be permissible to bleed in the heroic manner frequently recommended. A single bleeding, the amount regulated by the effect produced, is all that is ever likely to be of service.

[After the discovery of the uræmic origin of eclampsia in pregnant women the treatment by bleeding was very generally abandoned in the United States; but the more recent investigations of the causes of death have produced a reconsideration of this plan of treatment, and the tendency of the profession during the last ten or fifteen years has been toward venesection as a preventive of cerebral complications. In primiparæ with a full pulse and flushed face the rule with many of our obstetrical practitioners is to bleed the patient as early as practicable, and to do this at least once effectually, so as to produce, if possible, a noticeable impression. Where there are positive evidences of the existence of Bright's disease, of course this is inadmissible.—ED.]

Compression of the Carotids.—As a temporary expedient, having the same object in view, compression of the carotids during the paroxysms is worthy of trial. This was proposed by Trousseau in the eclampsia of infants, and in the single case of eclampsia in which I have tried it seemed to be decidedly beneficial. It is a simple measure, and it offers the advantage of not leading to any permanent deterioration of the blood, as in venesection.

Administration of Purgatives.—As a subsidiary means of diminishing vascular tension the administration of a strong purgative is desirable, and has the further effect of removing any irritant matter that may be lodged in the intestinal tract. If the patient be conscious, a full dose of the compound jalap powder may be given, or a few grains of calomel combined with jalap; and if she be comatose and unable to swallow, a drop of croton oil or a quarter of a grain of elaterium may be placed on the back of the tongue.

Administration of Sedatives and Narcotics.—The great indication in the management of eclampsia is the controlling of convulsive action by means of sedatives. Foremost amongst them must be placed the inhalation of chloroform—a remedy which is frequently remarkably useful, and which has the advantage of being applicable at all stages of the disease and whether the patient be comatose or not. Theoretical objections have been raised against its employment, as being likely to increase cerebral congestion: of this there is no satisfactory proof; on the contrary, there is reason to think that chloroform inhalation has rather the effect of lessening arterial tension, while it certainly controls the violent muscular action by which the hyperæmia is so much increased. Practically, no one who has used it can doubt its great value in diminishing the force and frequency of the convulsive paroxysms. Statistically, its usefulness is shown by Charpentier in his thesis on the effects of various methods of treatment in eclampsia, since, out of 63 cases in which it was used, in 48 it had the effect of diminishing or arresting the attacks, 1 only proving fatal. The mode of administration has varied. Some have given it almost continuously, keeping the patient in a more or less profound state of anæsthesia. Others have contented themselves with carefully watching the patient, and exhibiting the chloroform as soon as there were any indications of a recurring paroxysm, with the view of

controlling its intensity. The latter is the plan I have myself adopted, and of the value of which in most cases I have no doubt. Every now and again cases will occur in which chloroform inhalation is insufficient to control the paroxysm, or in which, from the very cyanosed state of the patient, its administration seems contraindicated. Moreover, it is advisable to have, if possible, some remedy more continuous in its action and requiring less constant personal supervision. Latterly, the internal administration of chloral has been recommended for this purpose. My own experience is decidedly in its favor, and I have used, as I believe, with marked advantage a combination of chloral with bromide of potassium, in the proportion of twenty grains of the former to half a drachm of the latter, repeated at intervals of from four to six hours. If the patient be unable to swallow, the chloral may be given in an enema or hypodermically, six grains being diluted in zj of water and injected under the skin. The remarkable influence of bromide of potassium in controlling the eclampsia of infants would seem to be an indication for its use in puerperal cases. Fordyce Barker is opposed to the use of chloral, which he thinks excites instead of lessens reflex irritability.¹ Another remedy, not entirely free from theoretical objections, but strongly recommended, is the subcutaneous injection of morphia, which has the advantage of being applicable when the patient is quite unable to swallow. It may be given in doses of one-third of a grain, repeated in a few hours, so as to keep the patient well under its influence. It is to be remembered that the object is to control muscular action, so as to prevent as much as possible the violent convulsive paroxysm, and therefore it is necessary that the narcosis, however produced, should be continuous. It is rational, therefore, to combine the intermittent action of chloroform with the more continuous action of other remedies, so that the former should supplement the latter when insufficient. Inhalation of the nitrite of amyl has been recommended on physiological grounds as likely to be useful, and is well worthy of trial, but of its action I have as yet no personal experience. Pilocarpine has recently been tried in the hope that the diaphoresis and salivation it produces might diminish arterial tension and free the blood of toxic matters. Braun² administered 3 centigrammes of the muriate of pilocarpine hypodermically, and reports favorably of the result; Fordyce Barker,³ however, is of opinion that it produces so much depression as to be dangerous.

Other remedies, supposed to act in the way of antidotes to uræmic poisoning, have been advised, such as acetic or benzoic acid, but they are far too uncertain to have any reliance placed on them, and they distract attention from more useful measures.

Precautions during the Paroxysm.—Precautions are necessary during the fits to prevent the patient injuring herself, especially to obviate laceration of the tongue; the latter can be best done by placing something between the teeth as the paroxysm comes on, such as the handle of a teaspoon enveloped in several folds of flannel.

Obstetric Management.—The obstetric management of eclampsia will naturally give rise to much anxiety, and on this point there has been

¹ *The Puerperal Diseases*, p. 120.

² *Berlin Klin. Woch.*, June 16, 1879.

³ *New York Med. Rec.*, March 1, 1879.

considerable difference of opinion. On the one hand, we have practitioners who advise the immediate emptying of the uterus, even when labor has commenced; on the other, those who would leave the labor entirely alone. Thus Gooch said, "Attend to the convulsions and leave the labor to take care of itself;" and Schroeder says, "Especially no kind of obstetric manipulation is required for the safety of the mother;" but he admits, however, that it is sometimes advisable to hasten the labor to ensure the safety of the child.

In cases in which the convulsions come on during labor the pains are often strong and regular, the labor progresses satisfactorily, and no interference is needful. In others we cannot but feel that emptying the uterus would be decidedly beneficial. We have to reflect, however, that any active interference might of itself prove very irritating and excite fresh attacks. The influence of uterine irritation is apparent by the frequency with which the paroxysms recur with the pains. If, therefore, the os be undilated and labor have not begun, no active means to induce it should be adopted, although the membranes may be ruptured with advantage, since that procedure produces no irritation. Forceful dilatation of the os, and especially turning, are strongly contraindicated.

The rule laid down by Tyler Smith seems that which is most advisable to follow—that we should adopt the course which seems least likely to prove a source of irritation to the mother. Thus, if the fits seem evidently induced and kept up by the pressure of the fœtus, and the head be within reach, the forceps, or even craniotomy, may be resorted to. But if, on the other hand, there be reason to think that the operation necessary to complete delivery is likely *per se* to prove a greater source of irritation than leaving the case to nature, then we should not interfere.

CHAPTER IV.

PUERPERAL INSANITY.

Classification.—Under the head of "*Puerperal Mania*" writers on obstetrics have indiscriminately classed all cases of mental disease connected with pregnancy and parturition. The result has been unfortunate, for the distinction between the various types of mental disorder has, in consequence, been very generally lost sight of. But little study of the subject suffices to show that the term "*puerperal mania*" is wrong in more ways than one, for we find that a large number of cases are not cases of "*mania*" at all, but of melancholia; while a considerable number are not, strictly speaking, "*puerperal*," as they either come on during pregnancy or long after the immediate risks of the puerperal period are over, being in the latter case associated with anæmia produced by over-lactation. For the sake of brevity the generic term "*Puerperal Insanity*" may be employed to cover all cases of mental disorders con-

nected with gestation, which may be further conveniently subdivided into three classes, each having its special characteristics—viz.:

I. The *Insanity of Pregnancy*.

II. *Puerperal Insanity*, properly so called—that is, insanity coming on within a limited period after delivery.

III. The *Insanity of Lactation*.

This division is a strictly natural one, and includes all the cases likely to come under observation. The relative proportion these classes bear to each other can only be determined by accurate statistical observations on a large scale, but these materials we do not possess. The returns from large asylums are obviously open to objection, for only the worst and most confirmed cases find their way into these institutions, while by far the greater proportion, both before and after labor, are treated in their own homes.

Proportion of these Forms of Insanity.—Taking such returns as only approximate, we find from Dr. Batty Tuke¹ that in the Edinburgh Asylum, out of 155 cases of puerperal insanity, 28 occurred before delivery, 73 during the puerperal period, and 54 during lactation. The relative proportions of each per hundred are as follows:

Insanity of Pregnancy,	18.06 per cent.
Puerperal Insanity,	47.09 “
Insanity of Lactation,	34.83 “

Marcé² collects together several series of cases from various authorities, amounting to 310 in all, and the results are not very different from those of the Edinburgh Asylum, except in the relatively smaller number of cases occurring before delivery. The percentage is calculated from his figures:

Insanity of Pregnancy,	8.06 per cent.
Puerperal Insanity,	58.06 “
Insanity of Lactation,	30.30 “

As each of these classes differs in various important respects from the others, it will be better to consider each separately.

Insanity of Pregnancy.—The insanity of pregnancy is, without doubt, the least common of the three forms. The intense mental depression which in many women accompanies pregnancy, and causes the patient to take a desponding view of her condition and to look forward to the result of her labor with the most gloomy apprehension, seems to be often only a lesser degree of the actual mental derangement which is occasionally met with. The relation between the two states is further borne out by the fact that a large majority of cases of insanity during pregnancy are well-marked types of melancholia: out of 28 cases reported by Tuke, 15 were examples of pure melancholia, 5 of dementia with melancholia. In many of these the attack could be traced as developing itself out of the ordinary hypochondriasis of pregnancy. In others the symptoms came on at a later period of pregnancy, the earlier months of which had not been marked by any unusual lowness of spirits. The age of the patient seems to have some influence, the proportion of cases between 30 and 40 years of age being much larger than in younger women. A larger proportion of cases occur in primiparæ

¹ *Edin. Med. Journ.*, vol. x.

² *Traité de la Folie des Femmes enceintes.*

than in multiparæ—a fact that no doubt depends on the greater dread and apprehension experienced by women who are pregnant for the first time, especially if not very young. Hereditary disposition plays an important part, as in all forms of puerperal insanity. It is not always easy to ascertain the fact of an hereditary taint, since it is often studiously concealed by the friends. Tuke, however, found distinct evidence of it in no less than 12 out of 28 cases. Fürstner¹ believes that other neuroses have an important influence in the causation of the disease. Out of 32 cases he found direct hereditary taint in 9, but in 11 more there was a family history of epilepsy, drunkenness, or hysteria.

Period of Pregnancy at which it Occurs.—The period of pregnancy at which mental derangement most commonly shows itself varies. Most generally, perhaps, it is at the end of the third or the beginning of the fourth month. It may, however, begin with conception, and even return with every impregnation. Montgomery relates an instance in which it recurred in three successive pregnancies. Marcé distinguishes between true insanity coming on during pregnancy and aggravated hypochondriasis by the fact that the latter usually lessens after the third month, while the former most commonly only begins after that date. It is unquestionable that in many cases no such distinction can be made, and that the two are often very intimately associated.

Form of Insanity.—The form of insanity does not differ from ordinary melancholia. The suicidal tendency is generally very strongly developed. Should the mental disorder continue after delivery, the patient may very probably experience a strong impulse to kill her child. Moral perversions have not been uncommonly observed. Tuke especially mentions a tendency to dipsomania in the early months, even in women who have not shown any disposition to excess at other times. He suggests that this may be an exaggeration of the depraved appetite or morbid craving so commonly observed in pregnant women, just as melancholia may be a further development of lowness of spirits. Laycock mentions a disposition to “kleptomania” as very characteristic of the disease. Casper² relates a curious case where this occurred in a pregnant lady of rank, and the influence of pregnancy in developing an irresistible tendency was pleaded in a criminal trial in which one of her petty thefts had involved her.

Prognosis.—The prognosis may be said to be, on the whole, favorable. Out of Dr. Tuke's 28 cases, 19 recovered within six months. There is little hope of a cure until after the termination of the pregnancy, as, out of 19 cases recorded by Marcé, only in 2 did the insanity disappear before delivery.

Transient Mania during Delivery.—There is a peculiar form of mental derangement sometimes observed during labor which is by some talked of as a temporary insanity. It may, perhaps, be more accurately described as a kind of acute delirium, produced, in the latter stage of labor, by the intensity of the suffering caused by the pains. According to Montgomery, it is most apt to occur as the head is passing through the os uteri, or, at a later period, during the expulsion of the child. It

¹ *Archiv für Psychiatrie*, Band v. Heft 2.

² Casper's *Forensic Medicine*, New Syd. Soc., vol. iv. p. 308.

may consist of merely a loss of control over the mind, during which the patient, unless carefully watched, might in her agony seriously injure herself or her child. Sometimes it produces actual hallucination, as in the case described by Tarnier in which the patient fancied she saw a spectre standing at the foot of her bed, which she made violent efforts to drive away. This kind of mania, if it may be so called, is merely transitory in its character, and disappears as soon as the labor is over. From a medico-legal point of view it may be of importance, as it has been held by some that in certain cases of infanticide the mother has destroyed the child when in this state of transient frenzy and when she was irresponsible for her acts. In the treatment of this variety of delirium we must, of course, try to lessen the intensity of the suffering, and it is in such cases that chloroform will find one of its most valuable applications.

Puerperal Insanity (proper).—True puerperal insanity has always attracted much attention from obstetricians, often to the exclusion of other forms of mental disturbance connected with the puerperal state. We may define it to be that form of insanity which comes on within a limited period after delivery, and which is probably intimately connected with that process. Out of 73 examples of the disease tabulated by Dr. Tuke, only 2 came on later than a month after delivery, and in these there were other causes present which might possibly remove them from this class.

Although a large number of these cases assume the character of acute mania, that is by no means the only kind of insanity which is observed, a not inconsiderable number being well-marked examples of melancholia. The distinction between them was long ago pointed out by Gooch, whose admirable monograph on the disease contains one of the most graphic and accurate accounts of puerperal insanity that has yet been written.

There are also some peculiarities as to the period at which these varieties of insanity show themselves, which, taken in connection with certain facts in their etiology, may eventually justify us in drawing a stronger line of demarcation between them than has been usual. It appears that cases of acute mania are apt to come on at a period much nearer delivery than melancholia. Thus, Tuke found that all the cases of mania came on within sixteen days after delivery, and that all cases of melancholia developed themselves after that period. We shall presently see that one of the most recent theories as to the causation of the disease attributes it to some morbid condition of the blood. Should further investigation confirm this supposition, inasmuch as septic conditions of the blood are most likely to occur a short time after labor, it would not be an improbable hypothesis that cases of acute mania occurring within a short time after labor may depend on such septic causes, while melancholia is more likely to arise from general conditions favoring the development of mental disease. This must, however, be regarded as a mere speculation requiring further investigation.

Causes.—Hereditary predisposition is very frequently met with, and a careful inquiry into the patient's history will generally show that other members of the family have suffered from mental derangement. Reid found that out of 111 cases in Bethlehem Hospital there was clear evi-

dence of hereditary taint in 45. Tuke made the same observation in 22 out of his 73 cases; and, indeed, it is pretty generally admitted by all alienist physicians that hereditary tendencies form one of the strongest predisposing causes of mental disturbance in the puerperal state. In a large proportion of cases circumstances producing debility and exhaustion or mental depression have preceded the attack. Thus, it is often found that patients attacked with it have had post-partum hemorrhage, or have suffered from some other conditions producing exhaustion, such as severe and complicated labor, or they may have been weakened by over-frequent pregnancies, or by lactation during the early months of pregnancy. Indeed, anaemia is always well marked in this disease. Mental conditions also are frequently traceable in connection with its production. Morbid dread during pregnancy, insufficient to produce insanity before delivery, may develop into mental derangement after it. Shame and fear of exposure in unmarried women not unfrequently lead to it, as is evidenced by the fact that out of 2281 cases, gathered from the reports of various asylums, above 64 per cent. were unmarried.¹ Sudden moral shocks or vivid mental impressions may be the determining cause in predisposed persons. Gooch narrates an example of this in a lady who was attacked immediately after a fright produced by a fire close to her house, the hallucinations in this case being all connected with light; and Tyler Smith, that of another whose illness dated from the sudden death of a relative. The age of the patient has some influence, and there seems to be a decidedly greater liability at advanced ages, especially when such women are pregnant for the first time.

Theory of its Dependence on a Morbid State of the Blood.—The possibility of the acute form of puerperal insanity, coming on shortly after delivery, being dependent on some form of septicæmia is one which deserves careful consideration. The idea originated with Sir James Simpson, who found albumen in the urine of four patients. He suggested that this might properly indicate the presence in the blood of certain urinary constituents which might have determined the attack, much in the same way as in eclampsia. Dr. Donkin subsequently wrote an important paper,² in which he warmly supported this theory, and arrived at the conclusion "that the acute dangerous class of cases are examples of uræmic blood-poisoning, of which the mania, rapid pulse, and other constitutional symptoms are merely the phenomena; and that the affection, therefore, ought to be termed uræmic or renal puerperal mania in contradistinction to the other form of the disease." He also suggests that the immediate poison may be carbonate of ammonia, resulting from the decomposition of urea retained in the blood. It will be observed, therefore, that the pathological condition producing puerperal mania would, supposing this theory to be correct, be precisely the same as that which at other times is supposed to give rise to puerperal eclampsia. There can be no doubt that the patient immediately after delivery is in a condition rendering her peculiarly liable to various forms of septic disease; and it must be admitted that there is no inherent improbability in the supposition that some morbid material circulating in the blood may be the effective cause of the attack in a person

¹ *Journ. of Mental Science*, 1870-71, p. 159.

² *Edin. Med. Journ.*, vol. vii.

otherwise predisposed to it. It is also certain, as I have already pointed out, that there are two distinct classes of cases, differing according to the period after delivery at which the attack comes on. Whether this difference depends on the presence in the blood of some septic matter—especially urinary excreta—is a question which our knowledge by no means justifies us in answering; it is, however, one which well merits further careful study.

Objections to this Theory.—It is only fair to point out some difficulties which appear to militate against the view which Dr. Donkin maintains. In the first place, the albuminuria is merely transient, while its supposed effects last for weeks or months. Sir James Simpson says, with regard to his cases, "I have seen all traces of albuminuria in puerperal insanity disappear from the urine within fifty hours of the access of the malady. The general rapidity of its disappearance is perhaps the principal—or indeed the only—reason why this complication has escaped the notice of those physicians among us who devote themselves with such ardor and zeal to the treatment of insanity in our public asylums." This apparent anomaly Simpson attempts to explain by the hypothesis that when once the uræmic poisoning has done its work and set the disease in progress, the mania progresses of itself. This, however, is pure speculation, and in the supposed analogous case of eclampsia the albuminuria certainly lasts as long as its effects. It is not easy to understand, also, why uræmic poisoning should in one case give rise to insanity and in another to convulsions. For all we know to the contrary, transient albuminuria may be much more common after delivery than has been generally supposed, and further investigation on this point is required. Albumen is by no means unfrequently observed in the urine, for a short time, in various conditions of the body, without any serious consequences, as, for example, after bathing; and we may too readily draw an unjustifiable conclusion from its detection in a few cases of mania. There are, however, many other kinds of blood-poisoning besides uræmia which may have an influence in the production of the disease, and it is to be hoped that future observations may enable us to speak with more certainty on this point.

Prognosis.—The prognosis of puerperal insanity is a point which will always deeply interest those who have to deal with so distressing a malady. It may resolve itself into a consideration of the immediate risk to life and of the chances of ultimate restoration of the mental faculties. It is an old aphorism of Gooch's—and one the correctness of which is justified by modern experience—that "mania is more dangerous to life, melancholia to reason." It has very generally been supposed that the immediate risk to life in puerperal mania is not great; and on the whole this may be taken as correct. Tuke found that death took place, from all causes, in 10.9 per cent. of the cases under observation; these, however, were all women who had been admitted into asylums, and in whom the attack may be assumed to have been exceptionally severe. Great stress was laid by Hunter and Gooch on extreme rapidity of the pulse as indicating a fatal tendency. There can be no doubt that it is a symptom of great gravity, but by no means one which need lead us to despair of our patient's recovery. The most dangerous class

of cases are those attended with some inflammatory complication; and if there be marked elevation of temperature, indicating the presence of some such concomitant state, our prognosis must be more grave than when there is mere excitement of the circulation.

Post-mortem Signs.—There are no marked post-mortem signs found in fatal cases to guide us in forming an opinion as to the nature of the disease. "No constant morbid changes," says Tyler Smith, "are found within the head, and most frequently the only condition found in the brain is that of unusual paleness and exsanguinity. Many pathologists have also remarked upon the extremely empty condition of the blood-vessels, particularly the veins."

Duration of the Disease.—The duration of the disease varies considerably. Generally speaking, cases of mania do not last so long as melancholia, and recovery takes place within a period of three months, often earlier. Very few of the cases admitted into the Edinburgh Asylum remained there more than six months, and after that time the chances of ultimate recovery greatly lessened. When the patient gets well, it often happens that her recollection of the events occurring during her illness is lost; at other times the delusions from which she suffered remain; as, for example, in a case which was under my care in which the personal antipathies which the patient formed when insane became permanently established.

Insanity of Lactation.—54 out of the 155 cases collected by Dr. Tuke were examples of the insanity of lactation, which would appear, therefore, to be nearly twice as common as that of pregnancy, but considerably less so than the true puerperal form. Its dependence on causes producing anæmia and exhaustion is obvious and well marked. In the large majority of cases it occurs in multiparæ who have been debilitated by frequent pregnancies and by length of nursing. When occurring in primiparæ, it is generally in women who have suffered from post-partum hemorrhage or other causes of exhaustion, or whose constitution was such as should have contraindicated any attempt at lactation. The bruit de diable is almost invariably present in the veins of the neck, indicating the impoverished condition of the blood.

The type is far more frequently melancholic than maniacal, and when the latter form occurs the attack is much more transient than in true puerperal insanity. The danger to life is not great, especially if the cause producing debility be recognized and at once removed.

There seems, however, to be more risk of the insanity becoming permanent than in the other forms. In 12 out of Dr. Tuke's cases the melancholia degenerated into dementia, and the patients became hopelessly insane.

Symptoms.—The symptoms of these various forms of insanity are practically the same as in the non-pregnant state.

In Cases of Mania.—Generally, in cases of mania there is more or less premonitory indication of mental disturbance, which may pass unperceived. The attack is often preceded by restlessness and loss of sleep, the latter being a very common and well-marked symptom, or, if the patient sleep, her rest is broken and disturbed by dreams. Causeless dislikes to those around her are often observed: the nurse, the husband,

the doctor, or the child becomes the object of suspicion, and, unless proper care be taken, the child may be seriously injured. As the disease advances the patient becomes incoherent and rambling in her talk, and in a fully-developed case she is incessantly pouring forth an unconnected jumble of sentences out of which no meaning can be made. Often some prevalent idea which is dwelling in the patient's mind can be traced running through her ravings, and it has been noticed that this is frequently of a sexual character, causing women of unblemished reputation to use obscene and disgusting language which it is difficult to understand their even having heard. The tendency of such patients to make accusations impugning their own chastity was specially insisted on by many eminent authorities in a recent celebrated trial, when Sir James Simpson stated that in his experience "the organ diseased gave a type to the insanity, so that with women suffering from affections of the genital organs the delusions would be more likely to be connected with sexual matters." Religious delusions—as a fear of eternal damnation or of having committed some unpardonable sin—are of frequent occurrence, but perhaps more often in cases which are tending to the melancholic type. There is generally intolerable restlessness, and the patient's whole manner and appearance are those of excessive excitement. She may refuse to remain in bed, may tear off her clothes, or may attempt to injure herself. The suicidal tendency is often very marked. In one case under my care the patient made incessant efforts to destroy herself, which were only frustrated by the most careful watching; she endeavored to strangle herself with the bedclothes, to swallow any article she could lay hold of, and even to gouge out her own eyes. Food is generally persistently refused, and the utmost coaxing may fail in inducing the patient to take nourishment. The pulse is rapid and small, and the more violent the excitement and furious the delirium the more excited is the circulation. The tongue is coated and furred, the bowels constipated and disordered, and the feces, as well as the urine, are frequently passed involuntarily. The urine is scanty and high-colored, and after the disease has lasted for some time it becomes loaded with phosphates. The lochia and the secretion of milk generally become arrested at the commencement of the disease. The waste of tissue, from the incessant restlessness and movement of the patient, is very great, and if the disease continue for some time she falls into a condition of marasmus, which may be so excessive that she becomes wasted to a shadow of her former size.

Symptoms of Melancholia.—When the insanity assumes the form of melancholia its advent is more gradual. It may commence with depression of spirits without any adequate cause, associated with insomnia, disturbed digestion, headache, and other indications of bodily derangement. Such symptoms, showing themselves in women who have been nursing for a length of time or in whom any other evident cause of exhaustion exists, should never pass unnoticed. Soon the signs of mental depression increase and positive delusions show themselves. These may vary much in their amount, but they are all more or less of the same type, and very often of a religious character. The amount of constitutional disturbance varies much. In some cases which approach in character those of mania there is considerable excitement, rapid pulse, furred

tongue, and restlessness. Probably cases of acute melancholia coming on during the puerperal state most often assume this form. In others, again, there is less of these general symptoms; the patients are profoundly dejected, sit for hours without speaking or moving; but there is not much excitement, and this is the form most generally characterizing the insanity of lactation. In all cases there is a marked disinclination to food. There is also, almost invariably, a disposition to suicide; and it should never be forgotten in melancholic cases that this may develop itself in an instant, and that a moment's carelessness on the part of the attendants may lead to disastrous results.

Treatment.—Bearing in mind what has been said of the essential character of puerperal insanity, it is obvious that the course of treatment must be mainly directed to maintain the strength of the patient, so as to enable her to pass through the disease without fatal exhaustion of the vital powers, while we endeavor at the same time to calm the excitement and give rest to the disturbed brain. Any over-active measures—for example, bleeding, blistering the shaven scalp, and the like—are distinctly contraindicated.

There is a general agreement on the part of alienist physicians that in cases of acute mania the two things most needed are a sufficient quantity of suitable food and sleep.

Importance of Administering Nourishment.—Every endeavor should be made to induce the patient to take plenty of nourishment, to remedy the defects of the excessive waste of tissue and support her strength until the disease abates. Dr. Blandford, who has especially insisted on the importance of this, says: ¹ “Now, with regard to the food, skilful attendants will coax a patient into taking a large quantity, and we can hardly give too much. Messes of minced meat with potato and greens, diluted with beef-tea, bread and milk, rum and milk, arrowroot, and so on, may be got down. Never give mere liquids so long as you can get down solids. As the malady progresses the tongue and mouth may become so dry and foul that nothing but liquids can be swallowed; but, reserving our beef-tea and brandy, let us give plenty of solid food while we can.”

Forcible Administration of Food.—The patient may in mania, as well as in melancholia—perhaps even more in the latter—obstinately refuse to take nourishment at all, and we may be compelled to use force. Various contrivances have been employed for this purpose. One of the simplest is introducing a dessert-spoon forcibly between the teeth, the patient being controlled by an adequate number of attendants, and slowly injecting into the mouth suitable nourishment by an india-rubber bottle with an ivory nozzle, such as is sold by all chemists. Care must be taken not to inject more than an ounce at a time, and to allow the patient to breathe between each deglutition. So extreme a measure will seldom be required if the patient have experienced attendants who can overcome her resistance to food by gentler means; but it may be essential, and it is far better to employ it than to allow the patient to become exhausted from want of nourishment. In one case I had to feed a patient in this way three times a day for several weeks, and used for the pur-

¹ Blandford, *Insanity and its Treatment*.

pose a contrivance known in asylums as Paley's feeding-bottle, which reduced the difficulty of the process to a minimum. Beef-tea or strong soup mixed with some farinaceous material, such as Revalenta Arabica, or wheaten flour or milk, forms the best mess for this purpose.

Stimulants.—In the early stages the patient is probably better without stimulants, which seem only to increase the excitement. As the disease progresses and exhaustion becomes marked, it may be necessary to have recourse to them. In melancholia they seem to be more useful, and may be administered with greater freedom.

State of the Bowels.—The state of the bowels requires especial attention. They are almost always disordered, the evacuations being dark and offensive in odor. In the early stages of the disease the prompt clearing of the bowels by a suitable purgative sometimes has the effect of cutting short an impending attack. A curious example of this is recorded by Gooch, in which the patient's recovery seemed to date from the free evacuation of the bowels. A few grains of calomel or a dose of compound jalap powder or of castor oil may generally be readily given. During the continuance of the illness the state of the primæ viæ should be attended to, and occasional aperients will be useful, but strong and repeated purgation is hurtful, from the debility it produces.

The Procuring Sleep.—The procuring sleep will necessarily form one of the most important points of treatment. For this purpose there is no drug so valuable as the hydrate of chloral, either alone or in combination with bromide of potassium, which has a distinct effect in increasing its hypnotic action. Given in a full dose at bedtime—say 15 grs. to 3ss—it rarely fails in procuring at least some sleep, and in an early stage of acute mania this may be followed by the best effects. It may be necessary to repeat this draught night after night during the acute stage of the malady. If we cannot induce the patient to swallow the medicine, it may be given in the form of enema.

Question of Administering Opiates.—It is generally admitted that in mania preparations of opium, formerly much relied on in the treatment of the disease, are apt to do more harm than good. Dr. Blandford gives a strong opinion on this point. He says: "In prolonged delirious mania I believe opium never does good, and may do great harm. We shall see the effects of narcotic poisoning if it be pushed, but none that are beneficial. This applies equally to opium given by the mouth and by subcutaneous injection. The latter, as it is more certain and effectual in producing good results, is also more deadly when it acts as a narcotic poison. After the administration of a dose of morphia by the subcutaneous method the patient will probably at once fall asleep, and we congratulate ourselves that our long-wished-for object is attained. But after half an hour or so the sleep suddenly terminates, and the mania and excitement are worse than before. Here you may possibly think that, had the dose been larger, instead of half an hour's sleep you would have obtained one of longer duration; and you may administer more, but with a like result. Large doses of morphia not merely fail to produce refreshing sleep; they poison the patient, and produce, if not the symptoms of actual narcotic poisoning, at any rate that typhoid condition which indicates prostration and approaching collapse. I believe

there is no drug the use of which more often becomes abused than that of opium." It is otherwise in cases of melancholia, especially in the more chronic forms. In these opiates, in moderate doses, not pushed to excess, may be given with great advantage. The subcutaneous injection of morphia is by far the best means of exhibiting the drug, from its rapidity of action and facility of administration.

Other Calmatives.—There are other methods of calming the excitement of the patient besides the use of medicines. The prolonged use of the warm bath, the patient being immersed in water at a temperature of 90° or 92° for at least half an hour, is highly recommended by some as a sedative. The wet pack serves the same purpose, and is more readily applied in refractory subjects.

Importance of Judicious Nursing.—Judicious nursing is of primary importance. The patient should be kept in a cool, well-ventilated, and somewhat darkened room. If possible she should remain in bed, or at least endeavors should be made to restrain the excessive restless motion which has so much effect in promoting exhaustion. The presence of relatives and friends, especially the husband, has generally a prejudicial and exciting effect; and it is advisable to place the patient under the care of nurses experienced in the management of the insane, who, as strangers, are likely to have more control over her. It is not too much to say that much of the success in treatment must depend on the manner in which this indication is met. Rough, unskilled nurses, who do not know how to use gentleness combined with firmness, will certainly aggravate and prolong the disorder. Inasmuch as no patient should be left unwatched by day or night, more than one nurse is essential.

Question of Removal to an Asylum.—The question of the removal of the patient to an asylum is one which will give rise to anxious consideration. As the fact of having been under such restraint of necessity fixes a certain lasting stigma upon a patient, this is a step which every one would wish to avoid if possible. In cases of acute mania, which will probably last a comparatively short time, home-treatment can generally be efficiently carried out. Much must depend on the circumstances of the patient. If these be of a nature which preclude the possibility of her obtaining thoroughly efficient nursing and treatment in her own home, it is advisable to remove her to a place where these essentials can be obtained, even at the cost of some subsequent annoyance. In cases of chronic melancholia, the management of which is on the whole more difficult, the necessity for such a measure is more likely to arise, and should not be postponed too late. Many examples of incurable dementia arising out of puerperal melancholia can be traced to unnecessary delay in placing the patients under the most favorable conditions for recovery.

Treatment during Convalescence.—When convalescence is commencing, change of air and scene will often be found of great value. Removal to some quiet country place, where the patient can enjoy abundance of air and exercise in the company of her nurses, without the excitement of seeing many people, is especially to be recommended. Great caution must be used in admitting the visits of relatives and friends. In two cases under my own care the patients relapsed, when apparently pro-

gressing favorably, because the husbands insisted, contrary to advice, on seeing them. On the other hand, Gooch has pointed out that when the patient is not recovering, when month after month has been passed in seclusion without any improvement, the visit of a friend or relative may produce a favorable moral impression and inaugurate a change for the better. It is probably in cases of melancholia, rather than in mania, that this is likely to happen. The experiment may under such circumstances be worth trying, but it is one the result of which we must contemplate with some anxiety.

CHAPTER V.

PUERPERAL SEPTICÆMIA.

Difference of Opinion as to Puerperal Fever.—There is no subject in the whole range of obstetrics which has caused so much discussion and difference of opinion as that to which this chapter is devoted. Under the name of *Puerperal Fever* the disease we have to consider has given rise to endless controversy. One writer after another has stated his view of the nature of the affection with dogmatic precision, often on no other grounds than his own preconceived notions and an erroneous interpretation of some of the post-mortem appearances. Thus, one states that puerperal fever is only a local inflammation, such as peritonitis; others declare it to be phlebitis, metritis, metro-peritonitis, or an essential zymotic disease, *sui generis*, which affects lying-in women only. The result has been a hopeless confusion, and the student rises from the study of the subject with little more useful knowledge than when he began. Fortunately, modern research is beginning to throw a little light upon this chaos.

Modern View of the Disease.—The whole tendency of recent investigation is daily rendering it more and more certain that obstetricians have been led into error by the special virulence and intensity of the disease, and that they have erroneously considered it to be something special to the puerperal state, instead of recognizing in it a form of septic disease practically identical with that which is familiar to surgeons under the name of pyæmia or septicæmia.

Objection to the Name "Puerperal Fever."—If this view be correct, the term "puerperal fever," conveying the idea of a fever such as typhus or typhoid, must be acknowledged to be misleading, and one that should be discarded as only tending to confusion. Before discussing at length the reasons which render it probable that the disease is in no way specific or peculiar to the puerperal state, it will be well to relate briefly some of the leading facts connected with it.

History of the Disease.—More or less distinct references to the existence of the so-called puerperal fever are met with in the classical authors,

proving, beyond doubt, that the disease was well known to them; and Hippocrates, besides relating several cases the nature of which is unquestionable, clearly recognizes the possibility of its originating in the retention and decomposition of portions of the placenta. Although Harvey and other writers showed that they were more or less familiar with it, and even made most creditable observations on its etiology, it was not until the latter half of the last century that it came prominently into notice. At that time the frightful mortality occurring in some of the principal lying-in hospitals, especially in the Hôtel Dieu at Paris, attracted attention; and ever since the disease has been familiar to obstetricians.

Mortality resulting from it in Lying-in Hospitals.—Its prevalence in hospitals in which lying-in women are congregated has been constantly observed both in this country and abroad, occasionally producing an appalling death-rate; the disease, when once it has appeared, frequently spreading from one patient to another in spite of all that could be done to arrest it. It would be easy to give many startling instances of this. Thus, it prevailed in London in the years 1760, 1768, and 1770 to such an extent that in some lying-in institutions nearly all the patients died. Of the Edinburgh Infirmary in 1773 it is stated that "almost every woman as soon as she was delivered, or perhaps about twenty-four hours after, was seized with it, *and all of them died*, though every method was used to cure the disorder." On the Continent, where the lying-in institutions are on a much larger scale, the mortality was equally great. Thus in the Maison d'Accouchements of Paris in a number of different years sometimes as many as 1 in 3 of the women delivered died, on one occasion 10 women dying out of 15 delivered. Similar results were observed in other great continental hospitals, as in Vienna, where, in 1823, 19 per cent. of the cases died, and, in 1842, 16 per cent.; and in Berlin in 1862 hardly a single patient escaped, the hospital being eventually closed.

Should Lying-in Hospitals be Abolished?—Such facts, the correctness of which is beyond any question, prove to demonstration the great risk which may accompany the aggregation of lying-in women. Whether they justify the conclusion that all lying-in hospitals should be abolished is another and a very wide question which can scarcely be satisfactorily discussed in a practical work. It is to be observed, however, that most of the cases in which the disease produced such disastrous results occurred before our more recent knowledge of its mode of propagation was acquired, when no sufficient hygienic precautions were adopted, when ventilation was little thought of, and when, in a word, every condition prevailed that would tend to favor the spread of a contagious disease from one patient to another. More recent experience proves that when the contrary is the case (as, for example, in such an institution as the Rotunda Hospital in Dublin), the occurrence of epidemics of this kind may be entirely prevented and the mortality approximated to that of home-practice.

The Assumption of a Puerperal Miasm is Unnecessary.—The more closely the history of these outbreaks in hospitals is studied, the more apparent does it become that they are not dependent on any miasm

necessarily produced by the aggregation of puerperal patients, but on the direct conveyance of septic matter from one patient to another.

In numerous instances the disease has been said to be generally epidemic in domiciliary practice, much in the same way as scarlet fever or any zymotic complaint might be. Such epidemics are described as having occurred in London in 1827–28, in Leeds in 1809–12, in Edinburgh in 1825, and many others might be cited. There is, however, no sufficient ground for believing that the disease has ever been epidemic in the strict sense of the word. That numerous cases have often occurred in the same place and at the same time is beyond question; but this can easily be explained without admitting an epidemic influence, knowing as we do how readily septic matter may be conveyed from one patient to another. In many of the so-called epidemics the disease has been limited to the patients of certain midwives or practitioners, while those of others have entirely escaped—a fact easily understood on the assumption of the disease being produced by septic matter conveyed to the patient, but irreconcilable with the view of general epidemic influence. We are not in possession of any reliable statistics of the mortality arising from puerperal septicæmia in ordinary general practice. It has, however, been well pointed out, in the Report on Puerperal Fever presented by the Obstetrical Society of Berlin to the Prussian Minister of Health,¹ that not only do the published returns of death from metria afford no reliable estimate of the actual mortality from this source, but that they are very far more numerous than deaths from any other cause in connection with pregnancy and childbirth.

Numerous Theories advanced regarding its Nature.—It would be a useless task to detail at length the theories that have been advanced to explain the disease. Indeed, it may safely be held that the supposed necessity of providing a theory which would explain all the facts of the disease has done more to surround it with obscurity than even the difficulties of the subject itself. If any real advance is to be made, it can only be by adopting a humble attitude, by admitting that we are only on the threshold of the inquiry, and by a careful observation of clinical facts, without drawing from them too positive deductions.

Theory of its Local Origin.—Many have taught that the disease is essentially a local inflammation producing secondary constitutional effects. This view doubtless originated from too exclusive attention to the morbid changes found on post-mortem examination. Excessive peritonitis, phlebitis, inflammation of the lymphatics or of the tissues of the uterus, are very commonly found after death; and each of these has, in its turn, been believed to be the real source of the disease. This view finds but little favor with modern pathologists, and is in so many ways inconsistent with clinical facts that it may be considered to be obsolete. No one of the conditions above mentioned is universally found, and in the worst cases definite signs of local inflammation may be entirely absent. Nor will this theory explain the conveyance of the disease from one patient to another, or the peculiar severity of the constitutional symptoms.

Theory of an Essential Zymotic Fever.—A more admissible theory,

¹ See *Edin. Med. Journ.*, Nov., 1878.

and one which has been extensively entertained, is, that there is an essential zymotic fever peculiar to, and only attacking, puerperal women, which is as specific in its nature as typhus or typhoid, and to which the local phenomena observed after death bear the same relation that the pustules on the skin do to small-pox or the ulcers in the intestinal glands to typhoid. This fever is supposed to spread by contagion and infection, and to prevail epidemically, both in private and in hospital practice. The most recent exponent of this view is Fordyce Barker, who in his excellent work on the *Puerperal Diseases* has entered at length into all the theories of the disease. He, like others who hold his opinions, has, I cannot but think, entirely failed to bring forward any conclusive evidence of the existence of such a specific fever. It is no doubt true that in typhus and typhoid and other undoubted examples of this class of disease there are well-marked local secondary phenomena; but then they are distinct and constant. He makes no attempt to prove that anything of the kind occurs in puerperal fever. On the contrary, probably there are no two cases in which similar local phenomena occur, nor is there any case in which the most practised obstetrician could foretell either the course and duration of the illness or the local phenomenon. Again, this theory altogether fails to explain the very important class of cases which can be distinctly traced to sources originating in the patient herself—viz. the absorption of septic matter from decomposing coagula and the like. Barker meets this difficulty by placing such cases of auto-infection under a separate category, admitting that they are examples of septicæmia. But he fails to show that there is any difference in symptomatology or post-mortem signs between them and the cases he believes to depend on an essential fever; nor would it be possible to distinguish the one from the other by either their clinical or pathological history.

Theory of its Identity with Surgical Septicæmia.—The modern view, which holds that the disease is, in fact, identical with the condition known as pyæmia or septicæmia, is by no means free from objections, and much patient clinical investigation is required to give a satisfactory explanation of certain peculiarities which the disease presents; but in spite of these difficulties, which time may serve to remove, it offers a far better explanation of the phenomena observed than any other that has yet been advanced.

According to this theory, the so-called puerperal fever is produced by the absorption of septic matter into the system through solutions of continuity in the generative tract, such as always exist after labor. It is not essential that the poison should be peculiar or specific, for, just as in surgical pyæmia, any decomposing organic matter, either originating within the generative organs of the patient herself or coming from without, may set up the morbid action.

In describing the disease under discussion I shall assume that, so far as our present knowledge goes, this view is the one most consonant with facts; but, bearing in mind that very little is yet known of surgical septicæmia, it must not be expected that obstetricians can satisfactorily explain all the phenomena they observe.

Basis of Description.—The best basis of description I know of is that given by John Sanderson when he says, "In every pyæmic process

you may trace a focus, a centre of origin, lines of diffusion or distribution, and secondary results from the distribution—in every case an initial process from which infection commences, from which the infection spreads, and secondary processes which come out of this primary one.”¹ Adopting this division, I shall first treat of the mode in which the infection may commence in obstetric cases, and point out the special difficulties which this part of the subject presents.

Channels through which Septic Matter may be Absorbed.—The fact that all recently-delivered women present lesions of continuity in the generative tract, through which septic matter, brought into contact with them, may be readily absorbed, has long been recognized. The analogy between the interior of the uterus after delivery and the surface of a stump after amputation was particularly insisted on by Cruveilhier, Simpson, and others—an analogy which was, to a great extent, based on erroneous conceptions of what took place, since they conceived that the whole interior of the uterus was bared. It is now well known that that is not the case; but the fact remains that at the placental site, at any rate, there are open vessels through which absorption may readily take place. That absorption of septic material occurs through this channel is probable in certain cases in which decomposing materials exist in the interior of the uterus, especially when, from defective uterine contraction, the venous sinuses are abnormally patulous and are not occluded by thrombi. It is difficult to understand how septic matter, introduced from without, can reach the placental site. Other sites of absorption are, however, always available. These exist in every case in the form of slight abrasions or lacerations about the cervix, or in the vagina, or, especially in primiparæ, about the fourchette and perineum. There is even some reason to think that absorption of septic matter may take place through the mucous membrane of the vagina or cervix without any breach of surface. This might serve to account for the occasional although rare cases in which symptoms of the disease develop themselves before delivery, or so soon after it as to show that the infection must have preceded labor; nor is there any inherent improbability in the supposition that septic material may be occasionally absorbed through the unbroken mucous membrane, as is certainly the case with some poisons—for example, that of syphilis. Hence there is no difficulty in recognizing the similarity of a lying-in woman to a patient suffering from a recent surgical lesion, or in understanding how septic matter conveyed to her during or shortly after labor may be absorbed. It is necessary, however, to suppose that absorption takes place immediately or very shortly after these lesions of continuity are formed, for it is well known that the power of absorption is arrested after they have commenced to heal. This fact may explain the cases in which sloughing about the perineum or vagina exists without any septicæmia resulting, or the far from uncommon cases in which an intensely fetid lochial discharge may be present a few days after delivery without any infection taking place.

The character and sources of the septic matter constitute one of the most obscure questions in connection with septicæmia, and that which is most open to discussion.

¹ *Clinical Transactions*, vol. viii. p. 108.

Division into Autogenetic and Heterogenetic Cases.—The most practical division of the subject is into cases in which the septic matter originates within the patient, so that she infects herself, the disease then being properly *autogenetic*; and into those in which the septic matter is conveyed from without, and brought into contact with absorptive surfaces in the generative tract, the disease then being *heterogenetic*.

Sources of Self-infection.—The sources of auto-infection may be various, but they are not difficult to understand. Any condition giving rise to decomposition, either of the tissues of the mother herself, of matters retained in the uterus or vagina that ought to have been expelled, or decomposing matter derived from a putrid fœtus, may start the septicæmic process. Thus it may happen that from continuous pressure on the maternal soft parts during labor sloughing has set in, or there may be already decomposing material present from some previous morbid state of the genital tracts, as in carcinoma. A more common origin is the retention of coagula or of small portions of membrane or of placenta in the interior of the uterus, which have putrefied from access of air, or in the decomposition of the lochia. That the retention of portions of the placental tissue has at all times been the cause of septicæmia may be illustrated by the case of the Duchesse d'Orleans (in the time of Louis XIII.), who had an easy labor, but died of childbed fever. An examination was made by the leading physicians of Paris, in their report of which it was stated: "On the right side of the womb was found a small portion of after-birth so firmly adherent that it could be hardly torn off by the finger-nails."¹ The reason why self-infection does not more often occur from such sources, since more or less decomposition is of necessity so often present, has already been referred to in the fact that absorption of such matters is not apt to occur when the lesions of continuity always existing after parturition have commenced to heal. This observation may also serve to explain how previous bad states of health, by interfering with the healthy reparative process occurring after delivery, may predispose to self-infection. It is interesting to note that puerperal septicæmia arising from such sources is not limited to the human race. In the debate on pyæmia at the Clinical Society, Mr. Hutchinson recorded several well-marked examples occurring in ewes, in whose uteri portions of retained placenta were found.

Sources of Heterogenetic Infection.—The sources of septic matter conveyed from without are much more difficult to trace, and there are many facts connected with heterogenetic infection which are very difficult to reconcile with theory, and of which, it must be admitted, we are not yet able to give a satisfactory explanation.

It is probable that any decomposing organic matter may infect, but that some forms operate with more certainty and greater virulence than others.

Influence of Cadaveric Poisoning.—One of these, which has attracted special attention, is what may be termed cadaveric poison, derived from dissection of the dead subject in the anatomical and post-mortem theatres, and conveyed to the genital tract by the hands of the accoucheur. Attention was particularly directed to this source of infection

¹ *Louise Bourgeois*, by Goodell.

by the observations of Semmelweiss, who showed that in the division of the Vienna Lying-in Hospital attended by medical men and students who frequented the dissecting-rooms the mortality was seldom less than 1 in 10, while in the division solely attended by women the mortality never exceeded 1 in 34; the number of deaths in the former division at once falling to that of the latter so soon as proper precautions and means of disinfection were used. Many other facts of a like nature have since been recorded which render this origin of puerperal septicæmia a matter of certainty. An interesting example is related by Simpson with characteristic candor: "In 1836 or 1837, Mr. Sidey of this city had a rapid succession of five or six cases of puerperal fever in his practice at a time when the disease was not known to exist in the practice of any other practitioners in the locality. Dr. Simpson, who had then no firm or proper belief in the contagious propagation of puerperal fever, attended the dissection of Mr. Sidey's patients and freely handled the diseased parts. The next four cases of midwifery which Dr. Simpson attended were all affected with puerperal fever, and it was the first time he had seen it in practice. Dr. Patterson of Leith examined the ovaries, etc. The three next cases which Dr. Patterson attended in that town were attacked with the disease."¹ Negative examples are of course brought forward of those who have attended post-mortem examinations without injury to their obstetric patients, which merely prove that the cadaveric poison does not, of necessity, attach itself to the hands of the dissector; and no amount of such testimony can invalidate such positive evidence as that just narrated. Barnes believes that there is not so much danger attending the dissection of patients who have died of any ordinary disease, but that the risk attending the dissection of those who have died of infectious or contagious complaints is very great indeed.² I presume there is no doubt that the risk is greater when the subject has died from zymotic disease, but the distinction is too delicate to rely on; and the attendant on midwifery will certainly err on the safe side by avoiding as much as possible having anything to do with the conduct of dissections or post-mortem examinations.

Infection from Erysipelas.—Another possible source of infection is erysipelatos disease in all its forms. The intimate connection between erysipelas and surgical pyæmia has long been recognized by surgeons, and the influence of erysipelas in producing puerperal septicæmia has been especially observed in surgical hospitals into which lying-in patients were also admitted. Trousseau relates instances of this kind occurring in Paris. The only instance that I know of in London was in the lying-in ward of King's College Hospital, where, in spite of every hygienic precaution, the mortality was so great as to necessitate the closure of the ward. Here the association of erysipelas with puerperal septicæmia was again and again observed, the latter proving fatal in direct proportion to the prevalence of the former in the surgical wards. The dependence of the two on the same poison was in one instance curiously shown by the fact of the child of a patient who died of puerperal septicæmia dying from erysipelas which started from a slight abrasion

¹ *Selected Obstet. Works*, p. 508.

² "Lectures on Puerperal Fever," *Lancet*, vol. ii., 1865.

produced by the forceps. A more recent and very remarkable example is related by Dr. Lombe Atthill.¹ A patient suffering from erysipelas was admitted into the Rotunda Hospital on February 15, 1877. The sanitary condition of the hospital was at the time excellent. The patient was removed next day, but of the next 10 patients confined in adjoining wards, 9 were attacked with puerperal peritonitis, the only one who escaped being a case of abortion. But the connection between erysipelas and puerperal septicæmia is not limited to hospitals, having been often observed in domiciliary practice. Some interesting facts have been collected by Dr. Minor,² who has shown that the two diseases have frequently prevailed together in various parts of the United States, and that during a recent outbreak of puerperal fever in Cincinnati it occurred chiefly in the practice of those physicians who attended cases of erysipelas. Many children also died from erysipelas whose mothers had died from puerperal fever.

Infection from other Zymotic Diseases.—There is good reason to believe that the contagium of other zymotic diseases may produce a form of disease indistinguishable from ordinary puerperal septicæmia and presenting none of the characteristic features of the specific complaint from which the contagium was derived. This is admitted to be a fact by the majority of our most eminent British obstetricians, although it does not seem to be allowed by continental authorities, and it is strongly controverted by some writers in this country. It is certainly difficult to reconcile this with the theory of septicæmia, and we are not in a position to give a satisfactory explanation of it. I believe, however, that the evidence in favor of the possibility of puerperal septicæmia originating in this way is too strong to be assailable.

Cases Produced by the Contagion of Scarlet Fever.—The scarlatinal poison is that regarding which the greatest number of observations have been made. Numerous cases of this kind are to be found scattered through our obstetric literature, but the largest number are to be met with in a paper by Dr. Braxton Hicks in the twelfth volume of the *Obstetrical Transactions*; and they are especially valuable from that gentleman's well-known accuracy as a clinical observer. Out of 68 cases of puerperal disease seen in consultation, no less than 37 were distinctly traced to the scarlatinal poison. Of these, 20 had the characteristic rash of the disease; but the remaining 17, although the history clearly proved exposure to the contagium of scarlet fever, showed none of its usual symptoms, and were not to be distinguished from ordinary typical cases of the so-called puerperal fever. On the theory that it is impossible for the specific contagious diseases to be modified by the puerperal state, we have to admit that one physician met with 17 cases of puerperal septicæmia in which, by a mere coincidence, the contagion of scarlet fever had been traced, and that the disease nevertheless originated from some other source—an hypothesis so improbable that its mere mention carries its own refutation.

Cases Produced by the Contagion of other Zymotic Diseases.—With regard to the other zymotic diseases the evidence is not so strong, proba-

¹ *Medical Press and Circular*, April, 1877.

² *Erysipelas and Childhood Fever*, Cincinnati, 1874.

bly from the comparative rarity of the diseases. Hicks mentions one case in which the diphtheritic poison was traced, although none of the usual phenomena of the disease were present. I lately saw a case in which a lady a few days after delivery had a very serious attack of septicæmia without any diphtheritic symptoms, her husband being at the same time attacked with diphtheria of a most marked type. Here it would be difficult not to admit the dependence of the two diseases on the same poison.

The Zymotic Diseases are not always Modified in the Puerperal State.—It is, however, certain that all the zymotic diseases may attack a newly-delivered woman and run their characteristic course without any peculiar intensity. Probably most practitioners have seen cases of this kind; and this is precisely one of the points of difficulty which we cannot at present explain, but on which future research may be expected to throw some light. It seems to me not improbable that the explanation of the fact that zymotic poison may in one puerperal patient run its ordinary course, and in another produce symptoms of intense septicæmia, may be found in the channel of absorption. It is at any rate comprehensible that if the contagium be absorbed through the skin or the ordinary channel it may produce its characteristic symptoms and run its usual course, while if brought into contact with lesions of continuity in the generative tract it may act more in the way of septic poison, or with such intensity that its specific symptoms are not developed.

It may reasonably be objected that if puerperal and surgical septicæmia be identical the zymotic poisons ought to be similarly modified when they infect patients after surgical operations. The subject of specific contagium as a cause of surgical pyæmia has been so little studied that I do not think any one would be justified in asserting that such an occurrence is not possible. Fritsch of Halle and other German physicians have recently shown how elaborate antiseptic precautions in lying-in hospitals may prevent the origin of the disease from such sources. Sir James Paget, in his *Clinical Lectures*, seems to believe in the possibility of such modification. He says: "I think it not improbable that in some cases results occurring with obscure symptoms within two or three days after operations have been due to scarlet-fever poison, hindered in some way from its usual progress." Sir Spencer Wells informs me that he has seen cases of surgical pyæmia which he had reason to believe originated in the scarlatinal poison; and his well-known success as an ovariologist is no doubt, in a great measure, to be attributed to his extreme care in seeing that no one likely to come in contact with his patients has been exposed to any such source of infection.

Sewer Gas and Defective Sanitary Arrangements.—Exposure to sewer gas may, I feel sure, produce the disease. In two cases of the kind I had the opportunity of closely watching an untrapped drain opened directly into the bedroom—in one instance into a bath, in the other into a water-closet. Both cases were indistinguishable from the ordinary form of the disease, and in both improvement commenced as soon as the patient was removed into another room.

In a case I saw some years ago in Notting Hill the patient, who had been confined within a week, had all the symptoms of a most intense

attack of septicæmia, but none of a diphtheritic character, while her husband lay in an adjoining room suffering from a diphtheritic sore throat. Here the waste-pipe of the bath was found to communicate directly with the sewer. In spite of her intense illness, I had the patient removed to another house, and from that moment she began to improve. In two other cases in which the same source of disease was detected the removal of the patient from the infected atmosphere was immediately followed by a marked amelioration in the symptoms. I know of three similar cases which ended fatally in which I have every reason to believe that the cause of the disease was poisoning by sewer gas. Frankenhauser has related a curious case of the poisoning of four puerperal women by sewer gas. In fact, the whole question of defective sanitary conditions on the puerperal state deserves much more serious study than it has ever yet received, and I have long been satisfied that they have often much to do with certain grave forms of illness in the lying-in state the origin of which cannot otherwise be traced.

Septicæmia from Contagion conveyed from other Puerperal Patients.—The last source from which septic matter may be conveyed is from a patient suffering from puerperal septicæmia—a mode of origin which has of late attracted special attention. That this is the explanation of the occasional endemic prevalence of the disease in lying-in hospitals can scarcely be doubted. The theory of a special puerperal miasm pervading the hospital is not required to account for the facts, for there are a hundred ways impossible to detect or avoid—on the hands of nurses or attendants, in sponges, bed-pans, sheets, or even suspended in the atmosphere—in which septic material derived from one patient may be carried to another.

The poison may be conveyed in the same manner from one private patient to another. Of this there are many lamentable instances recorded. Thus it was mentioned by a gentleman at the recent discussion at the Obstetrical Society that 5 out of 14 women he attended died, no other practitioner in the neighborhood having a case. This origin of the disease was clearly pointed out by Gordon¹ toward the end of last century, who stated that he himself "was the means of carrying the infection to a great number of women;" and he also traced the spread of the disease in the same way in the practice of certain midwives. In some remarkable instances the unhappy property of carrying contagion has clung to individuals in a way which is most mysterious, and which has led to the supposition that the whole system becomes saturated with the poison. One of the strangest cases of this kind was that of Dr. Rutter of Philadelphia, which caused much discussion. He had 45 cases of puerperal septicæmia in his own practice in one year, while none of his neighbors' patients were attacked. Of him it is related, "Dr. Rutter, to rid himself of the mysterious influence which seemed to attend upon his practice, left the city for ten days, and before waiting on the next parturient case had his hair shaved off and put on a wig, took a hot bath, and changed every article of his apparel, taking nothing with him that he had worn or carried to his knowledge on any former occasion; and mark the result. The lady, notwithstanding that she had

¹ See *Lectures on Puerperal Fever*, by Robert J. Lee, M. D.

an easy parturition, was seized the next day with childbed fever, and died on the eleventh day after the birth of the child. Two years later he made another attempt at self-purification, and the next case attended fell a victim to the same disease." No wonder that Meigs, in commenting on such a history, refused to believe that the doctor carried the poison, and rather thought that he was "merely unhappy in meeting with such accidents through God's providence." It appears, however, that Dr. Rutter was the subject of a form of ozæna, and it is quite obvious that under such circumstances his hands could never have been free from septic matter.¹ This observation is of peculiar interest, as showing that the sources of infection may exist in conditions difficult to suspect and impossible to obviate, and it affords a satisfactory explanation of a case which was for years considered puzzling in the extreme. It is quite possible that other similar cases—of which many are on record, although none so remarkable—may possibly have depended on some similar cause personal to the medical attendant.

The sources of septic poison being thus multifarious, a few words may be said as to the mode in which it may be conveyed to the patient.

Mode in which the Poison may be Conveyed to the Patient.—As on the view of puerperal septicæmia which seems most to agree with recorded facts the poison, from whatever source it may be derived, must come into actual contact with lesions of continuity in the generative tract, it is obvious that one method of conveyance may be on the hands of the accoucheur. That this is a possibility, and that the disease has often been unhappily conveyed in this way, no one can doubt. Still, it would be unfair in the extreme to conclude that this is the only way in which infection may arise. In town-practice, especially, there are many other ways in which septic matter may reach the patient. The nurse may be the means of communication, and, if she have been in contact with septic matter, she is even more likely than the medical attendant to convey it when washing the genitals during the first few days after delivery, the time that absorption is most apt to occur. Barnes relates a whole series of cases occurring in a suburb of London in the practice of different practitioners, every one of which was attended by the same nurse. Again, septic matter may be carried in sponges, linen, and other articles. What is more likely, for example, than that a careless nurse might use an imperfectly-washed sponge on which discharge has been allowed to remain and decompose? Nor do I see any reason to question the possibility of infection from septic matter suspended in the atmosphere; and in lying-in hospitals, where many women are congregated together, there can be little doubt that this is a common origin of the disease. It is certain, whatever view we may take of the character of the septic material,

¹ This is stated on the authority of an obstetrical contemporary of Dr. Rutter. (See *Amer. Journ. of Med. Science*, April, 1875, p. 474.)

The author quotes from the editor. Dr. Rutter had an ozæna which in time much disfigured him from its effect upon the contour of his nose. He was unfortunately inoculated in his index finger from a patient, and neglected the pustule. He had 95 cases of puerperal septicæmia in four years and nine months, with 18 deaths. The question of Dr. Meigs, who was a non-contagionist in regard to puerperal peritonitis, was remarkably apposite: "Did he distil a subtle essence which he carried with him?" —Harris's note to third American edition.

that it must be in a state of very minute subdivision, and there is no theoretical difficulty in the assumption of its being conveyed by the atmosphere.

Conduct of the Practitioner in Relation to the Disease.—This question naturally involves a reference to the duty of those who are unfortunately brought into contact with septic matter in any form, either in a patient suffering from puerperal septicæmia, zymotic disease, or offensive discharges. The practitioner cannot always avoid such contact, and it is practically impossible to relinquish obstetric work every time that he is in attendance on a case from which contagion may be carried. Nor do I believe, especially in these days when the use of antiseptics is so well understood, that it is essential. It was otherwise when antiseptics were not employed, but I can scarcely conceive any case in which the risk of infection cannot be prevented by proper care. The danger I believe to be chiefly in not recognizing the possible risk and in neglecting the use of proper precautions. It is impossible, therefore, to urge too strongly the necessity of extreme, and even exaggerated, care in this direction. The practitioner should accustom himself, as much as possible, to use the left hand only in touching patients suffering from infectious diseases, as that which is not used under ordinary circumstances in obstetric manipulations. He should be most careful in the frequent employment of antiseptics in washing his hands, such as Condyl's fluid, carbolic acid, or the 1-in-1000 solution of perchloride of mercury. Clothing should be changed on leaving an infectious case. Much more care than is usually practised should be taken by nurses, especially in securing perfect cleanliness in everything brought into contact with the patient. When, however, a practitioner is in actual and constant attendance on a case of puerperal septicæmia—when he is visiting his patient many times a day, especially if he be himself washing out the uterus with antiseptic lotions—it is certain that he cannot deliver other patients with safety; and he should secure the assistance of a brother-practitioner, although there seems no reason why he should not visit women already confined in whom he has not to make vaginal examinations.

Prophylaxis of Septicæmia.—If the views here inculcated as to the nature of, and mode of infection in, puerperal septicæmia be correct, it is obvious that much may be done in the way of prophylaxis. A perfectly aseptic management of puerperal women is practically impossible. In many lying-in institutions on the Continent, and in some in this country, very rigid rules have been laid down to prevent the possibility of infective matter being conveyed to the patient either on the hands of the attendants or on instruments, napkins, and the like, and, it is said, with very satisfactory results. As the risk is much greater when lying-in women are collected together, such precautions, which this is not the place to discuss, are absolutely indicated. They are not, however, applicable in ordinary private practice, but there are certain simple precautions, which every one might adopt without trouble, which will materially lessen the risk of septic poisoning. Amongst these may be indicated the use of antiseptic lotions, with which the practitioner and nurse should always wash their hands before attending any case or touching the genital organs; the use of carbolized oil, 1-in-8 for lubricating the fingers,

catheter, forceps, etc. ; syringing out the vagina night and morning with diluted Condyl's fluid ; rigid attention to cleanliness in bedding, napkins, etc. Precautions such as these, although they may appear to some frivolous and useless, indicate a recognition of danger and an endeavor to remove it, and, if they were generally inculcated on nurses (see *note*, p. 555) and others, might go far to prevent the occurrence of septic mischief.

Nature of the Septic Poison.—As to the precise character of the septic poison, although of late much has been said about it, and there is good reason to believe that further research may throw light on this obscure subject, too little is known to justify any positive statement. With regard to the influence of minute micro-organisms and their supposed connection with the production of the disease, this is especially the case. The recent researches of Heiberg, Von Recklinghausen, Steurer, and others have shown that in puerperal septicæmia, as in surgical fever, erysipelas, and other infectious diseases, chain-like micrococci in large numbers may be traced passing between the muscular and connective-tissue fibres through the lymphatics, and thus into the general circulation, and that they may be found in various organs and pathological products. These observations are of much importance, as tending to confirm by scientific observation the intimate relation between these various forms of disease which has long been believed to exist. It may be taken as certain that these bodies bear an intimate and important relation to the disease ; but whether they themselves form the septic matter or carry it, or whether they are mere accidental concomitants of the pyæmic processes, it is impossible, in the present state of our knowledge, to state, and I therefore prefer to dwell on that part of the subject which is of clinical importance, rather than enter into speculative theories which may to-morrow prove to be valueless.¹

Channels of Diffusion.—Passing on to the channels of diffusion through which the septic matter may act, we have to consider its effects on the structures with which it is brought into contact and the mode in which it may infect the system at large ; and this will include a consideration of the pathological phenomena.

Local changes consequent on the absorption of the poison are pretty constant, and of these we may form an intelligible idea by thinking of them as similar in character and causation to those which we have the opportunity of studying when septic matter is applied to a wound open to observation ; as, for example, in cases of blood-poisoning following a dissection wound. Distinct traces of local action are not of invariable occurrence, and in some of the worst class of cases, when the amount of septic matter is great and its absorption rapid, death may occur after an illness of short duration, but great intensity, and before appreciable local changes, either at the site of absorption or in the system at large, have had time to develop themselves. The fact that puerperal fever may prove fatal without leaving any tangible post-mortem signs has often been pointed out, such cases most frequently occurring during the en-

¹ For the latest information on this point see "Our Present Knowledge of the Relations between Micro-organisms and Puerperal Fever," by Carl Lomer, M. D., *Amer. Journ. of Obstet.*, July, 1884.

demic prevalence of the disease in lying-in hospitals. There can be little doubt, however, that in such cases of intense septicæmia marked pathological changes exist in the form of alterations of the blood and degenerations of tissue, but not of a character which can be detected by an ordinary post-mortem examination. In the great majority of cases indications of the disease exist at the site of absorption. These are described by pathologists as identical in their character with the inflammatory oedema which occurs in connection with phlegmonous erysipelas. If lacerations exist in the cervix or vagina, they take on unhealthy action, their edges swell, and their surfaces become covered with a yellowish coat similar in appearance to diphtheritic membrane. The mucous membrane of the uterus is also generally found to be affected, and in a degree varying with the intensity of the local septic process. There is evidence of severe endometritis, and very frequently the whole lining of the uterus is profoundly altered, softened, covered with patches of diphtheritic deposit, and, it may be, in a state of general necrosis. In the severer cases these changes affect the muscular tissue of the uterus, which is found to be swollen, soft, imperfectly contracted, and even partially necrosed—a condition which is likened by Heiberg to hospital gangrene. The connective tissue surrounding the generative tract is also swollen and oedematous, and the inflammation may in this way reach the peritoneum, although peritonitis, so often observed in puerperal septicæmia, does not necessarily depend on the direct transmission of inflammation from the pelvic connective tissue, but is more often a secondary phenomenon.

Channels through which Systemic Infection is Produced.—The channels through which general systemic infection may supervene are the lymphatics and the venous sinuses, the former being by far the most important. Recent researches have shown the great number and complexity of the lymphatics in connection with the pelvic viscera, and marked traces of the absorption of septic matter are almost always to be found, except in those very intense cases, already alluded to, in which no appreciable post-mortem signs are discoverable. The septic matter is probably absorbed from the lymph-spaces abounding in the connective tissue, and carried along the lymphatic canals to the nearest glands. The result is inflammation of their coats and thrombosis of their contents, which may be seen on section as a creamy, purulent substance. The absorption of septic material may, as Virchow has shown, be delayed by the local changes produced in the lymphatics and in the glands with which they communicate, which are therefore conservative in their action; and the further progress of the case may in this way be stopped and local inflammation alone result, such cases being believed by Heiberg to be examples of abortive pyæmia. On the other hand, the free septic material may be too abundant and intense to be so arrested; it may pass on through the lymph-canals and glands until it reaches the blood-current through the thoracic duct, and so produces a general blood-infection. This mode of absorption of septic matter, and the tendency of the glands to arrest its further progress, serve to explain the progressive character of many cases in which fresh exacerbations seem to occur from time to time, since fresh quantities of poison, generated at

its source of origin, may be absorbed as the case progresses. The uterine veins are supposed by D'Espine to be the channel of absorption in the intense form of disease which proves fatal very shortly after delivery, too soon for the more gradual process of lymphatic absorption to have become established. It is evident that the veins are not likely to act in this way, since they must, under ordinary circumstances, be completely occluded by thrombi, otherwise hemorrhage would occur. If, however, uterine contraction be incomplete, the occlusion of the venous sinuses may be imperfect, and absorption of septic material through them may then take place. Some writers have laid great stress on imperfect uterine contraction in predisposing to septicæmia, and its influence may thus be well explained. The veins may bear an important part in the production of septicæmia independent of the direct absorption of septic matter through them by means of the detachment of minute portions of their occluding thrombi in the form of emboli. If phlegmonous inflammation occur in the immediate vicinity of the veins, the thrombi they contain may become infected. When once blood-infection has occurred by any of these channels, general septicæmia, the so-called puerperal fever, is developed.

Pathological Phenomena observed after General Blood-Infection.—The variety of pathological phenomena found on post-mortem examination has had much to do with the prevalent confusion as to the nature of the disease. This has resulted in the description of many distinct forms of puerperal fever, the most marked pathological alteration having been taken to be the essential element of the disease. As a matter of fact, there is no doubt that various types of pathological change are met with. Heiberg describes four chief classes which are by no means distinctly separated from one another, are often found simultaneously in the same subject, and are certainly not to be distinguished by the symptoms during life.

Intense Cases without Marked Post-mortem Signs.—Of these, the first is the class of cases in which no appreciable morbid phenomena are found after death. This formidable and fatal form of the disease has long been well known, and is that described by some of our authors as adynamic or malignant puerperal fever. It is the variety which was so prevalent in our lying-in hospitals, and which Ramsbotham talks of as being second only to cholera in the severity and suddenness of its onset and in the rapidity with which it carried off its victims. It is quite erroneous to suppose that the existence of pathological changes in this form of disease has never been recognized. Even with the coarse methods of examination formerly used, the occurrence of a fluid and altered state of the blood and ecchymoses in connection with various organs—especially the lungs, spleen, and kidneys—were noticed and specially described by Copland in his *Dictionary of Medicine*. More recently it has been clearly proved by the microscope that there exists, in addition, the commencement of inflammation in most of the tissues, shown by cloudy swellings and granular infiltration and disintegration of the cell-elements; proving that the blood, heavily charged with septic matter, had set up morbid action wherever it circulated, the patient succumbing before this had time to develop.

Cases characterized by Inflammation of the Serous Membranes.—In the second type, and that perhaps most commonly met with, the morbid changes are more frequently found in the serous membranes, in the pleura, in the pericardium, but, above all, in the peritoneum, the alterations in which have long attracted notice, and have been taken by many writers as proving peritonitis to be the main element of the disease. Evidences of more or less peritonitis are very general. In the more severe cases there is little or no exudation of plastic lymph, such as is found in peritonitis unassociated with septicæmia. There is a greater or less quantity of brownish serum only, the coils of intestine, distended with flatus and highly congested, being surrounded by it. More often there are patchy deposits of fibrinous exudation over many of the viscera, the fundus uteri, the under surface of the liver, and the distended intestines. There is then also a considerable quantity of sero-purulent fluid in the abdominal cavity. The pleural cavities may also exhibit similar traces of inflammatory action, containing imperfectly-organized lymph and sero-purulent fluid. Schroeder states that pleurisy is more often the direct result of transmission of inflammation through the substance of the diaphragm or lung than a secondary consequence of the septicæmia. In like manner, evidences of pericarditis may exist, the surface of the pericardium being highly injected and its cavity containing serous fluid. Inflammation of the synovial membranes of the larger joints, occasionally ending in suppuration, is not uncommon, and may probably be best included under this class of cases.


Cases characterized by Changes in the Mucous Membranes.—In the third type the mucous membranes appear to bear the brunt of the disease. The pathological changes are most marked in the mucous membrane lining the intestines, which is highly congested, and even ulcerated in patches, with numerous small spots of blood extravasated in the sub-mucous tissue. Similar small apoplectic effusions have been observed in the substance of the kidneys and under the mucous membrane of the bladder. Pneumonia is of common occurrence. In most cases it is probably secondary to the impaction of minute emboli in the smaller branches of the pulmonary artery, but it may doubtless arise from independent inflammation of the lung-tissue, and will then be included in the class of cases now under consideration.

Cases characterized by the Impaction of Infected Emboli and Secondary Inflammation and Abscess.—The fourth class of pathological phenomena are those which are produced chiefly by the impaction of minute infected emboli in small vessels in various parts of the body. These are the cases which most closely resemble surgical pyæmia, both in their symptoms and post-mortem signs, and which by many writers are described under the name of puerperal pyæmia. The dependence of puerperal fever on phlebitis of the uterine veins was a favorite theory, and in a large proportion of cases the coats of the veins show signs of inflammation, their canals being occupied with thrombi in a more or less advanced state of disintegration. The mode in which these thrombi may become infected has been shown by Babnoff, who has proved that leucocytes may penetrate the coats of the vein, and, entering its contained coagulum, may set up disintegration and suppuration. This

to find that the symptoms are also very various in different cases. This can readily be explained by the amount and virulence of the poison absorbed, the channels of infection, and the organs which are chiefly implicated; but it renders it very difficult to describe the disease satisfactorily.

The symptoms generally show themselves within two or three days after delivery. As infection most often occurs during labor, or, in cases which are autogenetic, within a short time afterward, and before the lesions of continuity in the generative tract have commenced to cicatrize,

FIG. 194.




it can be understood why septicæmia rarely commences later than the fourth or fifth day.

The Early Symptoms are not Well Marked.—In the great majority of cases the disease begins insidiously. There are, generally, some chilliness and rigor, but by no means always, and even when present they frequently escape observation or are referred to some transient cause. The first symptom which excites attention is a rise in the pulse, which may vary from 100 to 140 or more, according to the severity of the attack; and the thermometer will also show that the temperature is raised to 102°, or in bad cases even to 104° or 106°. Still, it must be borne in mind that both the pulse and temperature may be increased in the puerperal state from transient cause, and do not, of themselves, justify the diagnosis of septicæmia.

Symptoms of Intense Septicæmia.—In the more intense class of cases, in which the whole system seems overwhelmed with the severity of the attack, the disease progresses with great rapidity and often without any appreciable indication of local complication. The pulse is very rapid, small, and feeble, varying from 120 to 140, and there is generally a temperature of 103° or 104° . In the worst form of cases the temperature is steadily high, without marked remissions. (See Figs. 193, 194, and 195.) There may be little or no pain, or there may be slight tenderness on pressure over the abdomen or uterus, and as the disease progresses the intestines get largely distended with flatus, so that intense tympanites often form a most distressing symptom. The countenance is sallow, sunken, and has a very anxious expression. As a rule, intelligence is unimpaired, and this may be the case even in the worst forms of the disease and up to the period of death. At other times there is a good deal of low muttering delirium, which often occurs at night alone, and alternates with intervals of complete consciousness, but is occasionally intensified for a short time into a more acute form. Diarrhœa and vomiting are of very frequent occurrence; by the latter dark, grumous, coffee-ground substances are ejected. The diarrhœa is occa-

FIG. 195.




sionally very profuse and uncontrollable; in mild cases it seems to relieve the severity of the symptoms. The tongue is moist and loaded with sordes, but sometimes it gets dark and dry, especially toward the termination of the disease. The lochia are generally suppressed or altered in character, and sometimes they have a highly offensive odor, especially

when the disease is autogenetic. The breathing is hurried and panting, and the breath itself has a very characteristic, heavy, sweetish odor. The secretion of milk is often, but not always, arrested.

Duration of the Disease.—With more or less of these symptoms the case goes on, and when it ends fatally it generally does so within a week, the fatal termination being indicated by more weakness, rapid, thread-like, or intermittent pulse, marked delirium, great tympanites, and sometimes a sudden fall of temperature, until at last the patient sinks with all the symptoms of profound exhaustion.

Variety of Symptoms in Different Cases.—In milder cases similar symptoms, variously modified and combined, are present. It is seldom

FIG. 196.




that two precisely similar cases are met with ; in some the rapid weak pulse is most marked ; in others, abdominal distension, vomiting, diarrhoea, or delirium.

Symptoms of Peritonitis.—Local complications variously modify the symptoms and course of the disease. The most common is peritonitis, so much so that with some authors puerperal fever and puerperal peritonitis are synonymous terms. Here the first symptom is severe abdominal pain, commencing at the lower part of the abdomen, where the uterus is felt enlarged and tender. As the abdominal pain and tenderness spread, the sufferings of the patient greatly increase, the intestines become enormously distended with flatus, and the breathing is entirely

racteristic rub; pleurisy by dulness on percussion; kidney affection by albuminuria and the presence of casts; liver complication by jaundice; and so on.

Pyæmic Forms of the Disease.—The course of the disease is not always so intense and rapid, being in some cases of a more chronic cha-

FIG. 198.



racter and lasting many weeks. The symptoms in the early stage are often indistinguishable from those already described, and it is generally only after the second week that indications of purulent infection develop themselves. Then we often have recurrent and very severe rigors, with marked elevations and remissions of temperature. At the same time there is generally an exacerbation of the general symptoms, a peculiar yellowish discoloration of the skin, and occasionally well-developed jaundice. Transient patches of erythema are not uncommonly observed on various parts of the skin, and such eruptions have often been mistaken for those of scarlet fever or other zymotic disease. Localized inflammations and suppuration may rapidly follow. Amongst the most common is inflammation or even suppuration of the joints—the knees, shoulders, or hips—which is preceded by difficulty of movement, swelling, and very acute pain. Large collections of pus in various parts of the muscles and connective tissues are not rare. Suppurative inflammation may also be found in connection with many organs, as in the eye, in the pleura, pericardium, or lungs; each of which will, of course,

give rise to characteristic symptoms, more or less modified by the type of the disease and the intensity of the inflammation.

Puerperal Malarial Fever.—There is a peculiar form of febrile disturbance which sometimes occurs in the puerperal state, and which is apt to be confounded with septicæmia, to which attention has been specially directed by Fordyce Barker¹ under the name of “puerperal malarial fever.” It is specially apt to be met with in women who have been exposed to malarial poison during their former lives, the recurrence of the fever being probably determined by the puerperal state. Of this I have seen several very well-marked examples in ladies who have formerly contracted fever or ague in India. One of my patients, who has long been in India and suffered from intermitten fever for years, is invariably attacked with it after delivery, and herself warned me of the fact the first time I attended her. The diagnosis is not always easy. Barker insists on the fact that puerperal malarial fever generally commences after the fifth day from delivery, while septicæmia almost always does so before that time. In the malarial fever, moreover, the intermissions are much more marked, while there are frequently-recurring chills or rigors, which is not the case in septicæmia.


Treatment.—In considering the all-important subject of treatment the views of the practitioner are naturally biassed by the theory he has adopted of the nature of the disease. If that here inculcated be correct, the indications we have to bear in mind are—1st, to discover, if possible, the source of the poison, in the hope of arresting further septic absorption; 2d, to keep the patient alive until the effects of the poison are worn off; and, 3d, to treat any local complications that may arise.

The Use of Antiseptic Injections.—The first is likely to be of great importance in cases of self-infection, as fresh quantities of septic matter may be from time to time absorbed. We, fortunately, are in possession of a powerful means of preventing further absorption by the application of antiseptics to the interior of the uterus and to the canal of the vagina.² This is especially valuable when the existence of decomposing coagula or other sources of septic matter is suspected in the uterine cavity, or when offensive discharges are present. Disinfection is readily

¹ “Puerperal Malarial Fever,” *Amer. Journ. of Obstet.*, April, 1880.

² My colleague, Dr. Hayes, has invented a silver tube for the purpose of administering such intra-uterine injections (Fig. 199), which answers its purpose admirably. The

FIG. 199.




Hayes's Tube for Intra-uterine Injections.

numerous apertures at its extremity allow of a number of minute streams of fluid being thrown out in the form of a spray over the interior of the uterus, the complete bathing of its surface and washing out of its cavity being thus ensured. It is, moreover, introduced more easily than the ordinary vaginal pipe, and can be attached to a Higginson syringe.


accomplished by washing out the uterine cavity at least twice daily by means of a Higginson's syringe with a long vaginal pipe attached. The

FIG. 200.



third day after delivery, when the temperature was found to be slightly increased. On the morning of the eighth day the temperature had risen to 105.8°. She was delirious, with a rapid, thready pulse, clammy per-

FIG. 201.



spiration, tympanitic abdomen, and her general condition indicated the most urgent danger. On vaginal examination a piece of compressed and putrid placenta was found in the os. This was removed by my colleague, Dr. Hayes, and the uterus thoroughly washed out with Cond's fluid and water. The same evening the temperature had sunk to 99°, and the general symptoms were much improved. The next day there was a slight return of offensive discharge and an aggravation of the symptoms. After again washing out the uterus the temperature fell, and from that date the patient convalesced without a single bad symptom (Fig. 201).

This is a very well-marked example of the value of local antiseptic treatment, and I have seen many cases of the same kind. It should therefore never be omitted in all cases in which self-infection is possible; and, indeed, even when there is no reason to suspect the presence of a local focus of infection the use of antiseptic lotions is advisable as a matter of precaution, since it can do no harm and is generally com-

forting to the patient. Any antiseptic may be used, such as a weak solution of carbolic acid, 1 in 50, or of tincture of iodine, or Cond's fluid

largely diluted. I generally use the two latter alternately, the one in the morning, the other in the evening. Occasionally I have employed a 1-in-50 solution of carbolic acid, with about 5 grains to the ounce of iodoform suspended in it. This has the advantage of not only being a very powerful antiseptic, but of acting more continuously in consequence of the powdered iodoform remaining partially attached to the uterine walls; or, as some have advised, an iodoform pessary may be placed in the uterine cavity. The nozzle of the syringe should be guided well through the cervix, and the cavity of the uterus thoroughly washed out until the fluid that issues from the vagina is no longer discolored. As the os is always patulous, there is no risk of producing the troublesome symptoms of uterine colic which occasionally follow the use of intra-uterine injections in the unimpregnated state. It is quite useless to entrust the injection to the nurse, and it should be performed at least twice daily by the practitioner himself in all cases in which the discharges are offensive. It is not advisable, however, that such injections should be used indiscriminately, since they are not entirely free from risk, nor should they be continued for more than a few days. The vulva should in all cases be carefully inspected, with the view of ascertaining if the source of infection be not some local slough or necrotic ulcer about the perineum or orifice of the vagina, in which case its surface should be freely covered with iodoform. I have seen more than one instance in which this simple procedure has sufficed to cut short symptoms of a very threatening character.

Administration of Food and Stimulants.—In a disease characterized by so marked a tendency to prostration the importance of sustaining the vital powers by an abundance of easily-assimilated nourishment cannot be overrated. Strong beef-tea or other forms of animal soup, milk, alone or mixed either with lime- or soda-water, and the yolk of eggs beat up with milk and brandy, should be given at short intervals, and in as large quantities as the patient can be induced to take; and the value of thoroughly efficient nursing will be especially apparent in the management of this important part of the treatment. As there is frequently a tendency to nausea, the patient may resist the administration of food, and the resources of the practitioner will be taxed in administering it in such form and variety as will prove least distasteful. Generally speaking, not more than one or two hours should be allowed to elapse without some nutriment being given. The amount of stimulant required will vary with the intensity of the symptoms and the indications of debility. Generally, stimulants are well borne, prove decidedly beneficial, and require to be given pretty freely. In cases of moderate severity a tablespoonful of good old brandy or whiskey every four hours may suffice; but when the pulse is very rapid and thready, when there is much low delirium, tympanites, or sweating (indicating profound exhaustion), it may be advisable to give them in much larger quantities and at shorter intervals. The careful practitioner will closely watch the effects produced, and regulate the amount by the state of the patient, rather than by any fixed rule; but in severe cases eight or twelve ounces of brandy, or even more, in the twenty-four hours may be given with decided benefit.

Venesection not Admissible.—Venesection, both general and local, was long considered a sheet-anchor in this disease. Modern views are, however, entirely opposed to its use; and in a disease characterized by so profound an alteration of the blood and so much prostration it is too dangerous a remedy to employ, although it is possible that it might alleviate temporarily the severity of some of the symptoms, especially in cases in which peritonitis is well marked and much local pain and tenderness are present.^[1]

Medicinal Treatment.—The rational indications in medicinal treatment are to lessen the force of the circulation as much as is possible without favoring exhaustion, and to diminish the temperature.

Use of Arterial Sedatives.—For the former purpose Barker strongly advocates the use of *veratrum viride*, in doses of five drops of the tincture every hour, until the pulse falls to below 100, when its effects are subsequently kept up by two or three drops every second hour. Of this drug I have no personal experience, but I have extensively used minute doses of tincture of aconite for the same purpose, and when carefully given I believe it to be a most valuable remedy. The way I have administered it is to give a single drop of the tincture, at first every half hour, increasing the interval of administration according to the effect produced. Generally, after giving four or five doses at intervals of half an hour, the pulse begins to fall, and afterward a few doses, at intervals of one or two hours, will suffice to prevent the heart's action rising to its former rapidity. The advantage of thus modifying cardiac action with the view of preventing excessive waste of tissue cannot be questioned. It is evident that so powerful a remedy must not be used without the most careful supervision, for, if continued too long or given at too frequent intervals, it may unduly depress the circulation, and do more harm than good. It is necessary, therefore, that the practitioner should constantly watch the effect of the drug, and stop it if the pulse become very weak or if it intermit. It is most likely to be useful at an early stage of the disease before much exhaustion is present, and then only when the pulse is of a certain force and volume. Barker says of the *veratrum viride*, what is also true of aconite, that "it should not be given in those cases in which rapid prostration is manifested by a feeble, thread-like, irregular pulse, profuse sweats, and cold extremities."

Reduction of Temperature.—The reduction of temperature must form an important part of our treatment, and for this purpose many agents are at our disposal.

Quinine.—Quinine in large doses, of from 10 to 30 grains, has been much used for this purpose, especially in Germany. After its exhibition

[¹ I believe that the entire abandonment of venesection has been a grave error, and that where there is early in the attack a high pulse, with great abdominal distension and tenderness and a decided elevation of temperature, we ought to bleed the patient, sitting, at once, and to such a degree as to produce a decided impression. One of the worst cases I ever saw was cured in this way. The woman was delivered at 3½ p. m. of one day, and the disease manifested itself in twenty hours. At 9 the next morning she was apparently doing well; at 1 she was in great suffering, and could not bear her abdomen to be touched; vs. f 5xvj: at 9 p. m., symptoms more grave; vs. f 5xl in a sitting posture until she felt sick. At 10 p. m. pulse 150: in twenty-four hours from this no fever and very little pain: in three days regarded as out of danger. Saw her in robust health, with her child living, a year later.—Ed.]

the temperature frequently falls one or two degrees. It may be given morning and evening. Unpleasant head-symptoms, deafness, and ringing in the ears often render its continuance for a length of time impossible. These may, however, be much lessened by the addition of 10 to 15 minims of hydrobromic acid to each dose.

Salicylic Acid.—Salicylic acid, in doses of from 10 to 20 grains, or the salicylate of soda in the same doses, is a valuable antipyretic which I have found on the whole more manageable than quinine. Under its use the temperature often falls considerably in a short space of time. It is, however, apt to depress the circulation, and thus requires to be carefully watched while it is being administered; and should the pulse become very small and feeble it should be discontinued.

Warburg's Tincture.—In some cases, especially when the fever has assumed a remittent type, I have administered with marked benefit a drug which is of high repute in India in the worst class of malarious remittent fevers, and the almost marvellous effects of which in such cases I had myself witnessed in India many years ago. This is the so-called "Warburg's tincture," the value of which has been testified to by many high authorities, among whom I may mention Dr. Maclean of Netley, Dr. Broadbent, and Sir Alexander Armstrong, the Director-General of the Medical Department of the Navy, who informs me that it is now supplied to all Her Majesty's ships in the tropics, because it is found to be of the utmost value in cases in which quinine has little or no effect.

Recently its composition has been made public by Dr. Maclean. The basis is quinine, in combination with various aromatics and bitters, some of which probably intensify its action. Be this as it may, the testimony in favor of the antipyretic action of the remedy is very strong. I have found its exhibition followed by a profuse diaphoresis (this being its almost invariable effect), and sometimes a rapid amelioration of the symptoms. In other cases in which I have tried it, like everything else, it has proved of no avail. Of its use in ten malarial cases above alluded to Dr. Fordyce Barker says: "For nearly two years past, in those cases where the stomach will tolerate it, I have found Warburg's tincture much more effective and speedy in producing the results desired than the largest doses of quinine."¹

Application of Cold.—Cold may be advantageously tried in suitable cases. The simplest mode of using it is by Thornton's ice-cap, by which a current of cold water is kept continuously running round the head. This has been found of great value in pyrexia after ovariectomy, and I have also found it useful as a means of reducing temperature in puerperal cases. It is a comforting application, and gives great relief to the throbbing headache, which often causes much suffering. Under its use the temperature often falls two or more degrees, and it is easily continued day and night.

In very serious cases, when the temperature reaches 105° and upward, the external application of cold to the rest of the body may be tried. I have elsewhere related² a case of puerperal septicæmia with hyper-

¹ *Op. cit.*, p. 278.

² "A Lecture on a Case of Puerperal Septicæmia, with Hyperpyrexia, treated by the Continuous Application of Cold," *Brit. Med. Journ.*, Nov. 17, 1877.

pyrexia, the temperature continuously ranging over 105° , in which I kept the patient for eleven days nearly continuously covered with cloths soaked in iced water, by which means only was the temperature kept within moderate bounds and life preserved. But this method of treatment is excessively troublesome and is in no way curative. It is only of use in moderating the temperature when it has reached a point at which it could not continue long without destroying the patient. I should therefore never think of employing it unless the temperature was over 105° , and then only as a temporary expedient, requiring incessant watching, to be desisted from as soon as the temperature had reached a more moderate height. It is clearly impossible to place a puerperal patient in a bath, as is practised in hyperpyrexia associated with acute rheumatism or typhoid fever. The same effect may, however, be obtained by placing her on macintosh sheeting, or, still better, on a water-bed, into which cold water is run from time to time, and covering the body with towels soaked in iced water, which are frequently renewed by the attendant nurses. During the application the temperature should be constantly taken, and as soon as it has fallen to 101° the cold applications should be discontinued.

Administration of Turpentine.—Amongst other remedies which have been used is turpentine, which was highly thought of by the Dublin school. In cases with much tympanitic distension and a small weak pulse it is sometimes of unquestionable value, and it probably acts as a strong nervine stimulant. Given in doses of 15 to 20 minims, rubbed up with mucilage, it can generally be taken in spite of its nauseous taste.

Evacuant Remedies.—Purgatives, diaphoretics, or even emetics, have often been employed as eliminants of the poison. The former are strongly recommended by Schroeder and other German authorities, and in this country they were formerly amongst the most favorite remedies, and there is a general concurrence of opinion amongst our older writers as to their value. In the first volume of the *Obstetrical Journal* there is a paper by Mr. Morton in which this practice is strongly advocated, and some interesting cases are recorded in which it apparently acted well. He administers calomel in doses of 3 or 4 grains with compound extract of colocynth, so as to keep up a free action of the bowels. It seems quite reasonable, when there is constipation, to promote a gentle action of the bowels by some mild aperient; but, bearing in mind that severe and exhausting diarrhoea is a common accompaniment of the disease, I should myself hesitate to run the risk of inducing it artificially, especially as there is no proof whatever that septic matter can really be eliminated in this way. At the commencement of the disease, however, I have often given one or two aperient doses of calomel with decided benefit.

Internal Antiseptic Remedies.—It is possible that further research will give us some means of counteracting the septic state of the blood; and the sulphites and carbolates have been given for this purpose, but as yet with no reliable results.

Tincture of Perchloride of Iron.—The tincture of the perchloride of iron naturally suggests itself from its well-known effects in surgical

pyæmia. In the less intense forms of the disease, especially when local suppurations exist, it is certainly useful, and may be given in doses of 10 to 20 minims every three or four hours. In very acute cases other remedies are more reliable, and the iron has the disadvantage of not unfrequently causing nausea or vomiting.

Opiates.—When restlessness, irritation, and want of sleep are prominent symptoms, sedatives may be required. Under such circumstances opiates may be given at night, and Battley's solution, nepenthe, or the hypodermic injection of morphia are the forms which answer best.

Treatment of Local Complications.—Pain and tenderness and local complications must be treated on general principles. The distress from them is most experienced when peritonitis is well marked. Then warm and moist applications, in the form of poultices or fomentations, are very useful. Relief is also sometimes obtained from turpentine stupes, and when the tympanites is distressing turpentine enemata are very serviceable. I have found the free application over the abdomen of the flexible collodium of the Pharmacopœia decidedly useful in alleviating the suffering from peritonitis.

Such are the remedies most used in this disease. It is needless to say that it is quite impossible to lay down fixed rules for the management of any individual case; and it is obvious that if puerperal septicæmia be not a special and distinct disease, its judicious treatment must depend on the general knowledge of the attendant and on a careful study of the symptoms each separate case presents.

CHAPTER VI.

PUERPERAL VENOUS THROMBOSIS AND EMBOLISM.

Puerperal Thrombosis, and its Results.—Under the head of *thrombosis* we may class several important diseases connected with the puerperal state which have received far less attention than they deserve. It is only of late years that some—we may probably safely say the majority—of those terribly sudden deaths which from time to time occur after delivery have been traced to their true cause—viz. obstruction of the right side of the heart and pulmonary arteries from a blood-clot, either carried from a distance or, as I shall hope to show, formed *in situ*. Although the result, and, to a great extent, the symptoms, are identical in both, still a careful consideration of the history of these two classes of cases tends to show that in their causation they are distinct and that they ought not to be confounded. In the former we have primarily a clotting of blood in some part of the peripheral venous system, and the separation of a portion of such a thrombus due to changes undergone during retrograde metamorphosis tending to its eventual absorption. In the latter we have a local deposition of fibrin, the result of blood-changes

consequent on pregnancy and the puerperal state. The formation of such a coagulum in vessels the complete obstruction of which is incompatible with life explains the fatal results. When, however, a coagulum chances to be formed in more distant parts of the circulation, the vital functions are not immediately interfered with, and we have other phenomena occurring due to the obstruction. The disease known as phlegmasia dolens, I shall presently attempt to show, is one result of blood-clot forming in peripheral vessels. But from the evident and tangible symptoms it produces it has long been considered an essential and special disease, and the general blood-dyscrasia which produces it, as well as other allied states, has not been studied separately. I shall hope to show that all these various conditions, dissimilar as they at first sight appear, are very closely connected, and that they are in fact due to a common cause; and thus, I think, we shall arrive at a clearer and more correct idea of their true nature than if we looked upon them as distinct and separate affections, as has been commonly done. I am aware that in phlegmasia dolens, the pathology of which has received perhaps more study than that of almost any other puerperal affection, something beyond simple obstruction of the venous system of the affected limb is probably required to account for the peculiar tense and shining swelling which is so characteristic. Whether this be an obstruction of the lymphatics, as Dr. Tilbury Fox and others have maintained with much show of reason, or whether it is some as yet undiscovered state, further investigation is required to show. But it is beyond any doubt that the important and essential part of the disease is the presence of a thrombus in the vessels; and I think it will not be difficult to prove that in its causation and history it is precisely similar to the more serious cases in which the pulmonary arteries are involved.

It will be well to commence the study of the subject by a consideration of the conditions which in the puerperal state render the blood so peculiarly liable to coagulation, and we may then proceed to discuss the symptoms and results of the formation of coagula in various parts of the circulatory system.

Conditions which Favor Thrombosis.—The researches of Virchow, Benjamin Ball, Humphrey, Richardson, and others have rendered us tolerably familiar with the conditions which favor the coagulation of the blood in the vessels. These are, chiefly—1. A stagnant or arrested circulation; as, for example, when the blood coagulates in the veins which draw blood from the gluteal region in old and bedridden people, or as in some forms of pulmonary thrombosis, in which the clots in the arteries are probably the result of obstruction in the circulation through the lung-capillaries, as in certain cases of emphysema, pneumonia, or pulmonary apoplexy. 2. A mechanical obstruction around which coagula form, as in certain morbid states of the vessels, or, a better example still, secondary coagula which form around a travelled embolus impacted in the pulmonary arteries. 3. And, most important of all, in which the coagulation is the result of some morbid state of the blood itself. Examples of this last condition are frequently met with in the course of various diseases, such as rheumatism or fever, in which the quantity of fibrin is increased and the blood itself is loaded with morbid

material. Thrombosis from this cause is of by no means infrequent occurrence after severe surgical operations, especially such as have been attended with much hemorrhage, or when the patient is in a weak and anæmic condition. This has been specially dwelt upon as a not infrequent source of death after operation by Fayrer and other surgeons.¹

Conditions which Favor Coagulation in the Puerperal State.—But little consideration is required to show why thrombosis plays so important a part in the puerperal state, for there most of the causes favoring its occurrence are present. Probably there is no other condition in which they exist in so marked a degree or are so frequently combined. The blood contains an excess of fibrin, which largely increases in the latter months of utero-gestation, until, as has been pointed out by Andral and Gavarret, it not unfrequently contains a third more than the average amount present in the non-pregnant state. As soon as delivery is completed other causes of blood-dyscrasia come into operation. Involution of the largely hypertrophied uterus commences, and the blood is charged with a quantity of effete material, which must be present, in greater or less amount, until that process is completed. It is an old observation that phlegmasia dolens is of very common occurrence in patients who have lost much blood during labor. Thus, Dr. Leishman says: "In no class of cases has it been so frequently observed as in women whose strength has been reduced to a low ebb by hemorrhage either during or after labor; and this, no doubt, accounts for the observation made by Merriman, that it is relatively a common occurrence after placenta prævia."² An examination of the cases in which death results from pulmonary thrombosis shows the same facts, as in a large proportion of them severe post-partum hemorrhage has occurred. The exhaustion following the excessive losses so common after labor must of itself strongly predispose to thrombosis, and, indeed, loss of blood has been distinctly pointed out by Richardson to be one of its most common antecedents. "There is," he observes, "a condition which has been long known to favor coagulation and fibrinous deposition. I mean loss of blood and syncope or exhaustion during impoverished states of the body."

Since, then, so many of the predisposing causes of thrombosis are present in the puerperal state, it is hardly a matter of astonishment that it should be of frequent occurrence or that it should lead to conditions of serious gravity. And yet the attention of the profession has been for the most part limited to a study of one only of the results of this tendency to blood-clotting after delivery, no doubt because of its comparative frequency and evident symptoms. True, the balance of professional opinion has lately held that phlegmasia dolens is chiefly the result of some morbid condition of the blood, producing plugging of the veins; but the wider view which I am attempting to maintain, which would bring this disease into close relation with the more rarely observed but infinitely important obstructions of the pulmonary arteries, has scarcely, if at all, been insisted on. Doubtless, further investigation will show that it is not in these parts of the venous system alone that puerperal

¹ *Edin. Med. Journ.*, March, 1861; *Indian Annals of Med.*, July, 1867.

² Leishman, *System of Obstetrics*, p. 710.

thrombosis occurs, but the symptoms and effects of venous obstruction elsewhere, important though they may be, are unknown.

Distinction between Thrombosis and Embolism.—I propose, then, to describe the symptoms and pathology of blood-clot in the right side of the heart and pulmonary artery. It may be useful here to repeat that this is essentially distinct from embolism of the same parts. The latter is obstruction due to the impaction of a separated portion of a thrombus formed elsewhere, and for its production it is essential that thrombosis should have preceded it. Embolism is, in fact, an accident of thrombosis, not a primary affection. The condition we are now discussing I hold to be primary, precisely similar in its causation to the venous obstruction which in other situations gives rise to phlegmasia dolens.

Is Primary Thrombosis in the Pulmonary Arteries Possible?—At the threshold of this inquiry we have to meet the objection, started by several who have written on this subject,¹ that spontaneous coagulation of the blood in the right side of the heart and pulmonary arteries is a mechanical and physiological impossibility. This was the view of Virchow, who, with his followers, maintained that whenever death from pulmonary obstruction occurred an embolus was of necessity the starting-point of the malady and the nucleus round which secondary deposition of fibrin took place. Virchow holds that the primary factor in thrombosis is a stagnant state of the blood, and that the impulse imparted to the blood by the right ventricle is of itself sufficient to prevent coagulation. It is to be observed that these objections are purely theoretical. Without denying that there is considerable force in the arguments adduced, I think that the clinical history of these cases strongly favors the view of spontaneous coagulation; and I would apply to the theoretical objections advanced the argument used by one of their strongest upholders with regard to another disputed point: "*Je préfère laisser la parole aux faits, car devant eux la théorie s'incline.*"²

The anatomical arrangement of the pulmonary arteries shows how spontaneous coagulation may be favored in them; for, as Dr. Humphry has pointed out,³ "the artery breaks up at once into a number of branches, which radiate from it, at different angles, to the several parts of the lungs. Consequently, a large extent of surface is presented to the blood, and there are numerous angular projections into the currents; both which conditions are calculated to induce the spontaneous coagulation of the fibrin." We know also that thrombosis generally occurs in patients of feeble constitution often debilitated by hemorrhage, in whom the action of the heart is much weakened. These facts of themselves go far to meet the objections of those who deny the possibility of spontaneous coagulation at the roots of the pulmonary arteries.

Results of Post-mortem Examinations.—The records of post-mortem examinations show also that in many of the cases the right side of the heart, as well as the larger branches of the pulmonary arteries, contained firm, leathery, decolorized, and laminated coagula, which could not have been recently formed. The advocates of the purely embolic theory maintain that these are secondary coagula formed round an embolus.

¹ See especially Bertin, *Des Embolies*, p. 46 et seq.

² *Ibid.*, p. 149.

³ Humphry, *On the Coagulation of the Blood in the Venous System during Life*.

But surely the mechanical causes which are sufficient to prevent spontaneous deposition of fibrin would also suffice to prevent its gathering round an embolus—unless, indeed, the obstruction was sufficient to arrest the circulation altogether, when death would occur before there was any time for a secondary deposit. Before we can admit the possibility of embolism we must have at least one factor—that is, thrombosis in a peripheral vessel—from which an embolus can come. In many of the recorded cases nothing of the kind was found, and although, as is argued, this may have been overlooked, yet such an oversight can hardly always have been made.

Clinical Facts Support this View.—The strongest argument, however, in favor of the spontaneous origin of pulmonary thrombosis is one which I originally pointed out in a series of papers "On Thrombosis and Embolism of the Pulmonary Artery as a Cause of Death in the Puerperal State."¹ I there showed, from a careful analysis of 25 cases of sudden death after delivery in which accurate post-mortem examinations had been made, that cases of spontaneous thrombosis and embolism may be divided from each other by a clear line of demarcation, depending on the period after delivery at which the fatal result occurs. In 7 out of these cases there was distinct evidence of embolism, and in them death occurred at a remote period after delivery; in none before the nineteenth day. This contrasts remarkably with the cases in which the post-mortem examination afforded no evidence of embolism. These amounted to 15 out of the 25, and in all of them, with one exception, death occurred before the fourteenth day, often on the second or third. The reason of this seems to be that in the former time is required to admit of degenerative changes taking place in the deposited fibrin leading to separation of an embolus, while in the latter the thrombosis corresponds in time, and to a great extent no doubt also in cause, to the original peripheral thrombosis from which in the former the embolus was derived. Many cases I have since collected illustrate the same rule in a very curious and instructive way.

Another clinical fact I have observed points to the same conclusion. In one or two cases distinct signs of pulmonary obstruction have shown themselves without proving immediately fatal, and shortly *afterward* peripheral thrombosis, as evidenced by phlegmasia dolens of one extremity, has commenced. Here the peripheral thrombosis obviously followed the central, both being produced by identical causes, and the order of events necessary to uphold the purely embolic theory was reversed.

I hold, then, that those who deny the possibility of spontaneous coagulation in the heart and pulmonary arteries do so on insufficient grounds, and that we may consider it to be an occurrence, rare no doubt, but still sufficiently often met with, and certainly of sufficient importance, to merit very careful study.

History.—Dr. Charles D. Meigs of Philadelphia was one of the first to direct attention to spontaneous coagulation of the blood in the right side of the heart and pulmonary arteries as a cause of sudden death in the puerperal state. The occurrence itself, however, has been carefully studied by Paget, whose paper was published in 1855, four years before

¹ *Lancet*, 1867.

gs wrote on the subject.¹ It is true that none of Paget's cases happened after delivery, but he none the less clearly apprehended the nature of the obstruction. In 1855, Hecker² attributed the majority of these cases to embolism proper; and since that date most authors have taken the same view, believing that spontaneous coagulation only occurs in exceptional cases, such as those in which, on account of some obstruction in the lung or in the debility of the last few hours before death, coagula form in the smaller ramifications of the pulmonary arteries and gradually creep backward toward the heart.

Symptoms of Pulmonary Obstruction.—The symptoms can hardly be mistaken, and there seems to be no essential difference between the symptomatology of spontaneous and embolic obstructions, so that the same description will suffice for both. In a larger proportion of cases the attack comes on with an appalling suddenness which forms one of its most striking characteristics. Nothing in the condition of the patient need have given rise to the least suspicion of impending mischief, when all at once an intense and horrible dyspnoea comes on; she gasps and struggles for breadth, tears off the coverings from her chest in a vain endeavor to get more air, and often dies in a few minutes, long before medical aid can be had, with all the symptoms of asphyxia. The muscles of the face and thorax are violently agitated in the attempt to oxygenate the blood, and an appearance closely resembling an epileptic convulsion may be presented. The face may be either pale or deeply mosed. Thus in one case I have elsewhere recorded, which was an undoubted example of true embolism, Mr. Pedler, the resident accoucheur at King's College Hospital, who was present during the attack, writes of the patient:³ "She was suffering from extreme dyspnoea, the countenance was excessively pale, her lips white, the face generally expressing deep anxiety." In another, which was probably an example of spontaneous thrombosis⁴ occurring on the twelfth day after delivery, it is stated, "The face had assumed a livid purple hue, which was so remarkable as to attract the attention both of the nurse and of her mother, who was with her." The extreme embarrassment of the circulation is shown by the tumultuous and irregular action of the heart in its endeavor to send the venous blood through the obstructed pulmonary arteries. Soon it gets exhausted, as shown by its feeble and fluttering beat. The pulse is thread-like and nearly imperceptible, the respirations short and hurried, but air may be heard entering the lungs freely. The intelligence during the struggle is unimpaired, and the dreadful consciousness of impending death adds not a little to the patient's sufferings and to the terror of the scene. Such is an imperfect account of the symptoms gathered from a record of what has been observed in fatal cases. It will be readily understood why, in the presence of so sudden and awful an attack, symptoms have not been recorded with the accuracy of ordinary clinical observation.

Is Recovery Possible?—A question of great practical interest, which has been entirely overlooked by writers on the subject, is. Have we any

¹ *Medical Clinic, Trans.*, vol. xxvii. p. 162 and vol. xxviii. p. 552; *Philadelphia Medical Examiner*, 1849.

² *Brit. Med. Journ.*, March 27, 1869.

³ *Deutsche Kliniker*, 1855.

⁴ *Obst. Trans.*, vol. xii. p. 194.

ground for supposing that there is a possibility of recovery after symptoms of pulmonary obstruction have developed themselves? That such a result must be of extreme rarity is beyond question, but I have little doubt that in some few cases, entirely inexplicable on any other hypothesis, life is prolonged until the coagulum is absorbed and the pulmonary circulation restored. In order to admit of this it is of course essential that the obstruction be not sufficient to prevent the passage of a certain quantity of blood to the lungs to carry on the vital functions. The history of many cases tends to show that the obstructing clot was present for a considerable time before death, and that it was only when some sudden exertion was made, such as rising from bed or the like, calling for an increased supply of blood which could not pass through the occluded arteries, that fatal symptoms manifested themselves. This was long ago pointed out by Paget,¹ who says: "The case proves that in certain circumstances a great part of the pulmonary circulation may be arrested in the course of a week (or a few days more or less) without immediate danger to life or any indication of what had happened;" and, after referring to some illustrative cases, "Yet in all these cases the characters of the clots by which the pulmonary arteries were obstructed showed plainly that they had been a week or more in the process of formation." If we admit the possibility of the continuance of life for a certain time, we must, I think, also admit the possibility, in a few rare cases, of eventual complete recovery. What is required is time for the absorption of the clot. In the peripheral venous system coagula are constantly removed by absorption. So strong, indeed, is the tendency to this that Humphrey observes with regard to it, "It appears that the blood is almost sure to revert to its natural channel in process of time."² If, then, the obstruction be only partial, if sufficient blood pass to keep the patient alive, and a sudden supply of oxygenated blood is not demanded by any exertion which the embarrassed circulation is unable to meet, it is not inconceivable that the patient may live until the obstruction is removed.

Illustrative Cases.—Such I believe to be the only explanation of certain cases, some of which on any other hypothesis it is impossible to understand. The symptoms are precisely those of pulmonary obstructions, and the description I have given above may be applied to them in every particular; and, after repeated paroxysms, each of which seems to threaten immediate dissolution, an eventual recovery takes place. What, then, I am entitled to ask, can the condition be if not that which I suggest? As the question I am considering has never, so far as I am aware, been treated of by any other writer, I may be permitted to state, very briefly, the facts of one or two of the cases on which I found my argument, some of which I have already published in detail elsewhere:

K. H., delicate young lady. Labor easy. First child. Profuse post-partum hemorrhage. Did well until the seventh day, during the whole of which she felt weak. Same day an alarming attack of dyspnoea came on. For several days she remained in a very critical condition, the slightest exertion bringing on the attacks. A slight blowing murmur heard for a few days at the base of the heart, and then

¹ *Op. cit.*, p. 358.

² *Med.-Chir. Trans.*, vol. xxvii. p. 14.

disappeared. For two months patient remained in the same state. As long as she was in the recumbent position she felt pretty comfortable, but any attempt at sitting up in bed or any unusual exertion immediately brought on the embarrassed respiration. During all this time it was found necessary to administer stimulants profusely to ward off the attacks. Eventually the patient recovered completely.

Q. F., æt. 44. Mother of twelve children. Confined on July 6. On the eleventh day she went to bed feeling well. There was no swelling or discomfort of any kind about the lower extremities at this time. About half-past 3 A. M. she was sitting up in bed, when she was suddenly attacked with an indescribable sense of oppression in the chest, and fell back in a semi-unconscious state, gasping for breath. She remained in a very critical condition, with the same symptoms of embarrassed respiration, for three days, when they gradually passed away. Two days *after* the attack phlegmasia dolens came on, the leg swelled, and remained so for several months.

This case is an example of the fact I have already referred to, of phlegmasia dolens coming on *after* the symptoms of pulmonary obstruction had manifested themselves; the inference being that both depended on similar causes operating on two distinct parts of the circulatory system.

C. H., æt. 24. Confined of her first child on August 20, 1867. Thirty hours after delivery she complained of great weakness and dyspnoea. This was alleviated by the treatment employed, but on the ninth day, after making a sudden exertion, the dyspnoea returned with increased violence, and continued unabated until I saw the patient on September 4, fourteen days after her confinement. The following are the notes of her condition, made at the time of the visit: "I found her sitting on the sofa propped up with pillows, as she said she could not breathe in the recumbent position. The least excitement or talking brought on the most aggravated dyspnoea, which was so bad as to threaten almost instant death. Her sufferings during these paroxysms were terrible to witness. She panted and struggled for breath, and her chest heaved with short, gasping respirations. She could not even bear any one to stand in front of her, waving them away with her hand and calling for more air. These attacks were very frequent, and were brought on by the most trivial causes. She talked in a low, suppressed voice, as if she could not spare breath for articulation. On auscultation air was found to enter the lungs freely in every direction, both in front and behind. Immediately over the site of the pulmonary arteries there was a distinct harsh, rasping murmur, confined to a very limited space and not propagated either upward or downward. The heart-sounds were feeble and tumultuous." These symptoms led me to diagnose pulmonary obstruction, and I of course gave a most unfavorable prognosis, but to my great surprise the patient slowly recovered. I saw her again six weeks later, when her heart-sounds were regular and distinct and the murmur had completely disappeared.

E. E., æt. 42, was confined for the first time on November 5, 1873, in the sixth month of utero-gestation. She had severe post-partum hemorrhage, depending on partially-adherent placenta, which was removed artificially. She did perfectly well until the fourteenth day after delivery, when she was suddenly attacked with intense dyspnoea, aggravated in paroxysms. Pulse pretty full, 130, but distinctly intermittent. Air entered lungs freely. The heart's action was fluttering and irregular, and at the juncture of the fourth and fifth ribs with the sternum there was a loud blowing, systolic murmur. This was certainly non-existent before, as the heart had been carefully auscultated before administering chloroform during labor. For two days the patient remained in the same state, her death being almost momentarily expected. On the 21st—that is, two days *after* the appearance of the chest symptoms—phlegmasia dolens of a severe kind developed itself in the right thigh and leg. She continued in the same state for many days, lying more or less tranquilly, but having paroxysms of the most intense apnoea, varying from two to six or eight in the twenty-four hours. No one who saw her in one of these could have expected her to live through it. Shortly after the first appearance of the paroxysms it was observed that the cellular tissue of the neck and part of the

face became swollen and oedematous, giving an appearance not unlike that of phlegmasia dolens. The attacks were always relieved by stimulants. These she incessantly called for, declaring that she felt they kept her alive. During all this time the mind was clear and collected. The pulse varied from 110 to 130; respirations about 60; temperature 101° to 102.5° . By slow degrees the patient seemed to be rallying. The paroxysms diminished in number, and after December 1st she never had another, and the breathing became free and easy. The pulse fell to 80 and the cardiac murmur entirely disappeared. The patient remained, however, very weak and feeble, and the debility seemed to increase. Toward the second week in December she became delirious, and died, apparently exhausted, without any fresh chest symptoms, on the 19th of that month. No post-mortem examination was allowed.

I have narrated this case, although it terminated fatally, because I hold it to be one of the class I am considering. The death was certainly not due to the obstruction, all symptoms of which had disappeared, but apparently to exhaustion from the severity of the former illness. It illustrates, too, the simultaneous appearance of symptoms of pulmonary obstruction and peripheral thrombosis. The swelling of the neck was a curious symptom which has not been recorded in any other cases, and may possibly be a further proof of the analogy between this condition and phlegmasia dolens.

Such Cases can only Depend on Pulmonary Obstruction.—Now, it may, of course, be argued that these cases do not prove my thesis, inasmuch as I only assume the presence of a coagulum. But I may fairly ask in return, What other condition could possibly explain the symptoms? They are precisely those which are noticed in death from undoubted pulmonary obstruction. No one seeing one of them, or even reading an account of the symptoms, while ignorant of the result, could hesitate a single instant in the diagnosis. Surely, then, the inference is fair that they depended on the same cause. In the very nature of things my hypothesis cannot be verified by post-mortem examination, but there is at least one case on record in which, after similar symptoms, a clot was actually found. The case is related by Dr. Richardson.¹ It was that of a man who for weeks had symptoms precisely similar to those observed in the cases I have narrated. In one of his agonizing struggles for breath he died, and after death it was found "that a fibrinous band, having its hold in the ventricle, extended into the pulmonary artery." This observation proves to a certainty that life may continue for weeks after deposition of a coagulum; and, moreover, this condition was precisely what we should anticipate, since, of course, the obstructing coagulum must necessarily be small, otherwise the vital functions would be immediately arrested.

Cardiac Murmurs in Pulmonary Obstruction.—There is a symptom noted in two of the above cases, and to a less extent in a third, which has not been mentioned in any account of fatal cases occurring after delivery—viz. a murmur over the site of the pulmonary arteries. It is a sign we should naturally expect, and very possibly it would be met with in fatal cases if attention were particularly directed to the point. In both these instances it was exceedingly well marked, and in both it entirely disappeared when the symptoms abated. The probability of

¹ *Clinical Essays*, p. 224 et seq.

a murmur being audible in cases of thrombosis of the pulmonary artery has been recognized by one of our highest authorities in cardiac disease, who actually observed it in a non-puerperal case. In the last edition of his work on diseases of the heart Dr. Walshe¹ says: "The only physical condition connected with the vessel itself would probably be systolic basic murmur following the course of the pulmonary main trunk and of its immediate divisions to the left and right of the sternum. This sign I most certainly heard in an old gentleman whose life was brought to a sudden close, in the course of an acute affection, by coagulation in the pulmonary artery, and to a moderate extent in the right ventricle."

Similar cases have probably been overlooked or misinterpreted. Many seem to have been attributed to shock, in the absence of a better explanation—a condition to which they bear no kind of resemblance.

Causes of Death.—The precise mode of death in pulmonary obstruction, whether dependent on thrombosis or embolism, has given rise to considerable difference of opinion. Virchow attributes it to syncope² depending on stoppage of the cardiac contraction. Panum,³ on the other hand, contests this view, maintaining that the heart continues to beat even after all signs of life have ceased. Certainly, tumultuous and regular pulsations of the heart are prominent symptoms in most of the recorded cases, and are not reconcilable with the idea of syncope. Panum's own theory is that death is the result of cerebral anæmia. Paget seems to think that the mode of death is altogether peculiar, in some respects resembling syncope, in others anæmia. Bertin, who has discussed the subject at great length, attributes the fatal result purely to asphyxia. The condition, indeed, is in all respects similar to that state, the oxygenation of the blood being prevented, not because air cannot get to the blood, but because blood cannot get to the air. The symptoms also seem best explained by this theory: the intense dyspnoea, the terrible struggle for air, the preservation of intelligence, the tumultuous action of the heart, are certainly not characteristic either of syncope or anæmia.

Post-mortem Appearances of Clots.—The anatomical character of the clots seems to vary considerably. Ball, by whom they have been most carefully described, believes that they generally commence in the smaller ramifications of the arteries, extending backward toward the heart and filling the vessels more or less completely. Toward its cardiac extremity the coagulum terminates in a rounded head, in which respect it resembles those spontaneously formed in the peripheral veins. It is non-adherent to the coats of the vessels, and the blood circulates, when it can do so at all, between it and the vascular walls. Such clots are white, dense, and of a homogeneous structure, consisting of layers of decolorized fibrin, firm at the periphery, where the fibrin has been most recently deposited, and softened in the centre, where amylaceous or fatty degeneration has commenced. Ball maintains that if the coagulum have commenced in the larger branches of the arteries, it must have first begun in the ventricle and extended into them. According to Humphrey, the

¹ Walshe, *On Diseases of the Heart*, 4th ed. 1873.

² *Gesamte Anatomie*, 1862, p. 316.

³ *Virchow's Archiv*, 1863.

same changes take place in pulmonary as in peripheral thrombi, and they may become adherent to the walls of the vessels or converted into threads or bands. When the obstruction is due to embolism, provided the case is a well-marked one and the embolus of some size, the appearances presented are different. We have no longer a laminated and decolorized coagulum with a rounded head, similar to a peripheral thrombus. The obstruction in this case generally takes place at the point of bifurcation of the artery, and we there meet with a grayish-white mass, contrasting remarkably with the more recently-deposited fibrin before and behind it. It may be that the form of the embolus shows that it has recently been separated from a clot elsewhere; and in many cases it has been possible to fit the travelled portion to the extremity of the clot from which it has been broken. We may also, perhaps, find that the embolus has undergone an amount of retrograde metamorphosis corresponding with that of the peripheral thrombus from which we suppose it to have come, but differing from that of the more recently-deposited fibrin around it. It must be admitted, however, that the anatomical peculiarities of the coagula will by no means always enable us to trace them to their true origin. In many cases emboli may escape detection from their smallness or from the quantity of fibrin surrounding them.

Treatment.—But few words need be said as to the treatment of pulmonary obstruction. In a large majority of cases the fatal result so rapidly follows the appearance of the symptoms that no time is given us even to make an attempt to alleviate the patient's sufferings. Should we meet with a case not immediately fatal, it seems that there are but two indications of treatment affording the slightest rational ground of hope :

1. To keep the patient alive by the administration of stimulants—brandy, ether, ammonia, and the like—to be repeated at intervals corresponding to the intensity of the paroxysms and the results produced. In the cases I have above narrated in which recovery ensued this took the place of all other medication. Possibly leeches or dry cupping to the chest might prove of some service in relieving the circulation.

2. To enjoin the most absolute and complete repose. The object of this is evident. The only chance for the patient seems to be that the vital functions should be carried on until the coagulum has been absorbed, or at least until it has been so much lessened in size as to admit of blood passing it to the lungs. The slightest movements may give rise to a fatal paroxysm of dyspnoea from the increased supply of oxygenated blood required. It must not be forgotten that in a large proportion of cases death immediately followed some exertion in itself trivial, such as rising out of bed. Too much attention, then, cannot be given to this point. The patient should be absolutely still; she should be fed with abundance of fluid food, such as milk, strong soups, and the like, and should on no account be permitted to raise herself in bed or attempt the slightest muscular exertion. If we are fortunate enough to meet with a case apparently tending to recovery, these precautions must be carried on long after the severity of the symptoms has lessened, for a moment's imprudence may suffice to bring them back in all their original intensity.

Bertin,¹ indeed, recommends a system of treatment very different from this. In the vain hope that the violent effort induced may cause the displacement of the impacted embolus (to which alone he attributes pulmonary obstruction) he recommends the administration of emetics. Few, I fancy, will be found bold enough to attempt so hazardous a plan of treatment.

Various drugs have been suggested in these cases. Richardson² recommended ammonia, a deficiency of which he at that time believed to be the chief cause of coagulation. He has since advised that liquor ammoniæ should be given in large doses, 20 minims every hour, in the hope of causing solution of the deposited fibrin; and he has stated that he has seen good results from the practice. Others advise the administration of alkalies, in the hope that they may favor absorption. The best that can be said for them is that they are not likely to do much harm.

Puerperal Pleuro-pneumonia.—This is perhaps the best place to mention an important but little-understood class of cases which I believe to be less uncommon than is generally supposed. I refer to severe pleuro-pneumonia occurring in connection with the puerperal state, but not distinctly associated with septicæmia. Two carefully-observed cases of this kind are recorded by MacDonald, occurring in his practice; I myself have met with three very marked examples within the past three years, one of which proved fatal, the other two giving rise to most serious illness, from which the patient recovered with difficulty.

Peculiarities of these Cases.—So far as my own observation goes, there are marked peculiarities in such cases which clearly differentiate them from the ordinary course of pneumonia. The onset is sudden and unconnected with exposure to cold or other cause of lung disease; there is no definite crisis, but a continuous pyrexia, of moderate intensity, lasting a variable time; and the physical signs differ from those of ordinary pneumonia.

Physical Signs.—In MacDonald's case, as well as in my own, they were peculiar in this respect, that there was very slight crepitation, marked rusty sputum, and a wooden dulness, much more intense than in ordinary pneumonia, extending over a large lung-space, with a very slight entrance of air into the lung-tissue. It is also remarkable that a very large proportion of the cases was associated with phlegmasia dolens. Thus, it existed in one of MacDonald's two cases, and in two out of my own three. Like phlegmasia dolens, moreover, the disease generally commenced some weeks after delivery; my own cases, for example, occurred respectively fifteen, twenty-eight, and thirty-five days after labor. It is difficult to believe that there is not some connection between these two conditions, and there is much in their peculiar history to lead to the belief that such forms of lung disease depend, in fact, on thrombotic or embolic obstruction of the minute branches of the pulmonary arteries, caused by conditions similar to those which have produced the phlebotic obstructions in the lower extremities. In the absence of careful post-mortem examination this hypothesis is clearly not susceptible of proof. MacDonald, while admitting that "a limited thrombosis of the pulmonary arteries would no doubt explain the facts of the cases,"

¹ *Op. cit.*, p. 393.

² *Heart Disease during Pregnancy*, p. 209.

is rather inclined to "seek the chief explanation of their occurrence in the alterations which the pregnant and puerperal conditions impress upon the blood and the blood-vascular system."

I confess that, to my mind, the former hypothesis is not only the most definite, but the one which most readily explains all the peculiarities of these cases. I cannot, however, do more than suggest it, in the hope that further observations, and especially carefully-conducted autopsies, may throw some light on this obscure and little-studied subject.

Treatment.—As regards treatment, it is obvious that it must be conducted on general principles, carefully avoiding over-severe measures, and supporting the patient through a trial to the system that must necessarily be severe.

CHAPTER VII.

PUERPERAL ARTERIAL THROMBOSIS AND EMBOLISM.

THE same condition of the blood which so strongly predisposes to coagulation in the vessels through which venous blood circulates tends to similar results in the arterial system. These, however, are by no means so common, and do not, as a rule, lead to such important consequences. The subject has been but little studied, and almost all our knowledge of it is derived from a very interesting essay by Sir James Simpson.¹ As I have devoted so much space to the consideration of venous thrombosis and embolism, I shall but briefly consider the effects of arterial obstruction.

Causes.—In a considerable number of recorded cases the obstruction has resulted from the detachment of vegetations deposited on the cardiac valves, the result of endocarditis, either produced by antecedent rheumatism or as a complication of the puerperal state. Sometimes the obstruction seems to depend on some general blood-dyscrasia, similar to that producing venous thrombosis, or on some local change in the artery itself. Thus, Simpson records a case apparently produced by local arteritis, which caused acute gangrene of both local extremities, ending fatally in the third week after delivery. In other cases it has been attributed to coagulation following spontaneous laceration and corrugation of the internal coat of the artery.

Symptoms.—The symptoms of puerperal arterial obstruction must of course vary with the particular arteries affected. Those with the obstruction of which we are most familiar are the cerebral, the brachial, and the femoral. The effects produced must also be modified by the size of the embolus and the more or less complete obstruction it produces. Thus, for example, if the middle cerebral artery be blocked up entirely, the functions of those portions of the brain supplied by it will be more or less completely arrested, and hemiplegia of the opposite side

¹ *Selected Obst. Works*, vol. i. p. 523.

of the body, followed by softening of the brain-texture, will probably result. If the nervous symptoms be developed gradually or increase in intensity after their first appearance, it may be that an obstruction, at first incomplete, has increased by the deposition of fibrin around it. So the occasional sudden supervention of blindness, with destruction of the eyeball—cases of which are recorded by Simpson—not improbably depends on the occlusion of the ophthalmic artery, the function of the organ depending on its supply through the single artery. The effects of obstruction of the visceral arteries in the puerperal state are entirely unknown, but it is far from unlikely that further investigation may prove them to be of great importance. In the extremities arterial obstruction produces effects which are well marked. They are classified by Simpson under the following heads: 1. *Arrest of pulse below the site of obstruction.*—This has been observed to come on either suddenly or gradually, and if the occlusion be in one of the large arterial trunks, it is a symptom which a careful examination will readily enable us to detect. 2. *Increased force of pulsation in the arteries above the seat of obstruction.* 3. *Fall in the temperature of the limb.*—This is a symptom which is easily appreciable by the thermometer, and when the main artery of the limb is occluded the coldness of the extremity is well marked. 4. *Lesions of motor and sensory functions, paralysis, neuralgia, etc. etc.*—Loss of power in the affected limb is often a prominent symptom, and when it comes on suddenly, and is complete, the main artery will probably be occluded. It may be diagnosed from paralysis depending on cerebral or spinal causes by the absence of head-symptoms, by the history of the attack, and by the presence of other indications of arterial obstruction, such as loss of pulsation in the artery, fall of temperature, etc. The sensory functions in these cases are generally also seriously disturbed—not so much by loss of sensation as by severe pain and neuralgia. Sometimes the pain has been excessive, and occasionally it has been the first symptom which directed attention to the state of the limb. 5. *Gangrene below or beyond the seat of arterial obstruction.*—Several interesting cases are recorded in which gangrene has followed arterial obstruction. Generally speaking, gangrene will not follow occlusion of the main arterial trunk of an extremity, as the collateral circulation becomes soon sufficiently developed to maintain its vitality. In many of the cases either thrombi have obstructed the channels of collateral circulation as well, or the veins of the limb have also been blocked up. When such extensive obstructions occur, they obviously cannot be embolic, but must depend on a local thrombosis traceable to some general blood-dyscrasia depending on the puerperal state.

Treatment.—Little can be said of such cases as to the treatment, which must vary with the gravity and nature of the symptoms in each. Beyond absolute rest (in the hope of eventual absorption of the thrombus or embolus), generous diet, attention to the general health of the patient, and sedative applications to relieve the local pain, there is little in our power. Should gangrene of an extremity supervene in a puerperal patient, the case must necessarily be wellnigh hopeless. Simpson, however, records one instance in which amputation was performed above the line of demarcation, the patient eventually recovering.

CHAPTER VIII.

OTHER CAUSES OF SUDDEN DEATH DURING LABOR AND THE PUERPERAL STATE.

Various Causes of Sudden Death.—A large number of the cases in which sudden death occurs during or after delivery find their explanation, as I have already pointed out, in thrombosis or embolism of the heart and pulmonary arteries. Probably many cases of the so-called *idiopathic asphyxia* were, in fact, examples of this accident, the true nature of which had been misunderstood. Besides these, there are no doubt many other conditions which may lead to a suddenly fatal result in connection with parturition.

Some of these are of an organic, others of a functional, nature.

Organic Causes.—Among the former may be mentioned cases in which the straining efforts of the second stage of labor have produced death in patients suffering from some pre-existent disease of the heart. Rupture of that organ has probably occurred from fatty degeneration of its walls. Dehous¹ narrates an instance in which the efforts of labor caused the rupture of an aneurism. Another case, from interference with the action of the heart in a patient who had pericardial effusion, is narrated by Ramsbotham. Dr. Devilliers relates an instance occurring in a young woman during the second stage of labor. The heart was found to be healthy, but the lungs were intensely congested and blood was extensively extravasated all through their texture. This was probably caused by pulmonary congestion and apoplexy, produced by the severe straining efforts. Many cases from effusion of blood into the brain-substance or on its surface are on record, no doubt in patients who, from arterial degeneration or other causes, were predisposed to apoplectic effusions. The so-called apoplectic convulsions, formerly described in most works on obstetrics as a variety of puerperal convulsions, are evidently nothing more than apoplexy coming on during or after labor. As regards their pathology, they do not seem to differ from ordinary cases of apoplexy in the non-pregnant condition. One example is recorded of death which was attributed to rupture of the diaphragm from excessive action in the second stage.

Functional Causes.—Among the causes of death which cannot be traced to some distinct organic lesion may be classed cases of syncope, shock, and exhaustion. Many instances of this kind are recorded. Thus, in some women of susceptible nervous organization the severity of the suffering appears to bring on a condition similar to that produced by excessive shock or exhaustion, which has not unfrequently proved fatal. Several examples of this kind have been cited by McClintock.² It is also not unlikely that sudden syncope sometimes produces a fatal result during or after labor. Most cases of death otherwise inexplicable used

¹ Dehous, *Sur les Morts subites*.

² *Union méd.*, 1853.

to be referred to this cause; but accurate autopsies were seldom made, and even when they were—the important effects of pulmonary coagula being unknown—it is more than probable that the true cause of death was overlooked. It has been supposed that the sudden removal of pressure from the veins of the abdomen by the emptying of the gravid uterus after delivery may favor an increased afflux of blood into the lower parts of the body, and thus tend to an anæmic condition of the brain and the production of syncope. However this may be, the possibility of its occurrence, and its manifest danger in a recently-delivered woman, are sufficient reasons for enforcing the recumbent position after labor is over. In some of the cases the syncope was evidently produced by the patient's suddenly sitting upright.

Death from Air in the Veins.—Some cases of sudden death immediately after labor seem to be due to the entrance of air into the veins. Six examples are cited by McClintock which were probably due to this cause. La Chapelle relates two. An interesting case is related by M. Lionet.¹ In this the patient died five and a half hours after an easy and natural labor, the chief symptoms being extreme pallor, efforts at vomiting, and dyspnœa. Air was found in the heart and in the arachnoid veins. There can be no question that the uterine sinuses after delivery are nearly as well adapted as the veins of the neck for allowing the entrance of air. They are firmly attached to the muscular walls of the uterus, so that they gape open when that organ is relaxed, and it is easy to understand how air might enter. Indeed, in the post-mortem examination in one of the cases occurring in the practice of Mme. La Chapelle it is stated that “the uterine sinuses opened in the interior of the uterus by large orifices (one line and a half in diameter), through which air could readily be blown as far as the iliac veins, and *vice versâ*.” The condition of the uterus after delivery also enables the air to have ready access to the mouths of the sinuses, for the alternate relaxation and contraction of the uterus occurring after the placenta is expelled would tend to draw in the air as by a suction-pump. Hence an additional reason for insisting on firm contraction of the uterus, as this will lessen the risk of this accident.

Cause of Death in such Cases.—The precise mechanism of death from air in the veins has been a subject of dispute among pathologists. By Bichat² it was referred to anæmia and syncope from want of blood in the vessels of the brain, which are occupied by air. Nysten³ attributed it to distension of the cavities of the heart by rarefied air, producing paralysis of its wall; Leroy, to a stoppage of the pulmonary circulation, and consequent want of proper blood-supply to the left heart; while Leroy d'Etoilles thought it might depend on any of these causes or a combination of all of them. These and many other hypotheses on the subject have been advanced, to all of which serious objection could be raised. The most recent theory is one maintained by Virchow and Oppolzer,⁴ and more recently by Feltz, which attributes the fatal results

¹ Dehous, *op. cit.*, p. 58.

² *Recherches sur la Vie et la Mort*, 1853.

³ Nysten, *Recherches de Phys. et Chem. Path.*, 1811.

⁴ *Claustiques des Embolies*; *Wiener Med. Woch.*, 1863; *Des Embolies capillaires*, 1868; and *op. cit.*, p. 115.

to impaction of the air-globules in the lesser divisions of the pulmonary arteries, where they form gaseous emboli, and cause death exactly in the same way as when the obstruction depends on a fibrinous embolus. The symptoms observed in fatal cases closely correspond to those of pulmonary obstruction, and it is not unlikely that some cases attributed to other causes may really depend on the entrance of air through the uterine sinuses. Such, for example, was most probably the explanation of a case referred to by Dr. Graily Hewitt in a discussion at the Obstetrical Society.¹ Death occurred shortly after the removal of an adherent placenta, during which, no doubt, air could readily enter the uterine cavity. The symptoms—viz. “severe pain in the cardiac region, distress as regards respiration, and pulselessness”—are identical with those of pulmonary obstruction. Dr. Hewitt refers the death to shock, which certainly does not generally produce such phenomena.

CHAPTER IX.

PERIPHERAL VENOUS THROMBOSIS (SYNS.: CRURAL PHLEBITIS, PHLEGMASIA DOLENS; ANASARCA SEROSA; OEDEMA LACTEUM; MILK LEG, ETC.).

Peripheral Thrombosis.—We now come to discuss the symptoms and pathology of the conditions associated with the formation of thrombi in the peripheral venous system, or rather in the veins of the lower extremities, since too little is known of their occurrence in other parts to enable us to say anything on the subject.

The most important of these is the well-known disease which under the name *phlegmasia dolens* has attracted much attention and given rise to numerous theories as to its nature and pathology. In describing it as a local manifestation of a general blood-dyscrasia, and not as an essential local disease, I am making an assumption as to its pathology that many eminent authorities would not consider justifiable. I have, however, already stated some of the reasons for so doing, and I shall shortly hope to show that this view is not incompatible with the most probable explanation of the peculiar state of the affected limb.

Symptoms.—The first symptom which usually attracts attention is severe pain in some part of the limb that is about to be affected. The character of the pain varies in different cases. In some it is extremely acute, and is most felt in the neighborhood and along the course of the chief venous trunks. It may begin in the groin or hip and extend downward, or it may commence in the calf and proceed upward toward the pelvis. The pain abates somewhat after swelling of the limb (which generally begins within twenty-four hours), but it is always a distressing symptom, and continues as long as the acute stage of the disease lasts.

¹ *Obstet. Trans.*, vol. x. p. 28.

restlessness, want of sleep, and suffering which it produces are sometimes excessive. Coincident with the pain, and sometimes preceding it, more or less *malaise* is experienced. The patient may for a day or two be restless, irritable, and out of sorts without any very definite cause, or the disease may be ushered in by a distinct rigor. Generally, there is constitutional disturbance, varying with the intensity of the case. The pulse is rapid and weak, 120 or thereabouts; the temperature elevated from 101° to 102° , with an evening exacerbation. The patient is thirsty, the tongue glazed or white and loaded, the bowels are constipated. In some few cases, when the local affection is slight, none of these constitutional symptoms are observed.

Condition of the Affected Limb.—The characteristic swelling rapidly follows the commencement of the symptoms. It generally begins in the groin, whence it extends downward. It may be limited to the thigh, or the whole limb, even to the feet, may be implicated. More rarely it commences in the calf of the leg, extending upward to the thigh and downward to the feet. The affected parts have a peculiar appearance which is pathognomonic of the disease. They are hard, tense, and brawny, of a shiny white color, and not yielding on pressure, except toward the beginning and end of the illness. The appearances presented are quite different from those of ordinary œdema. When the whole thigh is affected the limb is enormously increased in size. Frequently the venous trunks, especially the femoral and popliteal veins, are felt obstructed with coagula, and rolling under the finger. They are painful when handled, and in their course more or less redness is occasionally observed. Either leg may be attacked, but the left more frequently than the right. There is a marked tendency for the disease to spread, and we often find, in a case which is progressing apparently well, a rise of temperature and an accession of febrile symptoms followed by the swelling of the other limb.

Progress of the Disease.—After the acute stage has lasted from a week to a fortnight the constitutional disturbance becomes less marked, the pulse and temperature fall, the pain abates, and the sleeplessness and restlessness are less. The swelling and tension of the limb now begin to diminish, and absorption commences. This is invariably a slow process. It is always many weeks before the effusion has disappeared, and it may be many months. The limb retains for a length of time the peculiar *wooden* feeling, as Dr. Churchill terms it. Any imprudence, such as a too early attempt at walking, may bring on a relapse and fresh swelling of the limb. This gradual recovery is by far the most common termination of the disease. In some rare cases suppuration may take place, either in the subcutaneous cellular tissue, the lymphatic glands, or even in the joints, and death may result from exhaustion. The possibility of pulmonary obstruction and sudden death from separation of an embolus have already been pointed out: and the fact that this lamentable occurrence has generally followed some undue exertion should be borne in mind as a guide in the management of our patient.

Period of Commencement.—The disease usually begins within a short time after delivery, rarely before the second week. In 22 cases tabulated by Dr. Robert Lee, 7 were attacked between the fourth and twelfth

days, and 14 after the second week. Some cases have been described as commencing even months after delivery. It is questionable if these can be classed as puerperal, for it must not be forgotten that phlegmasia dolens is by no means necessarily a puerperal disease [or confined to the female sex]. There are many other conditions which may give rise to it, all of them, however, such as produce a septic and hyperinosed state of the blood, such as malignant disease, dysentery, phthisis, and the like. My own experience would lead me to think that cases of this kind are much more common than is generally believed.

History and Pathology.—The disease has long attracted the attention of the profession. Passing over more or less obscure notices by Hippocrates, De Castro, and others, we find the first clear account in the writings of Mauriceau, who not only gave a very accurate description of its symptoms, but made a guess at its pathology which was certainly more happy than the speculations of his successors: it is, he says, caused “by a reflux on the parts of certain humors which ought to have been evacuated by the lochia.” Puzos ascribed it to the arrest of the secretion of milk and its extravasation in the affected limb. This theory, adopted by Levret and many subsequent writers, took a strong hold on both professional and public opinion, and to it we owe many of the names by which the disease is known to this day, such as *œdema lacteum*, milk leg, etc. In 1784, Mr. White of Manchester attributed it to some morbid condition of the lymphatic glands and vessels of the affected parts; and this or some analogous theory, such as that of rupture of the lymphatics crossing the pelvic brim, as maintained by Tyre of Gloucester, or general inflammation of the absorbents, as held by Dr. Ferrier, was generally adopted.

Phlebitic Theory.—It was not until the year 1823 that attention was drawn to the condition of the veins. To Bouillaud belongs the undoubted merit of first pointing out that the veins of the affected limb were blocked up by coagula, although the fact had been previously observed by Dr. Davis of University College. Dr. Davis made dissections of the veins in a fatal case, and found, as Bouillaud had done, that they were filled with coagula, which he assumed to be the results of inflammation of their coats; hence the name of “*crural phlebitis*” which has been extensively adopted instead of *phlegmasia dolens*. Dr. Robert Lee did much to favor this view, and, finding that thrombi were present in the iliac and uterine as well as in the femoral veins, he concluded that the phlebitis commenced in the uterine branches of the hypogastric veins and extended downward to the femorals. He pointed out that *phlegmasia dolens* was not limited to the puerperal state, but that, when it did occur independently of it, other causes of uterine phlebitis were present, such as cancer of the os and cervix uteri. The inflammatory theory was pretty generally received, and even now is considered by many to be a sufficient explanation of the disease. Indeed, the fact that more or less thrombosis was always present could not be denied; and on the supposition that thrombosis could only be caused by phlebitis, as was long supposed to be the case, the inflammatory theory was the natural one. Before long, however, pathologists pointed out that thrombosis was by no means necessarily, or even generally, the result of inflamma-

tion of the vessels in which the clot was contained, but that the inflammation was more generally the result of the coagulum.

Theory of its Dependence on Septic Causes.—The late Dr. Mackenzie took a prominent part in opposing the phlebitic theory. He proved by numerous experiments on the lower animals that inflammation is not sufficient of itself to produce the extensive thrombi which are found to exist, and that inflammation originating in one part of a vein is not apt to spread along its canal, as the phlebitic theory assumes. His conclusion is, that the origin of the disease is rather to be sought in some septic or altered condition of the blood producing coagulation in the veins. Dr. Tyler Smith¹ pointed out an occasional analogy between the causes of phlegmasia dolens and puerperal fever, evidently recognizing the dependence of the former on blood-dyscrasia. "I believe," he says, "that contagion and infection play a very important part in the production of the disease. I look on a woman attacked with phlegmasia dolens as having made a fortunate escape from the greater dangers of diffuse phlebitis or puerperal fever." In illustration of this he narrates the following instructive history: "A short time ago a friend of mine had been in close attendance on a patient dying of erysipelatous sore throat with sloughing, and was himself affected with sore throat. Under these circumstances he attended, within the space of twenty-four hours, three ladies in their confinements, all of whom were attacked with phlegmasia dolens."

View of Tilbury Fox.—The latest important contribution to the pathology of the disease is contained in two papers by Dr. Tilbury Fox, published in the second volume of the *Obstetrical Transactions*. He maintained that something beyond the mere presence of coagula in the veins is required to produce the phenomena of the disease, although he admitted that to be an important, and even an essential, part of the pathological changes present. The thrombi he believed to be produced either by extrinsic or intrinsic causes—the former comprising all cases of pressure by tumor or the like; the latter, and the most important, being divisible into the heads of—

1. True inflammatory changes in the vessels, as seen in the epidemic form of the disease.

2. Simple thrombus, produced by rapid absorption of morbid fluid.

3. Virus action and thrombus conjoined, the phlegmasia dolens itself being the result of simple thrombus, and not produced by diseased (inflamed) coats of vessels; the general symptoms the result of the general blood-state.

He further pointed out that the peculiar swelling of the limbs cannot be explained by the mere presence of œdema, from which it is essentially different; the white appearance of the skin, the severe neuralgic pain, and the persistent numbness indicating that the whole of the cutaneous textures, the cutis vera, and even the epithelial layer, are infiltrated with fibrinous deposit. He concluded, therefore, that the swelling is the result of œdema *plus* something else, that something being obstruction of the lymphatics, by which the absorption of effused serum is prevented. The efficient cause which produces these changes he believes to be, in the

¹ Tyler Smith, *Manual of Obstetrics*, p. 538.

majority of cases, a septic action originating in the uterus, producing a condition similar to that in which phlegmasia dolens arises in the non-puerperal state.

There is no doubt much force in Dr. Fox's arguments, and it may, I think, be conceded that obstruction of the veins, *per se*, is not sufficient to produce the peculiar appearance of the limb. It is, moreover, certain that phlebitis alone is also an insufficient explanation not only of the symptoms, but even of the presence of thrombi so extensive as those that are found. The view which traces the disease solely to inflammation or obstruction of lymphatics is purely theoretical, has no basis of facts to support it, and finds now-a-days no supporters. The experiments of Mackenzie and Lee, as well as the vastly increased knowledge of the causes of thrombosis which the researches of modern pathologists have given us, seem to point strongly to the view already stated, that the disease can only be explained by a general blood-dyscrasia depending on the puerperal state. It by no means follows that we are to consider Dr. Fox's speculations as incorrect. It is far from improbable that the lymphatic vessels are implicated in the production of the peculiar swelling, only we are not as yet in a position to prove it. There is no inherent improbability in the supposition that the same morbid state of the blood which produces thrombosis in the veins may also give rise to such an amount of irritation in the lymphatics as may interfere with their functions, and even obstruct them altogether. The essential and all-important point in the pathology of the disease, however, seems undoubtedly to be thrombosis in the veins; and the probability of there being some as yet undetermined pathological changes in addition to this by no means militates against the view I have taken of the intimate connection of the disease with other results of thrombosis in different vessels.

Changes occurring in the Thrombi.—The changes which take place in the thrombi all tend to their ultimate absorption. These have been described by various authors as leading to organization or suppuration. It is probable, however, that the appearances which have led to such a supposition are fallacious, and that they are really due to retrograde metamorphosis of the fibrin, generally of an amylaceous or fatty character.

Detachment of Emboli.—The peculiarities of a clot that most favor detachment of an embolus are such a shape as admits of a portion floating freely in the blood-current, by the force of which it is detached and carried to its ultimate destination. When the accident has occurred it is often possible to recognize the peripheral thrombus from which the embolus has separated, by the fact of its terminal extremity presenting a freshly-fractured end, instead of the rounded head natural to it. Such detachment is unlikely to occur, even when favored by the shape of the clot, unless sufficient time have elapsed after its formation to admit of its softening and becoming brittle. The curious fact I have before mentioned, of true puerperal embolism occurring in the large majority of cases only after the nineteenth day from delivery, finds a ready explanation in this theory, which it remarkably corroborates.

Treatment.—On the supposition that phlegmasia dolens was the result

of inflammation of the veins of the affected limb, an antiphlogistic course of treatment was naturally adopted. Accordingly, most writers on the subject recommend depletion, generally by the application of leeches along the course of the affected vessels. We are told that if the pain continue the leeches should be applied a second or even a third time. If we admit the septic origin of the disease, we must, I think, see the impropriety of such a practice. The fact that it occurs in a large majority of cases in patients of a weakly and debilitated constitution, often in women who have suffered from hemorrhage, is a further reason for not adopting this routine custom. If local loss of blood be used at all, it should be strictly limited to cases in which there is much tenderness and redness across the course of the veins, and then only in patients of plethoric habits and strong constitution. Cases of this kind will form a very small minority of those coming under our observation.

Over-active Treatment Unadvisable.—What has been said of the pathology of the affection tends to the conclusion that active treatment of any kind in the hope of curing the disease is likely to be useless. Our chief reliance must be on time and perfect rest, in order to admit of the thrombi and the secondary effusion being absorbed, while we relieve the pain and other prominent symptoms and support the strength and improve the constitution of the patient.

Relief of Pain, etc.—The constant application of heat and moisture to the affected limb will do much to lessen the tension and pain. Wrapping the entire limb in linseed-meal poultices, frequently changed, is one of the best means of meeting this indication. If, as is sometimes the case, the weight of the poultices be too great to be readily borne, we may substitute warm flannel stupes covered with oiled silk. Local anodyne applications afford much relief, and may be advantageously used along with the poultices and stupes, either by sprinkling their surface freely with laudanum or chloroform and belladonna liniment, or by soaking the flannels in poppy-head fomentations. It is needless to say that the most absolute rest in bed should be enjoined, even in slight cases, and that the limb should be effectually guarded from undue pressure by a cradle or some similar contrivance. Local counter-irritation has been strongly recommended, and frequent blisters have been considered by some to be almost specific. I should myself hesitate to use blisters, as they certainly would not be soothing applications, and one hardly sees how they can be of much service in hastening the absorption of the effusion.

Constitutional Treatment.—During the acute stage of the disease the constitutional treatment must be regulated by the condition of the patient. Light but nutritious diet must be administered in abundance, such as milk, beef-tea, and soups. Should there be much debility, stimulants in moderation may prove of service. With regard to medicines, we shall probably find benefit from such as are calculated to improve the condition of the blood and the general health of the patient. Chlorate of potash, with diluted hydrochloric acid, quinine, either alone or in combination with sesquicarbonate of ammonia, the tincture of the perchloride of iron, are the drugs that are most likely to prove of service. Alkalies and other medicines, which have been recommended in

the hope of hastening the absorption of coagula, must be considered as altogether useless. Pain must be relieved and sleep procured by the judicious use of anodynes, such as Dover's powder, the subcutaneous injection of morphia, or chloral. Generally, no form answers so well as the hypodermic injection of morphia.

Subsequent Local Treatment.—When the acute symptoms have abated and the temperature has fallen, the poultices and stupes may be discontinued and the limb swathed in a flannel roller from the toes upward. The equable pressure and support thus afforded materially aid the absorption of the effusion and tend to diminish the size of the limb. At a still later stage very gentle inunctions of weak iodine ointment may be used with advantage once a day before the roller is applied. Shampooing and friction of the limb, generally recommended for the purpose of hastening absorption, should be carefully avoided, on account of the possible risk of detaching a portion of the coagulum and producing embolism. This is no merely imaginary danger, as the following fact narrated by Trousseau proves: "A phlegmasia alba dolens had appeared on the left side in a young woman suffering from peri-uterine phlegmon. The pain having ceased, a thickened venous trunk was felt on the upper and internal part of the thigh. Rather strong pressure was being made, when M. Demarquay felt something yield under his fingers. A few minutes afterward the patient was attacked with dreadful palpitation, tumultuous cardiac action, and extreme pallor, and death was believed to be imminent. After some hours, however, the oppression ceased, and the patient eventually recovered. A slightly-attached coagulum must have become separated and conveyed to the heart or pulmonary artery."¹ Warm douches of water—of salt water, if it can be obtained—may be advantageously used in the later stages of the disease, and they may be applied night and morning, the limb being bandaged in the interval. The occasional use of the electric current is said to promote absorption, and it would seem likely to be a serviceable remedy.

Change of Air, etc.—When the patient is well enough to be moved a change of air to the seaside will be of value. Great caution, however, should be recommended in using the limb, and it is far better not to run the risk of a relapse by any undue haste in this respect. It is well to warn the patient and her friends that a considerable time must of necessity elapse before the local signs of the disease have completely disappeared.

¹ Trousseau, "Clinique de l'Hôtel Dieu," in *Gaz. des Hôp.*, 1860, p. 577.

CHAPTER X.

PELVIC CELLULITIS AND PELVIC PERITONITIS.

FROM the earliest time the occurrence after parturition of severe forms of inflammatory disease in and about the pelvis, frequently ending in suppuration, has been well known. It is only of late years, however, that these diseases have been made the subject of accurate clinical and pathological investigation, and that their true nature has begun to be understood. Nor is our knowledge of them as yet by any means complete. They merit careful study on the part of the accoucheur, for they give rise to some of the most severe and protracted illnesses from which puerperal patients suffer. They are often obscure in their origin and apt to be overlooked, and they not rarely leave behind them lasting mischief.

These diseases are not limited to the puerperal state. On the contrary, many of the severest cases arise from causes altogether unconnected with childbearing. These will not be now considered, and this chapter deals solely with such forms as may be directly traced to childbirth.

Two Distinct Forms.—Modern researches have demonstrated that there are two distinct varieties of inflammatory disease met with after labor, which differ materially from each other in many respects. In one of these the inflammation affects chiefly the connective tissue surrounding the generative organs contained within the pelvis, or extends up from beneath the peritoneum and into the iliac fossæ. In the other it attacks that portion of the peritoneum which covers the pelvic viscera, and is limited to it.

So much is admitted by all writers, but great obscurity in description, and consequent difficulty in understanding satisfactorily the nature of these affections, have resulted from the variety of nomenclature which different authors have adopted.

Thus, the former disease has been variously described as pelvic cellulitis, peri-uterine phlegmon, parametritis, or pelvic abscess; while the latter is not unfrequently called perimetritis, as contradistinguished from parametritis. The use of the prefix *para* or *peri* to distinguish the cellular or peritoneal variety of inflammation, originally suggested by Virchow, has been pretty generally adopted in Germany, and has been strongly advocated in this country by Matthews Duncan. It has never, however, found much favor with English writers, and the similarity of the two names is so great as to lead to confusion. I have therefore selected the terms "*pelvic peritonitis*" and "*pelvic cellulitis*," as conveying in themselves a fairly accurate notion of the tissues mainly involved.

Importance of Distinguishing the Two Classes of Cases.—The important fact to remember is that there exist two distinct varieties of inflammatory disease, presenting many similarities in their course, symptoms, and results, often occurring simultaneously, but in the main distinct in their pathology and capable of being differentiated. Thomas compares

them—and, as serving to fix the facts on the memory, the illustration is a good one—to pleurisy and pneumonia. “Like them,” he says, “they are separate and distinct, like them affect different kinds of structure, and like them they generally complicate each other.” It might therefore be advisable, as most writers on the disease occurring in the non-puerperal state have done, to treat of them in two separate chapters. There is, however, more difficulty in distinguishing them as puerperal than as non-puerperal affections, for which reason, as well as for the sake of brevity, I think it better to consider them together, pointing out as I proceed the distinctive peculiarities of each.

Seat of Disease.—When attention was first directed to this class of diseases the pelvic cellular tissue was believed to be the only structure affected. This was the view maintained by Nonat, Simpson, and many modern writers. Attention was first prominently directed to the importance of localized inflammation of the peritoneum, and to the fact that many of the supposed cases of cellulitis were really peritonitic, by Bernutz. There can be no doubt that he here made an enormous step in advance. Like many authors, however, he rode his hobby a little too hard, and he erred in denying the occurrence of cellulitis in many cases in which it undoubtedly exists.

Etiology.—The great influence of childbirth in producing these diseases has long been fully recognized. Courty estimates that about two-thirds of all the cases met with occur in connection with delivery or abortion, and Duncan found that, out of 40 carefully-observed cases, 25 were associated with the puerperal state.

The Inflammation is Secondary, and never Idiopathic.—It is pretty generally admitted by most modern writers that both varieties of the disease are produced by the extension of inflammation from either the uterus, the Fallopian tubes, or the ovaries. This point has been especially insisted on by Duncan, who maintains that the disease is never idiopathic, and is “invariably secondary either to mechanical injury, or to the extension of inflammation of some of the pelvic viscera, or to the irritation of the noxious discharges through or from the tubes or ovaries.”

Often Intimately connected with Septicæmia.—Their intimate connection with puerperal septicæmia is also a prominent fact in the natural history of the diseases. Barker mentions a curious observation illustrative of this, that when puerperal fever is endemic in the Bellevue Hospital in New York cases of pelvic peritonitis and cellulitis are also invariably met with. Olshausen has also remarked that in the Lying-in Hospital at Halle during the autumn vacation, when the patients are not attended by practitioners, and when, therefore, the chance of septic infection being conveyed to them is less, these inflammations are almost always absent. As inflammation of the lining membrane of the uterus, of the vaginal mucous membrane, and of the pelvic connective tissue are of very constant occurrence as local phenomena of septic absorption, the connection between the two classes of cases is readily susceptible of explanation. Schroeder, indeed, goes farther, and includes his description of these diseases under the head of puerperal fever. They do not, however, necessarily depend upon it; for, although it must be admitted that cases of this kind form a large pro-

THE PUERPERAL STATE.

of those met with, others unquestionably occur which cannot be traced to such sources, but are the direct result of causes altogether unconnected with the inflammation attending on septic absorption, such as the exertion shortly after delivery or premature coition. Mechanical causes may beyond doubt excite the disease in a woman predisposed by the puerperal process, but they cannot fairly be included under the head of puerperal fever.

Seat of the Inflammation in Pelvic Cellulitis.—Abundance of areolar tissue exists in connection with the pelvic viscera which may be the seat of cellulitis. It forms a loose padding between the organs contained in the pelvis proper, surrounds the vagina, the rectum, and the bladder, and is found in considerable quantity between the folds of the broad ligaments. From these parts it extends upward to the iliac fossæ and the inner surface of the abdominal parietes. In any of these positions it may be the seat of the kind of inflammation we are discussing. The essential character of the inflammation is similar to that which accompanies areolar inflammation in other parts of the body. There is first an acute inflammatory œdema, followed by the infiltration of the areolæ of the connective tissue with exudation, and the consequent formation of appreciable swellings. These may form in any part of the pelvis. Thus, we may meet with them—and this is a common situation—between the folds of the broad ligaments, or distinct hard tumors connected with the uterus and extending to the pelvic walls, their rounded outlines being readily made out by bimanual examination. If the cellulitis be limited in extent, such a swelling may exist on one side of the uterus only, forming a rounded mass of varying size and apparently attached to it. At other times the exudation is more extensive, and may completely or partially surround the uterus, extending to the cellular tissue between the vagina and rectum or between the uterus and the bladder. In such cases the uterus is imbedded and firmly fixed in dense, hard exudation. At other times the inflammation chiefly affects the cellular tissue covering the muscles lining the iliac fossæ. There it forms a mass, easily made out by palpation, but on vaginal examination little or no trace of the exudation can be felt, or only a sense of thickness at the roof of the vagina on the same side as the swelling.

Seat of the Inflammation in Pelvic Peritonitis.—In pelvic peritonitis the inflammation is limited to that portion of the peritoneum which invests the pelvic viscera. Its extent necessarily varies with the intensity and duration of the attack. In some cases there may be little more than irritation, while more often it runs on to exudation of plastic material. The result is generally complete fixation of the uterus and hardening and swelling in the roof of the vagina, and the lymph poured out may mat together the surrounding viscera, so as to form swellings difficult, in some cases, to differentiate from those resulting from cellulitis. On post-mortem examination the pelvic viscera are found extensively adherent, and the agglutination may involve the coils of the intestine in the vicinity, so as sometimes to form tumors of considerable size.

Relative Frequency of the Two Forms of Disease.—The relative frequency of these two forms of inflammation as puerperal affections is not easy to ascertain. In the non-puerperal state the peritonitic variety is

much the more common, but in the puerperal state they very generally complicate each other, and it is rare for cellulitis to exist to any great extent without more or less peritonitis.

Symptomatology.—The earliest symptom is pain in the lower part of the abdomen, which is generally preceded by rigor or chilliness. The amount of pain varies much. Sometimes it is comparatively slight, and it is by no means rare to meet with patients, the subjects of very considerable exudations, who suffer little more than a certain sense of weight and discomfort at the lower part of the abdomen. On the other hand, the suffering may be excessive, and is characterized by paroxysmal exacerbations, the patient being comparatively free from pain for several successive hours, and then having attacks of the most acute agony. Schroeder says that pain is always a symptom of peritonitis, and that it does not exist in uncomplicated cellulitis. The swellings of cellulitis are certainly sometimes remarkably free from tenderness, and I have often seen masses of exudation in the iliac fossæ which could bear even rough handling. On the other hand, although this is certainly more often met with in non-puerperal cases, the tenderness over the abdomen is sometimes excessive, the patient shrinking from the slightest touch. The pulse is raised, generally from 100 to 120, and the thermometer shows the presence of pyrexia. During the entire course of the disease both these symptoms continue. The temperature is often very high, but more frequently it varies from 100° to 104°, and it generally shows more or less marked remissions. In some cases the temperature is said not to be elevated at all, or even to be subnormal, but this is certainly quite exceptional. Other signs of local and general irritation often exist. Among them—and most distinctly in cases of peritonitis—are nausea and vomiting and an anxious, pinched expression of the countenance, while the local mischief often causes distressing dysuria and tenesmus. The latter is especially apt to occur when there is exudation between the rectum and vagina, which presses on the bowel. The passage of feces, unless in a very liquid form, may then cause intolerable suffering.

The Symptoms often Insidious in their Onset.—Such symptoms may show themselves within a few days after delivery, and then they can barely fail to attract attention. On the other hand, they may not commence for some weeks after labor, and then they are often insidious in their onset and apt to be overlooked. It is far from rare to meet with cases six weeks or more after confinement in which the patient complains of little beyond a feeling of *malaise* and discomfort, and in which, on investigation, a considerable amount of exudation is detected which had previously entirely escaped observation.

Results of Physical Examination.—On introducing the finger into the vagina it will be found to be hot and swollen, in some cases distinctly œdematous, and on reaching the vaginal *cul-de-sac* the existence of exudation may generally be made out. The amount of this varies much. Sometimes, especially in the early stage of the disease, there is little more than a diffuse sense of thickness and induration at either side of, or behind, the uterus. More generally, careful bi-manual examination enables us to detect a distinct hardening and swelling, possibly a tumor

THE PUERPERAL STATE.

size, which may apparently be attached to the sides of the uterus, rise above the pelvic brim, or may extend quite to the umbilicus. The examination should be very carefully and systematically made with both hands, so as to explore the whole contour of the uterus before, behind, and on either side, as well as the iliac fossæ, in case a considerable exudation might readily escape detection. When the exudation is at all great more or less fixity of the uterus is sure to be observed and is a very characteristic symptom. The womb, instead of being movable by the examining finger, is firmly fixed by the surrounding exudation, and in severe forms of the disease is quite encased in it. More or less displacement of the organ is also of common occurrence. The swelling is limited to one side of the pelvis or to Douglas's space, the uterus is displaced in the opposite direction, so that it is no longer in its usual central position.

The Two Forms cannot Always be Distinguished.—The differential diagnosis of pelvic cellulitis and pelvic peritonitis cannot always be made, and indeed in many cases it is impossible, since both varieties of disease coexist. The elements of diagnosis generally insisted on are the greater general disturbance, the sea, etc. in pelvic peritonitis, and an earlier commencement of the symptoms after labor. The swellings in pelvic peritonitis are also more extensive, with less clearly-defined outlines than those of cellulitis. When the cellulitis involves the iliac fossa, the diagnosis is of course easy, the continuous retraction of the uterus on the affected side (an invagination position assumed with the view of deepening the muscles lining the iliac fossa at rest) is often observed. When the inflammation is chiefly limited to the cavity of the pelvis, the distinction between the two classes of cases cannot be made with any degree of certainty.

Terminations.—Both forms of disease may end either in resolution or in suppuration. In the former case, after the acute symptoms have existed for a variable time—it may be for a few days only, it may be for many weeks—their severity abates, the swellings become less tender and commence to contract, become harder, and are gradually absorbed, until at last the fixity of the uterus disappears, and it again resumes its central position in the pelvic cavity. This process is often very gradual. It is by no means rare to find a patient, even some months after the attack, when all acute symptoms have long disappeared, who is even able to move about without inconvenience, in whom the uterus is still immovably fixed in a mass of deposit or is at least adherent in some part of its contour. More or less permanent adhesions are of common occurrence, and give rise to symptoms of considerable obscurity, which are often not traced to their proper source.

Symptoms of Suppuration.—When the inflammation is about to terminate in suppuration, the pyrexial symptoms continue, and eventually well-marked hectic is developed, the temperature generally showing a distinct exacerbation at night. At the same time, rigors, loss of appetite, a peculiar yellowish discoloration of the face, and other signs of suppuration show themselves. The relative frequency of this termination is variously estimated by authors. Duncan quotes Simpson as calculating it as occurring in half the cases of pelvic cellulitis, but states his o

belief that it is much more frequent. West observed it in 23 out of 43 cases following delivery or abortion, and McClintock in 37 out of 70. Schroeder says that he has only once seen suppuration in 92 cases of distinctly demonstrable exudation—a result which is certainly totally opposed to common experience. Barker also states that in his experience suppuration in either pelvic peritonitis or cellulitis “is very rare, except when they are associated with pyæmia or puerperal fever.” It is certain that suppuration is more likely to occur in pelvic cellulitis than in pelvic peritonitis, but it unquestionably occurs, in this country at least, much more frequently than the statements of either of these authors would lead us to suppose.

Channels through which Pus may Escape.—The pus may find an exit through various channels. In pelvic cellulitis, more especially when the areolar tissue of the iliac fossa is implicated, the most common site of exit is through the abdominal wall. It may, however, open at other positions, and the pus may find its way through the cellular tissue and point at the side of the anus or in the vagina, or it may take even a more tortuous course and reach the inner surface of the thigh. Pelvic abscesses not uncommonly open into the rectum or bladder, causing very considerable distress from tenesmus or dysuria. According to Hervieux, it is chiefly the peritoneal varieties which open in this way. Not unfrequently more than one opening is formed, and when the pus has burrowed for any distance long fistulous tracts result, which secrete pus for a length of time and are very slow to heal. Rupture of an abscess into the peritoneal cavity, especially of a peritonitic abscess, is a possible (but fortunately a very rare) termination, and will generally prove fatal by producing general peritonitis. In one case, which I have recorded in the fifteenth volume of the *Obstetrical Transactions*, suppuration was followed by extensive necrosis of the pelvic bones. Two similar cases are related by Trousseau in his *Clinical Medicine*, but I have not been able to meet with any other examples of this rare complication, which was probably rather the result of some obscure septicæmic condition than of extension of the inflammation.

Prognosis.—The prognosis is favorable as regards ultimate recovery, but there is great risk of a protracted illness which may seriously impair the health of the patient, especially if suppuration result. Hence it is necessary to be guarded in an expression of opinion as to the consequences of the disease. Secondary mischief is also far from unlikely to follow from the physical changes produced by the exudation, such as permanent adhesions or malpositions of the uterus or organic alterations in the ovaries or Fallopian tubes.

Treatment.—In the treatment of both forms of disease the important points to bear in mind are the relief of pain and the necessity of absolute rest; and to these objects all our measures must be subordinate, since it is quite hopeless to attempt to cut short the inflammation by any active medication.

If the disease be recognized at a very early stage, the local abstraction of blood, by the application of a few leeches to the groin or to the hemorrhoidal veins, may give relief; but the influence of this remedy has been greatly exaggerated, and when the disease is of any standing it is

THE PUERPERAL STATE.

leeches to the uterus, often recommended, are, I believe, to do less harm than good (unless in very skilful hands), from irritation produced by passing the speculum. Opiates in large doses are said to be our sheet-anchor in treatment whenever the pain is severe, either by the mouth, in the form of morphia suppositories, or rectally subcutaneously. In the not uncommon cases in which pain is severe in paroxysms the opiates should be administered in sufficient quantity to lull the pain; and it is a good plan to give the full supply of morphia suppositories (which often act better than any other form of administering the drug), with directions to use them whenever the pain threatens to come on. When there is much pain, large doses of quinine may be given with great advantage along with the opiates. The state of the bowels requires careful attention. Opiates are apt to produce constipation, and the passage of hardened feces causes much suffering. Hence it is desirable to keep the bowels open. Nothing answers this purpose so well as small doses of castor oil, such as half a teaspoonful given every morning. Warmth and moisture, constantly applied to the lower part of the abdomen, either in the form of large poultices of linseed meal, or, if these prove too heavy, of spongio-piline soaked in boiling water, give great relief. The poultices may be advantageously sprinkled with laudanum or belladonna liniment. I say nothing of the use of mercurials, iodide of potassium, and other so-called absorbent remedies, since I believe them to be quite valueless and apt to divert attention from more useful plans of treatment.

Importance of Rest.—The most absolute rest in the recumbent position is essential, and it should be persevered in for some time after the intensity of the symptoms is lessened. The beneficial effect of rest in alleviating pain is often seen in neglected cases the nature of which has been overlooked, instant relief following the laying up of the patient.

Counter-irritation.—When the acute symptoms have lessened, absorption of the exudation may be favored and considerable relief obtained from counter-irritation, which should be gentle and long continued. The daily use of tincture of iodine until the skin peels perhaps best meets this indication, but frequently-repeated blisters are often very serviceable. This I believe to be a better plan than keeping up an open sore with savine ointment or similar irritating applications.

Opening of Pelvic Abscesses.—When suppuration is established the question of opening the abscess arises. When this points in the groin and the matter is superficial, a free incision may be made; and here, as in mammary abscess, the antiseptic treatment is likely to prove very serviceable. The abscess should, however, not be opened too soon, and it is better to wait until the pus is near the surface. The importance of not being in too great a hurry to open pelvic abscesses has been insisted on by West, Duncan, and other writers, and I have no doubt the rule is a good one. It is more especially applicable when the abscess is pointing in the vagina or rectum, where exploratory incisions are apt to be dangerous, and when the presence of pus should be positively ascertained before operating. We have in the aspirator a most useful instrument in the treatment of such cases, which enables us to remove

the greater part of the pus without any risk, and the use of which is not attended with danger even if employed prematurely. If it do not sufficiently evacuate the abscess, a free opening can afterward be safely made with the bistoury. The surgical treatment of pelvic abscess is, however, too wide a subject to admit of being satisfactorily treated here.

Diet and Regimen.—The diet should be abundant, but simple and nutritious. In the early stages of the disease milk, beef-tea, eggs, and the like will be sufficient. After suppuration a large quantity of animal food is necessary, and a sufficient amount of stimulants. The drain on the system is then often very great, and the amount of nourishment patients will require and assimilate when a copious purulent discharge is going on is often quite remarkable. A general tonic plan of medication is also indicated, and such drugs as iron, quinine, and cod-liver oil will prove useful.



INDEX.

ABDOMEN, adipose enlargement of, 157
 enlargement of, as a sign of pregnancy, 149
 state of, after delivery, 551
Abdominal pregnancy. See *Extra-uterine Pregnancy.*
Abortion, 242
 causes of, 244
 difficulty in procuring artificial, 248
 liability to recurrence of, 243
 production of, in vomiting of pregnancy, 201
 retention of secundines in, 249, 253
 symptoms of, 248
 treatment of, 249
 value of opium in prevention of, 249 [250]
Abscess of mammae. See *Mammary Abscess.*
Abscess, pelvic. See *Pelvic Cellulitis.*
After-pains, 552
 treatment of, 554
Age, influence of, in labor, 340
Albuminuria in pregnancy, 206
 relation of, to eclampsia, 576
 relation of, to puerperal insanity, 586
Allantois, 167
Amnii, liquor, 109
Amnion, formation of, 106
 pathology of, 236
 structure of, 109
Amputations (intra-uterine), 240
Anæmia in pregnancy, 206
Anæsthesia in labor, 295
 in forceps operations, 478
 value of, in difficult cases of turning, 470
Anasarca in pregnancy, 209
Anteversion of the gravid uterus, 217
Antiseptic midwifery, 604
Apoplexy during or after labor, 635
Arbor vitæ, 59
Area germinativa, 105
Area pellucida, 105
Areola, 80
 changes of, during pregnancy, 146
Arm, presentation of. See *Shoulder Presentation.*
 dorsal displacement of, 330
Arterial transfusion, 542
Artificial human milk, 571
Artificial respiration in cases of apparent still-birth, 558

Ascites as a cause of dystocia, 373
Asphyxia (idiopathic), 635
 of new-born children, 557
Atropine, hypodermic injection of, in rigidity of cervix, 352
Auscultatory signs of pregnancy, 152

BAGS (Barnes's). See *Dilators.*
Ballotement, 151
Basilyst, the, 508
Bi-lobed uterus, gestation in, 192
Binder, uses of, 294
 [application of, 290]
Bladder, distension of, as a cause of protracted labor, 340
 exfoliation of lining membrane of, 213
 state of, after delivery, 553
Blastodermic membrane, 99
 division and layers of, 105
Blood, alteration in, after delivery, 547
 changes of, during pregnancy, 139
Blood-diseases transmitted to fetus, 237
Blunt-hook in breech presentation, 309
Bowels, action of, after delivery, 556
Breech presentations. See *Pelvic Presentations.*
Broad ligaments of uterus, 68
Bronchitis as a cause of protracted labor, 340
Brow presentations, 318

CÆSAREAN section, 330, 359, 398, 511, 513
 [in America, 525]
 [in Great Britain, 525]
 causes of mortality after, 517
 causes requiring the operation, 575
 description of, 521
 history of, 511
 [improved methods of performing, 523]
 [mortality in cases of fibroid tumor, 357]
 [in pelvic exostosis, 388]
 post-mortem operation, 516
 [process of Cohnstein, 523]
 [of Frank, 522]
 [of Kehrer, 524]
 [of Säger, 524]
 results to child in, 574
 [results in Great Britain and the United States compared, 525]

INDEX.

section
 ics of,
 t Ame., 513]
 American dwarfs, 514]
 substitutes for, 526
 res in, 522
 ures in the United States, 522]
 transverse position of fetus, 330]
 ulus of bladder obstructing labor, 359
 succedaneum, 279
 aa in, unc, 223
 r, 353
 a pulmonary obstruc-
 teet. i pregnancy, 204
 a twins, birth of, 370]
 myrtiformes, 52
 er introduction of, 51
 264
 i See *Pelvic Cellulitis*.
 nipsy. See *Craniotomy*.
 uteri, 58
 alterations of, after childbirth, 58
 cavity of, 59
 dition of, in labor, 259
 trophic elongation of, 353
 pation of, before fetal head, 286
 of, for rigidity, 354
 is of, 439
 ion of, by pregnancy, 136
 mucous membrane of, 63
 rganic causes of rigidity of, 353
 rigidity of, as a cause of protracted
 labor, 351
 treatment of rigidity, 352
 villi of, 64
 Charlotte, Princess of Wales, death of, 348
 Child (the new-born). See *Infant*.
 Child, risks to, in forceps operations, 486
 Childbirth, mortality of, 546
 Chloral in labor, 295
 in rigidity of cervix, 352
 Chloroform in labor, 296
 [deaths from, 297]
 in difficult cases of turning, 462
 in rigidity of cervix, 352
 Chorea in pregnancy, 212
 Chorion, 110
 vesicular degeneration of, 229
 Circulation of fetus, 129
 Cleavage of yolk, 99
 Clitoris, 48
 Coccyx, 35
 ligaments of, 36
 ossification of, 36
 mobility of, 36
 Cold in the treatment of puerperal hyper-
 pyrexia, 619
 Colostrum, 559
 Complex presentations, 330
 Conception, signs of, 143
 Constipation in pregnancy, 202
 Constriction of uterus, tetanoid, 355
 Continued fever in pregnancy, 221

Convulsions puerperal. See *Eclampsia*.
 Corps reticulé, 108
 Corpus luteum, 84
 false, 85
 Cranioclast, 500
 Craniotomy, 497
 cases requiring, 502
 comparative merits of, and cephalo-
 tripsy, 505
 description of cephalotripsy, 506
 extraction of head by craniotomy for-
 ceps, 508
 method of perforating, 504
 perforators, 499
 perforation of after-coming head, 505
 religious objections to, 498
 Cran my forceps, 500
 Crotaceus, 499
 Cyclical theory of menstruation, 92
 Cystocele, obstructing labor, 359

DEATH, apparent, of new-born child.
 See *Infant*.
 sudden, during labor and the puer-
 peral state, 635
 from air in the veins, 636
 functional causes of, 635
 nic causes of, 635
 tion of fetus, 510
 , 100
 end of pregnancy and after deliv-
 ery, 104
 cavity between d. vera and reflexa,
 104
 divisions of, 100
 fatty degeneration of, as the cause of
 labor, 256
 formation of d. reflexa, 102
 structure of, 101
 Delivery, state of patient after, 547
 contraction of uterus after, 549
 management of patient after, 553
 nervous shock after, 547
 prediction of date of, 161
 signs of recent, 164
 state of pulse after, 547
 weight of uterus after, 550
 Diabetes, 143
 Diameters of fetal skull, 127
 of pelvis, 41
 Diarrhœa in pregnancy, 202
 [Diet, milk, in nursing mothers, 564]
 Diet of lying-in women, 554
 Dilators (cautheouc) in the induction of
 premature labor, 453
 in rigidity of cervix, 353
 Diphtheria in the puerperal state, 600
 Diseases of pregnancy, 198
 albuminuria, 206
 anæmia and chlorosis, 206
 carcinoma, 223
 cardiac diseases, 222
 chorea, 212
 constipation, 202

Diseases of pregnancy—
 diarrhoea, 202
 disorders of the nervous system, 211
 respiratory organs, 204
 teeth, 204
 urinary system, 213
 displacements of the gravid uterus, 216
 epilepsy, 223
 eruptive fevers, 221
 fibroid tumors, 225
 hæmorrhoids, 203
 icterus, 223
 leucorrhœa, 214
 ovarian tumor, 224
 palpitation, 205
 paralysis, 211
 pneumonia, 221
 pruritus, 215
 ptyalism, 204
 syncope, 205
 syphilis, 222
 varicose veins, 215
 vomiting (excessive), 198
Dropsies affecting the fœtus, 239
Ductus arteriosus, 129
 venosus, 129
Dystocia from fœtus, 363

ECLAMPSIA, 573
 cause of death in, 576
 condition of patient between the attacks, 575
 confusion from defective nomenclature, 573
 exciting causes of, 578
 obstetric management in, 581
 pathology of, 576
 premonitory symptoms of, 574
 relation of, to labor, 575
 results to mother and child in, 576
 symptoms of, 574
 transfusion in, 536
 Traube and Rosenstein's theory of, 577
 treatment of, 579
 uræmic theory of, 576
 venesection in, 579
 views of MacDonal, 578
Écraseur, use of, as a substitute for craniotomy, 501
Embolism. See Thrombosis.
Embryotomy, 509
Emotion, mental, as a cause of protracted labor, 340
Epiblast, 105
Epilepsy, in pregnancy, 223
Epileptic convulsions, 574
Ergot of rye, 342
 as a means of inducing labor, 452
 objections to use of, 343
 mode of administration, 342
 value of, after delivery, 294
Ergotin, hypodermic injection of, in post-partum hemorrhage, 422
Eruptive fevers in pregnancy, 221

Erysipelas as a cause of puerperal septicæmia, 599
Ether in labor, 297 [298]
Evisceration, 511
Exhaustion, importance of distinguishing between temporary and permanent in labor, 342
Expression, uterine (see Pressure); of the placenta, 291
Extra-uterine pregnancy, 171
 abdominal variety of, 182
 causes of, 173
 changes of the fœtus in, 185
 classification of, 172
 diagnosis of abdominal variety, 186
 diagnosis of tubal variety, 177
 [faradic current in, 180]
 gastrotomy in, 182, 187
 [non-removal of placenta in, 179]
 [laparotomy, primary and secondary, relative risks of, 189]
 pseudo-labor in, 184
 vaginal section in, 179
 [Mathieson's case of, 179]
 symptoms of rupture in, 176
 treatment after rupture, 182
 [Lawson Tait's operations, 182]
 treatment of abdominal variety, 186
 tubal variety, 175
 treatment of tubal variety, 178

FACE presentation, 310
 causes of, 310
 diagnosis of, 311
 difficulties connected with, 317
 erroneous views formerly entertained of, 310
 mechanism of delivery in, 312
 mento-posterior positions in, 314
 prognosis in, 316
 treatment of, 316
Fallopian tubes, 71
False corpus luteum, 85
False pains, character and treatment of, 282, 283
Faradization in apparent stillbirth, 558
 in destroying the vitality of the fœtus in abnormal pregnancies, 180
 in hemorrhage after delivery, 426
 in labor, 344
Fibroid tumor, in pregnancy, 225
 obstructing labor, 356 [357]
Fillet, 496
 in breech presentations, 309
 nature of the instrument, 496
 objections to its use, 497
Flattened pelvis, 378
Fœtal head, anatomy of, 120
 induction of premature labor, for large size of, 450
Fœtal heart, sounds of, in pregnancy, 153
Fœtus, anatomy and physiology of, 118
 [anencephalous, producing enuresis, 214]

INDEX.

appearance of a putrid, 241
 appearance of, at various stages of development, 118
 at term, 119
 circulation of, 129
 changes in circulation of, as cause of labor, 255
 changes in position of, during pregnancy
 without water, 559]

position in utero by palpation, 243
 primary viability of, 243
 excessive development of, as a cause of difficult labor, 374
 explanation of its position in utero, 125
 functions of, 127
 nutrition of, 127
 pathology of, 237
 position of, in utero, 123
 respiration of, 128
 signs and diagnosis of death of, 242
 tangles, 120
 diagnosis of, 301
 presentations. See *Pelvic Presentation*
 of, 129
 of, 476
 advantages of pelvic curve in, 473
 application of, to after-coming head in breech presentations, 307
 application of, within the cervix, 355
 [carried over abdomen to complete delivery of head, 494]
 [at inferior strait, 491]
 [at superior strait, 493]
 cases in which a straight instrument should be used, 473
 dangers of, 347, 485
 dangers of, to child, 486
 description of, 472
 description of the operation, 479
 difference between high and low operations, 478
 disadvantages of a weak instrument, 476
 frequent use of, in modern practice, 346, 472
 high operations, 484
 [in America, 486]
 long, 474
 preliminary considerations before using, 478
 short, 472
 use of anesthetics in forceps delivery, 479
 use of, in deformed pelvis, 395
 use of, in difficult occipito-posterior positions, 320
 use of, in protracted labor, 346
 [Bedford's, 489]
 [Davis's, 488]

Forceps—

[Elliot's, 490]
 [Hodge's, 488]
 [Meigs's craniotomy, 509]
 [Sawyer's, 490]
 [Wallace's, 488]
 [White's, 490]
 Forceps saw, 501
 Fossa navicularis, 53
 Funis. See *Umbilical Cord*.
 Funnel-shaped pelvis, 378

GALACTAGOGUES, 564

Galactorrhœa, 565
 Galvanism as a means of inducing labor, 152
 Gangrene of limbs from arterial obstruction, 634
 Gastrotomy, after rupture of uterus, 438
 in extra-uterine pregnancy, 182, 187
 Gastro-elytrotomy. See *Laparo-elytrotomy*.
 o-hysterotomy, possibilities of, 528]
 tive organs, in the female, 48
 division according to function, 48
 minimal vesicle, disappearance of, after impregnation, 98
 m. See *Pregnancy*.
 tria in pregnancy, 143
 lactation, 549
 human follicle, 75
 structure of, 77

HEMATOCELE, obstructing labor, 360

[Hand, introduction of in occipito-posterior positions, 322]
 Hand-feeding of infants, 569
 ass's milk in, 569
 artificial human milk in, 570
 causes of mortality in, 569
 cow's milk in, and its preparation, 570
 goat's milk in, 570
 method of, 572
 Head presentations, 268
 description of cranial positions in, 268, 269
 division of, 269
 frequency of first position, 268
 explanation of, 270
 mechanism of first position, 272
 second position, 276
 third position, 276
 fourth position, 278
 relative frequency of various positions, 269
 Heart, diseases of, in pregnancy, 222
 hypertrophy of, in pregnancy, 141
 Hemorrhage, accidental, 411
 causes and pathology of, 412
 concealed internal, 413
 diagnosis, prognosis, and treatment of concealed internal, 413, 414

Hemorrhage—

- prognosis of, 414
- symptoms and diagnosis of, 412
- treatment of, 414
- after delivery, 415
- causes of, 416
- constitutional predisposition to, 419
- curative treatment of, 421
- from laceration of maternal structures, 427
- nature's mode of preventing, 266, 415
- preventive treatment of, 420
- secondary causes of, 417
- secondary treatment of, 427
- symptoms of, 420
- transfusion of blood in, 428
- after delivery (secondary), 428
- distinction between, and profuse lochial discharge, 428
- local causes of, 429
- treatment of, 430
- unavoidable. See *Placenta Prævia*.
- Hemorrhoids, in pregnancy, 203
- Hernia, in labor, 360
- Hour-glass contraction of uterus, 417 [418]
- ante-partum, 355
- Hydatids of uterus, 229
- Hydramnios, 236
- Hydrocephalus of fœtus, as a cause of difficult labor, 371
- Hydorrhœa gravidarum, 228
- Hymen, 52
- Hypoblast, 105
- Hysteria during labor, 574

INDUCTION of premature labor. See *Premature Labor*.

Inertia of the uterus, frequent child-bearing as a cause of, 339

Infant, apparent death of, 557

- appearance of, in cases of apparent death, 557

- clothing of, 559
- evils of over-suckling, 560
- management of, 561
- management of, when food disagrees, 573

- treatment of apparent death of, 557
- various kinds of food of, 572
- washing and dressing of, 558

Infantile mortality, diminution of, as a reason for more frequent use of forceps, 347

Inflammatory diseases affecting the fœtus, 239

Injectons, uterine, of hot water, 424

Insanity (puerperal), 582

- classification of, 583
- of lactation, 588
- of pregnancy, 583
- predisposing causes of, 583
- puerperal (proper), 585
- causes of, 585

Insanity—

- form of, prognosis of, 584
- post-mortem signs of, 588
- symptoms of, 588
- transient mania during delivery, 584
- treatment of, 590
- treatment during convalescence, 592
- question of removal to an asylum, 592
- Insomnia in pregnancy, 211
- Intermittent fever affecting the fœtus, 238
- Intestines, disorders of, as influencing labor, 340
- Inversion of uterus. See *Uterus*.
- Irregular uterine contractions after labor, 417
- as a cause of lingering labor, 341
- Irritable bladder in pregnancy, 213
- Ischium, planes of the, 46

JAUNDICE in pregnancy, 223**K**IESTEIN, 142 [143]

- Knee presentation, 301
- Knots of the umbilical cord, 235
- Kyphotic deformity of pelvis, 386

LABIA majora, 49

Labia minora, 50

Labor, 255

- age, influence of, on, 340
- anæsthesia in, 295
- arrest of, 164
- causes of, 255
- causes of precipitate, 350
- causes of protracted, 337
- character and source of pain in, 261
- character of false pains, 263
- dilatation of cervix in, 257
- duration of, 266
- effect of uterine contractions in, 257
- evil effects of protracted, 337
- induction of. See *Premature Labor*.
- influence of stage of, in protracted, 337
- management of, in deformed pelvis, 394
- management of natural, 280
- management of third stage of, 289
- mechanism of, in head presentation, 268
- [missed, causes of, 195]
- obstructed by faulty condition of the soft parts, 351
- period of day at which labor commences, 267
- phenomena of, 255
- position of patient during, 285
- [premature, induction of, 456]
- preparatory treatment, 280
- precipitate, 350
- prolonged and precipitate, 337
- rupture of membranes in, 259
- stages of, 262

Labor—

- symptoms of protracted, 338
- treatment of protracted, 341

Lactation, defective secretion of milk in, 564

- diet of nursing women during, 562
- diseases of the eye during, 566
- evil results of prolonged, 560
- excessive flow of milk in, 565
- importance of, to mother, 560
- importance of wet-nursing to child, 560, 561
- insanity of, 588
- management of, 561
- means of arresting secretion of milk in, 563
- period of weaning in, 563

Lamine dorsales, 105

Laparo-elytrotomy, 529

- [latest statistics of, 530]

[Laparotomies, American puerperal, 438]

Lead-poisoning, affecting the fetus, 238

- as a cause of abortion, 247

Leucorrhœa, in pregnancy, 214

Lever. See *Vectis*.

Liquor amnii, 109

- uses of, 110
- source of, 109
- deficiency of, 237

Lithopædion, 185

Liver, acute yellow atrophy of, 223

Lochia, 551

- variation in amount and duration of, 552

- occasional fetor of, 552

Lying-in hospitals, mortality in, 594

Lypo hæmia, 146, 205

MALARIAL puerperal fever, 615

- Malpresentations, peculiar form of bag of membranes in, 301

Mammary abscess, 566

- antiseptic treatment of, 567
- signs and symptoms of, 566
- treatment of, 566
- changes during pregnancy, 148
- their diagnostic value, 79
- glands, 146
- their sympathetic relations with the uterus, 80

Mania, puerperal. See *Insanity, Puerperal*.

Measles, affecting the fetus, 238

- in pregnancy, 221

Meconium, 131

Membranes, artificial rupture of, 284

- puncture of, as a means of inducing labor, 451

Menstruation, 81

- cessation of, 93
- during pregnancy, 144
- changes in Graafian follicle after, 81
- period of, duration, and recurrence, 87

Menstruation—

- purpose of, 92
- source of blood in, 89
- theory of, 90
- quantity of blood lost in, 88
- vicarious, 92

[Menstrual life, duration of, 94]

Mesoblast, 105

Milk, artificial human, 570

- ass's, 569
- cow's, and its preparation, 570
- defective secretion of, 564
- excessive secretion of, 565
- goat's, 570
- means of arresting the secretion of, 563
- secretion of, after delivery, 560
- transfusion of, 539

Milk-fever, 548

Miscarriage. See *Abortion*.

Missed labor, 193

Moles, 245

Monstrosity (double), 367

- classification of, 367
- mechanism of delivery in, 367

Mons veneris, 49

Montgomery's cups, 102

Morning sickness, 145

Mortality of childbirth, 546

Mucous membrane of uterus. See *Uterus*.

Müller's operation, 526

Myxoma fibrosum, 232

NERVOUS shock after delivery, 547

Nervous system, changes in, during pregnancy, 141

- disorders of, in pregnancy, 211
- excitability of, in puerperal women, 578

Neuralgia in pregnancy, 211

Nipple, 80

Nipples, depressed, 564

- fissures and excoriations of, 564

Nursing. See *Lactation*.

Nutrition of fetus, 127

Nympheæ. See *Labia minora*.**O**BLIQUELY contracted pelvis, 386

Obstetric bag, 281

Occipito-posterior positions, difficult cases of, 319

- causes of face-to-pubes delivery in, 319

- [version by the vertex, 321]

- forceps in, 320

- treatment of, 319

- vectis or fillet in, 320

Omphalo-mesenteric artery and vein, 107

Opiates, use of, after delivery, 553

[Opium to arrest labor in threatened abortion, 250]

Os innominatum, 33

Osteomalacia, as a cause of deformity, 376

- Osteophytes**, formation of, during pregnancy, 141
- Os uteri**, constriction of internal, as a cause of dystocia, 355
dilatation of, as a means of inducing labor, 453
occlusion of, in labor, 354
- Ovarian pregnancy**. See *Extra-uterine Pregnancy*.
tumor in pregnancy, 224
- Ovariectomy** in pregnancy, 224
- Ovary**, 72
functions of, 81
structure of, 74
vascular arrangements of, 78
- Ovule**, 77
changes in, after impregnation, 98
changes in, when retained in utero after its death, 245
formation of, 75
- Ovum**, blighted, retained in utero, 245
- Oxytocic remedies**, 342
- PAINS**, after-, 552
false, 282
irregular and spasmodic, as a cause of protracted labor, 341
labor, 261
- Palpitation** in pregnancy, 205
- Pampiniform plexus**, 65
- Paralysis** in pregnancy, 211
from embolism of the cerebral arteries, 634
from embolism of the main arteries of the limb, 634
- Parovarium**, 69
- Parturient canal**, axis of, 45
- Pathology** of decidua and ovum, 227
- Pelvic cellulitis and peritonitis**, 644
etiology of, 645
importance of, distinguishing the two forms of disease, 644
connection with septicæmia, 645
opening of abscess in, 650
prognosis of, 649
relative frequency of the two forms of disease, 646
results of physical examination, 647
seat of inflammation in cellulitis, 646
seat of inflammation in peritonitis, 646
suppuration in, 648
symptomatology, 647
terminations of, 648
treatment of, 649
two distinct forms of disease, 644
- Pelvic presentations**, 299
application of forceps to the after-coming head in, 307
causes of, 299
danger to children in, 300
diagnosis of, 300
frequency of, 299
- Pelvic presentations**—
management of impacted breech in, 308
mechanism of, 302
prognosis in, 299
treatment of, 306
- Pelvimeters**, various forms of, 392
- Pelvis**, alterations in, articulations of, during pregnancy, 39
anatomy of, 33
articulations of, 36
axes of, 45
Cæsarean section in deformities of, 398
causes of deformity of, 375
comparative estimate of turning and forceps in deformity of, 397
craniotomy in deformity of, 398
diagnosis of deformity, 391
deformities of, 375
development of, 46
difference according to race, 47
differences in the two sexes, 41
division into true and false, 34
equally contracted, 377
equally enlarged, 377
flattened, 378
forceps in deformity of, 395
induction of premature labor in deformity of, 398
infantile, 46
[justo-minor in a large woman, 377]
kyphotic, 386
ligaments of, 37
funnel-shaped, 378
masculine, 378
mechanism of delivery in deformed, 390
movements of the articulations of, 38
obliquely-contracted, 385
planes of, 385
Robert's, 387
scoliotic, 379
soft parts connected with, 48
tumors of, 388
turning in deformity of, 396
undeveloped, 377
- Perchloride of iron**, injections of, in post-partum hemorrhage, 426
- Perforation** of after-coming head, 504
- Perforators**, 498
- Perineum**, distension of, in labor, 264, 286
incision of, 287
laceration of, 288
relaxation of, 287
rigidity of, as a cause of protracted labor, 356
- Peritonitis**, pelvic. See *Pelvic Cellulitis*.
[puerperal, venesection in, 618]
puerperal. See *Septicæmia*.
- Phlegmasia dolens**. See *Thrombosis, Puerperal, Venous*.
- Placenta**, adhesion of, after delivery, 419
degeneration of, 116
detachment of, in labor, 265

Placenta—

- expression of, 291
- fœtal portion of, 112
- form of, in man and animals, 111
- formation of, from chorion, 110
- functions of, 116
- maternal portion of, 114
- minute structure of, 112
- pathology of, 232
- sinus system of, 114
- sounds produced during separation of, 156
- treatment of adherent, 423
- treatment of, in extra-uterine fixation, 188
- Placenta membranacea, 232
- Placenta prævia, 400
 - causes of, 400
 - causes of hemorrhage in, 403
 - natural termination of labor in, 405
 - pathological changes of placenta in, 405
 - prognosis in, 406
 - sources of hemorrhage in, 403
 - summary of rules for treatment in, 410
 - symptoms of, 403
 - treatment of, 406
 - turning in, 408
 - [by Hicks' method, 411]
- Placentæ succenturiæ, 232
- Placentitis, 233
- Plugging of vagina, 252
- Plural births, 166, 363
 - arrangement of placenta and membranes in, 168
 - causes of, 167
 - diagnosis of, 169
 - relative frequency of, in different countries, 166
 - sex of children in, 167
 - treatment of, 363
- Pneumonia in pregnancy, 221
 - puerperal embolic, 632
- "Polar globule," 99
- [Polypus obstructing labor, 362]
- Porro's operation, 525
- [Porro-Cesarean statistics, the latest, 526]
- Position of cranium in head presentation. See *Head Presentations*.
- Post-partum hemorrhage. See *Hemorrhage*.
- Pregnancy, 132
 - abnormal, 166
 - affections of respiratory organs, 204
 - alteration of color of vaginal mucous membrane, as a sign of, 152
 - ballotement, as a sign of, 151
 - changes in the blood during, 139
 - changes in the liver, lymphatics, and spleen during, 141
 - changes in the urine during, 142
 - complicated with ovarian tumor, 224
 - deposits of pigmentary matter during, 148

Pregnancy—

- diabetes in, 210
 - differential diagnosis of, 157
 - dress of patient in, 280
 - duration of, 160
 - enlargement of abdomen as a sign of, 149
 - extra-uterine. See *Extra-uterine Pregnancy*.
 - fœtal movements in, 149
 - formation of osteophytes during, 141
 - hypertrophy of the heart during, 141
 - in cases of double uterus, 67
 - in the absence of menstruation, 145
 - intermittent uterine contractions, as a sign of, 150
 - ptyalism in, 204
 - prolapse of the uterus in, 216
 - protraction, 163
 - pruritus in, 215
 - quickening, 149
 - sickness of, 145
 - signs and diagnosis of, 144
 - sounds produced by the fœtal movements in, 156
 - spurious, 159
 - sympathetic disturbances of, 145
 - [tension of abdomen in, producing dyspnea at night, how treated, 205]
 - uterine fluctuation in, 151
 - vaginal pulsation in, 151
 - vaginal signs of, 151
- Premature labor, 242
- history of the operation of induction of, 449
 - induction of, 449 [456]
 - induction of, in deformed pelvis, 398
 - injection of carbonic acid gas as a means of inducing, 455
 - insertion of flexible bougie as a means of inducing, 455
 - objects of the operation of induction of, 449
 - oxytocics as a means of inducing, 452
 - period for the induction of, in deformed pelvis, 399
 - precautions as regards the child in the induction of, 456
 - puncture of the membranes as a means of inducing, 451
 - separation of the membranes as a means of inducing, 454
 - vaginal and uterine douches as a means of inducing, 454
- Pressure as a means of inducing uterine contractions, 344
- mode of applying, 345
- Prolapse of umbilical cord. See *Umbilical Cord*.
- Ptyalism in pregnancy, 204
- Puerperal convulsions. See *Eclampsia*.
- Puerperal fever. See *Septicæmia*.
- Puerperal mania. See *Insanity*.
- Puerperal state, 546
 - after-treatment in, 556

Puerperal state—

- diet and regimen in, 554
- diminution of uterus in, 549
- importance of prolonged rest in, 556
- pulse in, 547
- secretions and excretions in, 548
- temperature in, 548

Pulmonary arteries, anatomical arrangement of, as favoring thrombosis, 624

QUICKENING, 149

Quinine as an oxytocic, 343

RACE, as influencing the size of the fetal skull, 122

Recto-vaginal fistula, 439

Respiration of fetus, 123

Retroversion of the gravid uterus, 217

Rickets as a cause of pelvic deformity, 376

Rosenmüller, organ of. See Parovarium.

Round ligaments of the uterus, 70

Rules for monthly nurses, 556

Rupture of uterus. See Uterus.

SACRUM, anatomy of, 35

mechanical relations, 35

Salivation in pregnancy, 204

Scarlet fever affecting the fetus, 238

in pregnancy, 221

in the puerperal state, 600

Scoliotic deformity of pelvis, 379

Scybake in the rectum obstructing labor, 360

Septicæmia (puerperal), 593

bacteria in, 605

channels of diffusion in, 605

through which septic matter may be absorbed, 597

cold in treatment of, 619

conduct of practitioner in regard to, 604

contagion from other puerperal patients as a cause of, 602

description of, 609

division into auto-genetic and hetero-genetic forms, 598

epidemics of, 595

history of, 593

importance of antiseptic precautions in, 604

influence of cadaveric poison as a cause of, 598

influence of zymotic disease in causing, 600

its connection with pelvic cellulitis and peritonitis, 645

local changes in, 605

malarial, 615

mode in which the poison may be conveyed to patients in, 603

mortality in lying-in hospitals, 594

Septicæmia (puerperal)—

nature of septic poison, 605

pathological phenomena in, 607

prevention of, 604

pyæmic forms of, 614

sewer gas as a source of infection, 601

sources of auto-infection in, 598

of hetero-infection, 598

symptoms of the intense forms, 611

theory of an essential zymotic fever, 595

of identity with surgical septicæmia, 596

of local origin, 595

transfusion of blood in, 536

treatment of, 615

venesection in, 618

Warburg's tincture in the treatment of, 619

Sex, discovery of, of fetus during pregnancy, 153

of fetus as influencing the size of the skull, 122

Shoulder presentations, 322

diagnosis of, 325

division of, 323

mechanism of, 327

prognosis and frequency of, 325

spontaneous version in, 327

spontaneous evolution in, 328

treatment of, 329

Siamese twins, how born, 368

Sickness of pregnancy, 145

Smallpox affecting the fetus, 237

in pregnancy, 221

Smith's, Tyler, theory of labor, 257

Spondylolisthesis, 381 [382]

Spondylolizema, 382

Spontaneous evolution, 328

version, 327

Spurious pregnancy, 159

diagnosis of, 160

symptoms of, 159

[Stethoscope, Cammann's, 155]

Stillbirth, apparent, 557

Symphyseotomy, 527

[in Italy, revival of, 529]

Syncope during or after labor, 636

in pregnancy, 205

[postural treatment of, 426]

Syphilis affecting the fetus, 238

as a cause of abortion, 247

in pregnancy, 222

Sugar, in urine of pregnancy, 143

Super-fecundation and super-fetation, 169

Sutures of fetal head, 120

TEMPERATURE after delivery, 548

Thrombosis (peripheral venous), 637

changes in thrombi in, 641

condition of the affected limb, 638

detachment of emboli in, 641

history and pathology of, 639

progress of the disease, 638

Thrombosis (peripheral venous)—
 symptoms of, 637
 treatment of, 641
 (puerperal), 621
 arterial thrombosis and embolism, 633
 cardiac murmur in pulmonary, 629
 cases illustrating recovery from pulmonary, 627-629
 causes of death in pulmonary, 630
 clinical facts in favor of pulmonary, 625
 conditions which favor thrombosis in the puerperal state, 623
 distinction between thrombosis and embolism, 624
 phlegmasia dolens a consequence of, 622
 post-mortem appearance of clots in pulmonary, 630
 pulmonary, as a cause of pleuro-pneumonia, 632
 question of primary thrombosis in the pulmonary arteries, 624
 question of recovery from pulmonary, 626
 symptoms of arterial, 633
 of pulmonary obstruction in, 626
 treatment of arterial, 634
 of pulmonary, 631
Thrombosis of uterine vessels, 416
Thrombus. See Hematocoele.
Toothache in pregnancy, 204
Transfusion of blood, 534
 addition of chemical reagents to prevent coagulation of fibrin, 538
 cases suitable for the operation, 540
 dangers of the operation, 539
 defibrination of blood in, 543
 difficulties of the operation, 536
 effects of successful transfusion, 545
 history of the operation, 535
 immediate transfusion, 537
 method of injecting defibrinated blood, 544
 method of performing immediate transfusion, 541
 method of preparing defibrinated blood, 544
 nature and object of the operation, 536
 Schäfer's directions for immediate, 541
 secondary effects of, 545
 statistical results of, 539
Tropics, influence of residence in, on labor, 339
Trunk, presentation of. See Shoulder Presentation.
Tumors, diagnosis of uterine and ovarian, 158
 fetal, 239
 obstructing labor, 373
 (maternal) obstructing delivery, 358
Tunica albuginea, 74
Turning, 457

Turning—
 after perforation, 505
 anaesthesia in, 462
 [bi-manual, of Hicks, in placenta prævia, 411]
 by combined method, 462
 by external manipulation only, 459
 cases suitable for the operation, 459
 operating by combined method, 458
 cephalic, 457
 choice of hand to be used, 462
 history of the operation, 457
 in abdomino-anterior positions, 469
 in deformed pelvis, 396
 in placenta prævia, 408, 468
 method of cephalic, 460
 of performing by external manipulation, 459
 of podalic, 466
 object and nature of the operation, 458
 period when the operation should be performed, 462
 podalic, 461, 465
 position of patient in, 461
 statistics and dangers of, 458
 value of anaesthetics in difficult cases of, 470
Twins. See Plural Births.
Twins [Carolina, how born, 370]
 [Carolina, manner of birth of, 370]
 conjoined, 367
 [Hungarian, manner of birth of, 370]
 locked, 364
 [Siamese, manner of birth of, 368]

UMBILICAL cord, 117
 knots of, 117, 235
 ligature of, 289
 pathology of, 235
 prolapse of, 332
 cause of, 333
 prolapse of, diagnosis of, 333
 frequency of, 332
 postural treatment of, 334
 prognosis of, 333
 reposition of, 335
 treatment by laceration, 289
Umbilical souffle, 155
 vesicle, 107
Urachus, 108
Uræmia, in connection with eclampsia, 576
 in connection with puerperal insanity, 586
Urethra, 51
Urine, changes in, during pregnancy, 142
 retention of, after delivery, 553
Uterine fluctuation, as a sign of pregnancy, 151
 souffle, 155
Utero-sacral ligaments, 70
Uterus, 56

Uterus—

analogy of interior of, after delivery, and stump of an amputated limb, 105
 anomalies of, 67
 ante-partum hour-glass contraction, 355
 arrangement of muscular fibres of, 61
 axis of, during pregnancy, 135
 changes in cervix during pregnancy, 135, 136
 changes in form and dimensions of, during pregnancy, 133
 changes in mucous membranes of, after delivery, 550
 changes in mucous membranes of, after impregnation, 100
 changes in tissues of, during pregnancy, 138
 changes in the vessels of, after delivery, 550
 congestive hypertrophy of, 158
 contractions of, in labor, 259
 dimensions of, 58
 diminution in size of, after delivery, 549
 distension of, as a cause of labor, 256
 distension of, by retained menses, 158
 fatty transformation of, after delivery, 550
 hour-glass contraction, 418
 intermittent contractions of, during pregnancy, 150
 internal surface of, 59
 inversion of, 442
 [inverted, spontaneous reposition of, 447]
 [painful intermittent contractions of threatening abortion, 150]
 [partitioned, 68]
 differential diagnosis of, 444
 production of, 444
 results of physical examination in, 443
 [rupture of, rational treatment of, 441]
 symptoms of, 443
 treatment of, 446
 ligaments of, 68
 lymphatics of, 65
 malposition of, as a cause of protracted labor, 341
 mode of action in labor, 259
 mucous membrane of, 61
 muscular fibres of, 61
 nerves of, 67
 regional divisions of, 58
 relations of, 56
 retroversion of gravid, 217

Uterus—

rupture of, 431
 alterations of tissues in, 433
 causes of, 432
 comparative result of various methods of treatment in, 438
 prognosis of, 436
 seat of laceration in, 432
 symptoms of, 435
 treatment of, 437
 gastrotomy in, 438
 size of, at various periods of pregnancy, 134
 state of, in protracted labor, 339
 structures composing, 60
 utricular glands of, 62
 vessels of, 65
 weight of, after delivery, 550

VAGINA, 54

bands and cicatrices of, obstructing delivery, 353
 contraction of, after delivery, 551
 lacerations of, 439
 orifice of, 52
 structure of, 54
 Varicose veins in pregnancy, 215
 Vectis, 495
 action of, 496
 cases in which it is applicable, 496
 Veins, entrance of air into, as a cause of sudden death after delivery, 636
 Venesection for rigidity of cervix, 352
 Version. See *Turning*.
 Vesico-uterine ligaments, 71
 Vesico-vaginal fistula, 439
 Vestibule, 51
 Vicarious menstruation, 92
 Vomiting in pregnancy, 198
 Vulva, 48
 condition of, after delivery, 551
 œdema of, obstructing labor, 360
 vascular supply of, 53
 Vulvo-vaginal glands, 52

WARBURG'S tincture, 619

Weaning. See *Lactation*.

Wet-nurse, selection of, 561

Wolfian bodies, 66, 118

[Womanhood, precocious physical, 86]

[Woman, puerperal, treatment of, 618]

Wounds of the fœtus, 240

ZONA pellucida, 77

Zymotic disease, affecting the fœtus, 238
 as a cause of septicæmia, 600



THE AMERICAN JOURNAL of the MEDICAL SCIENCES.

(Continued from first page.)

But it is not only in the *Original Department* that the JOURNAL of the future will seek to eclipse all its efforts in the past. The mass of contributions to medical literature and science increases with such rapidity, that if the reader is to keep abreast with them the matter must be carefully sifted, and arranged so as to enable him to grasp it understandingly with the least possible expenditure of time. In the *Bibliographical Department*, therefore, separate reviews are devoted only to works of exceptional importance. As a rule, new books are considered in groups of cognate subjects, the reviewer setting forth tersely the merits of the individual volumes with a condensed statement of the views of the authors. In this manner the reader is kept advised of the products of the press in the most convenient manner.

A similar plan is adopted in the *Quarterly Summary of Progress*. The various branches of medical science have been assigned to the following gentlemen, who will furnish well-digested *résumés* of progress, paying special attention to clinical application: Anatomy, George D. Thane, M. R. C. S.; Physiology, Gerald F. Yeo, M. D.; Materia Medica, Therapeutics and Pharmacology, Roberts Bartholow, M. D., LL. D.; Medicine, William Osler, M. D.; Surgery, in America, R. J. Hall, M. D.; in Europe, Frederick Treves, F. R. C. S.; Ophthalmology, L. Webster Fox, M. D.; Otology, Charles H. Burnett, M. D.; Laryngology, J. Solis Cohen, M. D.; Dermatology, Louis A. Duhring, M. D., and H. W. Stelwagon, M. D.; Midwifery and Gynecology, D. Berry Hart, M. D.; Jurisprudence, Matthew Hay, M. D.; Public Health, Shirley F. Murphy, M. R. C. S.

The publishers feel an honest pride in thus being the means of bringing together the professions of the two great English-speaking peoples, and in laying before them a periodical which must be universally recognized as marking a new era in medical progress. Believing that it will be regarded as indispensable by all intelligent physicians on both sides of the Atlantic, they feel themselves warranted, by the expectation of a large increase in circulation, in maintaining the present very moderate subscription price, notwithstanding the greatly augmented expenditure entailed by the change.

The JOURNAL will continue to be published quarterly, as heretofore, on the first of January, April, July and October.

Price, FIVE DOLLARS Per Annum, in Advance.

THE MEDICAL NEWS.

A National Weekly Periodical, containing 28 to 32 Quarto Pages in Each Issue.

THE continually increasing appreciation of THE MEDICAL NEWS by the profession throughout the country, is a most gratifying recognition of the policy pursued by the managers of this journal in their unceasing efforts to enhance its value to the practitioner.

Possessing a most efficient organization THE NEWS unites the best features of the medical magazine and newspaper. Its large and able Editorial Staff discusses in each issue the important topics of the day in a thoughtful and scholarly manner, while its corps of qualified reporters and correspondents, covering every medical centre, insures that its readers shall be promptly and thoroughly posted upon all matters of interest in the world of medicine. On account of the position conceded to THE NEWS, it has become the medium chosen by the leading minds of the profession for the publication of their most important contributions to medical science. The valuable instruction afforded in Clinical Lectures, and the rich experience gained in the leading Hospitals of the world are constantly laid before the readers of THE NEWS, while prompt and authentic reports of Society Proceedings are received from special reporters in various sections of the country by mail and Telegraph. In the pages devoted to the progress of Medical Science are found early notes of all important advances, gleaned from the principal journals of both hemispheres. Ample space is devoted to Reviews, News Items, Correspondence and Notes and Queries. In short, every branch of medicine is adequately represented in THE NEWS, and the details of plan and typography have been carefully studied in order to economize the time and secure the comfort of the reader in every possible way.

Price, FIVE DOLLARS Per Annum, in Advance.

COMMUTATION RATE.

To subscribers paying in advance for 1887:—

AMERICAN JOURNAL OF THE MEDICAL SCIENCES (quarterly)	} To one address for \$9.00
THE MEDICAL NEWS (weekly)	


SPECIAL OFFERS.

Advance-paying subscribers to either or both of the above-named periodicals may take advantage of any one of the following offers:

(1). THE MEDICAL NEWS VISITING LIST for 1887, dated, either for 30 patients per week (1 vol.), or for 60 patients (2 vols.), or for 90 patients (3 vols.), will be sent

post-paid on receipt of 75 cents per volume (regular price \$1.25). (2). **THE YEAR-BOOK OF TREATMENT** for 1886 will be sent on receipt of 75 cents (regular price \$1.25). (3). An *advance* remittance of \$10 will procure **THE NEWS** and **THE JOURNAL** for one year, together with any one volume of **THE MEDICAL NEWS VISITING LIST** and **THE YEAR-BOOK OF TREATMENT** for 1886, as offered above. Subsequent volumes of **THE MEDICAL NEWS VISITING LIST** may be procured by *advance*-paying subscribers for 75 cents each (regular price \$1.25). Thumb-letter Index for quick use, 25 cents additional.

Subscribers can obtain, at the close of each volume, cloth covers for THE JOURNAL (one annually), and for THE NEWS (one annually), free by mail, by remitting Ten Cents for the JOURNAL cover, and Fifteen Cents for the NEWS cover.

 The safest mode of remittance is by bank check or postal money order, drawn to the order of the undersigned; where these are not accessible, remittances for subscriptions may be sent at the risk of the publishers by forwarding in *registered* letters. Address,

LEA BROTHERS & CO., 706 and 708 Sansom Street, PHILADELPHIA.

THE MEDICAL NEWS VISITING LIST FOR 1887.

Containing Calendar for two years. Obstetric diagrams. Scheme of Dentition. Tables of weights and measures and comparative scales. Instructions for examining the urine. List of disinfectants. Table of eruptive fevers. Lists of new remedies and remedies not generally used, Incompatibles, Poisons and Antidotes. Artificial respiration. Table of doses, prepared to accord with the last revision of the U. S. Pharmacopoeia, an extended table of Diseases and their remedies, and directions for Ligation of Arteries. Blanks for all records of practice and Erasable tablet. Handsomely bound in limp Morocco, with tuck, pencil, rubber and catheter scale.

In response to numerous requests, **THE MEDICAL NEWS VISITING LIST** for 1887 is issued in three sizes, viz.: Dated, for 30 patients per week, 1 vol.; dated, for 60 patients, 2 vols.; dated, for 90 patients, 3 vols.; Price per volume, \$1.25. Also, furnished with Ready Reference Thumb-letter Index for quick use, 25 cents additional. For special offers, including **VISITING LIST**, see above.

THE MEDICAL NEWS PHYSICIANS' LEDGER.

Containing 400 pages of fine linen "ledger" paper, ruled so that all the accounts of a large practice may be conveniently kept in it, either by single or double entry, for a long period. Strongly bound in leather, with cloth sides, and with a patent flexible back, which permits it to lie perfectly flat when opened at any place. Price, \$5.00. Also, a small special lot of same Ledger, with 300 pages. Price, \$4.00.

HARTSHORNE, HENRY, A. M., M. D., LL. D.,

Lately Professor of Hygiene in the University of Pennsylvania.

A Conspectus of the Medical Sciences; Containing Handbooks on Anatomy, Physiology, Chemistry, Materia Medica, Practice of Medicine, Surgery and Obstetrics. Second edition, thoroughly revised and greatly improved. In one large royal 12mo. volume of 1028 pages, with 477 illustrations. Cloth, \$4.25; leather, \$5.00.

The object of this manual is to afford a convenient work of reference to students during the brief moments at their command while in attendance upon medical lectures. It is a favorable sign that it has been found necessary, in a short space of time, to issue a new and carefully revised edition. The illustrations are very numerous and unusually clear, and each part seems to have received its due share of attention. We can conceive such a work to be useful, not only to students, but to practitioners as well. It reflects credit upon the

industry and energy of its able editor.—*Boston Medical and Surgical Journal*, Sept. 3, 1874.

We can say with the strictest truth that it is the best work of the kind with which we are acquainted. It embodies in a condensed form all recent contributions to practical medicine, and is therefore useful to every busy practitioner throughout our country, besides being admirably adapted to the use of students of medicine. The book is faithfully and ably executed.—*Charleston Medical Journal*, April, 1875.

NEILL, JOHN, M. D., and SMITH, F. G., M. D.,

Late Surgeon to the Penna. Hospital.

Prof. of the Institutes of Med. in the Univ. of Penna.

An Analytical Compendium of the Various Branches of Medical Science, for the use and examination of Students. A new edition, revised and improved. In one large royal 12mo. volume of 974 pages, with 374 woodcuts. Cloth, \$4; leather, \$4.75.

LUDLOW, J. L., M. D.,

Consulting Physician to the Philadelphia Hospital, etc.

A Manual of Examinations upon Anatomy, Physiology, Surgery, Practice of Medicine, Obstetrics, Materia Medica, Chemistry, Pharmacy and Therapeutics. To which is added a Medical Formulary. 3d edition, thoroughly revised, and greatly enlarged. In one 12mo. volume of 816 pages, with 370 illustrations. Cloth, \$3.25; leather, \$3.75.

The arrangement of this volume in the form of question and answer renders it especially suitable for the office examination of students and for the examination of students.

DUNGLISON, ROBLEY, M. D.,*Late Professor of Institutes of Medicine in the Jefferson Medical College of Philadelphia.*

MEDICAL LEXICON; A Dictionary of Medical Science: Containing a concise Explanation of the various Subjects and Terms of Anatomy, Physiology, Pathology, Hygiene, Therapeutics, Pharmacology, Pharmacy, Surgery, Obstetrics, Medical Jurisprudence and Dentistry, Notices of Climate and of Mineral Waters, Formulæ for Official, Empirical and Dietetic Preparations, with the Accentuation and Etymology of the Terms, and the French and other Synonymes, so as to constitute a French as well as an English Medical Lexicon. Edited by RICHARD J. DUNGLISON, M.D. In one very large and handsome royal octavo volume of 1139 pages. Cloth, \$6.50; leather, raised bands, \$7.50; very handsome half Russia, raised bands, \$8.

The object of the author, from the outset, has not been to make the work a mere lexicon or dictionary of terms, but to afford under each word a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of medical science. Starting with this view, the immense demand which has existed for the work has enabled him, in repeated revisions, to augment its completeness and usefulness, until at length it has attained the position of a recognized and standard authority wherever the language is spoken. Special pains have been taken in the preparation of the present edition to maintain this enviable reputation. The additions to the vocabulary are more numerous than in any previous revision, and particular attention has been bestowed on the accentuation, which will be found marked on every word. The typographical arrangement has been greatly improved, rendering reference much more easy, and every care has been taken with the mechanical execution. The volume now contains the matter of at least four ordinary octavos.

About the first book purchased by the medical student is the Medical Dictionary. The lexicon explanatory of technical terms is simply a *sine qua non*. In a science so extensive and with such collaterals as medicine, it is as much a necessity also to the practising physician. To meet the wants of students and most physicians the dictionary must be condensed while comprehensive, and practical while perspicacious. It was because Dunglison met these indications that it became at once the dictionary of general use wherever medicine was studied in the English language. In no former revision have the alterations and additions been so great. The chief terms have been set in black letter, while the derivatives follow in small caps; an arrangement which greatly facilitates reference. —*Cincinnati Lancet and Clinic*, Jan. 10, 1874.

A book of which every American ought to be proud. When the learned author of the work

passed away, probably all of us feared lest the book should not maintain its place in the advancing science whose terms it defines. Fortunately, Dr. Richard J. Dunglison, having assisted his father in the revision of several editions of the work, and having been, therefore, trained in the methods and imbued with the spirit of the book, has been able to edit it as a work of the kind should be edited—to carry it on steadily, without jar or interruption, along the grooves of thought it has travelled during its lifetime. To show the magnitude of the task which Dr. Dunglison has assumed and carried through, it is only necessary to state that more than six thousand new subjects have been added in the present edition. —*Philadelphia Medical Times*, Jan. 3, 1874.

It has the rare merit that it certainly has no rival in the English language for accuracy and extent of references. —*London Medical Gazette*.

HOBLYN, RICHARD D., M. D.

A Dictionary of the Terms Used in Medicine and the Collateral Sciences. Revised, with numerous additions, by ISAAC HAYS, M. D., late editor of *The American Journal of the Medical Sciences*. In one large royal 12mo. volume of 520 double-columned pages. Cloth, \$1.50; leather, \$2.00.

It is the best book of definitions we have, and ought always to be upon the student's table. —*Southern Medical and Surgical Journal*.

STUDENTS' SERIES OF MANUALS.

A Series of Fifteen Manuals, for the use of Students and Practitioners of Medicine and Surgery, written by eminent Teachers or Examiners, and issued in pocket-size 12mo. volumes of 300-540 pages, richly illustrated and at a low price. The following volumes are now ready: TREVES' *Manual of Surgery*, by various writers, in three volumes, each, \$2; BELL'S *Comparative Physiology and Anatomy*, \$2; GOULD'S *Surgical Diagnosis*, \$2; ROBERTSON'S *Physiological Physics*, \$2; BRUCE'S *Materia Medica and Therapeutics*, \$1.50; POWER'S *Human Physiology*, \$1.50; CLARKE and LOCKWOOD'S *Dissectors' Manual*, \$1.50; RALFE'S *Clinical Chemistry*, \$1.50; TREVES' *Surgical Applied Anatomy*, \$2; PEPPER'S *Surgical Pathology*, \$2; and KLEIN'S *Elements of Histology*, \$1.50. The following are in press: BELLAMY'S *Operative Surgery*, PEPPER'S *Forensic Medicine*, and CURNOW'S *Medical Applied Anatomy*. For separate notices see index on last page.

SERIES OF CLINICAL MANUALS.

In arranging for this Series it has been the design of the publishers to provide the profession with a collection of authoritative monographs on important clinical subjects in a cheap and portable form. The volumes will contain about 550 pages and will be freely illustrated by chromo-lithographs and woodcuts. The following volumes are now ready: HUTCHINSON on *Syphilis*, \$2.25; MARSH on the *Joints*, \$2; OWEN on *Surgical Diseases of Children*, \$2; MORRIS on *Surgical Diseases of the Kidney*, \$2.25; PICK on *Fractures and Dislocations*, \$2; BUTLIN on the *Tongue*, \$3.50; TREVES on *Intestinal Obstruction*, \$2; and SAVAGE on *Insanity and Allied Neuroses*, \$2. The following are in active preparation: CARTER and FROST on *Ulnic Surgery*, BRYANT on the *Breast*, BROADBENT on the *Pulse*, LUCAS on *Urethra*, and BALL on the *Rectum and Anus*. For separate notices see index

GRAY, HENRY, F. R. S.,*Lecturer on Anatomy at St. George's Hospital, London.*

Anatomy, Descriptive and Surgical. The Drawings by H. V. CARTER, M. D., and Dr. WESTMACOTT. The dissections jointly by the AUTHOR and Dr. CARTER. With an Introduction on General Anatomy and Development by T. HOLMES, M. A., Surgeon to St. George's Hospital. Edited by T. PICKERING PICK, F. R. C. S., Surgeon to and Lecturer on Anatomy at St. George's Hospital, London, Examiner in Anatomy, Royal College of Surgeons of England. A new American from the eleventh enlarged and improved London edition, thoroughly revised and re-edited by WILLIAM W. KEEN, M. D., Professor of Anatomy in the Pennsylvania Academy of the Fine Arts, etc. To which is added the second American from the latest English edition of **LANDMARKS, MEDICAL AND SURGICAL**, by LUTHER HOLDEN, F. R. C. S. In one imperial octavo volume of about 1100 pages, with about 650 large and elaborate engravings on wood. Price of edition in black (see below): Cloth, \$6; leather, \$7; half Russia, \$7.50. The price of edition in colors will be announced shortly.

This work covers a more extended range of subjects than is customary in the ordinary text-books, giving not only the details necessary for the student, but also the application to those details to the practice of medicine and surgery. It thus forms both a guide for the learner and an admirable work of reference for the active practitioner. The engravings form a special feature in the work, many of them being the size of nature, nearly all original, and having the names of the various parts printed on the body of the cut, in place of figures of reference with descriptions at the foot. In this edition a new departure will be taken by the issue of the work with the arteries, veins and nerves distinguished by different colors. The engravings thus form a complete and splendid series, which will greatly assist the student in forming a clear idea of Anatomy, and will also serve to refresh the memory of those who may find in the exigencies of practice the necessity of recalling the details of the dissecting-room. Combining, as it does, a complete Atlas of Anatomy with a thorough treatise on systematic, descriptive and applied Anatomy, the work will be found of great service to all physicians who receive students in their offices, relieving both preceptor and pupil of much labor in laying the groundwork of a thorough medical education.

For the convenience of those who cannot afford the slight increase in cost necessitated by the use of colors, the volume will be published also in black alone, and maintained in this style at the price of former editions, notwithstanding the largely increased size of the work.

Landmarks, Medical and Surgical, by the distinguished Anatomist, Mr. Luther Holden, has been appended to the present edition as it was to the previous one. This work gives in a clear, condensed and systematic way all the information by which the practitioner can determine from the external surface of the body the position of internal parts. Thus complete, the work, it is believed, will furnish all the assistance that can be rendered by type and illustration in anatomical study.

ALSO FOR SALE SEPARATE—**HOLDEN, LUTHER, F. R. C. S.,***Surgeon to St. Bartholomew's and the Foundling Hospitals, London.*

Landmarks, Medical and Surgical. Second American from the latest revised English edition, with additions by W. W. KEEN, M. D., Professor of Artistic Anatomy in the Pennsylvania Academy of the Fine Arts, formerly Lecturer on Anatomy in the Philadelphia School of Anatomy. In one handsome 12mo. volume of 148 pages. Cloth, \$1.00.

This little book is all that can be desired within its scope, and its contents will be found simply invaluable to the young surgeon or physician, since they bring before him such data as he requires at every examination of a patient. It is written in language so clear and concise that one ought almost to learn it by heart. It teaches diagnosis by external examination, ocular and palpable, of the body, with such anatomical and physiological facts as directly bear on the subject. It is eminently the student's and young practitioner's book.—*Physician and Surgeon*, Nov. 1881.

The study of these Landmarks by both physi-

cians and surgeons is much to be encouraged. It inevitably leads to a progressive education of both the eye and the touch, by which the recognition of disease or the localization of injuries is vastly assisted. One thoroughly familiar with the facts here taught is capable of a degree of accuracy and a confidence of certainty which is otherwise unattainable. We cordially recommend the Landmarks to the attention of every physician who has not yet provided himself with a copy of this useful, practical guide to the correct placing of all the anatomical parts and organs.—*Canada Medical and Surgical Journal*, Dec. 1881.

WILSON, ERASMUS, F. R. S.

A System of Human Anatomy, General and Special. Edited by W. H. GOBRECHT, M. D., Professor of General and Surgical Anatomy in the Medical College of Ohio. In one large and handsome octavo volume of 616 pages, with 397 illustrations. Cloth, \$4.00; leather, \$5.00.

CLELAND, JOHN, M. D., F. R. S.,*Professor of Anatomy and Physiology in Queen's College, Galway.*

A Directory for the Dissection of the Human Body. In one 12mo. volume of 178 pages. Cloth, \$1.25.

ALLEN, HARRISON, M. D.,*Professor of Physiology in the University of Pennsylvania.*

A System of Human Anatomy, Including Its Medical and Surgical Relations. For the use of Practitioners and Students of Medicine. With an Introductory Section on Histology. By E. O. SHAKESPEARE, M. D., Ophthalmologist to the Philadelphia Hospital. Comprising 813 double-columned quarto pages, with 380 illustrations on 109 full page lithographic plates, many of which are in colors, and 241 engravings in the text. In six Sections, each in a portfolio. Section I. HISTOLOGY. Section II. BONES AND JOINTS. Section III. MUSCLES AND FASCLE. Section IV. ARTERIES, VEINS AND LYMPHATICS. Section V. NERVOUS SYSTEM. Section VI. ORGANS OF SENSE, OF DIGESTION AND GENITO-URINARY ORGANS, EMBRYOLOGY, DEVELOPMENT, TERATOLOGY, SUPERFICIAL ANATOMY, POST-MORTEM EXAMINATIONS, AND GENERAL AND CLINICAL INDEXES. Price per Section, \$3.50; also bound in one volume, cloth, \$23.00; very handsome half Russia, raised bands and open back, \$25.00. *For sale by subscription only. Apply to the Publishers.*

Extract from Introduction.

It is the design of this book to present the facts of human anatomy in the manner best suited to the requirements of the student and the practitioner of medicine. The author believes that such a book is needed, inasmuch as no treatise, as far as he knows, contains, in addition to the text descriptive of the subject, a systematic presentation of such anatomical facts as can be applied to practice.

A book which will be at once accurate in statement and concise in terms; which will be an acceptable expression of the present state of the science of anatomy; which will exclude nothing that can be made applicable to the medical art, and which will thus embrace all of surgical importance, while omitting nothing of value to clinical medicine,—would appear to have an excuse for existence in a country where most surgeons are general practitioners, and where there are few general practitioners who have no interest in surgery.

It is to be considered a study of applied anatomy in its widest sense—a systematic presentation of such anatomical facts as can be applied to the practice of medicine as well as of surgery. Our author is concise, accurate and practical in his statements, and succeeds admirably in infusing an interest into the study of what is generally considered a dry subject. The department of Histology is treated in a masterly manner, and the ground is travelled over by one thoroughly familiar with it. The illustrations are made with great

care, and are simply superb. There is as much of practical application of anatomical points to the every-day wants of the medical clinician as to those of the operating surgeon. In fact, few general practitioners will read the work without a feeling of surprised gratification that so many points, concerning which they may never have thought before are so well presented for their consideration. It is a work which is destined to be the best of its kind in any language.—*Medical Record*, Nov. 25, 1882.

CLARKE, W. B., F.R.C.S. & LOCKWOOD, C. B., F.R.C.S.*Demonstrators of Anatomy at St. Bartholomew's Hospital Medical School, London.*

The Dissector's Manual. In one pocket-size 12mo. volume of 396 pages, with 49 illustrations. Limp cloth, red edges, \$1.50. See *Students' Series of Manuals*, page 4.

Meers, Clarke and Lockwood have written a book that can hardly be rivalled as a practical aid to the dissector. Their purpose, which is "how to describe the best way to display the anatomical structure," has been fully attained. They excel in a lucidity of demonstration and graphic terseness of expression, which only a long training and

intimate association with students could have given. With such a guide as this, accompanied by so attractive a commentary as Treves' *Surgical Applied Anatomy* (same series), no student could fail to be deeply and absorbingly interested in the study of anatomy.—*New Orleans Medical and Surgical Journal*, April, 1884.

TREVES, FREDERICK, F. R. C. S.,*Senior Demonstrator of Anatomy and Assistant Surgeon at the London Hospital.*

Surgical Applied Anatomy. In one pocket-size 12mo. volume of 540 pages, with 61 illustrations. Limp cloth, red edges, \$2.00. See *Students' Series of Manuals*, page 4.

He has produced a work which will command a larger circle of readers than the class for which it was written. This class of a thorough, practical acquaintance with these fundamental branches,

quickened by daily use as a teacher and practitioner, has enabled our author to prepare a work which it would be a most difficult task to equal.—*The American Practitioner*, Feb. 1884.

CURNOW, JOHN, M. D., F. R. C. P.,*Professor of Anatomy at King's College, Physician at King's College Hospital.*

Medical Applied Anatomy. In one pocket-size 12mo. volume. Papering. See *Students' Series of Manuals*, page 4.

BELLAMY, EDWARD, F. R. C. S.,*Senior Assistant-Surgeon at the Charing-Cross Hospital, London.*

The Student's Guide to Surgical Anatomy: Being a Description of the most Important Surgical Regions of the Human Body, and intended as an Introduction to operative Surgery. In one 12mo. volume of 300 pages, with 50 illustrations. Cloth, \$2.25.

HARTSHORN'S HANDBOOK OF ANATOMY AND PHYSIOLOGY. Second edition, revised. In one royal 12mo. volume of 326 pages, with 226 woodcuts. Cloth, \$1.75.

BOEKER'S SPECIAL ANATOMY AND PHYSIOLOGY. Eighth edition, extensively revised and modified. In two octavo volumes of 236 pages, with 224 woodcuts. Cloth, \$5.50.

DRAPER, JOHN C., M. D., LL. D.,*Professor of Chemistry in the University of the City of New York.***Medical Physics.** A Text-book for Students and Practitioners of Medicine. In one octavo volume of 734 pages, with 376 woodcuts, mostly original. Cloth, \$4.**FROM THE PREFACE.**

The fact that a knowledge of Physics is indispensable to a thorough understanding of Medicine has not been as fully realized in this country as in Europe, where the admirable works of Desplats and Gariel, of Robertson and of numerous German writers constitute a branch of educational literature to which we can show no parallel. A full appreciation of this the author trusts will be sufficient justification for placing in book form the substance of his lectures on this department of science, delivered during many years at the University of the City of New York.

Broadly speaking, this work aims to impart a knowledge of the relations existing between Physics and Medicine in their latest state of development, and to embody in the pursuit of this object whatever experience the author has gained during a long period of teaching this special branch of applied science.

This elegant and useful work bears ample testimony to the learning and good judgment of the author. He has fitted his work admirably to the exigencies of the situation by presenting the reader with brief, clear and simple statements of such propositions as he is by necessity required to master. The subject matter is well arranged, liberally illustrated and carefully indexed. That it will take rank at once among the text-books is certain, and it is to be hoped that it will find a place upon the shelf of the practical physician, where, as a book of reference, it will be found useful and agreeable.—*Louisville Medical News*, September 26, 1885.

Certainly we have no text-book as full as the excellent one he has prepared. It begins with a statement of the properties of matter and energy. After these the special departments of physics are

explained, acoustics, optics, heat, electricity and magnetism, closing with a section on electrobiology. The applications of all these to physiology and medicine are kept constantly in view. The text is amply illustrated and the many difficult points of the subject are brought forward with remarkable clearness and ability.—*Medical and Surgical Reporter*, July 18, 1885.

That this work will greatly facilitate the study of medical physics is apparent upon even a mere cursory examination. It is marked by that scientific accuracy which always characterizes Dr. Draper's writings. Its peculiar value lies in the fact that it is written from the standpoint of the medical man. Hence much is omitted that appears in a mere treatise on physical science, while much is inserted of peculiar value to the physician.—*Medical Record*, August 22, 1885.

ROBERTSON, J. MCGREGOR, M. A., M. B.,*Muirhead Demonstrator of Physiology, University of Glasgow.***Physiological Physics.** In one 12mo. volume of 537 pages, with 219 illustrations. Limp cloth, \$2.00. See *Students' Series of Manuals*, page 4.

The title of this work sufficiently explains the nature of its contents. It is designed as a manual for the student of medicine, an auxiliary to his text-book in physiology, and it would be particularly useful as a guide to his laboratory experi-

ments. It will be found of great value to the practitioner. It is a carefully prepared book of reference, concise and accurate, and as such we heartily recommend it.—*Journal of the American Medical Association*, Dec. 6, 1884.

DALTON, JOHN C., M. D.,*Professor Emeritus of Physiology in the College of Physicians and Surgeons, New York.***Doctrines of the Circulation of the Blood.** A History of Physiological Opinion and Discovery in regard to the Circulation of the Blood. In one handsome 12mo. volume of 293 pages. Cloth, \$2.

Dr. Dalton's work is the fruit of the deep research of a cultured mind, and to the busy practitioner it cannot fail to be a source of instruction. It will inspire him with a feeling of gratitude and admiration for those plodding workers of olden times, who laid the foundation of the magnificent temple of medical science as it now stands.—*New Orleans Medical and Surgical Journal*, Aug. 1885.

In the progress of physiological study no fact was of greater moment, none more completely

revolutionized the theories of teachers, than the discovery of the circulation of the blood. This explains the extraordinary interest it has to all medical historians. The volume before us is one of three or four which have been written within a few years by American physicians. It is in several respects the most complete. The volume, though small in size, is one of the most creditable contributions from an American pen to medical history that has appeared.—*Med. & Surg. Rep.*, Dec. 6, 1884.

BELL, F. JEFFREY, M. A.,*Professor of Comparative Anatomy at King's College, London.***Comparative Physiology and Anatomy.** In one 12mo. volume of 561 pages, with 229 illustrations. Limp cloth, \$2.00. See *Students' Series of Manuals*, page 4.

The manual is preëminently a student's book—clear and simple in language and arrangement. It is well and abundantly illustrated, and is readable and interesting. On the whole we consider

it the best work in existence in the English language to place in the hands of the medical student.—*Bristol Medico-Chirurgical Journal*, Mar., 1886.

ELLIS, GEORGE VINER,*Emeritus Professor of Anatomy in University College, London.***Demonstrations of Anatomy.** Being a Guide to the Knowledge of the Human Body by Dissection. From the eighth and revised London edition. In one very handsome octavo volume of 716 pages, with 249 illustrations. Cloth, \$4.25; leather, \$5.25.**ROBERTS, JOHN B., A. M., M. D.,***Prof. of Applied Anat. and Oper. Surg. in Phila. Polytechnic and Coll. for Graduates in Medicine.***The Compend of Anatomy.** For use in the dissecting-room and in preparing for examinations. In one 16mo. volu

75 cents.

CHAPMAN, HENRY C., M. D.,

Professor of Institutes of Medicine and Medical Jurisprudence in the Jefferson Medical College of Philadelphia.

A Treatise on Human Physiology. In one handsome octavo volume of about 1000 pages, profusely illustrated. *Shortly.*

DALTON, JOHN C., M. D.,

Professor of Physiology in the College of Physicians and Surgeons, New York, etc.

A Treatise on Human Physiology. Designed for the use of Students and Practitioners of Medicine. Seventh edition, thoroughly revised and rewritten. In one very handsome octavo volume of 722 pages, with 252 beautiful engravings on wood. Cloth, \$5.00; leather, \$6.00; very handsome half Russia, raised bands, \$6.50.

This edition of Dr. Dalton's well-known work bears evidence of having been thoroughly and carefully revised. From the first appearance of the book it has been a favorite, owing as well to the author's renown as an oral teacher as to the charm of simplicity with which, as a writer, he always succeeds in venturing even intricate subjects. It must be gratifying to him to observe the frequency with which his work, written for students and practitioners, is quoted by other writers on physiology. This fact attests its value, and, in great measure, its originality. It now needs no such seal of approbation, however, for the thousands who have studied it in its various editions have never been in any doubt as to its sterling worth.—*N. Y. Medical Journal and Obstetrical Review*, Oct. 1882.

Professor Dalton's well-known and deservedly appreciated work has long passed the stage at

which it could be reviewed in the ordinary sense. The work is eminently one for the medical practitioner, since it treats most fully of those branches of physiology which have a direct bearing on the diagnosis and treatment of disease. The work is one which we can highly recommend to all our readers.—*Dublin Journal of Medical Science*, Feb. '83.

Certainly no physiological work as ever issued from the press that presented its subject-matter in a clearer and more attractive light. Almost every page bears evidence of the exhaustive revision that has taken place. The material is placed in a more compact form, yet its delightful charm is retained, and no subject is thrown into obscurity. Altogether this edition is far in advance of any previous one, and will tend to keep the profession posted as to the most recent additions to our physiological knowledge.—*Michigan Medical News*, April, 1882.

FOSTER, MICHAEL, M. D., F. R. S.,

Prefector in Physiology and Fellow of Trinity College, Cambridge, England.

Text-Book of Physiology. Third American from the fourth English edition, with notes and additions by E. T. REICHERT, M. D., Professor of Physiology in the University of Pennsylvania. In one handsome royal 12mo. volume of 908 pages, with 271 illustrations. Cloth, \$3.25; leather, \$3.75.

Dr. Foster's work upon physiology is so well-known as a text-book in this country, that it needs but little to be said in regard to it. There is scarcely a medical college in the United States where it is not in the hands of the students. The author, more than any other writer with whom we are acquainted, seems to understand what portions of the science are essential for students

to know and what may be passed over by them as not important. From the beginning to the end, physiology is taught in a systematic manner. To this third American edition numerous additions, corrections and alterations have been made, so that in its present form the usefulness of the book will be found to be much increased.—*Cincinnati Medical News*, July 1885.

POWER, HENRY, M. B., F. R. C. S.,

Examiner in Physiology, Royal College of Surgeons of England.

Human Physiology. In one handsome pocket-size 12mo. volume of 396 pages, with 47 illustrations. Cloth, \$1.50. See *Students' Series of Manuals*, page 4.

The prominent character of this work is that of judicious condensation, in which an able and successful effort appears to have been made by its accomplished author to teach the greatest number of facts in the fewest possible words. The result is a specimen of concentrated intellectual pabulum seldom surpassed, which ought to be carefully ingested and digested by every practitioner who desires to keep himself well informed upon this most progressive of the medical sciences. The volume is one which we cordially recommend

to every one of our readers.—*The American Journal of the Medical Sciences*, October, 1884.

This little work is deserving of the highest praise, and we can hardly conceive how the main facts of this science could have been more clearly or concisely stated. The price of the work is such as to place it within the reach of all, while the excellence of its text will certainly secure for it most favorable commendation.—*Cincinnati Lancet and Clinic*, Feb. 16, 1884.

SIMON, W., Ph. D., M. D.,

Professor of Chemistry and Toxicology in the College of Physicians and Surgeons, Baltimore, and Professor of Chemistry in the Maryland College of Pharmacy.

Manual of Chemistry. A Guide to Lectures and Laboratory work for Beginners in Chemistry. A Text-book, specially adapted for Students of Pharmacy and Medicine. In one 8vo. vol. of 410 pp., with 16 woodcuts and 7 plates, mostly of actual deposits, with colors illustrating 56 of the most important chemical reactions. Cloth, \$3.00; also without plates, cloth, \$2.50.

This book supplies a want long felt by students of medicine and pharmacy, and is a concise but thorough treatise on the subject. The long experience of the author as a teacher in schools of medicine and pharmacy is conspicuous in the perfect adaptation of the work to the special needs of the student of these branches. The colored

plates, beautifully executed, illustrating precipitates of various reactions, form a novel and valuable feature of the book, and cannot fail to be appreciated by both student and teacher as a help over the hard places of the science.—*Maryland Medical Journal*, Nov. 22, 1884.

Wöhler's Outlines of Organic Chemistry. Edited by FITTIG. Translated by IRA REMSEN, M. D., Ph. D. In one 12mo. volume of 550 pages. Cloth, \$3.

LEHMANN'S MANUAL OF CHEMICAL PHYSIOLOGY. In one octavo volume of 327 pages, with 41 illustrations. Cloth, \$2.25.

CARPENTER'S HUMAN PHYSIOLOGY. Edited

CARPENTER'S PRIZE ESSAY ON THE USE AND ABUSE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE. With explanations of scientific words. Small 12mo. 178 pages. Cloth, 60 cents.

FRANKLAND, E., D. C. L., F. R. S., & JAPP, F. R., F. I. C.,*Professor of Chemistry in the Normal School of Science, London.**Assist. Prof. of Chemistry in the Normal School of Science, London.*

Inorganic Chemistry. In one handsome octavo volume of 677 pages with 51 woodcuts and 2 plates. Cloth, \$3.75; leather, \$4.75.

This work should supersede other works of its class in the medical colleges. It is certainly better adapted than any work upon chemistry, with which we are acquainted, to impart that clear and full knowledge of the science which students of medicine should have. Physicians who feel that their chemical knowledge is behind the times, would do well to devote some of their leisure time to the study of this work. The descriptions and demonstrations are made so plain that there is no difficulty in understanding them.—*Cincinnati Medical News*, January, 1886.

This excellent treatise will not fail to take its place as one of the very best on the subject of which it treats. We have been much pleased with the comprehensive and lucid manner in which the difficulties of chemical notation and nomenclature have been cleared up by the writers. It shows on every page that the problem of rendering the obscurities of this science easy of comprehension has long and successfully engaged the attention of the authors.—*Medical and Surgical Reporter*, October 31, 1885.

FOWNES, GEORGE, Ph. D.

A Manual of Elementary Chemistry; Theoretical and Practical. Embodying WATTS' Physical Inorganic Chemistry. New American edition. In one large royal 12mo. volume of 1061 pages, with 168 illustrations on wood and a colored plate. Cloth, \$2.75; leather, \$3.25.

Fownes' Chemistry has been a standard text-book upon chemistry for many years. Its merits are very fully known by chemists and physicians everywhere in this country and in England. As the science has advanced by the making of new discoveries, the work has been revised so as to keep it abreast of the times. It has steadily maintained its position as a text-book with medical students. In this work are treated fully: Heat, Light and Electricity, including Magnetism. The influence exerted by these forces in chemical action upon health and disease, etc., is of the most important kind, and should be familiar to every medical practitioner. We can commend the work as one of the very best text-books upon

chemistry extant.—*Cincinnati Medical News*, October, 1885.

Of all the works on chemistry intended for the use of medical students, *Fownes' Chemistry* is perhaps the most widely used. Its popularity is based upon its excellence. This last edition contains all of the material found in the previous, and it is also enriched by the addition of *Watts' Physical and Inorganic Chemistry*. All of the matter is brought to the present standpoint of chemical knowledge. We may safely predict for this work a continuance of the fame and favor it enjoys among medical students.—*New Orleans Medical and Surgical Journal*, March, 1886.

ATTFIELD, JOHN, Ph. D.,*Professor of Practical Chemistry to the Pharmaceutical Society of Great Britain, etc.*

Chemistry, General, Medical and Pharmaceutical; Including the Chemistry of the U. S. Pharmacopœia. A Manual of the General Principles of the Science, and their Application to Medicine and Pharmacy. A new American, from the tenth English edition, specially revised by the Author. In one handsome royal 12mo. volume of 728 pages, with 87 illustrations. Cloth, \$2.50; leather, \$3.00.

A text-book which passes through ten editions in sixteen years must have good qualities. This remark is certainly applicable to Attfield's Chemistry, a book which is so well known that it is hardly necessary to do more than note the appearance of this new and improved edition. It seems, however, desirable to point out that feature of the book which, in all probability, has made it so popular. There can be little doubt that it is its thoroughly practical character, the expression being used in its best sense. The author understands what the student ought to learn, and is able

to put himself in the student's place and to appreciate his state of mind.—*American Chemical Journal*, April, 1884.

It is a book on which too much praise cannot be bestowed. As a text-book for medical schools it is unsurpassable in the present state of chemical science, and having been prepared with a special view towards medicine and pharmacy, it is alike indispensable to all persons engaged in those departments of science. It includes the whole chemistry of the last Pharmacopœia.—*Pacific Medical and Surgical Journal*, Jan. 1884.

BLOXAM, CHARLES L.,*Professor of Chemistry in King's College, London.*

Chemistry, Inorganic and Organic. New American from the fifth London edition, thoroughly revised and much improved. In one very handsome octavo volume of 727 pages, with 292 illustrations. Cloth, \$3.75; leather, \$4.75.

Comment from us on this standard work is almost superfluous. It differs widely in scope and aim from that of Attfield, and in its way is equally beyond criticism. It adopts the most direct methods in stating the principles, hypotheses and facts of the science. Its language is so terse and lucid, and its arrangement of matter so logical in sequence that the student never has occasion to complain that chemistry is a hard study. Much attention is paid to experimental illustrations of chemical principles and phenomena, and the mode of conducting these experiments. The book maintains the position it has always held as one of

the best manuals of general chemistry in the English language.—*Detroit Lancet*, Feb. 1884.

Professor Bloxam's book is a very satisfactory one. We know of no treatise on chemistry which contains so much practical information in the same number of pages. The book can be readily adapted not only to the needs of those who desire a tolerably complete course of chemistry, but also to the needs of those who desire only a general knowledge of the subject. We take pleasure in recommending this work both as a satisfactory text-book, and as a useful book of reference.—*Boston Medical and Surgical Journal*, June 19, 1884.

REMSEN, IRA, M. D., Ph. D.,*Professor of Chemistry in the Johns Hopkins University, Baltimore, Md.*

Principles of Theoretical Chemistry. A Treatise on the Principles of Chemical Compounds. New 12mo. volume of about 250 pages.

CHARLES, T. CRANSTOUN, M. D., F. C. S., M. S.,*Formerly Asst. Prof. and Demonstrator of Chemistry and Chemical Physics, Queen's College, Belfast.*

The Elements of Physiological and Pathological Chemistry. A Handbook for Medical Students and Practitioners. Containing a general account of Nutrition, Foods and Digestion, and the Chemistry of the Tissues, Organs, Secretions and Excretions of the Body in Health and in Disease. Together with the methods for preparing or separating their chief constituents, as also for their examination in detail, and an outline syllabus of a practical course of instruction for students. In one handsome octavo volume of 463 pages, with 38 woodcuts and 1 colored plate. Cloth, \$3.50.

This is, we believe, the first complete work of the kind in the English language, and may well serve to show what progress is being made in medicine. The student will find condensed in one volume such a store of knowledge as would formerly have cost him much reading to gather. The book will fully repay anyone to read, and every library should have it for reference.—*New Orleans Medical and Surgical Journal*, April, 1885.

Dr. Charles is fully impressed with the importance and practical reach of his subject, and he

has treated it in a competent and instructive manner. We cannot recommend a better book than the present. In fact, it fills a gap in medical textbooks, and that is a thing which can rarely be said nowadays. Dr. Charles has devoted much space to the elucidation of urinary mysteries. He does this with much detail, and yet in a practical and intelligible manner. In fact, the author has filled his book with many practical hints.—*Medical Record*, December 20, 1884.

HOFFMANN, F., A.M., Ph.D., & POWER F.B., Ph.D.,*Public Analyst to the State of New York.**Prof. of Anal. Chem. in the Phil. Coll. of Pharmacy.*

A Manual of Chemical Analysis, as applied to the Examination of Medicinal Chemicals and their Preparations. Being a Guide for the Determination of their Identity and Quality, and for the Detection of Impurities and Adulterations. For the use of Pharmacists, Physicians, Druggists and Manufacturing Chemists, and Pharmaceutical and Medical Students. Third edition, entirely rewritten and much enlarged. In one very handsome octavo volume of 621 pages, with 179 illustrations. Cloth, \$4.25.

We congratulate the author on the appearance of the third edition of this work, published for the first time in this country also. It is admirable and the information it undertakes to supply is both extensive and trustworthy. The selection of processes for determining the purity of the substances of which it treats is excellent and the descrip-

tion of them singularly explicit. Moreover, it is exceptionally free from typographical errors. We have no hesitation in recommending it to those who are engaged either in the manufacture or the testing of medicinal chemicals.—*London Pharmaceutical Journal and Transactions*, 1883.

CLOWES, FRANK, D. Sc., London,*Senior Science-Master at the High School, Newcastle-under-Lyme, etc.*

An Elementary Treatise on Practical Chemistry and Qualitative Inorganic Analysis. Specially adapted for use in the Laboratories of Schools and Colleges and by Beginners. Third American from the fourth and revised English edition. In one very handsome royal 12mo. volume of 387 pages, with 55 illustrations. Cloth, \$2.50.

We may simply repeat the favorable opinion which we expressed after the examination of the previous edition of this work. It is practical in its aims, and accurate and concise in its statements.—*American Journal of Pharmacy*, August, 1885.

The style is clear, the language terse and vigorous. Beginning with a list of apparatus necessary

for chemical work, he gradually unfolds the subject from its simpler to its more complex divisions. It is the most readable book of the kind we have yet seen, and is without doubt a systematic, intelligible and fully equipped laboratory guide and text-book.—*Medical Record*, July 18, 1885.

RALFE, CHARLES H., M. D., F. R. C. P.,*Assistant Physician at the London Hospital.*

Clinical Chemistry. In one pocket-size 12mo. volume of 314 pages, with 16 illustrations. Limp cloth, red edges, \$1.50.

This is one of the most instructive little works that we have met with in a long time. The author is a physician and physiologist, as well as a chemist, consequently the book is unqualifiedly practical, telling the physician just what he ought to know, of the applications of chemistry in medi-

cine. Dr. Ralfe is thoroughly acquainted with the latest contributions to his science, and it is quite refreshing to find the subject dealt with so clearly and simply, yet in such evident harmony with the modern scientific methods and spirit.—*Medical Record*, February 2, 1884.

CLASSEN, ALEXANDER,*Professor in the Royal Polytechnic School, Aix-la-Chapelle.*

Elementary Quantitative Analysis. Translated, with notes and additions, by EDGAR F. SMITH, Ph. D., Assistant Professor of Chemistry in the Towne Scientific School, University of Penna. In one 12mo. volume of 324 pages, with 36 illust. Cloth, \$2.00.

It is probably the best manual of an elementary nature extant, inasmuch as its methods are the best. It teaches by examples, commencing with single determinations, followed by separations,

and then advancing to the analysis of minerals and such products as are met with in applied chemistry. It is an indispensable book for students in chemistry.—*Boston Journal of Chemistry*, Oct. 1878.

GREENE, WILLIAM H., M. D.,*Demonstrator of Chemistry in the Medical Department of the University of Pennsylvania.*

A Manual of Medical Chemistry. For the use of Students. Based upon Bowman's Medical Chemistry. In one 12mo. volume of 310 pages, with 74 illus. Cloth, \$1.75.

It is a concise manual of three hundred pages, giving an excellent summary of the best methods of analyzing the liquids and solids of the body, both for the estimation of their normal constituents and

the recognition of compounds due to pathological conditions. The detection of poisons is treated with sufficient fullness for the purpose of the student or practitioner.—*Boston J. of Chem.* June, '80.

BRUNTON, T. LAUDER, M.D., D.Sc., F.R.S., F.R.C.P.,

Lecturer on Materia Medica and Therapeutics at St. Bartholomew's Hospital, London, etc.

A Text-book of Pharmacology, Therapeutics and Materia Medica; Including the Pharmacy, the Physiological Action and the Therapeutical Uses of Drugs. In one handsome octavo volume of 1033 pages, with 188 illustrations. Cloth, \$5.50; leather, \$6.50.

It is a scientific treatise worthy to be ranked with the highest productions in physiology, either in our own or any other language. Everything is practical, the dry, hard facts of physiology being pressed into service and applied to the treatment of the commonest complaints. The information is so systematically arranged that it is available for immediate use. The index is so carefully compiled that a reference to any special point is at once obtainable. Dr. Brunton is never satisfied with vague generalities, but gives clear and precise directions for prescribing the various drugs and preparations. We congratulate students on being at last placed in possession of a scientific treatise of enormous practical importance.—*The London Lancet*, June 27, 1886.

Of all the numerous publications of this year

upon the subject of Pharmacology the comprehensive work of Brunton is clearly the most important, and is beyond question the foremost English handbook of *Materia Medica* and Therapeutics since the appearance of Pereira's *Elements of Materia Medica* in 1842. It is original both in the arrangement of the subjects and in the mode of treatment, and develops in a comprehensive manner the foundation principles of the science of Pharmacology without leaving the needs of the practitioner out of sight for an instant. In fact, the author has written a book which deserves to be known far beyond the boundaries of England, and can serve as a model for the pharmacological works of the continent on account of its happy combination of theory with practice.—*Virchow's Jahresbericht*, Berlin, 1886.

BARTHOLOW, ROBERTS, A. M., M.D., LL. D.,

Professor of Materia Medica and General Therapeutics in the Jefferson Medical College of Philadelphia.

New Remedies of Indigenous Source: Their Physiological Actions and Therapeutical Uses. In one octavo volume of about 300 pages. *Preparing.*

PARRISH, EDWARD,

Late Professor of the Theory and Practice of Pharmacy in the Philadelphia College of Pharmacy.

A Treatise on Pharmacy: designed as a Text-book for the Student, and as a Guide for the Physician and Pharmacist. With many Formule and Prescriptions. Fifth edition, thoroughly revised, by THOMAS S. WIEGAND, Ph.G. In one handsome octavo volume of 1093 pages, with 256 illustrations. Cloth, \$5; leather, \$6.

No thorough-going pharmacist will fail to possess himself of so useful a guide to practice, and no physician who properly estimates the value of an accurate knowledge of the remedial agents employed by him in daily practice, so far as their miscibility, compatibility and most effective methods of combination are concerned, can afford to leave this work out of the list of their works of reference. The country practitioner, who must always be in a measure his own pharmacist, will find it indispensable.—*Louisville Medical News*, March 29, 1884.

This well-known work presents itself now based upon the recently revised new Pharmacopoeia.

Each page bears evidence of the care bestowed upon it, and conveys valuable information from the rich store of the editor's experience. In fact, all that relates to practical pharmacy—apparatus, processes and dispensing—has been arranged and described with clearness in its various aspects, so as to afford aid and advice alike to the student and to the practical pharmacist. The work is judiciously illustrated with good woodcuts.—*American Journal of Pharmacy*, January, 1884.

There is nothing to equal Parrish's *Pharmacy* in this or any other language.—*London Pharmaceutical Journal*.

HERMANN, Dr. L.,

Professor of Physiology in the University of Zurich.

Experimental Pharmacology. A Handbook of Methods for Determining the Physiological Actions of Drugs. Translated, with the Author's permission, and with extensive additions, by ROBERT MEADE SMITH, M.D., Demonstrator of Physiology in the University of Pennsylvania. In one handsome 12mo. volume of 199 pages, with 32 illustrations. Cloth, \$1.50.

MAISCH, JOHN M., Phar. D.,

Professor of Materia Medica and Botany in the Philadelphia College of Pharmacy.

A Manual of Organic Materia Medica; Being a Guide to *Materia Medica* of the Vegetable and Animal Kingdoms. For the use of Students, Druggists, Pharmacists and Physicians. Third edition. In one handsome royal 12mo. volume of about 550 pages, with about 250 illustrations. *Shortly.*

BRUCE, J. MITCHELL, M.D., F.R.C.P.,

Physician and Lecturer on Materia Medica and Therapeutics at Charing Cross Hospital, London.

Materia Medica and Therapeutics. An Introduction to Rational Treatment. Fourth edition. In one pocket-size 12mo. volume of 591 pages. Limp cloth, \$1.50. See *Students' Series of Manuals*, page 4.

GRIFFITH, ROBERT EGLINTON, M.D.

A Universal Formulary, containing the Pharmaceutical and Administrative Official and other Medicines, as used by Physicians and Pharmacists. Third edition, thoroughly revised, by ROBERT GRIFFITH, Phar.D., Professor of Materia Medica in the University of London. In one octavo volume of

STILLÉ, A., M. D., LL. D., & MAISCH, J. M., Phar. D.,

Professor Emeritus of the Theory and Practice of Medicine and of Clinical Medicine in the University of Pennsylvania.

Prof. of Mat. Med. and Botany in Philadelphia College of Pharmacy, Sec'y to the American Pharmaceutical Association.

NEW (FOURTH) EDITION.

The National Dispensatory.

CONTAINING THE NATURAL HISTORY, CHEMISTRY, PHARMACY, ACTIONS AND USES OF MEDICINES, INCLUDING THOSE RECOGNIZED IN THE PHARMACOPEIAS OF THE UNITED STATES, GREAT BRITAIN AND GERMANY, WITH NUMEROUS REFERENCES TO THE FRENCH CODEX.

Fourth edition, revised to October, 1886, and covering the new British Pharmacopœia.

In one magnificent imperial octavo volume of 1794 pages, with 311 elaborate engravings. Price in cloth, \$7.25; leather, raised bands, \$8.00; very handsome half Russia, raised bands and open back, \$9.00.

* * * This work will be furnished with Patent Ready Reference Thumb-letter Index for \$1.00 in addition to the price in any style of binding.

In this new edition of THE NATIONAL DISPENSATORY, all important changes in the recent British Pharmacopœia have been incorporated throughout the volume, while in the Addenda will be found, grouped in a convenient section of 24 pages, all therapeutical novelties which have been established in professional favor since the publication of the third edition two years ago. Detailed information is thus given of the following among the many drugs treated: Antipyrin, Cocaine Hydrochlorate, Cascara Sagrada, Fabiana, Franciscea, various new Glycerins, Gymnocladus, Hydroquinon, Hyppone, Iodol, Jacaranda, Lanolin, Menthol, Phormium, Sulphophenol, Thallin and Urethan. In this edition, as always before, THE NATIONAL DISPENSATORY may be said to be the representative of the most recent state of American, English, German and French Pharmacology, Therapeutics and Materia Medica.

The National Dispensatory is so well and favorably known on both sides of the Atlantic that scarcely anything else is left to the reviewer than to call attention to the fact that another edition of this valuable work has made its appearance. This last edition surpasses even its predecessors in thoroughness and accuracy. The fact that in 1884, when the third edition was published, no revision of the British Pharmacopœia had been made for seventeen years, has necessitated a thorough going over of the whole work in order that references might correspond to the latest revision of the work of our British cousins. In addition to these changes is a fairly full *résumé*, in the form of *addenda*, of the more important drugs which have come into general use in the last two or three years.—*The American Journal of the Medical Sciences*, April, 1887.

This, the most comprehensive of the several commentaries on the Pharmacopœias of the United States, Great Britain and Germany which has yet appeared, has by this last revision been brought

fully up to the existing knowledge upon the subject treated. Its references to the British Pharmacopœia have been amended for the late and much-changed new edition of that work; an "addenda" of twenty-five pages has been appended, treating of the latest and most important additions to the materia medica. This work should be in the hands of every physician and pharmacist.—*Boston Medical and Surgical Journal*, Feb. 10, 1887.

We think it a matter for congratulation that the profession of medicine and that of pharmacy have shown such appreciation of this great work as to call for four editions within the comparatively brief period of eight years. The matters with which it deals are of so practical a nature that neither the physician nor the pharmacist can do without the latest text-books on them, especially those that are so accurate and comprehensive as this one. The book is in every way creditable both to the authors and to the publishers.—*New York Medical Journal*, May 21, 1887.

FARQUHARSON, ROBERT, M. D.,

Lecturer on Materia Medica at St. Mary's Hospital Medical School.

A Guide to Therapeutics and Materia Medica. Third American edition, specially revised by the Author. Enlarged and adapted to the U. S. Pharmacopœia by FRANK WOODBURY, M. D. In one handsome 12mo. volume of 524 pages. Cloth, \$2.25.

Dr. Farquharson's Therapeutics is constructed upon a plan which brings before the reader all the essential points with reference to the properties of drugs. It impresses these upon him in such a way as to enable him to take a clear view of the actions of medicines and the disordered conditions in which they must prove useful. The double-col-

umn pages—one side containing the recognized physiological action of the medicine, and the other the disease in which observers (who are nearly always mentioned) have obtained from it good results—make a very good arrangement. The early chapter containing rules for prescribing is excellent.—*Canada Med. and Surg. Journal*, Dec. 1882.

EDES, ROBERT T., M. D.,

Jackson Professor of Clinical Medicine in Harvard University, Medical Department.

A Text-Book of Materia Medica and Therapeutics. In one octavo volume of about 600 pages, with illustrations. *In press.*

STILLÉ, ALFRED, M. D., LL. D.,

Professor of Theory and Practice of Med. and of Clinical Med. in the Univ. of Penna.

Therapeutics and Materia Medica. A Systematic Treatise on the Action and Uses of Medicinal Agents, including their Description and History. Fourth edition, revised and enlarged. In two large and handsome octavo volumes, containing 1936 pages.

COATS, JOSEPH, M. D., F. F. P. S.,*Pathologist to the Glasgow Western Infirmary.*

A Treatise on Pathology. In one very handsome octavo volume of 829 pages, with 339 beautiful illustrations. Cloth, \$5.50; leather, \$6.50.

The work before us treats the subject of Pathology more extensively than it is usually treated in similar works. Medical students as well as physicians, who desire a work for study or reference, that treats the subjects in the various departments in a very thorough manner, but without prolixity, will certainly give this one the preference to any with which we are acquainted. It sets forth the most recent discoveries, exhibits, in an interesting manner, the changes from a normal condition effected in structures by disease, and points out the characteristics of various morbid agencies, so that they can be easily recognized. But, not limited to morbid anatomy, it explains fully how the functions of organs are disturbed by abnormal conditions. There is nothing belonging to its de-

partment of medicine that is not as fully elucidated as our present knowledge will admit.—*Cincinnati Medical News*, Oct. 1883.

One of the best features of this treatise consists in the judicious admixture of foreign observation with private experience. Thus the subject is presented in a harmonious manner, facilitating the study of single topics and making the entire volume profitable and pleasant reading. The author includes in his descriptions, general pathology as well as the special pathological histology of the different systems and organs. He has succeeded in offering to students and practitioners a thoroughly acceptable work.—*Medical Record*, Dec. 22, 1883.

GREEN, T. HENRY, M. D.,*Lecturer on Pathology and Morbid Anatomy at Charing-Cross Hospital Medical School, London.*

Pathology and Morbid Anatomy. Fifth American from the sixth revised and enlarged English edition. In one very handsome octavo volume of 482 pages, with 150 fine engravings. Cloth, \$2.50.

The fact that this well-known treatise has so rapidly reached its sixth edition is a strong evidence of its popularity. The author is to be congratulated upon the thoroughness with which he has prepared this work. It is thoroughly abreast with all the most recent advances in pathology. No work in the English language is so admirably adapted to the wants of the student and practitioner as this, and we would recommend it most earnestly to every one.—*Nashville Journal of Medicine and Surgery*, Nov. 1884.

An extended review of such a well-known book is unnecessary. We had already regarded the book as a model of its kind, and the author's assurance that he believes the present edition to be fully up to date will be received as sufficient proof that nothing of importance has been omitted. The book has been most carefully revised, and bears upon every page the marks of the care and accuracy that have won for it an international reputation.—*New York Medical and Surgical Journal*, July 18, 1885.

WOODHEAD, G. SIMS, M. D., F. R. C. P. E.,*Demonstrator of Pathology in the University of Edinburgh.*

Practical Pathology. A Manual for Students and Practitioners. In one beautiful octavo volume of 497 pages, with 136 exquisitely colored illustrations. Cloth, \$6.00.

It forms a real guide for the student and practitioner who is thoroughly in earnest in his endeavor to see for himself and do for himself. To the laboratory student it will be a helpful companion, and all those who may wish to familiarize themselves with modern methods of examining morbid tissues are strongly urged to provide

themselves with this manual. The numerous drawings are not fancied pictures, or merely schematic diagrams, but they represent faithfully the actual images seen under the microscope. The author merits all praise for having produced a valuable work.—*Medical Record*, May 31, 1884.

SCHÄFER, EDWARD A., F. R. S.,*Assistant Professor of Physiology in University College, London.*

The Essentials of Histology. In one octavo volume of 246 pages, with 281 illustrations. Cloth, \$2.25.

This admirable work was greatly needed. To those who are familiar with the author's former "Course of Practical Histology," the book needs no recommendation. It has been written with the object of supplying the student with directions for the microscopical examination of the tissues, which are given in a clear and understandable way. Although especially adapted for laboratory work, at the same time it is intended to serve as

an elementary text-book of histology, comprising all the essential facts of the science, but omitting unimportant details. The author has recommended only those methods upon which long experience has proved that full dependence can be placed. The strict observance of this plan permits of no doubt, and makes the work eminently satisfactory.—*The Physician and Surgeon*, July, 1887.

KLEIN, E., M. D., F. R. S.,*Joint Lecturer on General Anat. and Phys. in the Med. School of St. Bartholomew's Hosp., London.*

Elements of Histology. In one pocket-size 12mo. volume of 360 pages, with 181 illus. Limp cloth, red edges, \$1.50. See *Students' Series of Manuals*, page 4.

This little volume, originally intended by its able author as a manual for medical students, contains much valuable information, systematically arranged, that will be acceptable to the general practitioner. It gives a graphic and lucid description of every tissue and organ in the hu-

man body; and, while small in size, it is full to overflowing with important facts in regard to these multifarious and complex structures. We know of no book of its size that will prove of greater value to medical students and practitioners of medicine.—*The Southern Practitioner*, Nov. 1883.

PEPPER, A. J., M. B., M. S., F. R. C. S.,*Surgeon and Lecturer at St. Mary's Hospital, London.*

Surgical Pathology. In one pocket-size 12mo. volume of 511 pages, with 81 illustrations. Limp cloth, red edges, \$2.00. See *Students' Series of Manuals*, page 4.

It is not pretentious, but it will serve exceedingly well as a book of reference. It embodies a great deal of matter, referring over the whole field of surgical pathology. Its form is practical, its language is clear, and the information set forth is well-arranged, well-indexed and well-

illustrated. The student will find in it nothing that is unnecessary. The list of subjects covers the whole range of surgery. The book supplies a very manifest want and should meet with success.—*New York Medical Journal*, May 31, 1884.

Cornil and Ranvier's Pathological Histology.—Translated by E. O. SHAKESPEARE, M. D., and J. HENRY C. SIMES, M. D. Octavo, 800 pp., 360 illustrations.

FLINT, AUSTIN, M. D.,*Prof. of the Principles and Practice of Med. and of Clin. Med. in Bellevue Hospital Medical College, N. Y.*

A Treatise on the Principles and Practice of Medicine. Designed for the use of Students and Practitioners of Medicine. New (sixth) edition, thoroughly revised and rewritten by the Author, assisted by WILLIAM H. WELCH, M.D., Professor of Pathology, Johns Hopkins University, Baltimore, and AUSTIN FLINT, JR., M.D., LL.D., Professor of Physiology, Bellevue Hospital Medical College, N. Y. In one very handsome octavo volume of about 1170 pages, with illustrations. Cloth, \$5.50; leather, \$6.50; very handsome half Russia, raised bands, \$7.00.

A new edition of a work of such established reputation as Flint's Medicine needs but few words to commend it to notice. It may in truth be said to embody the fruit of his labors in clinical medicine, ripened by the experience of a long life devoted to its pursuit. America may well be proud of having produced a man whose indefatigable industry and gifts of genius have done so much to advance medicine; and all English-reading students must be grateful for the work which he has left behind him. It has few equals, either in point of literary excellence, or of scientific learning, and no one can fail to study its pages without being struck by the lucidity and accuracy which characterize them. It is qualities such as these which render it so valuable for its purpose, and give it a foremost place among the text-books of this generation.—*The London Lancet*, March 12, 1887.

No text-book on the principles and practice of medicine has ever met in this country with such

general approval by medical students and practitioners as the work of Professor Flint. In all the medical colleges of the United States it is the favorite work upon Practice; and, as we have stated before in alluding to it, there is no other medical work that can be so generally found in the libraries of physicians. In every state and territory of this vast country the book that will be most likely to be found in the office of a medical man, whether in city, town, village, or at some cross-roads, is Flint's *Practice*. We make this statement to a considerable extent from personal observation, and it is the testimony also of others. An examination shows that very considerable changes have been made in the sixth edition. The work may undoubtedly be regarded as fairly representing the present state of the science of medicine, and as reflecting the views of those who exemplify in their practice the present stage of progress of medical art.—*Cincinnati Medical News*, Oct. 1886.

HARTSHORNE, HENRY, M. D., LL. D.,*Lately Professor of Hygiene in the University of Pennsylvania.*

Essentials of the Principles and Practice of Medicine. A Handbook for Students and Practitioners. Fifth edition, thoroughly revised and rewritten. In one royal 12mo. volume of 669 pages, with 144 illustrations. Cloth, \$2.75; half bound, \$3.00.

Within the compass of 600 pages it treats of the history of medicine, general pathology, general symptomatology, and physical diagnosis (including laryngoscope, ophthalmoscope, etc.), general therapeutics, nosology, and special pathology and practice. There is a wonderful amount of information contained in this work, and it is one of the best of its kind that we have seen.—*Glasgow Medical Journal*, Nov. 1882.

An indispensable book. No work ever exhibited a better average of actual practical treatment than

this one; and probably not one writer in our day had a better opportunity than Dr. Hartshorne for condensing all the views of eminent practitioners into a 12mo. The numerous illustrations will be very useful to students especially. These essentials, as the name suggests, are not intended to supersede the text-books of Flint and Bartholow, but they are the most valuable in affording the means to see at a glance the whole literature of any disease, and the most valuable treatment.—*Chicago Medical Journal and Examiner*, April, 1882.

BRISTOWE, JOHN SYER, M. D., F. R. C. P.,*Physician and Joint Lecturer on Medicine at St. Thomas' Hospital, London.*

A Treatise on the Practice of Medicine. Second American edition, revised by the Author. Edited, with additions, by JAMES H. HUTCHINSON, M.D., physician to the Pennsylvania Hospital. In one handsome octavo volume of 1085 pages, with illustrations. Cloth, \$5.00; leather, \$6.00; very handsome half Russia, raised bands, \$6.50.

The book is a model of conciseness, and combines, as successfully as one could conceive it to be possible, an encyclopædic character with the smallest dimensions. It differs from other admirable text-books in the completeness with which it covers the whole field of medicine.—*Michigan Medical News*, May 10, 1880.

His accuracy in the portraiture of disease, his care in stating subtle points of diagnosis, and the faithfully given pathology of abnormal processes have seldom been surpassed. He embraces many diseases not usually considered to belong to theory

and practice, as skin diseases, syphilis and insanity, but they will not be objected to by readers, as he has studied them conscientiously, and drawn from the life.—*Medical and Surgical Reporter*, December 20, 1879.

The reader will find every conceivable subject connected with the practice of medicine ably presented, in a style at once clear, interesting and concise. The additions made by Dr. Hutchinson are appropriate and practical, and greatly add to its usefulness to American readers.—*Buffalo Medical and Surgical Journal*, March, 1880.

WATSON, SIR THOMAS, M. D.,*Late Physician in Ordinary to the Queen.*

Lectures on the Principles and Practice of Physio. A new American from the fifth English edition. Edited, with additions, and 190 illustrations, by HENRY HARTSHORNE, A. M., M. D., late Professor of Hygiene in the University of Pennsylvania. In two large octavo volumes of 1840 pages. Cloth, \$9.00; leather, \$11.00.

LECTURES ON THE STUDY OF FEVER. By A. HUDSON, M. D., M. R. I. A. In one octavo volume of 308 pages. Cloth, \$2.50.

STOKES' LECTURES ON FEVER. Edited by John William Moore, M. D., F. R. C. P. In one octavo volume of 280 pages. Cloth, \$2.00.

A TREATISE ON FEVER. By ROBERT D. LYONS, K. C. C. In one 8vo. vol. of 354 pp. Cloth, \$2.25.

LA ROCHE ON YELLOW FEVER, considered in its Historical, Pathological, Etiological and Therapeutical Relations. In two large and handsome octavo volumes of 1468 pp. Cloth, \$7.00.

A CENTURY OF AMERICAN MEDICINE, 1776-1876. By Drs. E. H. CLARK, H. J. BIGLOW, S. D. GROSS, T. G. THOMAS, and J. S. BILLINGS. In one 12mo. volume of 370 pages. Cloth, \$2.25.

For Sale by Subscription Only.

A System of Practical Medicine.

BY AMERICAN AUTHORS.

EDITED BY WILLIAM PEPPER, M. D., LL. D.,

PROVOST AND PROFESSOR OF THE THEORY AND PRACTICE OF MEDICINE AND OF CLINICAL MEDICINE IN THE UNIVERSITY OF PENNSYLVANIA,

Assisted by LOUIS STARR, M. D., Clinical Professor of the Diseases of Children in the Hospital of the University of Pennsylvania.

The complete work, in five volumes, containing 5573 pages, with 198 illustrations, is just ready. Price per volume, cloth, \$5; leather, \$6; half Russia, raised bands and open back, \$7.

In this great work American medicine is for the first time reflected by its worthiest teachers, and presented in the full development of the practical utility which is its pre-eminent characteristic. The most able men—from the East and the West, from the North and the South, from all the prominent centres of education, and from all the hospitals which afford special opportunities for study and practice—have united in generous rivalry to bring together this vast aggregate of specialized experience.

The distinguished editor has so apportioned the work that to each author has been assigned the subject which he is peculiarly fitted to discuss, and in which his views will be accepted as the latest expression of scientific and practical knowledge. The practitioner will therefore find these volumes a complete, authoritative and unflinching work of reference, to which he may at all times turn with full certainty of finding what he needs in its most recent aspect, whether he seeks information on the general principles of medicine, or minute guidance in the treatment of special disease. So wide is the scope of the work that, with the exception of midwifery and matters strictly surgical, it embraces the whole domain of medicine, including the departments for which the physician is accustomed to rely on special treatises, such as diseases of women and children, of the genito-urinary organs, of the skin, of the nerves, hygiene and sanitary science, and medical ophthalmology and otology. Moreover, authors have inserted the formulas which they have found most efficient in the treatment of the various affections. It may thus be truly regarded as a COMPLETE LIBRARY OF PRACTICAL MEDICINE, and the general practitioner possessing it may feel secure that he will require little else in the daily round of professional duties.

In spite of every effort to condense the vast amount of practical information furnished, it has been impossible to present it in less than 5 large octavo volumes, containing about 5600 beautifully printed pages, and embodying the matter of about 15 ordinary octavos. Illustrations are introduced wherever requisite to elucidate the text.

A detailed prospectus will be sent to any address on application to the publishers.

These two volumes bring this admirable work to a close, and fully sustain the high standard reached by the earlier volumes; we have only therefore to echo the eulogium pronounced upon them. We would warmly congratulate the editor and his collaborators at the conclusion of their laborious task on the admirable manner in which, from first to last, they have performed their several duties. They have succeeded in producing a work which will long remain a standard work of reference, to which practitioners will look for guidance, and authors will resort to for facts. From a literary point of view, the work is without any serious blemish, and in respect of production, it has the beautiful finish that Americans always give their works.—*Edinburgh Medical Journal*, Jan. 1887.

* * * The greatest distinctively American work on the practice of medicine, and, indeed, the superlative adjective would not be inappropriate were even all other productions placed in comparison. An examination of the five volumes is sufficient to convince one of the magnitude of the enterprise, and of the success which has attended its fulfilment.—*The Medical Age*, July 26, 1886.

This huge volume forms a fitting close to the great system of medicine which in so short a time has won so high a place in medical literature, and has done such credit to the profession in this country. Among the twenty-three contributors are the names of the leading neurologists in America, and most of the work in the volume is of the highest order.—*Boston Medical and Surgical Journal*, July 21, 1887.

We consider it one of the grandest works on Practical Medicine in the English language. It is a work of which the profession of this country can feel proud. Written exclusively by American

physicians who are acquainted with all the varieties of climate in the United States, the character of the soil, the manners and customs of the people, etc., it is peculiarly adapted to the wants of American practitioners of medicine, and it seems to us that every one of them would desire to have it. It has been truly called a "Complete Library of Practical Medicine," and the general practitioner will require little else in his round of professional duties.—*Cincinnati Medical News*, March, 1886.

Each of the volumes is provided with a most copious index, and the work altogether promises to be one which will add much to the medical literature of the present century, and reflect great credit upon the scholarship and practical acumen of its authors.—*The London Lancet*, Oct. 3, 1885.

The feeling of proud satisfaction with which the American profession sees this, its representative system of practical medicine issued to the medical world, is fully justified by the character of the work. The entire caste of the system is in keeping with the best thoughts of the leaders and followers of our home school of medicine, and the combination of the scientific study of disease and the practical application of exact and experimental knowledge to the treatment of human maladies, makes every one of us share in the pride that has welcomed Dr. Pepper's labors. Sheared of the prolixity that wearies the readers of the German school, the articles glean these same fields for all that is valuable. It is the outcome of American brains, and is marked throughout by much of the sturdy independence of thought and originality that is a national characteristic. Yet nowhere is there lack of study of the most advanced views of the day.—*North Carolina Medical Journal*, Sept. 1886.

FOTHERGILL, J. M., M. D., Edin., M. R. C. P., Lond.,*Physician to the City of London Hospital for Diseases of the Chest.*

The Practitioner's Handbook of Treatment; Or, The Principles of Therapeutics. New (third) edition. In one octavo volume of 661 pages. Cloth, \$3.75; leather, \$4.75. *Just ready.*

Dr. Fothergill is always interesting and instructive, and in this standard work he shows his peculiar power as a writer on therapeutics to the best advantage. Everything he undertakes is done conscientiously. The book well sustains the favorable impression which it created at the beginning of its career, and in its present improved form it will be welcomed more than ever by the busy practitioner and the scientific student of medicine.—*The Medical News*, July 23, 1887.

To have a description of the normal physiological processes of an organ and of the methods of treatment of its morbid conditions brought together in a single chapter, and the relations between the two clearly stated, cannot fail to prove a great convenience to many thoughtful but busy physicians. The practical value of the volume is

greatly increased by the introduction of many prescriptions. That the profession appreciates that the author has undertaken an important work and has accomplished it is shown by the demand for this third edition.—*New York Medical Journal*, June 11, 1887.

This is a wonderful book. If there be such a thing as "medicine made easy," this is the work to accomplish this result. It imparts information so agreeably, so smoothly, that the reader almost thinks as he lays the book down that he "knew that before," when in reality he did not, or else he could before have reconciled facts which now become clear as daylight. The author deals with the "Principles of Therapeutics" the study of which will give great vantage to the physician.—*Virginia Medical Monthly*, June, 1887.

REYNOLDS, J. RUSSELL, M. D.,*Professor of the Principles and Practices of Medicine in University College, London.*

A System of Medicine. With notes and additions by HENRY HARTSHORNE, A. M., M. D., late Professor of Hygiene in the University of Pennsylvania. In three large and handsome octavo volumes, containing 3056 double-columned pages, with 317 illustrations. Price per volume, cloth, \$5.00; sheep, \$6.00; very handsome half Russia, raised bands, \$6.50. Per set, cloth, \$15; leather, \$18; half Russia, \$19.50. *Sold only by subscription.*

STILLÉ, ALFRED, M. D., LL. D.,*Professor Emeritus of the Theory and Practice of Med. and of Clinical Med. in the Univ. of Penna.*

Cholera: Its Origin, History, Causation, Symptoms, Lesions, Prevention and Treatment. In one handsome 12mo. volume of 163 pages, with a chart. Cloth, \$1.25. *Just ready.*

FINLAYSON, JAMES, M. D., Editor,*Physician and Lecturer on Clinical Medicine in the Glasgow Western Infirmary, etc.*

Clinical Manual for the Study of Medical Cases. With Chapters by Prof. Gairdner on the Physiognomy of Disease; Prof. Stephenson on Diseases of the Female Organs; Dr. Robertson on Insanity; Dr. Gemmell on Physical Diagnosis; Dr. Coats on Laryngoscopy and Post-Mortem Examinations, and by the Editor on Case-taking, Family History and Symptoms of Disorder in the Various Systems. New edition. In one 12mo. volume of 682 pages, with 158 illustrations. Cloth, \$2.50. *Just ready.*

The profession cannot but welcome the second edition of this very valuable work of Finlayson and his collaborators. The size of the book has been increased and the number of illustrations nearly doubled. The manner in which the subject is treated is a most practical one. Symptoms alone and their diagnostic indications form the basis of discussion. The text explains clearly and thoroughly the methods of examination and the conclusions to be drawn from the physical signs.—*The Medical News*, April 23, 1887.

This manual is one of the most complete and perfect of its kind. It goes thoroughly into the question of diagnosis from every possible point. It must lead to a thoroughness of observation, an examination in detail of every scientific appliance,

and a study of means to the end which cannot fail in laying an excellent foundation for the student for future success as an able diagnostician.—*Medical Record*, August 13, 1887.

The second edition of this manual is a very considerable improvement upon the first. Much new matter has been introduced and the work has been brought up to the present time in all respects. As it stands it is one of the best manuals of diagnosis in the English language for beginners. The whole work is so complete and so simply written, and yet contains such an amount of valuable information, that it should be a part of the library of every practitioner.—*New York Medical Journal*, July 23, 1887.

FENWICK, SAMUEL, M. D.,*Assistant Physician to the London Hospital.*

The Student's Guide to Medical Diagnosis. From the third revised and enlarged English edition. In one very handsome royal 12mo. volume of 328 pages, with 87 illustrations on wood. Cloth, \$2.25.

HABERSHON, S. O., M. D.,*Senior Physician to and late Lect. on Principles and Practice of Med. at Guy's Hospital, London.*

On the Diseases of the Abdomen; Comprising those of the Stomach, and other parts of the Alimentary Canal, (Esophagus, Cæcum, Intestines and Peritoneum. Second American from third enlarged and revised English edition. In one handsome octavo volume of 554 pages, with illustrations. Cloth, \$3.50.

TANNER, THOMAS HAWKES, M. D.

A Manual of Clinical Medicine and Physical Diagnosis. Third American from the second London edition. Revised and enlarged by TILBURY FOX, M. D. In one small 12mo. volume of 362 pages, with illustrations. Cloth, \$1.50.

BARTHOLOW, ROBERTS, A. M., M. D., LL. D.,

Prof. of Materia Medica and General Therapeutics in the Jefferson Med. Coll. of Phila., etc.

Medical Electricity. A Practical Treatise on the Applications of Electricity to Medicine and Surgery. New (third) edition. In one very handsome octavo volume of 308 pages, with 110 illustrations. Cloth, \$2.50. *Just ready.*

The fact that this work has reached its third edition in six years, and that it has been kept fully abreast with the increasing use and knowledge of electricity, demonstrates its claim to be considered a practical treatise of tried value to the profession. The matter added to the present edition embraces the most recent advances in electrical treatment. The illustrations are abundant and clear, and the work constitutes a full, clear and concise manual well adapted to the needs of both student and practitioner.—*The Medical News*, May 14, 1887.

This "practical treatise on the applications of electricity to medicine and surgery" has grown to be so important a work that every practitioner

should read it, especially when it is recalled what possibilities lie in the path of the further study of the therapeutics of electricity. Dr. Bartholow has here presented the profession with a concise work that, beginning with elementary descriptions and principles, gradually grows, page by page, into a magnificently practical treatise, describing operations in detail, and giving records of successes that prove electricity to be marvellous as a curative agent in many forms of disease. The doctor cannot now do better than to possess himself of Dr. Bartholow's treatise, just as it is.—*Virginia Medical Monthly*, June, 1887.

RICHARDSON, B. W., M.A., M.D., LL. D., F.R.S., F.S.A.

Fellow of the Royal College of Physicians, London.

Preventive Medicine. In one octavo volume of 729 pages. Cloth, \$4; leather, \$5; very handsome half Russia, raised bands, \$5.50.

Dr. Richardson has succeeded in producing a work which is elevated in conception, comprehensive in scope, scientific in character, systematic in arrangement, and which is written in a clear, concise and pleasant manner. He evinces the happy faculty of extracting the pith of what is known on the subject, and of presenting it in a most simple, intelligent and practical form. There is perhaps no similar work written for the general public that contains such a complete, reliable and instructive

collection of data upon the diseases common to the race, their origins, causes, and the measures for their prevention. The descriptions of diseases are clear, chaste and scholarly; the discussion of the question of disease is comprehensive, masterly and fully abreast with the latest and best knowledge on the subject, and the preventive measures advised are accurate, explicit and reliable.—*The American Journal of the Medical Sciences*, April, 1884.

HARTSHORNE, HENRY, M. D., LL. D.,

Formerly Professor of Hygiene in the University of Pennsylvania, and Professor of Physiology and Diseases of Children in the Woman's Medical College of Pennsylvania.

A Household Manual of Medicine, Surgery, Nursing and Hygiene: For Daily Use in the Preservation of Health and Care of the Sick and Injured, with an Introductory Outline of Anatomy and Physiology. In one very handsome royal octavo volume of 946 pages, with 8 plates and 283 engravings. Cloth, \$4.00; very handsome half Morocco, \$5.00.

THE YEAR-BOOK OF TREATMENT FOR 1886.

A Comprehensive and Critical Review for Practitioners of Medicine. In one 12mo. volume of 309 pages, bound in limp cloth, \$1.25.

This "review" includes every department of medical and surgical as well as obstetrical practice. It attempts nothing in the way of etiology, diagnosis or symptoms, but limits itself to the advances made in the treatment of diseases, injuries, etc. The work seems to us to be invaluable to

every practitioner, whether he be a general one or a specialist. It is a book to be kept on the office table for continuous reference. An excellent index to subjects, as well as to authors quoted, is appended.—*Virginia Medical Monthly*, April, 1887.

* * For special commutations with periodicals see page 3.

THE YEAR-BOOK OF TREATMENT FOR 1885.

Similar to that of 1886 above. 12mo., 320 pages. Limp cloth, \$1.25.

SCHREIBER, DR. JOSEPH.

A Manual of Treatment by Massage and Methodical Muscle Exercise. Translated by WALTER MENDELSON, M. D., of New York. In one handsome octavo volume of 278 pages, with 117 fine engravings. *Just ready.* Cloth, \$2.75.

The present volume will do much to establish mechanical therapeutics upon a scientific basis. The work is a very welcome addition to the library, and we heartily recommend it to our readers as a step in the right direction.—*New York Medical Journal*, July 16, 1887.

As a thorough and satisfactory exposition of the

science of mechanical therapeutics, adapted to the use of the general practitioner, this volume leaves nothing to be desired. The text is fully illustrated by well-drawn woodcuts, leaving no room for obscurity in the description of the various manipulations recommended.—*Atlanta Medical and Surgical Journal*, Aug. 1887.

STURGES' INTRODUCTION TO THE STUDY OF CLINICAL MEDICINE. Being a Guide to the Investigation of Disease. In one handsome 12mo. volume of 127 pages. Cloth, \$1.25.

DAVIS' CLINICAL LECTURES ON VARIOUS IMPORTANT DISEASES. By N. S. Davis, M. D. Edited by FRANK H. Davis, M. D. Second edition. 12mo. 287 pages. Cloth, \$1.75.

TODD'S CLINICAL LECTURES ON CERTAIN ACUTE DISEASES. In one octavo volume of 320 pages. Cloth, \$2.50.

PAVY'S TREATISE ON THE FUNCTION OF DIGESTION; its Disorders and their Treatment. From the second London edition. In one octavo volume of 238 pages. Cloth, \$2.00.

BARLOW'S MANUAL OF THE PRACTICE OF MEDICINE. With additions by D. F. CONNIE, M. D. 1 vol. 8vo., pp. 603. Cloth, \$2.50.

CHAMBERS' MANUAL OF DIET AND REGIMEN IN HEALTH AND SICKNESS. In one handsome octavo volume of 302 pp. Cloth, \$2.75.

HOLLAND'S MEDICAL NOTES AND REFLECTIONS. 1 vol. 8vo., pp. 483. Cloth, \$2.50.

FLINT, AUSTIN, M. D.,*Professor of the Principles and Practice of Medicine in Bellevue Hospital Medical College, N. Y.*

A Manual of Auscultation and Percussion; Of the Physical Diagnosis of Diseases of the Lungs and Heart, and of Thoracic Aneurism. Fourth edition. In one handsome royal 12mo. volume of 278 pages, with 14 illustrations. Cloth, \$1.75.

This admirable little book is too well known to require any extended notice. That a third and large edition has been exhausted in little more than two years, is evidence that the book is appreciated. We ourselves have used a former edition as a text-book in teaching the physical examination of the chest, and can consequently speak from experience.—*Boston Medical and Surgical Journal*, Feb. 11, 1886.

The student needs a first-class text-book in which the subject is fully explained for him to

study. Dr. Flint's work is just such a book. It contains the substance of the lessons which the author has for many years given in connection with practical instruction in auscultation and percussion to private classes, composed of medical students and practitioners. The fact that within a little more than two years a large edition of this manual has been exhausted, is proof of the favor with which it has been regarded by the medical profession.—*Cincinnati Medical News*, Feb. 1886.

BY THE SAME AUTHOR.

Physical Exploration of the Lungs by Means of Auscultation and Percussion. Three lectures delivered before the Philadelphia County Medical Society, 1882-83. In one handsome small 12mo. volume of 83 pages. Cloth, \$1.00.

A Practical Treatise on the Physical Exploration of the Chest and the Diagnosis of Diseases Affecting the Respiratory Organs. Second and revised edition. In one handsome octavo volume of 591 pages. Cloth, \$4.50.

Phthisis: Its Morbid Anatomy, Etiology, Symptomatic Events and Complications, Fatality and Prognosis, Treatment and Physical Diagnosis; In a series of Clinical Studies. In one handsome octavo volume of 442 pages. Cloth, \$3.50.

A Practical Treatise on the Diagnosis, Pathology and Treatment of Diseases of the Heart. Second revised and enlarged edition. In one octavo volume of 550 pages, with a plate. Cloth, \$4.

Essays on Conservative Medicine and Kindred Topics. In one very handsome royal 12mo. volume of 210 pages. Cloth, \$1.38.

GROSS, S. D., M.D., LL.D., D.C.L. Oxon., LL.D. Cantab.

A Practical Treatise on Foreign Bodies in the Air-passages. In one octavo volume of 452 pages, with 59 illustrations. Cloth, \$2.75.

COHEN, J. SOLIS, M. D.,*Lecturer on Laryngoscopy and Diseases of the Throat and Chest in the Jefferson Medical College.*

Diseases of the Throat and Nasal Passages. A Guide to the Diagnosis and Treatment of Affections of the Pharynx, Esophagus, Trachea, Larynx and Nares. Third edition, thoroughly revised and rewritten, with a large number of new illustrations. In one very handsome octavo volume. *Preparing.*

SEILER, CARL, M. D.,*Lecturer on Laryngoscopy in the University of Pennsylvania.*

A Handbook of Diagnosis and Treatment of Diseases of the Throat, Nose and Naso-Pharynx. Second edition. In one handsome royal 12mo. volume of 294 pages, with 77 illustrations. Cloth, \$1.75.

It is one of the best of the practical text-books on this subject with which we are acquainted. The present edition has been increased in size, but its eminently practical character has been maintained. Many new illustrations have also been introduced, a case-record sheet has been added,

and there are a valuable bibliography and a good index of the whole. For any one who wishes to make himself familiar with the practical management of cases of throat and nose disease, the book will be found of great value.—*New York Medical Journal*, June 9, 1883.

BROADBENT, W. H., M. D., F. R. C. P.,*Physician to and Lecturer on Medicine at St. Mary's Hospital.*

The Pulse. In one 12mo. volume. *Preparing.* See *Series of Clinical Manuals*, page 4.

FULLER ON DISEASES OF THE LUNGS AND AIR-PASSAGES. Their Pathology, Physical Diagnosis, Symptoms and Treatment. From the second and revised English edition. In one octavo volume of 476 pages. Cloth, \$3.50.

BLADE ON DIPHTHERIA; its Nature and Treatment, with an account of the History of its Prevalence in various Countries. Second and revised edition. In one 12mo. vol., pp. 158. Cloth, \$1.25.

WALSHE ON THE DISEASES OF THE HEART AND GREAT VESSELS. Third American edition. In 1 vol. 8vo., 416 pp. Cloth, \$3.00.

SMITH ON CONSUMPTION; its Early and Removable Stages. 1 vol. 8vo., pp. 253. Cloth, \$2.25.

LA ROCHE ON PNEUMONIA. 1 vol. 8vo. of 490 pages. Cloth, \$3.00.

WILLIAMS ON PULMONARY CONSUMPTION; its Nature, Varieties and Treatment. With an analysis of one thousand cases to exemplify its duration. In one 8vo. vol. of 303 pp. Cloth, \$2.50.

JONES CLINICAL OBSERVATIONS ON FUNCTIONAL NERVOUS DISORDERS. Second American edition. In one handsome octavo volume of 340 pages. Cloth, \$3.25.

ROSS, JAMES, M.D., F.R.C.P., LL.D.,*Senior Assistant Physician to the Manchester Royal Infirmary.***A Handbook on Diseases of the Nervous System.** In one octavo volume of 725 pages, with 184 illustrations. Cloth, \$4.50; leather, \$5.50.

This admirable work is intended for students of medicine and for such medical men as have not time for lengthy treatises. In the present instance the duty of arranging the vast store of material at the disposal of the author, and of abridging the description of the different aspects of nervous diseases, has been performed with singular skill, and the result is a concise and philosophical guide to

the department of medicine of which it treats. Dr. Ross holds such a high scientific position that any writings which bear his name are naturally expected to have the impress of a powerful intellect. In every part this handbook merits the highest praise, and will no doubt be found of the greatest value to the student as well as to the practitioner.—*Edinburgh Medical Journal*, Jan. 1887.

MITCHELL, S. WEIR, M.D.,*Physician to Orthopaedic Hospital and the Infirmary for Diseases of the Nervous System, Phila., etc.***Lectures on Diseases of the Nervous System; Especially in Women.** Second edition. In one 12mo. volume of 288 pages. Cloth, \$1.75.

No work in our language develops or displays more features of that many-sided affection, hysteria, or gives clearer directions for its differentiation, or sounder suggestions relative to its general management and treatment. The book is particularly valuable in that it represents in the main the author's own clinical studies, which have been so extensive and fruitful as to give his

teachings the stamp of authority all over the realm of medicine. The work, although written by a specialist, has no exclusive character, and the general practitioner above all others will find its perusal profitable, since it deals with diseases which he frequently encounters and must essay to treat.—*American Practitioner*, August, 1885.

HAMILTON, ALLAN McLANE, M.D.,*Attending Physician at the Hospital for Epileptics and Paralytics, Blackwell's Island, N. Y.***Nervous Diseases; Their Description and Treatment.** Second edition, thoroughly revised and rewritten. In one octavo volume of 598 pages, with 72 illustrations. Cloth, \$4.

When the first edition of this good book appeared we gave it our emphatic endorsement, and the present edition enhances our appreciation of the book and its author as a safe guide to students of clinical neurology. One of the best and most critical of English neurological journals, *Brain*, has

characterized this book as the best of its kind in any language, which is a handsome endorsement from an exalted source. The improvements in the new edition, and the additions to it, will justify its purchase even by those who possess the old.—*Alienist and Neurologist*, April, 1882.

TUKE, DANIEL HACK, M.D.,*Joint Author of The Manual of Psychological Medicine, etc.***Illustrations of the Influence of the Mind upon the Body in Health and Disease.** Designed to elucidate the Action of the Imagination. New edition. Thoroughly revised and rewritten. In one handsome octavo volume of 467 pages, with two colored plates. Cloth, \$3.00.

It is impossible to peruse these interesting chapters without being convinced of the author's perfect sincerity, impartiality, and thorough mental grasp. Dr. Tuke has exhibited the requisite amount of scientific address on all occasions, and the more intricate the phenomena the more firmly has he adhered to a physiological and rational

method of interpretation. Guided by an enlightened deduction, the author has reclaimed for science a most interesting domain in psychology, previously abandoned to charlatans and empirics. This book, well conceived and well written, must commend itself to every thoughtful understanding.—*New York Medical Journal*, September 6, 1884.

CLOUSTON, THOMAS S., M.D., F.R.C.P., L.R.C.S.,*Lecturer on Mental Diseases in the University of Edinburgh.***Clinical Lectures on Mental Diseases.** With an Appendix, containing an Abstract of the Statutes of the United States and of the Several States and Territories relating to the Custody of the Insane. By CHARLES F. FOLSOM, M.D., Assistant Professor of Mental Diseases, Med. Dep. of Harvard Univ. In one handsome octavo volume of 541 pages, with eight lithographic plates, four of which are beautifully colored. Cloth, \$4.

The practitioner as well as the student will accept the plain, practical teaching of the author as a forward step in the literature of insanity. It is refreshing to find a physician of Dr. Clouston's experience and high reputation giving the bedside notes upon which his experience has been founded and his mature judgment established. Such clinical observations cannot but be useful to

the general practitioner in guiding him to a diagnosis and indicating the treatment, especially in many obscure and doubtful cases of mental disease. To the American reader Dr. Folsom's Appendix adds greatly to the value of the work, and will make it a desirable addition to every library.—*American Psychological Journal*, July, 1884.

Dr. Folsom's *Abstract* may also be obtained separately in one octavo volume of 108 pages. Cloth, \$1.50.

SAVAGE, GEORGE H., M.D.,*Lecturer on Mental Diseases at Guy's Hospital, London.***Insanity and Allied Neuroses, Practical and Clinical.** In one 12mo. vol. of 551 pages, with 18 illus. Cloth, \$2.00. See *Series of Clinical Manuals*, page 4.**PLAYFAIR, W. S., M.D., F.R.C.P.****The Systematic Treatment of Nerve Prostration and Hysteria.** In one handsome small 12mo. volume of 97 pages. Cloth, \$1.00.**Blandford on Insanity and its Treatment: Lectures on the Treatment, Medical and Legal, of Insane Patients.** In one very handsome octavo volume.

ASHHURST, JOHN, Jr., M. D.,*Professor of Clinical Surgery, Univ. of Penna., Surgeon to the Episcopal Hospital, Philadelphia.*

The Principles and Practice of Surgery. New (fourth) edition, enlarged and revised. In one large and handsome octavo volume of 1114 pages, with 597 illustrations. Cloth, \$6; leather, \$7; half Russia, \$7.50.

As with Erichsen so with Ashhurst, its position in professional favor is established, and one has now but to notice the changes, if any, in theory and practice, that are apparent in the present as compared with the preceding edition, published three years ago. The work has been brought well up to date, and is larger and better illustrated than before, and its author may rest assured that it will certainly have a "continuance of the favor with which it has heretofore been received."—*The American Journal of the Medical Sciences*, Jan. 1886.

Every advance in surgery worth notice, chronicled in recent literature, has been suitably recognized and noted in its proper place. Suffice it to say, we regard Ashhurst's Surgery, as now presented in the fourth edition, as the best single volume on surgery published in the English language, valuable alike to the student and the practitioner, to the one as a text-book, to the other as a manual of practical surgery. With pleasure we give this volume our endorsement in full.—*New Orleans Medical and Surgical Journal*, Jan., 1886.

GROSS, S. D., M. D., LL. D., D. C. L. Oxon., LL. D. Cantab.,*Emeritus Professor of Surgery in the Jefferson Medical College of Philadelphia.*

A System of Surgery: Pathological, Diagnostic, Therapeutic and Operative. Sixth edition, thoroughly revised and greatly improved. In two large and beautifully-printed imperial octavo volumes containing 2382 pages, illustrated by 1623 engravings. Strongly bound in leather, raised bands, \$15; half Russia, raised bands, \$16.

Dr. Gross' *System of Surgery* has long been the standard work on that subject for students and practitioners.—*London Lancet*, May 10, 1884.

The work as a whole needs no commendation. Many years ago it earned for itself the enviable reputation of the leading American work on surgery, and it is still capable of maintaining that standard. A considerable amount of new material has been introduced, and altogether the distinguished author has reason to be satisfied that he has placed the work fully abreast of the state of our knowledge.—*Med. Record*, Nov. 18, 1882.

His *System of Surgery*, which, since its first edition in 1859, has been a standard work in this country as well as in America, in "the whole domain of surgery," tells how earnest and laborious and wise a surgeon he was, how thoroughly he appreciated the work done by men in other countries, and how much he contributed to promote the science and practice of surgery in his own. There has been no man to whom America is so much indebted in this respect as the Nestor of surgery.—*British Medical Journal*, May 10, 1884.

DRUITT, ROBERT, M. R. C. S., etc.

Manual of Modern Surgery. Twelfth edition, thoroughly revised by STANLEY BOYD, M.B., B.S., F.R.C.S. In one 8vo. volume of 965 pages, with 373 illustrations. Cloth, \$4; leather, \$5.

FROM THE EDITOR'S PREFACE TO THE TWELFTH EDITION.

Few works have been more widely known and appreciated than Dr. Robert's "Surgery." In England I am informed that 50,000 copies have been sold, whilst in America it has been so highly appreciated that a copy was issued by the Government to each surgeon serving in the Federal Army during the great Civil War. The twelfth edition differs much from the eleventh; scarcely a paragraph of the latter remains unaltered. In spite of my utmost endeavors to compress, the book has increased considerably in size. This is due to many causes, especially to the greatly increased range of subjects with which I have had to deal, to the replacement of many old illustrations by a number of considerably larger ones, and to the addition of a copious index.

GIBNEY, V. P., M. D.,*Surgeon to the Orthopædic Hospital, New York, etc.*

Orthopædic Surgery. For the use of Practitioners and Students. In one handsome octavo volume, profusely illustrated. *Preparing.*

ROBERTS, JOHN B., A. M., M. D.,*Lecturer on Anatomy and on Operative Surgery, at the Philadelphia School of Anatomy.*

The Principles and Practice of Modern Surgery. For the use of Students and Practitioners of Medicine and Surgery. In one very handsome octavo volume of about 500 pages, with many illustrations. *Preparing.*

BELLAMY, EDWARD, F. R. C. S.,*Surgeon and Lecturer on Surgery at Charing Cross Hospital, London.*

Operative Surgery. *Shortly.* See *Students' Series of Manuals*, page 4.

BALL, CHARLES B., M. Ch., Dub., F. R. C. S. E.,*Surgeon and Lecturer at St. P. David's Hospital, Dublin.*

Diseases of the Rectum and Anus. In one 12mo. volume of 550 pages. *Preparing.* See *Series of Clinical Manuals*, page 4.

MILLER'S PRACTICE OF SURGERY. Fourth and revised American from the last Edinburgh edition. In one large 8vo. vol. of 682 pages, with 320 illustrations. Cloth, \$5.

MILLER'S PRINCIPLES OF SURGERY. Fourth American from the third Edinburgh edition. In one 8vo. vol. of 638 pages, with 320 illustrations. Cloth, \$5.

ERICHSEN, JOHN E., F. R. S., F. R. C. S.,

Professor of Surgery in University College, London, etc.

The Science and Art of Surgery; Being a Treatise on Surgical Injuries, Diseases and Operations. From the eighth and enlarged English edition. In two large and beautiful octavo volumes of 2316 pages, illustrated with 984 engravings on wood. Cloth, \$9; leather, raised bands, \$11; half Russia, raised bands, \$12.

In noticing the eighth edition of this well-known work, it would appear superfluous to say more than that it has, like its predecessors, been brought fully up to the times, and is in consequence one of the best treatises upon surgery that has ever been penned by one man. We have always regarded "The Science and Art of Surgery" as one of the best surgical text-books in the English language, and this eighth edition only confirms our previous opinion. We take great pleasure in cordially commending it to our readers.—*The Medical News*, April 11, 1885.

After being before the profession for thirty

years and maintaining during that period a reputation as a leading work on surgery, there is not much to be said in the way of comment or criticism. That it still holds its own goes without saying. The author infuses into it his large experience and ripe judgment. Wedded to no school, committed to no theory, biassed by no hobby, he imparts an honest personality in his observations, and his teachings are the rulings of an impartial judge. Such men are always safe guides, and their works stand the tests of time and experience. Such an author is Erichsen, and such a work is his *Surgery*.—*Medical Record*, Feb. 21, 1885.

BRYANT, THOMAS, F. R. C. S.,

Surgeon and Lecturer on Surgery at Guy's Hospital, London.

The Practice of Surgery. Fourth American from the fourth and revised English edition. In one large and very handsome imperial octavo volume of 1040 pages, with 727 illustrations. Cloth, \$6.50; leather, \$7.50; half Russia, \$8.00.

The fourth edition of this work is fully abreast of the times. The author handles his subjects with that degree of judgment and skill which is attained by years of patient toil and varied experience. The present edition is a thorough revision of those which preceded it, with much new matter added. His diction is so graceful and logical, and his explanations are so lucid, as to place the work among the highest order of text-

books for the medical student. Almost every topic in surgery is presented in such a form as to enable the busy practitioner to review any subject in every-day practice in a short time. No time is lost with useless theories or superfluous verbiage. In short, the work is eminently clear, logical and practical.—*Chicago Medical Journal and Examiner*, April, 1886.

By the same Author.

Diseases of the Breast. In one 12mo. volume. *Preparing.* See *Series of Clinical Manuals*, page 4.

TREVES, FREDERICK, F. R. C. S.,

Hunterian Professor at the Royal College of Surgeons of England.

A Manual of Surgery. In Treatises by Various Authors. In three 12mo. volumes, containing 1866 pages, with 213 engravings. Price per volume, cloth, \$2. See *Students' Series of Manuals*, page 4.

We have here the opinions of thirty-three authors, in an encyclopedic form for easy and ready reference. The three volumes embrace every variety of surgical affections likely to be met with, the paragraphs are short and pithy, and the salient points and the beginnings of new subjects are always printed in extra-heavy type, so that a person may find whatever information he may be in need of at a moment's glance.—*Cincinnati Lancet-Clinic*, August 21, 1886.

The hand of Mr. Treves is evident throughout

in the choice, arrangement and logical sequence of the subjects. Every topic, as far as observed, is treated with a fulness of essential detail, which is somewhat surprising. Another characteristic of the work is the well-nigh universal acceptance of modern and progressive views of pathology and treatment. The entire work is conceived and executed in a scientific spirit. It contains the bone and marrow of modern surgery.—*Annals of Surgery*, Oct. 1886.

BUTLIN, HENRY T., F. R. C. S.,

Assistant Surgeon to St. Bartholomew's Hospital, London.

Diseases of the Tongue. In one 12mo. volume of 456 pages, with 8 colored plates and 3 woodcuts. Cloth, \$3.50. See *Series of Clinical Manuals*, page 4.

The language of the text is clear and concise. The author has aimed to state facts rather than to express opinions, and has compressed within the compass of this small volume the pathology, etiology, etc., of diseases of the tongue that are incon-

veniently scattered through general works on surgery and the practice of medicine. The physician and surgeon will appreciate its value as an aid and guide.—*Physician and Surgeon*, Sept. 1886.

TREVES, FREDERICK, F. R. C. S.,

Surgeon to and Lecturer on Surgery at the London Hospital.

Intestinal Obstruction. In one pocket-size 12mo. volume of 522 pages, with 60 illustrations. Limp cloth, blue edges, \$2.00. See *Series of Clinical Manuals*, page 4.

A standard work on a subject that has not been so comprehensively treated by any contemporary English writer. Its completeness renders a full review difficult, since every chapter deserves minute attention, and it is impossible to do thorough

justice to the author in a few paragraphs. *Intestinal Obstruction* is a work that will prove of equal value to the practitioner, the student, the pathologist, the physician and the operating surgeon.—*British Medical Journal*, Jan. 31, 1885.

GOULD, A. PEARCE, M. S., M. B., F. R. C. S.,

Assistant Surgeon to Middlesex Hospital.

Elements of Surgical Diagnosis. In one pocket-size 12mo. volume of 589 pages. Cloth, \$2.00. See *Students' Series of Manuals*, page 4.

PIRRIE'S PRINCIPLES AND PRACTICE OF SURGERY. Edited by JOHN NEILL, M. D. In one 8vo. vol. of 784 pp. with 316 illus. Cloth, \$3.75.

KEY'S OPERATIVE SURGERY. In one vol. 8vo. of 661 pages, with 81 woodcuts. Cloth, \$3.25.

HOLMES, TIMOTHY, M. A.,

Surgeon and Lecturer on Surgery at St. George's Hospital, London.

A System of Surgery; Theoretical and Practical. IN TREATISES BY VARIOUS AUTHORS. AMERICAN EDITION, THOROUGHLY REVISED AND RE-EDITED by JOHN H. PACKARD, M. D., Surgeon to the Episcopal and St. Joseph's Hospitals, Philadelphia, assisted by a corps of thirty-three of the most eminent American surgeons. In three large and very handsome imperial octavo volumes containing 3137 double-columned pages, with 979 illustrations on wood and 13 lithographic plates, beautifully colored. Price per set, cloth, \$18.00; leather, \$21.00; half Russia, \$22.50. *Sold only by subscription.*

HAMILTON, FRANK H., M. D., LL. D.,

Surgeon to Bellevue Hospital, New York.

A Practical Treatise on Fractures and Dislocations. Seventh edition thoroughly revised and much improved. In one very handsome octavo volume of 998 pages, with 379 illustrations. Cloth, \$5.50; leather, \$6.50; very handsome half Russia, open back, \$7.00.

It is about twenty-five years ago since the first edition of this great work appeared. The edition now issued is the seventh, and this fact alone is enough to testify to the excellence of it in all particulars. Books upon special subjects do not usually command extended sale, but this one is without a rival in any language. It is essentially a practical treatise, and it gathers within its covers almost everything valuable that has been written about fractures and dislocations. The principles and methods of treatment are very fully given. The book is so well known that it does not require

any lengthened review. We can only say that it is still unapproached as a treatise, and that it is a proof of the zeal and industry and great ability of its distinguished author.—*The Dublin Journal of Medical Science*, Feb. 1886.

His famous treatise on *Fractures and Dislocations*, published first in 1860, is justly regarded as the best book on that subject in existence. It has now run through seven editions, and has been translated into French and German.—*Medical Record*, Aug. 14, 1886.

SMITH, STEPHEN, M. D.,

Professor of Clinical Surgery in the University of the City of New York.

The Principles and Practice of Operative Surgery. New (second) and thoroughly revised edition. In one very handsome octavo volume of 892 pages, with 1005 illustrations. Cloth, \$4.00; leather, \$5.00. *Just ready.*

This work is too well and too favorably known to require any words of commendation, and its merits effectually protect it from adverse criticism. It is a treatise upon the principles as well as the practice of mechanical surgery. The subject matter is brought down to the very latest period, hence we shall find the work to be a faithful exponent of the art of surgery as practised now. Stephen Smith's Operative Surgery is one of the most com-

plete works in the English language, and is a fit companion to Malgaigne's magnificent treatise. The woodcuts are good, and are very numerous. The descriptions of operative procedures are plain, and the opinions expressed are conservative and judicious. The work reflects great credit upon the author and upon American surgical literature.—*The American Journal of the Medical Sciences*, April, 1887.

STIMSON, LEWIS A., B. A., M. D.,

Professor of Pathological Anatomy at the University of the City of New York, Surgeon and Curator to Bellevue Hospital, Surgeon to the Presbyterian Hospital, New York, etc.

A Manual of Operative Surgery. New (second) edition. In one very handsome royal 12mo. volume of 503 pages, with 342 illustrations. Cloth, \$2.50.

There is always room for a good book, so that while many works on operative surgery must be considered superfluous, that of Dr. Stimson has held its own. The author knows the difficult art of condensation. Thus the manual serves as a work of reference, and at the same time as a handy guide. It teaches what it professes, the steps of operations. In this edition Dr. Stimson has sought to indicate the changes that have been

effected in operative methods and procedures by the antiseptic system, and has added an account of many new operations and variations in the steps of older operations. We do not desire to extol this manual above many excellent standard British publications of the same class, still we believe that it contains much that is worthy of imitation.—*British Medical Journal*, Jan. 22, 1887.

By the same Author.

A Practical Treatise on Fractures. In one very handsome octavo volume of 598 pages, with 360 beautiful illustrations. Cloth, \$4.75; leather, \$5.75.

The author has given to the medical profession in this treatise on fractures what is likely to become a standard work on the subject. It certainly not surpassed by any work written in the English, or, for that matter, any other language. The author tells us in a short, concise and comprehensive manner, all that is known about his subject. There is nothing scanty or superficial about it, as in most other treatises; on the contrary, everything is thor-

ough. The chapters on repair of fractures and their treatment show him not only to be a profound student, but likewise a practical surgeon and pathologist. His mode of treatment of the different fractures is eminently sound and practical. We consider this work one of the best on fractures; and it will be welcomed not only as a text-book, but also by the surgeon in full practice.—*N. O. Medical and Surgical Journal*, March, 1883.

MARSH, HOWARD, F. R. C. S.,

Senior Assistant Surgeon to and Lecturer on Anatomy at St. Bartholomew's Hospital, London.

Diseases of the Joints. In one 12mo. volume of 468 pages, with 64 woodcuts and a colored plate. Cloth, \$2.00. See *Series of Clinical Manuals*, page 4.

PICK, T. PICKERING, F. R. C. S.,

Surgeon to and Lecturer on Surgery at St. George's Hospital, London.

Fractures and Dislocations. In one 12mo. volume of 530 pages, with 98 illustrations. Limp cloth, 2.00. See *Series of Clinical Manuals*, page 4.

BURNETT, CHARLES H., A. M., M. D.,*Professor of Otolaryngology in the Philadelphia Polyclinic; President of the American Otological Society.*

The Ear, Its Anatomy, Physiology and Diseases. A Practical Treatise for the use of Medical Students and Practitioners. New (second) edition. In one handsome octavo volume of 580 pages, with 107 illustrations. Cloth, \$4.00; leather, \$5.00.

We note with pleasure the appearance of a second edition of this valuable work. When it first came out it was accepted by the profession as one of the standard works on modern aural surgery in the English language; and in his second edition Dr. Burnett has fully maintained his reputation, for the book is replete with valuable information and suggestions. The revision has been carefully

carried out, and much new matter added. Dr. Burnett's work must be regarded as a very valuable contribution to aural surgery, not only on account of its comprehensiveness, but because it contains the results of the careful personal observation and experience of this eminent aural surgeon. —*London Lancet*, Feb. 21, 1885.

POLITZER, ADAM,*Imperial-Royal Prof. of Aural Therap. in the Univ. of Vienna.*

A Text-Book of the Ear and its Diseases. Translated, at the Author's request, by JAMES PATTERSON CASSELLS, M. D., M. R. C. S. In one handsome octavo volume of 800 pages, with 257 original illustrations. Cloth, \$5.50.

The work itself we do not hesitate to pronounce the best upon the subject of aural diseases which has ever appeared, systematic without being too diffuse on obsolete subjects, and eminently practical in every sense. The anatomical descriptions of each separate division of the ear are admirable, and profusely illustrated by woodcuts. They are followed immediately by the physiology of the

section, and this again by the pathological physiology, an arrangement which serves to keep up the interest of the student by showing the direct application of what has preceded to the study of disease. The whole work can be recommended as a reliable guide to the student, and an efficient aid to the practitioner in his treatment. —*Boston Medical and Surgical Journal*, June 7, 1883.

JULER, HENRY E., F. R. C. S.,*Senior Asst Surgeon, Royal Westminster Ophthalmic Hosp.; late Clinical Asst, Moorfields, London.*

A Handbook of Ophthalmic Science and Practice. In one handsome octavo volume of 460 pages, with 125 woodcuts, 27 colored plates, selections from the Test-types of Jaeger and Snellen, and Holmgren's Color-blindness Test. Cloth, \$4.50; leather, \$5.50.

This work is distinguished by the great number of colored plates which appear in it for illustrating various pathological conditions. They are very beautiful in appearance, and have been executed with great care to accuracy. An examination of the work shows it to be one of high standing, one that will be regarded as an authority among ophthalmologists. The treatment recommended is such as the author has learned from actual experience to be the best. —*Cincinnati Medical News*, Dec. 1884.

It presents to the student concise descriptions

and typical illustrations of all important eye affections, placed in juxtaposition, so as to be grasped at a glance. Beyond a doubt it is the best illustrated handbook of ophthalmic science which has ever appeared. Then, what is still better, these illustrations are nearly all original. We have examined this entire work with great care, and it represents the commonly accepted views of advanced ophthalmologists. We can most heartily commend this book to all medical students, practitioners and specialists. —*Detroit Lancet*, Jan. 1885.

NORRIS, WM. F., M. D., and OLIVER, CHAS. A., M. D.*Clin. Prof. of Ophthalmology in Univ. of Pa.*

A Text-Book of Ophthalmology. In one octavo volume of about 500 pages, with illustrations. *Preparing.*

CARTER, R. BRUDENELL, & FROST, W. ADAMS,**F. R. C. S.,****F. R. C. S.,***Ophthalmic Surgeon to and Lecturer on Ophthalmic Surgery at St. George's Hospital, London.**Assistant Ophthalmic Surgeon to and Joint Lecturer on Ophthalmic Surgery at St. George's Hospital, London.*

Ophthalmic Surgery. In one 12mo. volume of about 400 pages. *Preparing.* See *Series of Clinical Manuals*, page 4.

WELLS, J. SOELBERG, F. R. C. S.,*Professor of Ophthalmology in King's College Hospital, London, etc.*

A Treatise on Diseases of the Eye. New (fifth) American from the third London edition. In one large octavo volume. *Preparing.*

NETTLESHIP, EDWARD, F. R. C. S.,*Ophthalmic Surg. and Lect. on Ophth. Surg. at St. Thomas' Hospital, London.*

The Student's Guide to Diseases of the Eye. New (third) edition, thoroughly revised. With a chapter on the Detection of Color-Blindness, by WILLIAM THOMSON, M. D., Professor of Ophthalmology in the Jefferson Medical College. In one royal 12mo. volume of about 450 pages, with about 150 illustrations. Cloth, \$2. *Shortly.*

BROWNE, EDGAR A.,*Surgeon to the Liverpool Eye and Ear Infirmary and to the Dispensary for Skin Diseases.*

How to Use the Ophthalmoscope. Being Elementary Instructions in Ophthalmoscopy, arranged for the use of Students. In one small royal 12mo. volume of 116 pages, with 35 illustrations. Cloth, \$1.00.

LAURENCE AND MOON'S HANDY BOOK OF OPHTHALMIC SURGERY, for the use of Practitioners. Second edition. In one octavo volume of 227 pages, with 65 illust. Cloth, \$2.75.

LAWSON ON INJURIES TO THE EYE, ORBIT AND EYELIDS: Their Immediate and Remote Effects. 8 vo., 404 pp., 92 illus. Cloth, \$3.50.

ROBERTS, WILLIAM, M. D.,

Lecturer on Medicine in the Manchester School of Medicine, etc.

A Practical Treatise on Urinary and Renal Diseases, including Urinary Deposits. Fourth American from the fourth London edition. In one handsome octavo volume of 609 pages, with 81 illustrations. Cloth, \$3.50.

The previous editions of this book have made it so familiar to and so highly esteemed by the medical public, that little more is necessary than a mere announcement of the appearance of this, their successor. But it is pleasant to be able to say that, good as those were, this is still better. In fact, we think it may be said to be the best book in print on the subject of which it treats.—*The American Journal of the Medical Sciences*.—Jan. 1886.

The peculiar value and finish of the book are in a measure derived from its resolute maintenance of a clinical and practical character. It is an unrivalled exposition of everything which relates directly or indirectly to the diagnosis, prognosis and treatment of urinary diseases, and possesses a completeness not found elsewhere in our language in its account of the different affections.—*The Manchester Medical Chronicle*, July, 1885.

PURDY, CHARLES W., M. D.

Bright's Disease and Allied Affections of the Kidneys. In one octavo volume of 288 pages, with illustrations. Cloth, \$2. Just ready.

The object of this work is to "furnish a systematic, practical and concise description of the pathology and treatment of the chief organic diseases of the kidney associated with albuminuria, which shall represent the most recent advances in our knowledge on these subjects;" and this definition of the object is a fair description of the book. The work is a useful one, giving in a

short space the theories, facts and treatments, and going more fully into their later developments. On treatment the writer is particularly strong, steering clear of generalities, and seldom omitting what text-books usually do, the unimportant items which are all important to the general practitioner.—*The Manchester Medical Chronicle*, Oct., 1886.

MORRIS, HENRY, M. B., F. R. C. S.,

Surgeon to and Lecturer on Surgery at Middlesex Hospital, London.

Surgical Diseases of the Kidney. In one 12mo. volume of 554 pages, with 40 woodcuts, and 6 colored plates. Limp cloth, \$2.25. See *Series of Clinical Manuals*, page 4.

In this manual we have a distinct addition to surgical literature, which gives information not elsewhere to be met with in a single work. Such a book was distinctly required, and Mr. Morris has very diligently and ably performed the task

he took in hand. It is a full and trustworthy book of reference, both for students and practitioners in search of guidance. The illustrations in the text and the chromo-lithographs are beautifully executed.—*The London Lancet*, Feb. 26, 1886.

LUCAS, CLEMENT, M. B., B. S., F. R. C. S.,

Senior Assistant Surgeon to Guy's Hospital, London.

Diseases of the Urethra. In one 12mo. volume. Preparing. See *Series of Clinical Manuals*, page 4.

THOMPSON, SIR HENRY,

Surgeon and Professor of Clinical Surgery to University College Hospital, London.

Lectures on Diseases of the Urinary Organs. Second American from the third English edition. In one 8vo. volume of 203 pp., with 25 illustrations. Cloth, \$2.25.

By the Same Author.

On the Pathology and Treatment of Stricture of the Urethra and Urinary Fistulæ. From the third English edition. In one octavo volume of 350 pages, with 47 cuts and 3 plates. Cloth, \$3.50.

THE AMERICAN SYSTEM OF DENTISTRY.

In Treatises by Various Authors. Edited by WILBUR F. LITCH, M. D., D. D. S., Professor of Prosthetic Dentistry, Materia Medica and Therapeutics in the Pennsylvania College of Dental Surgery. In three very handsome octavo volumes of about 1000 pages each, richly illustrated. Per volume, cloth, \$6; leather, \$7; half Morocco, gilt top, \$8. Volume I., containing 1015 pages, with 6 plates and 537 woodcuts, just ready. For sale by subscription only.

As an encyclopedia of Dentistry it has no superior. It should form a part of every dentist's library, as the information it contains is of the greatest value to all engaged in the practice of dentistry.—*American Journal of Dental Science*, September, 1886.

A grand system, big enough and good enough and handsome enough for a monument (which

doubtless it is), to mark an epoch in the history of dentistry. Dentists will be satisfied with it and proud of it—they must. It is sure to be precisely what the student needs to put him and keep him in the right track, while the profession at large will receive incalculable benefit from it.—*Odontographic Journal*, Jan. 1887.

COLEMAN, A., L. R. C. P., F. R. C. S., Exam. L. D. S.,

Senior Dent. Surg. and Lect. on Dent. Surg. at St. Bartholomew's Hosp. and the Dent. Hosp., London.

A Manual of Oral Surgery and Pathology. Thoroughly revised and adapted to the use of American Students, by THOMAS C. STELLWAGEN, M. A., M. D., D. D. S., Prof. of Physiology at the Philadelphia Dental College. In one handsome octavo volume of 412 pages, with 331 illustrations. Cloth, \$3.25.

ESMARCH, Dr. FRIEDRICH,

Professor of Surgery at the University of Kiel, etc.

Early Aid in Injuries and Accidents. Five Ambulance Lectures. Translated by H. R. H. PRINCESS CHRISTIAN. In one handsome small 12mo. volume of 100 pages, with 24 illustrations. Cloth, 75 cents.

RASHAM ON RENAL DISEASES: A Clinical | one 12mo. vol. of 304 pages, with 21 Illustrations.

GROSS, SAMUEL W., A. M., M. D., LL. D.,*Professor of the Principles of Surgery and of Clinical Surgery in the Jefferson Medical College of Phila.*

A Practical Treatise on Impotence, Sterility, and Allied Disorders of the Male Sexual Organs. New (third) edition, thoroughly revised. In one very handsome octavo volume of 163 pages, with 16 illustrations. Cloth, \$1.50. *Just ready.*

We must congratulate the author that another edition has been made necessary. The tone of the book is healthy, and a cheerful prognosis is given of many of the affections of which it treats. We feel confident that the book will continue to sell on its merits.—*N. Y. Med. Journal*, June 18, 1887.

It must be gratifying to both author and publishers that large first and second editions of this little work were so soon exhausted, while the fact

that it has been translated into Russian may indicate that it filled a void even in foreign literature. His is a careful and physiological study of the sexual act, so far as concerns the male, and all his conclusions are scientifically reached. The book has a place by itself in our literature, and furnishes a large fund of information concerning important matters that are too often passed over in silence.—*The Medical Press*, June, 1887.

BUMSTEAD, F. J., and**M. D., LL. D.,***Late Professor of Venereal Diseases at the College of Physicians and Surgeons, New York, etc.***TAYLOR, R. W.,****A. M., M. D.,***Surgeon to Charity Hospital, New York, Prof. of Venereal and Skin Diseases in the University of Vermont, Pres. of the Am. Dermatological Ass'n.*

The Pathology and Treatment of Venereal Diseases. Including the results of recent investigations upon the subject. Fifth edition, revised and largely rewritten, by Dr. Taylor. In one large and handsome octavo volume of 898 pages with 139 illustrations, and thirteen chromo-lithographic figures. Cloth, \$4.75; leather, \$5.75; very handsome half Russia, \$6.25.

It is a splendid record of honest labor, wide research, just comparison, careful scrutiny and original experience, which will always be held as a high credit to American medical literature. This is not only the best work in the English language upon the subjects of which it treats, but also one which has no equal in other tongues for its clear, comprehensive and practical handling of its themes.—*Am. Jour. of the Med. Sciences*, Jan, 1884.

It is certainly the best single treatise on venereal in our own, and probably the best in any language.—*Boston Med. and Surg. Journal*, April 3, 1884.

The character of this standard work is so well

known that it would be superfluous here to pass in review its general or special points of excellence. The verdict of the profession has been passed; it has been accepted as the most thorough and complete exposition of the pathology and treatment of venereal diseases in the language. Admirable as a model of clear description, an exponent of sound pathological doctrine, and a guide for rational and successful treatment, it is an ornament to the medical literature of this country. The additions made to the present edition are eminently judicious, from the standpoint of practical utility.—*Journal of Cutaneous and Venereal Diseases*, Jan, 1884.

CORNIL, V.,*Professor to the Faculty of Medicine of Paris, and Physician to the Lourcine Hospital.*

Syphilis, its Morbid Anatomy, Diagnosis and Treatment. Specially revised by the Author, and translated with notes and additions by J. HENRY C. SIMES, M. D., Demonstrator of Pathological Histology in the University of Pennsylvania, and J. WILLIAM WHITE, M. D., Lecturer on Venereal Diseases and Demonstrator of Surgery in the University of Pennsylvania. In one handsome octavo volume of 461 pages, with 84 very beautiful illustrations. Cloth, \$3.75.

The anatomy, the histology, the pathology and the clinical features of syphilis are represented in this work in their best, most practical and most instructive form, and no one will rise from its

perusal without the feeling that his grasp of the wide and important subject on which it treats is a stronger and surer one.—*The London Practitioner*, Jan, 1882.

HUTCHINSON, JONATHAN, F. R. S., F. R. C. S.,*Consulting Surgeon to the London Hospital.*

Syphilis. In one 12mo. volume of 542 pages, with 8 chromo-lithographs. Cloth, \$2.25. *Just ready.* See *Series of Clinical Manuals*, page 4.

FROM THE PREFACE.

In the following pages I have aimed less at systematic completeness than at clinical exposition. To the latter subject I have devoted my best efforts, and my hope is that those who may honor this work by their attentive perusal will obtain from its pages clear impressions on the state of our knowledge on the topics which it concerns.

GROSS, S. D., M. D., LL. D., D. C. L., etc.

A Practical Treatise on the Diseases, Injuries and Malformations of the Urinary Bladder, the Prostate Gland and the Urethra. Third edition, thoroughly revised by SAMUEL W. GROSS, M. D. In one octavo volume of 574 pages, with 170 illustrations. Cloth, \$4.50.

CULLERIER, A., & BUMSTEAD, F. J., M. D., LL. D.,*Surgeon to the Hôpital du Midi.**Late Professor of Venereal Diseases in the College of Physicians and Surgeons, New York.*

An Atlas of Venereal Diseases. Translated and edited by FREEMAN J. BUMSTEAD, M. D. In one imperial 4to. volume of 328 pages, double-columns, with 26 plates, containing about 150 figures, beautifully colored, many of them the size of life. Strongly bound in cloth, \$17.00. A specimen of the plates and text sent by mail, on receipt of 25 cts.

HILL ON SYPHILIS AND LOCAL CONTAGIOUS FORMS OF LOCAL DISEASE AFFECTING DISORDERS. In one 8vo. vol. of 479 p. Cloth, \$3.25. **PRINCIPALLY THE ORGANS OF GENERAL LEE'S LECTURES ON SYPHILIS AND SOME TION.** In one 8vo. vol. of 246 pages. Cloth, \$2.25.

KAPOSI, MORIZ,

Of Vienna.

The Pathology and Treatment of Diseases of the Skin. For the use of Practitioners and Students. Translated, with the author's permission, by W. XAVIER SUDDUTH, M. D., F. R. M. S. In one octavo volume of about 600 pages, with 74 engravings and 8 colored plates. *Preparing.*

HYDE, J. NEVINS, A. M., M. D.,

Professor of Dermatology and Venereal Diseases in Rush Medical College, Chicago.

A Practical Treatise on Diseases of the Skin. For the use of Students and Practitioners. In one handsome octavo volume of 570 pages, with 66 beautiful and elaborate illustrations. Cloth, \$4.25; leather, \$5.25.

The author has given the student and practitioner a work admirably adapted to the wants of each. We can heartily commend the book as a valuable addition to our literature and a reliable guide to students and practitioners in their studies and practice.—*Am. Journ. of Med. Sci.*, July, 1883.

The aim of the author has been to present to his readers a work not only expounding the most modern conceptions of his subject, but presenting what is of standard value. He has more especially devoted its pages to the treatment of disease, and by his detailed descriptions of the therapeutic measures he has adapted them to the needs of the physician in active practice. In dealing with these questions the author leaves nothing to the presumed knowledge of the reader, but enters thoroughly into the most minute description, so that one is not only told what should be done under given conditions but how to do it as well. It is therefore in the best sense "a practical treatise." That it is comprehensive, a glance at the index will show.—*Maryland Medical Journal*, July 7, 1883.

Professor Hyde has long been known as one of the most intelligent and enthusiastic representatives of dermatology in the west. His numerous contributions to the literature of this specialty have gained for him a favorable recognition as a careful, conscientious and original observer. He has attempted, as he informs us, the task of presenting in a condensed form the results of the latest observation and experience. A careful examination of the work convinces us that he has accomplished his task with painstaking fidelity and with a creditable result.—*Journal of Cutaneous and Venereal Diseases*, June, 1883.

The several diseases are described very concisely, but at the same time with unusual clearness. The treatment is given in the simplest manner and apparently with great honesty. It is a good book, remarkably adapted to the needs of those for whom it was written. It should be at once placed in the library of every general practitioner.—*Detroit Lancet*, April, 1883.

FOX, T., M. D., F. R. C. P., and FOX, T. C., B. A., M. R. C. S.,

Physician to the Department for Skin Diseases, University College Hospital, London.

Physician for Diseases of the Skin to the Westminster Hospital, London.

An Epitome of Skin Diseases. With Formulæ. For Students and Practitioners. Third edition, revised and enlarged. In one very handsome 12mo. volume of 238 pages. Cloth, \$1.25.

The third edition of this convenient handbook calls for notice owing to the revision and expansion which it has undergone. The arrangement of skin diseases in alphabetical order, which is the method of classification adopted in this work, becomes a positive advantage to the student. The book is one which we can strongly recommend, not only to students but also to practitioners who require a compendious summary of the present state of dermatology.—*British Medical Journal*, July 2, 1883.

We cordially recommend Fox's *Epitome* to those whose time is limited and who wish a handy

manual to lie upon the table for instant reference. Its alphabetical arrangement is suited to this use, for all one has to know is the name of the disease, and here are its description and the appropriate treatment at hand and ready for instant application. The present edition has been very carefully revised and a number of new diseases are described, while most of the recent additions to dermal therapeutics find mention, and the formulæ at the end of the book has been considerably augmented.—*The Medical News*, December, 1883.

MORRIS, MALCOLM, F. R. C. S.

Joint Lecturer on Dermatology at St. Mary's Hospital Medical School, London.

Skin Diseases; including their Definitions, Symptoms, Diagnosis, Prognosis, Morbid Anatomy and Treatment. A Manual for Students and Practitioners. In one 12mo. volume of 316 pages, with illustrations. Cloth, \$1.75.

For a student who would like to know something about skin diseases, or that who, as a practitioner, seeks to refresh his memory, or to make a ready diagnosis and prescribe a rational treatment, we unhesitatingly recommend this book. Dr. Morris's *Skin Diseases* is a work that has been used in a series of lectures and has been found to be a most valuable and reliable guide. It is easy to read and contains a large amount of experience and practical information. It is a volume that every student and practitioner should have on his shelf.—*British Medical Journal*, April, 1883.

This is a volume that every student and practitioner should have on his shelf. It is a volume that every student and practitioner should have on his shelf. It is a volume that every student and practitioner should have on his shelf.

for clearness of expression and method; all arrangement is better adapted to promote a rational treatment of dermatology—a branch confessedly hitherto an unexplored territory to the beginner.—*St. Louis Medical Journal*, April, 1883.

The writer has given in a small compass a large amount of well-digested information, and has done so in a manner favorable with any other who has emanated from England, while many points are illustrated by himself from the study of a large number of others of his countrymen. It is a certainly excellent material in the book, and will be very useful.—*British Medical Journal*, March, 1883.

WILSON, ERASMUS, F. R. S.

The Student's Book of Cutaneous Medicine and Diseases of the Skin. In one handsomely bound octavo volume of 145 pages. Cloth, \$1.00.

HILLIER, THOMAS, M. D.

Physician to the Skin Diseases at the University College, London.

Handbook of Skin Diseases for Students and Practitioners. Second American edition. In one 12mo. volume of 145 pages, with plates. Cloth, \$2.25.

The American Systems of Gynecology and Obstetrics.

Systems of Gynecology and Obstetrics, in Treatises by American Authors. Gynecology edited by MATTHEW D. MANN, A. M., M. D., Professor of Obstetrics and Gynecology in the Medical Department of the University of Buffalo; and Obstetrics edited by BARTON COOKE HIRST, M. D., Obstetrician to the Philadelphia and to the Maternity Hospitals, Philadelphia. In four very handsome octavo volumes of about 800 pages each, fully illustrated by wood engravings and colored plates. Prices per volume: Cloth, \$6.00; leather, \$7.00; half Russia, \$8.00. Volume I. of the Gynecology, containing 784 pages, with 201 engravings on wood and 3 colored plates, is just ready. The subsequent volumes are to follow at regular intervals. *For sale by subscription only. Address the Publishers.* Full descriptive circular free on application.

LIST OF CONTRIBUTORS.

WILLIAM H. BAKER, M. D.,
FORDYCE BARKER, M. D., LL. D., Edin.,
ROBERT BATTEY, M. D.,
SAMUEL C. BUSEY, M. D.,
JAMES C. CAMERON, M. D.,
HENRY C. COE, A. M., M. D.,
E. C. DUDLEY, A. B., M. D.,
EDWARD S. DUNSTER, M. D.,
B. McE. EMMET, M. D.,
GEORGE J. ENGELMANN, M. D.,
HAROLD C. ERNST, M. D.,
HENRY J. GARRIGUES, A. M., M. D.,
WILLIAM GOODELL, A. M., M. D.,
EGBERT H. GRANDIN, A. M., M. D.,
CHARLES M. GREEN, M. D.,
SAMUEL W. GROSS, M. D.,
ROBERT P. HARRIS, M. D.,
STEPHEN Y. HOWELL, M. D.,
JAMES B. HUNTER, M. D.,
A. REEVES JACKSON, A. M., M. D.,
EDWARD W. JENKS, M. D., LL. D.,
JOSEPH TABER JOHNSON, M. D.,

CHARLES CARROLL LEE, M. D.,
WILLIAM T. LUSH, M. D., LL. D.,
MATTHEW D. MANN, A. M., M. D.,
H. NEWELL MARTIN, F. R. S., M. D.,
D. Sc. M. A.,
RICHARD B. MAURY, M. D.,
PAUL F. MUNDE, M. D.,
C. D. PALMER, M. D.,
ROSSELL PARK, M. D.,
THEOPHILUS PARVIN, M. D., LL. D.,
R. A. F. PENROSE, M. D., LL. D.,
THADDEUS A. REAMY, A. M., M. D.,
J. C. REEVE, M. D.,
WILLIAM L. RICHARDSON, M. D.,
A. D. ROCKWELL, A. M., M. D.,
ALEXANDER J. C. SKENE, M. D.,
J. LEWIS SMITH, M. D.,
R. STANSBURY SUTTON, A. M., M. D.,
LL. D.,
T. GAILLARD THOMAS, M. D., LL. D.,
ELY VAN DE WARKER, M. D.,
W. GILL WYLIE, M. D.

It is exceedingly fitting that a System of Gynecology should be given forth as the product of American pens. America is the fountain of modern gynecology, and has contributed more than any other country to its development. All things considered, the best treatise on gynecology must be expected from this country, and when the gynecologists of this country unite their efforts in

a "system," the result would be disappointing were it not to excel anything of the kind heretofore attempted. An examination of the first volume of the work before us justifies every expectation formed of it. We must commend it to every reader interested in the division of medicine of which it treats.—*The Medical Age*, July 26, 1887.

THOMAS, T. GAILLARD, M. D.,

Professor of Diseases of Women in the College of Physicians and Surgeons, N. Y.

A Practical Treatise on the Diseases of Women. Fifth edition, thoroughly revised and rewritten. In one large and handsome octavo volume of 810 pages, with 266 illustrations. Cloth, \$5.00; leather, \$6.00; very handsome half Russia, raised bands, \$6.50.

The words which follow "fifth edition" are in this case no mere formal announcement. The alterations and additions which have been made are both numerous and important. The attraction and the permanent character of this book lie in the clearness and truth of the clinical descriptions of diseases; the fertility of the author in therapeutic resources and the fulness with which the details of treatment are described; the definite character of the teaching; and last, but not least, the evident candor which pervades it. We would also particularize the fulness with which the history of the subject is gone into, which makes the

book additionally interesting and gives it value as a work of reference.—*London Medical Times and Gazette*, July 30, 1881.

That the previous editions of the treatise of Dr. Thomas were thought worthy of translation into German, French, Italian and Spanish, is enough to give it the stamp of genuine merit. At home it has made its way into the library of every obstetrician and gynecologist as a safe guide to practice. No small number of additions have been made to the present edition to make it correspond to recent improvements in treatment.—*Pacific Medical and Surgical Journal*, Jan. 1881.

EDIS, ARTHUR W., M. D., Lond., F. R. C. P., M. R. C. S.,

Asst. Obstetric Physician to Middlesex Hospital, late Physician to British Lying-in Hospital.

The Diseases of Women. Including their Pathology, Causation, Symptoms, Diagnosis and Treatment. A Manual for Students and Practitioners. In one handsome octavo volume of 576 pages, with 148 illustrations. Cloth, \$3.00; leather, \$4.00.

It is a pleasure to read a book so thoroughly good as this one. The special qualities which are conspicuous are thoroughness in covering the whole ground, clearness of description and conciseness of statement. Another marked feature of the book is the attention paid to the details of many minor surgical operations and procedures, as, for instance, the use of tents, application of leeches, and use of hot water injections. These

are among the more common methods of treatment, and yet very little is said about them in many of the text-books. The book is one to be warmly recommended especially to students and general practitioners, who need a concise but complete résumé of the whole subject. Specialists, too, will find many useful hints in its pages.—*Boston Med. and Surg. Journ.*, March 2, 1882.

BARNES, ROBERT, M. D., F. R. C. P.,

Obstetric Physician to St. Thomas' Hospital, London, etc.

A Clinical Exposition of the Medical and Surgical Diseases of Women. In one handsome octavo volume, with numerous illustrations. New edition. *Preparing.*

WEST, CHARLES, M. D.

Lectures on the Diseases of Women. Third American from the third London edition. In one octavo volume of 543 pages. Cloth, \$3.75; leather, \$4.75.

EMMET, THOMAS ADDIS, M. D., LL. D.,*Surgeon to the Woman's Hospital, New York, etc.*

The Principles and Practice of Gynecology; For the use of Students and Practitioners of Medicine. New (third) edition, thoroughly revised. In one large and very handsome octavo volume of 880 pages, with 150 illustrations. Cloth, \$5; leather, \$6; very handsome half Russia, raised bands, \$6.50.

We are in doubt whether to congratulate the author more than the profession upon the appearance of the third edition of this well-known work. Embodying, as it does, the life-long experience of one who has conspicuously distinguished himself as a bold and successful operator, and who has devoted so much attention to the specialty, we feel sure the profession will not fail to appreciate the privilege thus offered them of perusing the views and practice of the author. His earnestness of purpose and conscientiousness are manifest. He gives not only his individual experience but endeavors to represent the actual state of gynecological science and art.—*British Medical Journal*, May 16, 1885.

No jot or tittle of the high praise bestowed upon the first edition is abated. It is still a book of marked personality, one based upon large clinical experience, containing large and valuable additions to our knowledge, evidently written not only with honesty of purpose, but with a conscientious sense of responsibility, and a book that is at

once a credit to its author and to American medical literature. We repeat that it is a book to be studied, and one that is indispensable to every practitioner giving any attention to gynecology.—*American Journal of the Medical Sciences*, April, 1886.

The time has passed when Emmet's *Gynecology* was to be regarded as a book for a single country or for a single generation. It has always been his aim to popularize gynecology, to bring it within easy reach of the general practitioner. The originality of the ideas, aside from the perfect confidence which we feel in the author's statements, compels our admiration and respect. We may well take an honest pride in Dr. Emmet's work and feel that his book can hold its own against the criticism of two continents. It represents all that is most earnest and most thoughtful in American gynecology. Emmet's work will continue to reflect the individuality, the sterling integrity and the kindly heart of its honored author long after smaller books have been forgotten.—*American Journal of Obstetrics*, May, 1885.

DUNCAN, J. MATTHEWS, M.D., LL. D., F. R. S. E., etc.

Clinical Lectures on the Diseases of Women; Delivered in Saint Bartholomew's Hospital. In one handsome octavo volume of 175 pages. Cloth, \$1.50.

They are in every way worthy of their author; indeed, we look upon them as among the most valuable of his contributions. They are all upon matters of great interest to the general practitioner. Some of them deal with subjects that are not, as a

rule, adequately handled in the text-books; others of them, while bearing upon topics that are usually treated of at length in such works, yet bear such a stamp of individuality that they deserve to be widely read.—*N. Y. Medical Journal*, March, 1886.

MAY, CHARLES H., M. D.*Late House Surgeon to Mount Sinai Hospital, New York.*

A Manual of the Diseases of Women. Being a concise and systematic exposition of the theory and practice of gynecology. In one 12mo. volume of 342 pages. Cloth, \$1.75.

Medical students will find this work adapted to their wants. Also practitioners of medicine will find it exceedingly convenient to consult for the purpose of refreshing their minds upon the leading points of a gynecological subject. By systematic condensation, the omission of disputed ques-

tions, and the presentation only of accepted views, it constitutes a very satisfactory exposition of the leading principles of gynecology as they are understood at the present time.—*Cincinnati Medical News*, Nov. 1885.

HODGE, HUGH L., M. D.,*Emeritus Professor of Obstetrics, etc., in the University of Pennsylvania.*

On Diseases Peculiar to Women; Including Displacements of the Uterus. Second edition, revised and enlarged. In one beautifully printed octavo volume of 519 pages, with original illustrations. Cloth, \$4.50.

By the Same Author.

The Principles and Practice of Obstetrics. Illustrated with large lithographic plates containing 159 figures from original photographs, and with numerous woodcuts. In one large quarto volume of 542 double-columned pages. Strongly bound in cloth, \$14.00. Specimens of the plates and letter-press will be forwarded to any address, free by mail, on receipt of six cents in postage stamps.

RAMSBOTHAM, FRANCIS H., M. D.

The Principles and Practice of Obstetric Medicine and Surgery; In reference to the Process of Parturition. A new and enlarged edition, thoroughly revised by the Author. With additions by W. V. KEATING, M. D., Professor of Obstetrics, etc., in the Jefferson Medical College of Philadelphia. In one large and handsome imperial octavo volume of 640 pages, with 64 full-page plates and 43 woodcuts in the text, containing in all nearly 200 beautiful figures. Strongly bound in leather, with raised bands, \$7.

WINCKEL, F.

A Complete Treatise on the Pathology and Treatment of Childbed, For Students and Practitioners. Translated, with the consent of the Author, from the second German edition, by J. R. CHADWICK, M. D. Octavo 484 pages. Cloth, \$4.00.

ASHWELL'S PRACTICAL TREATISE ON THE DISEASES PECULIAR TO WOMEN. Third American from the third and revised London edition. In one 8vo. vol., pp. 520. Cloth, \$3.50.

CHURCHILL ON THE PUERPERAL FEVER

AND OTHER DISEASES PECULIAR TO WOMEN. In one 8vo. vol. of 464 pages. Cloth, \$2.50.

MEIGS ON THE NATURE, SIGNS AND TREATMENT OF CHILDBED FEVER. In one 8vo. volume of 346 pages. Cloth, \$2.00.

PARVIN, THEOPHILUS, M. D., LL. D.,

Prof. of Obstetrics and the Diseases of Women and Children in Jefferson Med. Coll., Phila.

The Science and Art of Obstetrics. In one handsome 8vo. volume of 697 pages, with 214 engravings and a colored plate. Cloth, \$4.25; leather, \$5.25.

It is a ripe harvest that Dr. Parvin offers to his readers. There is no book that can be more safely recommended to the student or that can be turned to in moments of doubt with greater assurance of aid, as it is a liberal digest of safe counsel that has been patiently gathered.—*The American Journal of the Medical Sciences*, July, 1887.

There is not in the language a treatise on the subject which so completely and intelligently gleams the whole field of obstetric literature, giving the reader the winnowed wheat in concise and well-jointed phrase, in language of exceptional purity and strength. The arrangement of the matter of this work is unique and exceedingly

favorable for an agreeable unfolding of the science and art of obstetrics. This new book is the easy superior of any single work among its predecessors for the student or practitioner seeking the best thought of the day in this department of medicine.—*The American Practitioner and Nurse*, April 2, 1887.

This treatise may be defined as exact, concise and scholarly. Parvin's distinguished position as a teacher, his scholarly attainments, and his honest endeavor to do his best by both the student and the physician, will secure for his treatise favorable recognition.—*American Journal of Obstetrics*, May, 1887.

BARNES, ROBERT, M. D., and FANCOURT, M. D.,

Phys. to the General Lying-in Hosp., Lond.

Obstetric Phys. to St. Thomas' Hosp., Lond.

A System of Obstetric Medicine and Surgery, Theoretical and Clinical. For the Student and the Practitioner. The Section on Embryology contributed by Prof. Milnes Marshall. In one handsome octavo volume of 872 pages, with 231 illustrations. Cloth, \$5; leather, \$6.

This system will be eagerly sought for, not only on account of its intrinsic merit, but also because the reputation which the elder Barnes, in particular, has secured, carries with it the conviction that any book emanating from him is necessarily sound in teaching and conservative in practice. It is indeed eminently fitting that a man who has done so much towards systematizing the obstetric art, who for so many years has been widely known as a capa-

ble teacher and trusted accoucheur, should embody within a single treatise the system which he has taught and in practice tested, and which is the outcome of a lifetime of earnest labor, careful observation and deep study. The result of this arrangement is the production of a work which rises above criticism and which in no respect need yield the palm to any obstetrical treatise hitherto published.—*American Journal of Obstetrics*, Feb. 1886.

PLAYFAIR, W. S., M. D., F. R. C. P.,

Professor of Obstetric Medicine in King's College, London, etc.

A Treatise on the Science and Practice of Midwifery. New (fourth) American, from the fifth English edition. Edited, with additions, by ROBERT P. HARRIS, M. D. In one handsome octavo volume of 654 pages, with 3 plates and 201 engravings. Cloth, \$4; leather, \$5; half Russia, \$5.50.

This still remains a favorite in America, not only because the author is recognized as a safe guide and eminently progressive man, but also as sparing no effort to make each successive edition a faithful mirror of the latest and best practice. A work so frequently noticed as the present requires no further review. We believe that this edition is simply the forerunner of many others, and that the demand will keep pace with the

supply.—*American Journal of Obstetrics*, Nov. 1885. Since its first publication, only eight years ago, it has rapidly become the favorite text-book, to the practical exclusion of all others. A large measure of its popularity is due to the clear and easy style in which it is written. Few text-books for students have very much to boast of in this respect.—*Medical Record*.

KING, A. F. A., M. D.,

Professor of Obstetrics and Diseases of Women in the Medical Department of the Columbian University, Washington, D. C., and in the University of Vermont, etc.

A Manual of Obstetrics. New (third) edition. In one very handsome 12mo. volume of 376 pages, with 102 illustrations. Cloth, \$2.25.

This little manual, certainly the best of its kind, fully deserves the popularity which has made a third edition necessary. Clear, practical, concise, its teachings are so fully abreast with recent advances in obstetric science that but few points can be criticised.—*American Journal of Obstetrics*, March, 1887.

This volume deserves commendation. It is not

bulky—it is concise. The chapters are divided with sub-headings, which aid materially in the finding of any particular subject, and the definitions are clear and explicit. It fulfils its purpose admirably, and we know of no better work to place in the student's hands. The illustrations are good.—*Archives of Gynecology*, January, 1887.

BARKER, FORDYCE, A. M., M. D., LL. D. Edin.,

Clinical Professor of Midwifery and the Diseases of Women in the Bellevue Hospital Medical College, New York, Honorary Fellow of the Obstetrical Societies of London and Edinburgh, etc., etc.

Obstetrical and Clinical Essays. In one handsome 12mo. volume of about 300 pages. Preparing.

BARNES, FANCOURT, M. D.,

Obstetric Physician to St. Thomas' Hospital, London.

A Manual of Midwifery for Midwives and Medical Students. In one royal 12mo. volume of 197 pages, with 50 illustrations. Cloth, \$1.25.

PARRY, JOHN S., M. D.,

Obstetrician to the Philadelphia Hospital, Vice-President of the Obstet. Society of Philadelphia.

Extra-Uterine Pregnancy: Its Clinical History, Diagnosis, Prognosis and Treatment. In one handsome octavo volume of 272 pages. Cloth, \$2.50.

TANNER ON PREGNANCY. Octavo, 490 pages, 4 colored plates, 16 cuts. Cloth, \$4.25.

LEWIS, WILLIAM, M. D.,

Regius Professor of Midwifery in the University of Glasgow, etc.

A System of Midwifery, Including the Diseases of Pregnancy and the Puerperal State. Third American edition, revised by the Author, with additions by JOHN H. PARRY, M. D., Obstetrician to the Philadelphia Hospital, etc. In one large and very handsome octavo volume of 740 pages, with 205 illustrations. Cloth, \$4.50; leather, \$6.50; very handsome half Russia, raised bands, \$6.00.

The author is broad in his teachings, and discusses briefly the comparative anatomy of the pelvis and the mobility of the pelvic articulations. The second chapter is devoted especially to the study of the pelvis, while in the third the female organs of generation are introduced. The structure and development of the ovum are admirably described. Then follow chapters upon the various subjects embraced in the study of midwifery. The descriptions throughout the work are plain and pleasing. It is sufficient to state that in this, the last edition of this well known work, every recent advancement in this field has been brought forward. *Physician and Surgeon*, Jan. 1880.

To the American student the work before us

must prove admirably adapted. Complete in all its parts, essentially modern in its teachings, and with demonstrations noted for clearness and precision, it will gain in favor and be recognized as a work of standard merit. The work cannot fail to be popular and is cordially recommended.—*N. O. Med. and Surg. Journ.*, March, 1880.

It has been well and carefully written. The views of the author are broad and liberal, and indicate a well-balanced judgment and a matured mind. We observe no spirit of dogmatism, but the earnest teaching of the thoughtful observer and lover of true science. Take the volume as a whole, and it has few equals.—*Maryland Medical Journal*, Feb. 1880.

LANDIS, HENRY G., A. M., M. D.,

Professor of Obstetrics and the Diseases of Women in Starting Medical College, Columbus, O.

The Management of Labor, and of the Lying-in Period. In one handsome 12mo. volume of 334 pages, with 28 illustrations. Cloth, \$1.75.

The author has designed to place in the hands of the young practitioner a book in which he can find necessary information in an instant. As far as we can see, nothing is omitted. The advice is sound, and the procedures are safe and practical. *Centralblatt für Gynäkologie*, December 4, 1880.

This is a book we can heartily recommend. The author goes much more practically into the details of the management of labor than most text books, and is so readable throughout as to

tempt any one who should happen to commence the book to read it through. The author presupposes a theoretical knowledge of obstetrics, and has consistently excluded from this little work everything that is not of practical use in the lying-in room. We think that if it is as widely read as it deserves, it will do much to improve obstetric practice in general.—*New Orleans Medical and Surgical Journal*, Mar. 1886.

SMITH, J. LEWIS, M. D.,

Clinical Professor of Diseases of Children in the Bellevue Hospital Medical College, N. Y.

A Treatise on the Diseases of Infancy and Childhood. New (sixth) edition, thoroughly revised and rewritten. In one handsome octavo volume of 867 pages, with 10 illustrations. Cloth, \$4.50; leather, \$5.50; half Russia, \$6.00.

Rarely does a pleasanter task fall to the lot of the bibliographer than to announce the appearance of a new edition of a medical classic like Prof. J. Lewis Smith's *Treatise on the Diseases of Infancy and Childhood*. For years it has stood high in the confidence of the profession, and with the addition of the changes now made it may be said to be the best book in the language on the subject of which it treats. The experience of the author fully sustains the confidence of the profession, and the present edition is a most valuable addition to the literature of the subject. The author has revised the text, and has added many new facts and observations, and has rewritten the whole work in a new and more readable form. The work is a most valuable addition to the literature of the subject, and is a most valuable addition to the library of every practitioner of medicine.

The author has revised the text, and has added many new facts and observations, and has rewritten the whole work in a new and more readable form. The work is a most valuable addition to the literature of the subject, and is a most valuable addition to the library of every practitioner of medicine.

stands alone amongst other works on its subject, of recommending treatment in accordance with the most recent therapeutical views.—*British and Foreign Medical-Chirurgical Review*.

It is a pleasure to the busy practitioner—interested in the advancement of his profession—to meet, fresh from the hands of its author, a medical classic such as Smith's *Diseases of Infancy and Childhood*. These familiar with former editions of the work will certainly recognize the taste-making with which this revision has been made. Many of the articles have been entirely rewritten. The whole work is enriched with a research and reasoning which is a most valuable addition to the literature of the subject. The author has spared neither time nor labor in bringing it to its present appearance, and the result is a most valuable addition to the literature of the subject, and is a most valuable addition to the library of every practitioner of medicine.

OWEN, EDMUND, M. B., F. R. C. S.,

Surgeon, St. George's Hospital, London.

Surgical Diseases of Children. In one 12mo. volume of 355 pages, with 4 illustrations. Cloth, \$2.50; leather, \$3.50. See *Series of Clinical Monographs*.

This is a most valuable addition to the literature of the subject, and is a most valuable addition to the library of every practitioner of medicine. The author has revised the text, and has added many new facts and observations, and has rewritten the whole work in a new and more readable form. The work is a most valuable addition to the literature of the subject, and is a most valuable addition to the library of every practitioner of medicine.

WEST, CHARLES, M. D.,

Professor of Diseases of the Nervous System in the University of Michigan, etc.

On Some Disorders of the Nervous System in Childhood. In one small 12mo. volume of 100 pages, with 10 illustrations. Cloth, \$1.50; leather, \$2.50.

This is a most valuable addition to the literature of the subject, and is a most valuable addition to the library of every practitioner of medicine. The author has revised the text, and has added many new facts and observations, and has rewritten the whole work in a new and more readable form. The work is a most valuable addition to the literature of the subject, and is a most valuable addition to the library of every practitioner of medicine.

TIDY, CHARLES MEYMOTT, M. B., F. C. S.,*Professor of Chemistry and of Forensic Medicine and Public Health at the London Hospital, etc.*

Legal Medicine. VOLUME II. Legitimacy and Paternity, Pregnancy, Abortion, Rape, Indecent Exposure, Sodomy, Bestiality, Live Birth, Infanticide, Asphyxia, Drowning, Hanging, Strangulation, Suffocation. Making a very handsome imperial octavo volume of 529 pages. Cloth, \$6.00; leather, \$7.00.

VOLUME I. Containing 664 imperial octavo pages, with two beautiful colored plates. Cloth, \$6.00; leather, \$7.00.

The satisfaction expressed with the first portion of this work is in no wise lessened by a perusal of the second volume. We find it characterised by the same fulness of detail and clearness of expression which we had occasion so highly to commend in our former notice, and which render it so valuable to the medical jurist. The copious

tables of cases appended to each division of the subject must have cost the author a prodigious amount of labor and research, but they constitute one of the most valuable features of the book, especially for reference in medico-legal trials.—*American Journal of the Medical Sciences*, April, 1884.

TAYLOR, ALFRED S., M. D.,*Lecturer on Medical Jurisprudence and Chemistry in Guy's Hospital, London.*

A Manual of Medical Jurisprudence. Eighth American from the tenth London edition, thoroughly revised and rewritten. Edited by JOHN J. REESE, M. D., Professor of Medical Jurisprudence and Toxicology in the University of Pennsylvania. In one large octavo volume of 937 pages, with 70 illustrations. Cloth, \$5.00; leather, \$6.00; half Russia, raised bands, \$6.50.

The American editions of this standard manual have for a long time laid claim to the attention of the profession in this country; and the eighth comes before us as embodying the latest thoughts and emendations of Dr. Taylor upon the subject to which he devoted his life with an assiduity and success which made him *facile princeps* among English writers on medical jurisprudence. Both the author and the book have made a mark too deep to be affected by criticism, whether it be censure or praise. In this case, however, we should

only have to seek for laudatory terms.—*American Journal of the Medical Sciences*, Jan. 1881.

This celebrated work has been the standard authority in its department for thirty-seven years, both in England and America, in both the professions which it concerns, and it is improbable that it will be superseded in many years. The work is simply indispensable to every physician, and nearly so to every liberally-educated lawyer, and we heartily commend the present edition to both professions.—*Albany Law Journal*, March 26, 1881.

By the Same Author.

The Principles and Practice of Medical Jurisprudence. Third edition. In two handsome octavo volumes, containing 1416 pages, with 188 illustrations. Cloth, \$10; leather, \$12.

For years Dr. Taylor was the highest authority in England upon the subject to which he gave especial attention. His experience was vast, his judgment excellent, and his skill beyond cavil. It is therefore well that the work of one who, as Dr. Stevenson says, had an "enormous grasp of all

matters connected with the subject," should be brought up to the present day and continued in its authoritative position. To accomplish this result Dr. Stevenson has subjected it to most careful editing, bringing it well up to the times.—*American Journal of the Medical Sciences*, Jan. 1884.

By the Same Author.

Poisons in Relation to Medical Jurisprudence and Medicine. Third American, from the third and revised English edition. In one large octavo volume of 788 pages. Cloth, \$5.50; leather, \$6.50.

PEPPER, AUGUSTUS J., M. S., M. B.,*Examiner in Forensic Medicine at the University of London.*

Forensic Medicine. In one pocket-size 12mo. volume *Series of Manuals*, page 4.

LEA, HENRY C.

Superstition and Force: Essays on The Battle, The Ordeal and Torture. Third revised edition. In one large octavo volume of 552 pages. Cloth, \$5.00; leather, \$6.00.


This valuable work is in reality a history of civilisation as interpreted by the progress of jurisprudence. . . In "Superstition and Force" we have a philosophic survey of the long period intervening between primitive barbarity and civilized enlightenment. There is not a chapter in the work

C. S.,*Students'*

Studies in Church History
Life of Henry VIII.—Excommunication
octavo volume of 605 pages. Cloth, \$5.00; leather, \$6.00.

The author is pre-eminently a specialist, and he has brought up every topic allied with the history of the Church, and traces it out to the minutest details of knowledge and impartiality, which will compel admiration. The whole is compressed into the book, and is a volume of other single volumes.

By the Same Author.
Studies in Church History
Life of Henry VIII.—Excommunication
octavo volume of 605 pages. Cloth, \$5.00; leather, \$6.00.



Allen's Anatomy	6	*Holmes' System of Surgery	22
American Journal of the Medical Sciences	3	Horner's Anatomy and Histology	6
American Systems of Gynecology	27	Hudson on Fever	4
American System of Practical Medicine	15	Hutchinson on Syphilis	4, 23
An American System of Dentistry	24	Hyde on the Diseases of the Skin	26
*Ashhurst's Surgery	20	Jones (C. Handfield) on Nervous Disorders	18
Ashwell on Diseases of Women	28	Juler's Ophthalmic Science and Practice	23
Attfield's Chemistry	9	Kaposi on Skin Diseases	26
Bail on the Rectum and Anus	4, 20	King's Manual of Obstetrics	33
Barker's Obstetrical and Clinical Essays	29	Klein's Histology	4, 13
Barlow's Practice of Medicine	17	Landis on Labor	30
Barnes' Midwifery	29	La Roche on Pneumonia, Malaria, etc.	18
*Barnes on Diseases of Women	27	La Roche on Yellow Fever	14
Barnes' System of Obstetric Medicine	29	Laurence and Moon's Ophthalmic Surgery	23
Bartholow on Electricity	17	Lawson on the Eye, Orbit and Eyelid	31
Bartholow's New Remedies and their Uses	11	Lea's Studies in Church History	31
Basham on Renal Diseases	24	Lea's Superstition and Force	25
Bell's Comparative Physiology and Anatomy	4, 7	Lee on Syphilis	8
Bellamy's Operative Surgery	4, 20	Lehmann's Chemical Physiology	8
Bellamy's Surgical Anatomy	6	*Leishman's Midwifery	30
Blandford on Insanity	19	Lucas on Diseases of the Urethra	4, 24
Bloxam's Chemistry	9	Ludlow's Manual of Examinations	3
*Bristowe's Practice of Medicine	14	Lyons on Fever	14
Brodbent on the Pulse	4, 18	Malsch's Organic Materia Medica	11
Browne on the Ophthalmoscope	23	Marsh on the Joint	4, 21
Bruce's Materia Medica and Therapeutics	22	May on Diseases of Women	28
Brunton's Materia Medica and Therapeutics	11	Medical News	1
Bryant on the Breast	4, 21	Medical News Visiting List	3
*Bryant's Practice of Surgery	21	Medical News Physicians' Ledger	3
*Bumstead on Venereal Diseases	25	Meligs on Childbed Fever	28
*Burnett on the Ear	23	Miller's Practice of Surgery	19
Bullin on the Tongue	4, 21	Miller's Principles of Surgery	20
Carpenter on the Use and Abuse of Alcohol	8	Mitchell's Nervous Diseases of Women	21
*Carpenter's Human Physiology	8	Morris on Diseases of the Kidney	4, 24
Carter & Frost's Ophthalmic Surgery	4, 23	Morris on Skin Diseases	26
Century of American Medicine	14	Nell and Smith's Compendium of Med. Sci.	3
Chambers on Diet and Regimen	17	Nettleship on Diseases of the Eye	23
Chapman's Human Physiology	8	Norris and Oliver on the Eye	23
Charles' Physiological and Pathological Chem.	10	Owen on Diseases of Children	4, 30
Churchill on Puerperal Fever	28	*Farrish's Practical Pharmacy	29
Clarke and Lockwood's Dissectors' Manual	4, 6	Perry on Extra-Uterine Pregnancy	29
Classes' Quantitative Analysis	10	Parvins Midwifery	17
Cleland's Dissector	5	Pavy on Digestion and its Disorders	17
Clouston on Insanity	19	Pepper's System of Medicine	15
Clowes' Practical Chemistry	10	Pepper's Forensic Medicine	4, 31
Coats' Pathology	13	Pepper's Surgical Pathology	4, 13
Cohen on the Throat	18	Pick on Fractures and Dislocations	4, 23
Coleman's Dental Surgery	24	Pirie's System of Surgery	29
Condie on Diseases of Children	30	Playfair on Nerve Prostration and Hysteria	19
Cornil on Syphilis	25	*Playfair's Midwifery	29
*Cornil and Ranvier's Pathological Histology	13	Pollitzer on the Ear and its Diseases	24
Cullerier's Atlas of Venereal Diseases	25	Power's Human Physiology	4, 8
Curnow's Medical Anatomy	4, 6	Purdy on Bright's Disease and Allied Affections	24
Dalton on the Circulation	7	Ralfs's Clinical Chemistry	4, 10
*Dalton's Human Physiology	8	Ramsbottom on Parturition	28
Davis' Clinical Lectures	17	Rensen's Theoretical Chemistry	9
Draper's Medical Physics	7	*Reynolds' System of Medicine	16
Druitt's Modern Surgery	20	Richardson's Preventive Medicine	17
Duncan on Diseases of Women	28	Roberts on Urinary Diseases	24
*Dunglison's Medical Dictionary	4	Roberts' Compend of Anatomy	7
Edes' Materia Medica and Therapeutics	12	Roberts' Principles and Practice of Surgery	20
Edis on Diseases of Women	27	Robertson's Physiological Physics	4, 7
Ellis' Demonstrations of Anatomy	7	Ross on Nervous Diseases	19
Emmet's Gynecology	21	*Sayce on Insanity, including Hysteria	4, 19
*Erichsen's System of Surgery	21	Schäfer's Essentials of Histology	13
Esmarch's Early Aid in Injuries and Accidents	24	Schreiber on Massage	17
Farquharson's Therapeutics and Mat. Med.	12	Seiler on the Throat, Nose and Naso-Pharynx	18
Fenwick's Medical Diagnosis	16	Series of Clinical Manuals	4
Finlayson's Clinical Diagnosis	16	Simon's Manual of Chemistry	9
Flint on Auscultation and Percussion	18	Skey's Operative Surgery	21
Flint on Phthisis	18	Slade on Diphtheria	18
Flint on Physical Exploration of the Lungs	18	Smith (Edward) on Consumption	18
Flint on Respiratory Organs	18	*Smith (J. Lewis) on Children	20
Flint on the Heart	18	Smith's Operative Surgery	32
Flint's Essays	18	Stille on Cholera	16
*Flint's Practice of Medicine	14	*Stille & Malsch's National Dispensary	12
Folsom's Laws of U. S. on Custody of Insane	19	*Stille's Therapeutics and Materia Medica	12
Foster's Physiology	8	Stimson on Fractures	22
*Fothergill's Handbook of Treatment	16	Stimson's Operative Surgery	22
Fownes' Elementary Chemistry	9	Stokes on Fever	14
Fox on Diseases of the Skin	26	Students' Series of Manuals	4
Frankland and Japp's Inorganic Chemistry	9	Sturges' Clinical Medicine	17
Fuller on the Lungs and Air Passages	18	Tanner on Signs and Diseases of Pregnancy	29
Galloway's Analysis	8	Tanner's Manual of Clinical Medicine	16
Gibney's Orthopedic Surgery	20	Taylor on Poisons	31
Gould's Surgical Diagnosis	4, 21	*Taylor's Medical Jurisprudence	31
*Gray's Anatomy	5	Taylor's Principles of Med. Jurisprudence	31
Greene's Medical Chemistry	10	*Thomas on Diseases of Women	24
Green's Pathology and Morbid Anatomy	13	Thompson on Stricture	24
Griffith's Universal Formulary	11	Thompson on Urinary Organs	24
Gross on Foreign Bodies in Air-Passages	15	Tidy's Legal Medicine	31
Gross on Impotence and Sterility	25	Todd on Acute Diseases	17
Gross on Urinary Organs	25	Treves' Manual of Surgery	21
*Gross' System of Surgery	20	Treves' Surgical Applied Anatomy	4, 6
Habershon on the Abdomen	16	Treves' Uterine Obstruction	4, 27
*Hamilton on Fractures and Dislocations	22	Take on the Influence of Mind on the Body	19
Hamilton on Nervous Diseases	19	Visiting List, The Medical News	8
Hartshorne's Anatomy and Physiology	6	Walsh on the Heart	19
Hartshorne's Conspectus of the Med. Sciences	3	Watson's Practice of Physic	14
Hartshorne's Essentials of Medicine	14	*Wells on the Eye	23
Hartshorne's Household Medicine	17	West on Diseases of Childhood	30
Hermann's Experimental Pharmacology	11	West on Diseases of Women	27
Hill on Syphilis	25	West on Nervous Disorders in Childhood	30
Hillier's Handbook of Skin Diseases	26	Williams on Consumption	18
Hoblyn's Medical Dictionary	4	Wilson's Handbook of Cutaneous Medicine	26
Hodge on Women	28	Wilson's Human Anatomy	5
Hodge's Obstetrics	28	Winckel on Pathol. and Treatment of Childbed	28
Hoffmann and Power's Chemical Analysis	10	Wöhler's Organic Chemistry	8
Holiden's Landmarks	5	Woodhead's Practical Pathology	13
Holland's Medical Notes and Reflections	17	Year-Books of Treatment for 1885 and 1886	17

Books marked * are also bound in half Russia.

LEA BROTHERS & CO., Philadelphia.



LANE MEDICAL LIBRARY

—
To avoid fine, this book should be returned on
or before the date last stamped below.

--	--	--

0124 Playfair, W.S.
P72 A treatise on the sci-
1885 ence and practice of
NAME DATE DUE
midwifery. 18735

